**Phase equilibria constraints on the melt fertility of crustal
source rocks: The effect of sub-solidus water loss**

Journal:	<i>Journal of Metamorphic Geology</i>
Manuscript ID:	JMG-14-0034
Manuscript Type:	Original Article
Date Submitted by the Author:	01-May-2014
Complete List of Authors:	Webb, Gordon; University of Melbourne, School of Earth Sciences Powell, Roger; University of Melbourne, School of Earth Sciences McLaren, Sandra; University of Melbourne, School of Earth Sciences
Keywords:	prograde metamorphism, melting, water, anatexis, THERMOCALC

1 Phase equilibria constraints on the melt fertility of
2 crustal source rocks: The effect of sub-solidus water
3 loss

4 G. Webb, R. Powell and S. McLaren

5 School Earth Sciences, The University of Melbourne, Vic 3010, Australia

6 *Corresponding author: gwebb@student.unimelb.edu.au

7 Short title: Prograde water loss and crustal melt fertility

ABSTRACT

During prograde metamorphism, free H₂O generated by ongoing dehydration reactions is likely to be continuously lost from a rock primarily driven by compaction. Classical melting experiments cannot easily simulate this phenomena because all run products are ideally conserved within the experimental charge even though significant equilibration and H₂O generation may occur during heating. Phase equilibria modelling is used to model the effect of subsolidus water loss on subsequent melting relationships of felsic lithologies (including metapelite, metagreywacke and metatonalite) in the suprasolidus. Subsolidus water loss drives the bulk composition towards the minimum saturation point within the wet melting field and results in significantly reduced subsequent melt generation when compared to melting experiments involving conservation of H₂O in an experimental charge from some temperature well below the solidus. This effect is most obvious at P - T conditions just above the solidus. The reduction in melt fertility due to subsolidus water loss causes rheologically critical thresholds to be shifted to higher temperatures (e.g. melt connectivity threshold, melt escape threshold and the solid-to-liquid transition). Subsolidus water loss is also likely to impact on crystallising melts as they cross the solidus. As a result, the H₂O content of granites may not reflect the H₂O contents of the magma from which they were derived. Instead they record the H₂O contents in the vicinity of the minimum saturation point at the P - T conditions at which they crossed the solidus.

keywords: prograde metamorphism, melting, water, anatexis, THERMOCALC

29 INTRODUCTION

30 Understanding crustal melt fertility is important for developing robust models for
 31 migmatite formation, granite petrogenesis and the differentiation and evolution of the
 32 continental crust (Brown & Fyfe, 1970; Huppert & Sparks, 1988; Clemens, 1990;
 33 Thompson, 1996; Sawyer, 1998; Annen *et al.*, 2008). Hydrous felsic lithologies melt
 34 incongruently at mid- to lower-crustal depths (10–60 km) at temperatures attainable during
 35 high-grade metamorphism (650–1000 °C; Thompson & Connolly, 1995; Thompson, 1999;
 36 White *et al.*, 2001). The amount of melt produced and its composition is determined by the
 37 composition of the source rock, the pressure and temperature conditions and any open
 38 system behaviour including the loss or gain of fluid, melt or solid phases. Of these,
 39 H₂O content is one of the key compositional variables affecting melting behaviour in the
 40 crust (Tuttle & Bowen, 1958; Holtz *et al.*, 1995; Johannes & Holtz, 1996).

41 Two dominant modes of melting have been identified to explain the observed
 42 characteristics of natural granites, migmatites and granulites. ‘Fluid-present melting’ or
 43 ‘wet melting’ describes the situation where a system is saturated with respect to a
 44 H₂O-dominated fluid phase as it crosses the wet solidus with increasing T and/or P . In
 45 fluid-present melting the activity of H₂O is effectively unity ($a_{\text{H}_2\text{O}} \approx 1$), neglecting the
 46 proportions of other constituents in the fluid that are likely to be small, at least for most
 47 crustal conditions. The resulting melt phase is saturated with respect to H₂O. Fluid-present
 48 melting may occur in some migmatite terrains (especially at low P ; e.g. Sawyer, 2010).
 49 However it is not generally considered to be the dominant mode of melting in the
 50 continental crust (Clemens, 1984). By contrast, ‘fluid-absent’, ‘vapour-absent’ or
 51 ‘dehydration’ melting proceeds via reactions involving the breakdown of hydrous minerals

(mainly muscovite, biotite and hornblende) in the absence of a free fluid ($a_{\text{H}_2\text{O}} < 1$) at higher temperatures than the wet solidus. It has been argued that fluid-absent melting is the dominant mechanism for the formation of most granite magmas, which are generally thought to have formed at high T and are undersaturated with respect H_2O (Clemens, 1984; Pickering & Johnston, 1998).

Estimates of the amount of melt generated during partial melting have largely come from melting experiments on a variety of starting compositions including metapelites (Vielzeuf & Holloway, 1988; Carrington & Harley, 1995; Gardien *et al.*, 1995; Pickering & Johnston, 1998), metagreywackes (Patino-Douce & Beard, 1996; Vielzeuf & Montel, 1994; Montel & Vielzeuf, 1997), tonalites (Huang & Wyllie, 1986; Rutter & Wyllie, 1988; Skjerlie & Johnston, 1993; Patino-Douce, 1997; 2004) and amphibolites (Rushmer, 1991; Wolf & Wyllie, 1994). The majority of recent experimental studies have focussed on fluid-absent melting because of its perceived importance in magma genesis and crustal differentiation. The results of these fluid-absent melting studies indicate that different crustal lithologies display markedly different melting behaviour at similar P – T conditions. Furthermore, different melt fractions have been reported for compositionally-similar starting materials (e.g. Gardien *et al.*, 1995).

Fluid-absent melting experiments are often described as having ‘no added water’ or being ‘vapour-absent’. This is generally achieved by pre-heating the starting material above $\sim 150^\circ\text{C}$ for a sustained period (*ca.* 24 hours), causing any adsorbed H_2O to escape, prior to sealing the experimental charge (e.g. Patino-Douce, 2004). It is then assumed that any H_2O retained within the starting material is structurally bound within hydrous minerals (e.g. muscovite, biotite and amphibole). However, hydrous minerals that are stable up to 150°C at ambient pressures may begin to break down, via subsolidus dehydration reactions,

76 as the experimental charge is heated (and pressurised) to the desired conditions of the
 77 experimental run. Because of the intentional ‘closed system’ nature of most high P melting
 78 experiments, any H_2O generated via subsolidus dehydration will be effectively trapped
 79 within the experimental charge. In this way a ‘vapour absent’ starting material can evolve,
 80 becoming ‘fluid-present’ at the experimental P – T conditions. Additionally, London *et al.*
 81 (2012) suggested that powdered samples loaded in air may absorb between 1 and 4 weight
 82 percent H_2O before the experimental charge is sealed. It has also been suggested that
 83 water is often lost from the experimental charge via diffusion of molecular H_2O through the
 84 metal casing at temperatures above $\sim 950^\circ C$ as suggested by decreasing melt modes and
 85 increasing modal proportions of solid phases (e.g. plagioclase and garnet) with increasing T
 86 during experimental runs (Patino-Douce & Beard, 1994). The combined effects of
 87 absorption during handling and diffusive loss at high temperature work in opposite
 88 directions, thereby increasing the uncertainty in the actual H_2O content inside the
 89 experimental charge during high temperature melting experiments. Furthermore, if H^+ is
 90 lost through the capsule rather than molecular H_2O , as has been suggested by some
 91 authors, this may affect the oxidation state during a melting experiment (Patino-Douce &
 92 Beard, 1994; White *et al.*, 2011).

93 An important constraint on the amount of H_2O available for melting reactions during
 94 regional metamorphism can be obtained by considering the evidence for the nature of the
 95 fluid regime prior to the onset of melting. During prograde metamorphism of rocks with an
 96 initially high H_2O content (e.g. metapelites and metagreywackes), free H_2O is likely to be
 97 continuously generated via devolatilisation reactions up to the point at which the wet
 98 solidus is crossed with increasing T and/or P . Above this point, H_2O is consumed via wet
 99 melting reactions and decreases until it is no longer stable as a free phase at progressively

100 higher T. Several authors have argued that any H₂O generated during subsolidus
 101 dehydration has a strong tendency to escape the source rock due to the combined effects of
 102 compaction and increasing fluid pressure (Thompson & Connolly, 1990; Connolly, 1997;
 103 Yardley, 2009). Decreasing H₂O content with increasing metamorphic grade is observed in
 104 most metamorphic terrains providing evidence that much of the H₂O generated during
 105 subsolidus dehydration escapes the system prior to the onset of melting (Miyashiro, 1961).
 106 Indeed, the loss of fluid from metamorphic terrains may be a requirement for preserving
 107 metamorphic isograds during retrograde cooling and/or decompression (Thompson, 1983;
 108 Guiraud *et al.*, 2001). Subsolidus H₂O loss (SWL) may be especially significant at the
 109 pressure conditions of the middle to lower crust where rock porosity is thought to be small
 110 (<0.1–0.5 %), and positive excursions in hydrostatic pressure above the prevailing
 111 lithostatic pressure cannot be maintained for significant time periods (Norton & Knapp,
 112 1977; Connolly, 1997; 2010).

113 Importantly, there are few existing datasets to constrain sub-solidus open system
 114 processes involving H₂O, and experimentally determined phase relationships do not
 115 adequately constrain melting relationships, particularly with respect to H₂O. Quantitative
 116 thermodynamic calculations using internally-consistent datasets can be readily employed in
 117 multicomponent systems that approach the compositional complexity of natural rocks (e.g.
 118 White & Powell, 2002; Powell *et al.*, 2005; White *et al.*, 2005; White & Powell, 2010). *P-X*
 119 or *T-X* pseudosections can be used to visualise the effect of changes in bulk composition, as
 120 a result of the loss of a fluid and/or solid component from the system. This kind of
 121 thermodynamic modelling has the potential to complement experimental studies to provide
 122 a clearer picture of open-system processes that are thought to occur in the continental
 123 crust. Here we use pseudosections to investigate the effect of SWL on melting relationships

for a variety of bulk compositions representing common crustal rock types (metapelites, metagreywackes, tonalites, granodiorites). The impact of SWL on melt fertility is modelled at temperatures appropriate to partial melting during regional metamorphism at mid- to lower-crustal pressures. The modelling results are compared to the results derived from melting experiments to provide improved constraints on the nature of melting during crustal anatexis.

CALCULATED PHASE RELATIONS

Mineral equilibria calculations were undertaken using THERMOCALC 3.37i (Powell & Holland, 1988; updated 2013) with an updated version of the internally consistent data set of Holland & Powell (1998; data set `tcds55`, file created 22 November 2003). The calculations were undertaken in the chemical system $\text{Na}_2\text{O}-\text{CaO}-\text{K}_2\text{O}-\text{FeO}-\text{MgO}-\text{Al}_2\text{O}_3-\text{SiO}_2-\text{H}_2\text{O}-\text{TiO}_2-\text{O}$ (NCKFMASHTO) of White *et al.* (2007). The phases considered in the calculations, and references to the activity-composition models used are: garnet, biotite and silicate melt (White *et al.*, 2007), cordierite and epidote (Holland & Powell, 1998), orthopyroxene and magnetite (White & Powell, 2002), muscovite (Coggon & Holland, 2002), plagioclase - K-feldspar (Holland & Powell, 2003), ilmenite (White *et al.*, 2000), and hornblende (Diener *et al.*, 2007). Sillimanite, kyanite and quartz are pure end-member phases. Mineral abbreviations used in diagrams and in the text are: opx – orthopyroxene; g – garnet; cd – cordierite; hb – hornblende; bi – biotite; mu – muscovite; ky – kyanite; sill – sillimanite; ksp – K-feldspar; pl – plagioclase; ilm – ilmenite; mt – magnetite; q – quartz; liq – silicate liquid/melt.

Bulk compositions for phase equilibria modelling were derived from starting

compositions of published melting experiments for which melt fraction (Φ) estimates are available (Table 1). The four starting compositions were chosen to represent common felsic rock types and have compositions that are comparable to the average compositions of the middle and upper crust (Fig. 1). Two metapelitic bulk compositions are from Vielzeuf & Holloway (1988) and Gardien *et al.* (1995) and denoted VH88P and G95P, respectively. VH88P is modelled on a natural metapelite from the Cabo Ortegal metamorphic complex (Galicia, NW Spain) composed of quartz (39 wt%), plagioclase (19 wt%), kyanite (7 wt%), muscovite (9 wt%), biotite (21 wt%), garnet (2 wt%), staurolite (1 wt%), secondary chlorite (1 wt%) and accessory minerals (1 wt%). VH88P is strongly peraluminous and has comparatively high molar K_2O/Na_2O and high FeO contents. G95P is derived from a model two-mica metapelite generated from a mixture of quartz (39 wt%), plagioclase (30 wt%), muscovite (15 wt%) and biotite (15 wt%). It is weakly peraluminous and has comparatively low K_2O/Na_2O , high SiO_2 and low CaO. A metagreywacke bulk composition (denoted VM94G) is derived from melting experiments on a natural metagreywacke from the vallée de la Beaume (Ardèche, France) composed of quartz (40 wt%), plagioclase (32 wt%), and biotite (25 wt%), and accessory minerals (3 wt%; Vielzeuf & Montel, 1994; Montel & Vielzeuf, 1997). This composition is weakly peraluminous with high K_2O/Na_2O , SiO_2 and low FeO. A tonalitic bulk composition (denoted PD97T) was derived from melting experiments on a natural tonalite starting material from the Sierra Nevada Batholith (Lee Vining Canyon area; Patino-Douce, 1997). The sample was reported to contain quartz (20 wt%), plagioclase (45 wt%), K-feldspar (7 wt%), hornblende (13 wt%), biotite (13 wt%) and accessory minerals (2 wt%). PD97T is metaluminous with low K_2O/Na_2O , SiO_2 and high CaO and Na_2O .

Some of the bulk compositions contain low concentrations of F, P, Mn and CO_2 . Small

170 amounts P_2O_5 , CO_2 and MnO are unlikely to have a measurable effect on the melting
 171 behaviour of felsic rocks, whereas F contents in the range 0.01–4 wt% may have the effect of
 172 lowering the solidus of granitic melts (Manning, 1981) and stabilising biotite to higher
 173 temperatures (Peterson *et al.*, 1991). Fluorine is not currently incorporated into the
 174 activity-composition models for biotite or melt. As a result, the models presented here may
 175 differ slightly in terms of melting behaviour from the more complex starting compositions of
 176 the corresponding experiments. Furthermore, it should be noted that the melt model used
 177 here does not incorporate TiO_2 or ferric iron. TiO_2 is readily incorporated into biotite and
 178 oxide minerals (e.g. ilmenite and magnetite) and is important for stabilising biotite at high
 179 temperature (White *et al.*, 2007). Hayden & Watson (2007) investigated the incorporation
 180 of TiO_2 into haplogranitic melt. Their results indicated that even for rutile-saturated bulk
 181 compositions the amount of TiO_2 incorporated into the melt phase was small and as such,
 182 is unlikely to have a first order effect on melt fertility. The ferric iron content of all of the
 183 bulk compositions in this study are fixed at low values such that the only stable iron oxide
 184 minerals are ilmenite $\pm$ magnetite.

185 For each starting composition the H_2O content was varied to produce a low- H_2O and a
 186 high- H_2O end-member (Table 2). The high and low values were chosen to encompass the
 187 transition from H_2O undersaturated to H_2O saturated conditions at the wet solidus. Each
 188 end-member pair were then used to construct T – M_{H_2O} pseudosections (Guiraud *et al.*,
 189 2001) for each model starting composition. Diagrams were constructed at 10 kbar for the
 190 starting compositions VH88P, G95P and VM94G and at 8 kbar for composition PD97T to
 191 facilitate comparison with the reported experimental results. Models were also generated at
 192 5 kbar for VH88P, G95P and VM94G and at 4 kbar for PD97T to investigate melting
 193 equilibria at low pressure.

Pseudosections

T – $M_{\text{H}_2\text{O}}$ pseudosections calculated for compositions VH88P, G95P, VM94G at 10 kbar, and PD97T at 8 kbar over the temperature range 450 to 1100 °C are shown in Fig. 2. The pseudosections for all four compositions have a sub horizontal H_2O -saturated solidus (wet-solidus). VH88P has the highest T wet solidus at $\sim 680^\circ\text{C}$. The wet solidi for G95P, VM94G and PD97T are all similar, occurring at $\sim 645^\circ\text{C}$. In each case the wet solidus extends beyond the range of the high H_2O end-member but is limited at low H_2O contents by the stability of free H_2O . The point at which the solidus is minimally saturated with respect to H_2O is referred to here as the minimum saturation point (MSP; Fig. 2). The narrow fields immediately above the wet-solidus in all four T – $M_{\text{H}_2\text{O}}$ pseudosections are low variance fields related to the underlying NCKASH univariant reaction,

The H_2O -out boundaries of these ‘wet-melting’ fields are intersected within 10–40 °C above the wet-solidus. The upper boundaries increase in T only slightly as they trend from the minimum saturation point towards the high H_2O end-member for each diagram.

At H_2O contents less than the value of the minimum saturation point the temperature of the solidus increases steeply as the equilibrium composition becomes increasingly undersaturated with respect to H_2O . Inflections in the solidus curves at low H_2O contents on each of the T – $M_{\text{H}_2\text{O}}$ pseudosections correspond to changes in the modal mineralogy of low variance fields emanating from ‘vapour absent’ univariant reactions in the underlying NCKASH and NCKFMASH systems including,

For all compositions, the subsolidus H₂O-out boundary trends to lower bulk H₂O contents as it approaches the solidus (at the minimum saturation point) with increasing T (Fig. 2). The H₂O-out boundary for VH88P rises steeply from <450 °C up to the plagioclase-in boundary (~615 °C) and then migrates towards a lower bulk H₂O value as it approaches the solidus. For compositions G95P and VM94G the H₂O-out boundary is steep and approximately linear from <450 °C up to the wet-solidus. The H₂O-out boundary for PD97T trends gradually towards lower bulk H₂O, with distinct inflections corresponding to muscovite-out, K-feldspar-in and hornblende-in boundaries, as it approaches the minimum saturation point at the wet solidus with increasing temperature.

The pseudosections for VH88P and G95P both contain sillimanite and kyanite. For VH88P, sillimanite is stable from the sill-ky phase transition (~800 °C) up to >1000 °C at high H₂O contents and is stable above 1100 °C at low H₂O contents (Fig. 2 a & b). Kyanite is stable from 800 °C down to below 450 °C over the entire range of modelled H₂O contents (0.11–14.02 mol%). Sillimanite is stable for G95P from the sill-ky phase transition to above 1100 °C near the low-H₂O end-member, but its upper limit of stability decreases to ~900 °C at the high-H₂O end-member. Kyanite is stable from 800 °C down to 450 °C at H₂O contents below ~3 mol%. Above this value the stability is limited by the kyanite-out boundary along the low T sides of the quadrivariant bi-mu-ksp-pl-liq-ilm-q-ky and the univariant bi-mu-pl-liq-ilm-q-ky fields. Neither sillimanite nor kyanite are stable for

233 VM94G and PD97T over the range modelled (Fig. 2 c & d).

234 K-feldspar is unstable at temperatures above the solidus at high bulk H₂O contents for
 235 all four modelled compositions. At bulk H₂O contents close to the minimum saturation
 236 point, K-feldspar is stable over the ranges ~790–1000 °C, ~700–980 °C, ~810–990 °C and
 237 <450–990 °C for compositions VH88P, G95P, VM94G and PD97T, respectively.

238 It is important to note that even at high T (>1000 °C) none of the bulk compositions
 239 reach their liquidus and form pure melts (including the metatonalite - PD97T; Fig. 2).
 240 Even for the high H₂O end-members the high T equilibrium assemblages comprise liq-g-ilm,
 241 liq-opx-ilm, liq-opx-ilm-q and liq-hb-pl-ilm for VH88P, G95P, VM94G and PD97T,
 242 respectively. Lower variance high T assemblages become increasingly stable at lower bulk
 243 H₂O contents in each pseudosection.

244 Ilmenite is stable for all for starting compositions over the entire modelled ranges of
 245 temperature and H₂O contents. The pseudosection for composition G95P contains rutile at
 246 low temperature and magnetite is stable for intermediate H₂O contents between
 247 800–880 °C. Staurolite is stable from just below to just above the solidus for composition
 248 VM94G. While these phases are involved in the NCKFMASHTO equilibria, their modal
 249 abundances (1-oxide molar basis) are always low (<0.01) and are, therefore, unlikely to
 250 exert a significant influence on the melting behaviour of the system modelled.

251 Prograde metamorphism and sub-solidus water loss

252 The contours of melt and free H₂O corresponding to the pseudosections in Fig. 2 are shown
 253 in Figure 3. While each T – $M_{\text{H}_2\text{O}}$ pseudosection displays distinct phase relationships in
 254 detail, the relationships between the free H₂O and melt bearing regions in each case are

255 broadly similar. The proportion of free H₂O present along a particular prograde heating
 256 path in the subsolidus (vertical line) can be described in terms of df/dT (representing the
 257 change in free H₂O productivity with increasing T). Similarly, the amount of melt produced
 258 along a path in the suprasolidus can be described in terms of dm/dT (i.e. change in melt
 259 productivity with increasing T ; Johnson *et al.*, 2008).

260 At sub-solidus conditions df/dT is positive from $<450^\circ\text{C}$ up to the solidus at bulk
 261 H₂O values above the H₂O-out boundary for each of the four compositions (grey regions in
 262 Fig. 3). As a consequence, all four modelled compositions will produce free H₂O along a
 263 prograde path, with fixed H₂O contents, that traverses the ‘free H₂O-stable’ region (i.e.
 264 bulk H₂O contents greater than the value of the minimum saturation point). At
 265 supra-solidus conditions, H₂O & melt are both stable within the narrow wet-melting fields.
 266 Within this region, df/dT becomes negative and the amount of free H₂O decreases rapidly
 267 towards the H₂O-out boundary with increasing T . This region also corresponds to an initial
 268 increase in melt production with melt contours that are perpendicular to the T -axis
 269 ($dm/dT > 10 \text{ mol\%/}10^\circ\text{C}$). Above the H₂O+melt region the melt contours have a slope
 270 influenced by the shape of the H₂O-undersaturated solidus such that increases in T and/or
 271 bulk H₂O correspond to increased melt production.

272 An idealised melting experiment involves conserving all of the starting components
 273 (including the initial H₂O contents) throughout the experimental run. This behaviour is
 274 represented by the conservation of subsolidus water (CSW) scenario, shown by the vertical
 275 grey arrows in Fig. 3. The water contents for the CSW scenario for VH88P, G95P and
 276 VM94G are derived from the reported water contents for the bulk starting material of the
 277 original experiments (2.15 w%, 1.9 w% and 1.44 w%, respectively; Table 2). The
 278 H₂O contents of PD97T was not reported. The CSW path for this composition was chosen

so as to intersect the observed melt fraction in the experiments (i.e. $\Phi = 0.30$ melting at 950 °C; Patino-Douce, 1997).

If subsolidus water loss occurs, free H₂O will be incrementally lost along the prograde heating path (Guiraud *et al.*, 2001), continuously driving the bulk composition towards the H₂O-out boundary. To reduce the bulk H₂O contents of the source rock below the value of the H₂O-out boundary would require the liberation of structurally bound H₂O from hydrous minerals. As there is no viable mechanism to achieve this, the subsolidus H₂O-out boundary effectively limits the minimum H₂O-contents of the source rock. As the rock crosses the solidus with increasing temperature any remaining free H₂O is rapidly consumed within the wet-melting fields and the newly generated melt becomes increasingly undersaturated with respect to H₂O. Because such melts are effectively ‘water-starved’ as soon as they cross the solidus, the H₂O contents of the system become fixed at the value of the minimum saturation point, if H₂O loss is efficient and the porosity is small, as generally suggested. As a result an isobaric prograde heating path for a rock composition affected by SWL will involve:

1. A sub-solidus path that traverses the free H₂O region at bulk H₂O contents just greater than the H₂O-out boundary as it approaches the minimum saturation point;
2. A supra-solidus path with the bulk H₂O fixed at the value of the minimum saturation point up to the maximum temperature of the system.

In natural systems, H₂O may additionally be lost via incorporation into escaping melt or as part of hydrous peritectic minerals that are themselves entrained into escaping melt (Clemens & Stevens, 2012). However, both of these mechanisms will lead to other changes in the bulk composition that cannot be determined by varying any single compositional

variable. The appropriate diagrams for modelling such phenomena are more complex and involve varying the equilibrium bulk composition at each increment of melt- and/or mineral-loss.

The mineral mode changes along the subsolidus sections of the SWL paths were calculated along the H₂O-out boundary so that the bulk H₂O contents continuously changes along this segment. The supra-solidus portion of the SWL path was modelled at a fixed H₂O value (corresponding to the minimum saturation point; Table 2). Changes in the modal abundances of the equilibrium assemblages along the modelled CSW and SWL paths for the four starting compositions are shown in Fig. 4.

For composition VH88P the SWL heating path results in significantly less melt generation when compared with the CSW path (especially at low T; Fig. 4a & 4b). The reduction in melt fraction along the SWL path corresponds to increases in the modal proportions of quartz, plagioclase and biotite. K-feldspar is stabilised at temperatures above $\sim 790^\circ\text{C}$, whereas K-feldspar is completely absent in the CSW scenario. Very high dm/dT melting (20 mol%/10 $^\circ\text{C}$) along the CSW path ($\Phi < 0.01$ to ~ 0.11) corresponds to the H₂O-out boundary just above the solidus, at $\sim 680^\circ\text{C}$. This feature is not observed along the SWL path. Moderate dm/dT melting (~ 3 mol%/10 $^\circ\text{C}$) over the interval $790\text{--}880^\circ\text{C}$ on the SWL path corresponds to breakdown of biotite and increased growth of garnet. Melting at higher temperatures continues to increase in a broadly monotonic fashion reaching melt fractions of ~ 0.65 and ~ 0.44 at 1000°C for the CSW and SWL paths, respectively ($dm/dT \sim 1\text{--}2$ mol%/10 $^\circ\text{C}$). Interestingly, muscovite begins to breakdown below the solidus (at $\sim 615^\circ\text{C}$, corresponding to the first appearance of plagioclase) and completely disappears just above the solidus, at $\sim 720^\circ\text{C}$, for both the SWL and CSW paths; however, there is no significant increase in the melt fraction corresponding to the

disappearance of muscovite from the equilibrium assemblage along either path.

The SWL path for G95P corresponds to reduced melt fertility and slight increases in the modal proportions of quartz and plagioclase at temperatures just above the solidus ($\sim 640^\circ\text{C}$) and an increase in the proportion of K-feldspar at temperatures above 760°C when compared to the CSW path (Fig. 4c & 4d). Two steep increases in melting occur along the CSW path at low melt fraction. The first corresponds to the H_2O -out boundary just above the solidus ($dm/dT \sim 6\text{--}8 \text{ mol\%/}10^\circ\text{C}$). The second step corresponds to the abrupt disappearance of muscovite and the first appearance of kyanite and K-feldspar (at $\sim 760^\circ\text{C}$; $dm/dT \sim 10\text{--}20 \text{ mol\%/}10^\circ\text{C}$). Due to the negligible amount of H_2O at the solidus in the SWL scenario the melt fraction remains low ($\Phi < 0.02$) until muscovite disappears at $\sim 760^\circ\text{C}$ at which point it increases abruptly to ($\Phi \sim 0.09$). Melting at higher temperatures continues to increase steeply with melt fractions reaching ~ 0.69 and ~ 0.51 at 1000°C for the CSW and SWL paths, respectively ($dm/dT \sim 3\text{--}5 \text{ mol\%/}10^\circ\text{C}$).

For composition VM94G, the SWL scenario results in significantly reduced melt fertility, moderate increases in the modal proportions of quartz and plagioclase at temperatures just above the solidus (Fig. 4e & 4f). K-feldspar increases in modal abundance and becomes stabilised over a broader temperature range ($\sim 805\text{--}1000^\circ\text{C}$) when compared to the CSW path ($\sim 905\text{--}930^\circ\text{C}$). A sharp increase in melting along the CSW path corresponds to the H_2O -out boundary just above the solidus (at $\sim 650^\circ\text{C}$; $dm/dT \sim 8\text{--}10 \text{ mol\%/}10^\circ\text{C}$). Again, this feature is absent from the SWL path. Immediately above the solidus the melt fraction remains low ($\Phi < 0.01$) along the SWL path with an initial, gradual increase in melt fraction, beginning at $\sim 755^\circ\text{C}$, corresponding to the first appearance of garnet and the initial breakdown of biotite (dm/dT increasing from < 0.1 up to $2 \text{ mol\%/}10^\circ\text{C}$). A steep increase in melting, from $\Phi \approx 0.09$ to $\Phi \approx 0.19$, corresponds to

the final disappearance of biotite and the first appearance of orthopyroxene across the temperature interval $\sim 895\text{--}910\text{ }^{\circ}\text{C}$ for the SWL path ($dm/dT \sim 7\text{ mol\%/10 }^{\circ}\text{C}$). A similar step occurs along the CSW path over the same temperature interval with melting abruptly increasing from $\Phi \approx 0.28$ to $\Phi \approx 0.36$. Above the biotite-out boundary, melt fractions climb steeply to values of ~ 0.56 and ~ 0.34 at $1000\text{ }^{\circ}\text{C}$ for the CSW and SWL paths, respectively ($dm/dT \sim 3\text{--}4\text{ mol\%/10 }^{\circ}\text{C}$).

The SWL scenario for composition PD97T results in decreased melt fertility and slight increases in the modal proportions of quartz, plagioclase and K-feldspar when compared to the CSW path (Fig. 4g & 4h). A small, moderate increase in melt fraction, from 0.0 to ~ 0.02 , occurs just above the solidus, at $\sim 640\text{ }^{\circ}\text{C}$, along the CSW path ($dm/dT \sim 5\text{ mol\%/10 }^{\circ}\text{C}$). Again, this initial step in melt fraction is absent from the SWL path. The melt fraction along the CSW path increases gently to 0.09 over the interval $\sim 645\text{--}840\text{ }^{\circ}\text{C}$ ($dm/dT \sim 0.1\text{--}0.2\text{ mol\%/10 }^{\circ}\text{C}$) and then abruptly increases to 0.17 between $\sim 840\text{--}850\text{ }^{\circ}\text{C}$ ($dm/dT \sim 10\text{ mol\%/10 }^{\circ}\text{C}$). A similar sharp increase in melt fraction occurs along the SWL path (from $\Phi \sim 0.03$ to 0.10) over the same temperature interval. For both paths this step corresponds to the first appearance of orthopyroxene, the breakdown and disappearance of biotite and an increase in the modal abundance of K-feldspar. Above the biotite-out boundary melt fractions increase moderately steeply to reach ~ 0.45 and ~ 0.36 at $1000\text{ }^{\circ}\text{C}$ for the CSW and SWL paths, respectively ($dm/dT \sim 10\text{ mol\%/10 }^{\circ}\text{C}$). Interestingly, hornblende modes do not noticeably decrease for either path over the calculated melting interval ($\sim 640\text{--}1100\text{ }^{\circ}\text{C}$).

Melting at low pressure

The minimum saturation point for each composition, and the corresponding SWL melting path, increases as a function of pressure. Therefore, it is not possible to create a single P - T pseudosection for a bulk composition affected by SWL. Rather it is necessary to determine T - $M_{\text{H}_2\text{O}}$ phase relationships and H_2O value of the minimum saturation point at each pressure of interest. As a result this method is only applicable to the consideration of isobaric melting paths. To examine the effect of pressure on the phase equilibria, temperature-mode diagrams were generated for the SWL scenario along prograde isobaric heating paths for VH88P, G95P and VM94G at 5 kbar and PD97T at 4 kbar (Fig. 5).

At 5 kbar, VH88P (SWL) produces very little melt ($\Phi < 0.03$) between the solidus, at $\sim 690^\circ\text{C}$, and $\sim 775^\circ\text{C}$ ($dm/dT \ll 0.1 \text{ mol\%/}10^\circ\text{C}$). The melt fraction then rises sharply to $\Phi \approx 0.07$ at $\sim 780^\circ\text{C}$ ($dm/dT \sim 10\text{--}20 \text{ mol\%/}10^\circ\text{C}$). This step corresponds to the disappearance of sillimanite from the equilibrium assemblage, a sharp decrease in the modal abundance of biotite, an increase in the modal abundance of cordierite and the first appearance of garnet. This feature is absent from the SWL scenario for composition VH88P at 10 kbar. Above $\sim 780^\circ\text{C}$ the melt fraction increases moderately steeply reaching $\Phi \approx 0.67$ at $\sim 1000^\circ\text{C}$ ($dm/dT \sim 2\text{--}3 \text{ mol\%/}10^\circ\text{C}$; *c.f.* 0.44 for VH88P at 10 kbar).

The melt fraction for composition G95P (SWL) at 5 kbar remains low (< 0.01) between the solidus ($\sim 655^\circ\text{C}$) and $\sim 675^\circ\text{C}$ ($dm/dT \ll 0.1 \text{ mol\%/}10^\circ\text{C}$). Between $675\text{--}680^\circ\text{C}$ the melt fraction jumps abruptly to ~ 0.07 ($dm/dT \sim 10\text{--}20 \text{ mol\%/}10^\circ\text{C}$). This step corresponds to the disappearance of muscovite and the first appearance of sillimanite in the equilibrium assemblage. A similar high dm/dT step is observed at $\sim 760^\circ\text{C}$ in the SWL melting curve for G95P at 10 kbar. From $680\text{--}835^\circ\text{C}$ the melt fraction increases gradually

394 to 0.19 ($dm/dT \sim 0.5\text{--}0.7 \text{ mol\%/10 }^\circ\text{C}$). From 835 $^\circ\text{C}$ the melting curve begins to increase
 395 steeply corresponding to the breakdown and disappearance of biotite and the first
 396 appearance of orthopyroxene in the equilibrium assemblage ($dm/dT \sim 8 \text{ mol\%/10 }^\circ\text{C}$).
 397 Melting then proceeds with a steep slope reaching 0.70 at $\sim 1000^\circ\text{C}$ (dm/dT
 398 $\sim 4\text{--}5 \text{ mol\%/10 }^\circ\text{C}$; *c.f.* 0.51 for G95P at 10 kbar).

399 At 5 kbar, the melting curve for composition VM94G (SWL) increases slightly, from
 400 $\Phi = 0.0$ to 0.02, between the solidus (675 $^\circ\text{C}$) and the first appearance of garnet in the
 401 equilibrium assemblage at 785 $^\circ\text{C}$ ($dm/dT \ll 0.1 \text{ mol\%/10 }^\circ\text{C}$). Between 785–840 $^\circ\text{C}$ the melt
 402 fraction increases sharply to 0.22 corresponding to the first appearance of orthopyroxene
 403 and the breakdown and disappearance of biotite from the equilibrium assemblage (dm/dT
 404 $\sim 8\text{--}10 \text{ mol\%/10 }^\circ\text{C}$). Above 840 $^\circ\text{C}$ the melting curve rises steeply to reach 0.60 at $\sim 1000^\circ\text{C}$
 405 ($dm/dT \sim 4 \text{ mol\%/10 }^\circ\text{C}$; *c.f.* 0.34 for G95P at 10 kbar).

406 At 4 kbar PC97T (SWL) produces very little melt (< 0.02) between the solidus, at
 407 $\sim 655^\circ\text{C}$, and 795 $^\circ\text{C}$ corresponding to the first appearance of orthopyroxene (dm/dT
 408 $\ll 0.1 \text{ mol\%/10 }^\circ\text{C}$). Between 795–805 $^\circ\text{C}$ the melt fraction increases sharply to 0.11 (dm/dT
 409 $\sim 10\text{--}15 \text{ mol\%/10 }^\circ\text{C}$). This interval corresponds to the rapid breakdown and disappearance
 410 of biotite and an increase in the abundance of K-feldspar and orthopyroxene in the
 411 equilibrium assemblage. Above 805 $^\circ\text{C}$, the melt fraction increases moderately steeply to
 412 reach 0.47 at $\sim 1000^\circ\text{C}$ (*c.f.* 0.36 for G95P at 10 kbar).

CRUSTAL MELT FERTILITY

Comparisons with melting experiments

The experimentally derived melt fractions corresponding to VH88P, G95P, VM94G and PD97T are shown in Figs. 3 & 6. As previously noted, the starting H₂O contents for the melting experiments reported by Patino-Douce (1997) are not known, so it is not possible to make a comparison with the modelled CSW melting path for PD97T. Nevertheless, it is still useful to compare an inferred CSW path from matching the reported experimental melt fraction with the modelled SWL scenario.

Vielzeuf & Holloway (1988) infer a very steep increase in melt fraction, from $\Phi = 0.10$ to 0.60 over the temperature range 850–900 °C corresponding to a biotite-out “reaction” (i.e. $\text{bi} + \text{sill/ky} + \text{pl} + \text{q} \rightarrow \text{liq} + \text{g} + \text{ksp}$; Fig. 6a). Above 900 °C, these authors describe a melting ‘plateau’ corresponding to the gradual breakdown of sillimanite and garnet. Neither of these features are observed for the CSW path for VH88P which displays broadly monotonic melting behaviour above 800 °C. At temperatures higher than ~950 °C, they report lower melt fractions than the VH88P() melting curve. A similar misfit is observed between the the experimental results of Vielzeuf & Montel (1994) and the modelled CSW path for VM94 at 1000 °C (Fig. 6c). Experiments above 950 °C were not reported in Gardien *et al.* (1995) or Patino-Douce (1997), so any high T effect cannot be assessed for these studies. Nevertheless, the generation of lower than predicted melt fractions at high T is consistent with observations from other experiments that melt modes often decrease at temperatures above 950 °C (possibly due to diffusive H₂O-loss through the wall of the experimental capsule; Patino-Douce & Beard, 1994).

The experimental melt fraction estimates reported by Gardien *et al.* (1995) show a

436 general trend of increasing melting with increasing T that broadly corresponds to the CSW
 437 melting curve for composition G95P (Fig. 6b). Interestingly, both the CSW and SWL
 438 melting curves for composition G95P record a steep increase in melt fraction at $\sim 760^\circ\text{C}$,
 439 corresponding to the mu-out boundary with increasing T , while the reported experimental
 440 melt fraction actually decreases by 5% between 750°C and 800°C before increasing at
 441 higher T . The melting estimates of Vielzeuf & Montel (1994) are slightly above the
 442 corresponding VM94G() melting curve at temperatures below $\sim 900^\circ\text{C}$ (Fig. 6b).

443 Importantly, the experimentally-derived melt fractions for all starting compositions are
 444 higher than the corresponding SWL melting curves as a function of T . This general result is
 445 consistent with the assumption that SWL constrained melting represents a minimum
 446 melting end-member for a particular bulk composition.

447 The general advantages and limitations of melting experiments and phase equilibria
 448 modelling have been discussed in detail elsewhere (e.g. Johnson *et al.*, 2008; White *et al.*,
 449 2011), so the following discussion focusses on the issues that are specific to estimating
 450 crustal melt fertility. The differences between the experimental estimates and the modelled
 451 CSW melting curves can be understood in terms of the different limitations inherent for
 452 each approach.

453 A key assumption of most melting experiments is that the analysed composition of the
 454 starting material is representative of the equilibrating bulk composition at experimental run
 455 conditions. As previously discussed above, the combined effects of H_2O absorption into
 456 powders during handling as well as diffusive H_2O loss through the experimental casing at
 457 high temperature may result in significant changes in the actual H_2O content of a given
 458 melting experiment. The CSW scenarios reported here represent idealised melting

experiments that are unaffected by these undesirable experimental phenomena.

Detailed melting experiments are labour and time intensive with individual experiments often taking several weeks to approach equilibration (e.g. Patino-Douce, 2004). As a result, the total number of data points generated is usually low and performing repeat experiments at the same P – T conditions is uncommon making it difficult to assess the statistical significance or the reproducibility of any single melting estimate. By contrast, phase equilibria modelling allows the determination of melt modes over a range of P – T – X conditions for which the constituent phase models are calibrated. This allows for the calculation of continuous melting curves along a particular P – T path of interest.

Melt fractions for each experimental charge are typically estimated either by mass balance calculations or by graphical assessment of the proportion of melt in two-dimensional back-scattered electron scans. Both methods have high analytical uncertainties that may be difficult to quantify (e.g. Vielzeuf & Montel, 1994). Quantitative determination of melt fractions from experimental data is particularly problematic at low melt fractions due to the small size of melt/glass volumes that may be observed and analysed within the matrix of the experimental charge. As a result, experimental studies can often only approximate the degree of melting occurring at temperatures near the solidus. This constraint does not apply to phase equilibria modelling where the degree of melting can be calculated in detail from the solidus up to high temperatures.

The NCKFMASH melt model used in this study is constrained by well calibrated experimental data on unary, binary and ternary subsystems (Holland & Powell, 2001). It can, therefore, be expected to satisfactorily describe the topology of the melt bearing fields in NCKFMASHTO, if not, for example, the precise PT position of fields. As previously

discussed, the effects of TiO_2 and Fe^{3+} on the modelled melt fertility are likely to be small and certainly less than the uncertainties inherent in the determination of experimental estimates on the corresponding natural rock compositions. The influence of other minor components (e.g. Mn, F, Cl, P, B) on melting behaviour cannot be assessed with the models used here and in any case it is unlikely that any of these components (with the possible exception of F) will significantly affect melting behaviour at the low concentrations reported for the experimental starting compositions (shown in Table 1). We, therefore, argue that the modelled melting curves presented here represent reliable estimates of the melt fertility of the starting compositions under conditions of CSW and SWL.

Estimates of crustal melt fertility

It is clear that the overall ‘shapes’ of the melting curves for the CSW and SWL scenarios for each starting composition are similar and reflect similar underlying melt-producing equilibria (Fig. 6). Inflections in the melting curves occur at similar T indicating that they are largely independent of bulk H_2O content over the ranges modelled. The key difference between the CSW and SWL curves, in each case, is that the CSW melting curve contains an initial steep increase, just above the solidus, corresponding to the disappearance of free H_2O from the equilibrium assemblage. This initial melting step effectively increases the baseline for subsequent melting but does not significantly affect the shape of the curve at higher T . The most obvious observation that can be made from comparing these curves is that CSW paths that incorporate excess free H_2O will yield more melt than the corresponding SWL path.

The SWL melting curves for each starting composition at high and low pressure are

shown in Figure 7. With decreasing pressure the solidus migrates to higher T (shown by the grey and black arrows in Fig. 7), while at the same time the H_2O solubility of the melt phase increases. At low pressure, SWL constrained heating paths intersect key melting equilibria at lower T compared to the corresponding high P paths (cf. SWL versus CSW paths at constant P ; Fig. 6). This is because the underlying fluid-absent melting reactions have a positive slope in P – T space. The resulting melting curves show increased melt generation in the lower pressure scenarios for all starting compositions. This is likely to be a general result that is applicable to most felsic rock compositions.

As noted above there is no obvious geological mechanism that would cause the bulk composition to contain less H_2O than the value corresponding to the minimum saturation point. As a result the SWL path results in the minimum melt fertility scenario for a given source rock. Increasing melt fertility above the minimum defined by efficient SWL may be achieved via the retention of sub-solidus H_2O or the addition of H_2O to the system at temperatures above the solidus (compare paths A, B & C in Fig. 8).

H_2O may be released from a source rock with a higher T solidus into an adjacent rock with a lower T solidus (White *et al.*, 2005). For example, at 10 kbar composition VH88P intersects its wet-solidus at $\sim 680^\circ\text{C}$ whereas the solidus for composition G95P occurs at $\sim 640^\circ\text{C}$. If these two compositions represent hypothetical adjacent source regions subjected to the same heating path, VH88P could provide up to 3% H_2O to G95P over the temperature interval between their respective solidi. assuming the source rocks were in equal proportions, this would result in a significant increase in the melt fertility of the G95P source rock ($\sim 10\%$ more melt at 800°C , $\sim 15\%$ more melt at 1000°C). Water may also be added to a particular source rock via dewatering of migrating melts. During cooling and crystallisation water is released as the melt composition intersects the H_2O saturated fields

near the solidus (e.g. paths D & E in Fig. 8). H₂O saturation in melt is strongly dependent
on pressure (Tuttle & Bowen, 1958; Johannes & Holtz, 1996). Therefore, the volume of
water released by a crystallising melt will depend on the initial H₂O content of the melt and
the pressure at which the melt intersects the H₂O-in boundary. These three mechanisms
may result in greater melt fertility than is defined by a SWL-moderated minimum melting
curve. However, under most geological conditions SWL paths and melt fertility are likely to
apply.

The modelling presented here does not take into consideration the effect of melt loss on
melt fertility of felsic rocks as this process involves modification of several compositional
vectors the cannot be represented on a simple T - M_{H_2O} diagram. The loss of melt will cause
the bulk composition to become more residual with the largest compositional change likely
to be a decrease in water content due to the hydrous nature of the melt derived from
melting of felsic rocks (White & Powell, 2002; 2010).

DISCUSSION and CONCLUSIONS

General conclusions

The water content of the rock comprises the sum of free H₂O and structurally-bound
H₂O residing in hydrous minerals (e.g. biotite, muscovite and hornblende). While free H₂O
can be lost from a source rock (due to compaction, deformation, migrating porosity, etc.),
structurally-bound H₂O is only made mobile when the hydrous host mineral becomes
unstable due to changes in the P - T - X conditions at equilibrium. This may occur at sub
solidus and/or suprasolidus conditions. Above the solidus, any H₂O liberated by the
breakdown of hydrous phases is incorporated into the melt phase (until it is H₂O

saturated). The H₂O contents of the systems described here cannot evolve to be less than the value of the minimum saturation point without also losing melt and/or solid phases (Fig. 8). In this way subsolidus water loss can be viewed as defining the minimum H₂O condition for a particular bulk composition that has undergone prograde heating. A corollary of this is that the subsolidus water loss path defines the minimum melt fertility of a rock, at least prior to melt loss.

Most melting experiments are likely to conserve any H₂O generated by devolatilisation of hydrous minerals as the experimental charge is heated and pressurised from ambient temperatures up to the conditions of the experimental run. Given the likelihood that subsolidus water loss is a significant phenomena affecting prograde metamorphic rocks, most experimentally derived melting curves will tend to overestimate the melt fertility of the starting material at the onset of melting within the continental crust. The modelled curves presented here allow subsolidus water loss to be taken into account and, therefore, allow determination of the minimum melting relationships for compositions of interest.

Implications for melt connectivity, melt loss and the rheology of the crust during anatexis

An important consequence of the reduced melt fertility implied by the subsolidus water loss model is that critical melt thresholds will be intersected at significantly higher temperatures (Fig. 6). At low-melt fractions melt will initially form in isolated domains along grain edges and corners between reactant phases (Mehnert *et al.*, 1973). As the melt fraction increases these isolated melt volumes begin to coalesce eventually forming an interconnected network through the rock volume (Vigneresse *et al.*, 1996). Rosenberg & Handy (2005) identified a

572 melt connectivity threshold which occurs at $\Phi \approx 0.07$ and corresponds to $\sim 80\%$ of grain
 573 boundaries containing melt. Experimental data suggests that rocks experience significant
 574 loss of strength between the onset of melting and the melt connectivity threshold (from
 575 ~ 800 MPa down to ~ 200 MPa; Rosenberg & Handy, 2005). It follows that rocks having
 576 experienced subsolidus water loss will remain comparatively stronger at higher
 577 temperatures than rocks that have not. For the bulk compositions modelled here the
 578 differences in the temperature of the melt connectivity threshold range from $<10^\circ\text{C}$ to
 579 $>150^\circ\text{C}$. Such a relationship between degree of melting, temperature and strength is likely
 580 to significantly influence the rheological behaviour of crust undergoing regional
 581 metamorphism and anatexis.

582 Higher degrees of melting are associated with critical thresholds governing melt
 583 segregation and the transition from a solid to a liquid rheology. The melt escape threshold
 584 refers to the melt fraction at which melt is able to segregate from the source (Vigneresse
 585 *et al.*, 1996). The melt escape threshold is likely to depend on a number of factors including
 586 viscosity and strain rate but is likely to occur at $\Phi \sim 0.2\text{--}0.25$ (Sawyer, 1994; Vigneresse
 587 *et al.*, 1996). The ‘solid-to-liquid transition’ is the melt content at the transition from a
 588 solid-supported to a liquid-supported structure (Rosenberg & Handy, 2005). The
 589 solid-to-liquid transition occurs at $\Phi \sim 0.4\text{--}0.6$. Both the melt escape threshold and the
 590 solid-to-liquid transition will be shifted to higher temperatures for a rock that has
 591 experienced subsolidus water loss.

Subsolidus water loss during magma crystallisation

The discussion so far has concerned the consequences of subsolidus water loss during

prograde heating, however crystallising magmas may also experience subsolidus water loss.

Large amounts of H_2O may be generated as a magma approaches and then crosses the

solidus if a magma had an initially high H_2O content. The magma is rapidly converted to

solids + H_2O across the narrow melt + H_2O fields just above the wet solidus (Paths D & E

in Fig. 8). Even at upper crustal conditions the inter-grain porosity is likely to be small and

the tendency for free H_2O to escape the now solid rock volume is likely to be high (Connolly,

1997). The escape of free H_2O will cause the bulk H_2O contents to decrease towards the

minimum saturation point. Further cooling will take the rock across the H_2O -out boundary

and the H_2O contents will become fixed. An important consequence of this is that the final

H_2O contents of the crystallised solid (e.g. a tonalite) will not reflect its H_2O content when

it was a magma but rather reflects the water contents in the vicinity of the minimum

saturation point. This would mean that analysing the H_2O contents of granites collected in

the field may tell us nothing about the H_2O contents of the original magma.

ACKNOWLEDGEMENTS

RP acknowledges support from Australian Research Council DP0987731. SM acknowledges

support from the Australian Research Council DP0987765

REFERENCES

References

- Annen, C., Blundy, J. D., Stephen, R. & Sparks, R. S. J., 2008. The sources of granitic melt in Deep Hot Zones. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, **97**, 297–309.
- Brown, G. C. & Fyfe, W. S., 1970. The production of granitic melts during ultrametamorphism. *Contributions to Mineralogy and Petrology*, **28**(4), 310–318.
- Carrington, D. P. & Harley, S. L., 1995. Partial melting and phase relations in high-grade metapelites: An experimental petrogenetic grid in the KFMASH system. *Contributions to Mineralogy and Petrology*, **120**(3-4), 270–291.
- Clemens, J. D., 1984. Water contents of silicic to intermediate magmas. *Lithos*, **17**, 273–287.
- Clemens, J. D., 1990. The granulite — granite connexion, *Granulites and crustal evolution*, pp. 25–36. Springer Netherlands.
- Clemens, J. D. & Stevens, G., 2012. What controls chemical variation in granitic magmas? *Lithos*, pp. 1–13.
- Coggon, R. & Holland, T. J. B., 2002. Mixing properties of phengitic micas and revised garnet-phengite thermobarometers. *Journal of Metamorphic Geology*, **20**(7), 683–696.
- Connolly, J. A. D., 1997. Devolatilization-generated fluid pressure and deformation-propagated fluid flow during prograde regional metamorphism. *Journal of Geophysical Research: Solid Earth*, **102**(B8), 18149–18173.

- 630 Connolly, J. A. D., 2010. The mechanics of metamorphic fluid expulsion.
631 *Elements*, **6**(3), 165–172.
- 632 Diener, J. F. A., Powell, R., White, R. W. & Holland, T. J. B., 2007. A new
633 thermodynamic model for clino- and orthoamphiboles in the system
634 Na₂O–CaO–FeO–MgO–Al₂O₃–SiO₂–H₂O–O. *Journal of Metamorphic
635 Geology*, **25**(6), 631–656.
- 636 Gardien, V., Thompson, A. B., Grujic, D. & Ulmer, P., 1995. Experimental melting of
637 biotite + plagioclase + quartz $\pm$ muscovite assemblages and implications for crustal
638 melting. *Journal of Geophysical Research: Solid Earth*, **100**(B8), 15581–15591.
- 639 Guiraud, M., Powell, R. & Rebay, G., 2001. H₂O in metamorphism and unexpected
640 behaviour in the preservation of metamorphic mineral assemblages. *Journal of
641 Metamorphic Geology*, **19**(4), 445–454.
- 642 Hayden, L. A. & Watson, E. B., 2007. Rutile saturation in hydrous siliceous melts and its
643 bearing on Ti-thermometry of quartz and zircon. *Earth and Planetary Science
644 Letters*, **258**(3-4), 561–568.
- 645 Holland, T. J. B. & Powell, R., 1998. An internally consistent thermodynamic data set for
646 phases of petrological interest. *Journal of Metamorphic Geology*, **16**(3), 309–343.
- 647 Holland, T. J. B. & Powell, R., 2001. Calculation of phase relations involving haplogranitic
648 melts using an internally consistent thermodynamic dataset. *Journal of
649 Petrology*, **42**(4), 673–683.
- 650 Holland, T. J. B. & Powell, R., 2003. Activity-composition relations for phases in

- 651 petrological calculations: An asymmetric multicomponent formulation. *Contributions to*
652 *Mineralogy and Petrology*, **145**(4), 492–501.
- 653 Holtz, F., Behrens, H., Dingwell, D. B. & Johannes, W., 1995. H₂O solubility in
654 haplogranitic melts: Compositional, pressure, and temperature dependence. *American*
655 *Mineralogist*, **80**, 94–108.
- 656 Huang, W. L. & Wyllie, P. J., 1986. Phase-relationships of gabbro-tonalite-granite-water at
657 15 kbar with applications to differentiation and anatexis. *American*
658 *Mineralogist*, **71**, 301–316.
- 659 Huppert, H. E. & Sparks, R. S. J., 1988. The generation of granitic magmas by intrusion of
660 basalt into continental crust. *Journal of Petrology*, **29**(3), 599–624.
- 661 Johannes, W. & Holtz, F., 1996. *Petrogenesis and experimental petrology of granitic rocks*,
662 Springer, Berlin.
- 663 Johnson, T. E., White, R. W. & Powell, R., 2008. Partial melting of metagreywacke: a
664 calculated mineral equilibria study. *Journal of Metamorphic Geology*, **26**(8), 837–853.
- 665 London, D., Morgan, VI, G. B. & Acosta-Vigil, A., 2012. Experimental simulations of
666 anatexis and assimilation involving metapelite and granitic melt. *Lithos*, **153**, 292–307.
- 667 Manning, D., 1981. The effect of fluorine on liquidus phase relationships in the system
668 Qz-Ab-Or with excess water at 1 kb. *Contributions to Mineralogy and Petrology*, .
- 669 Mehnert, K. R., Büsch, W. & Schneider, G., 1973. Initial melting at grain boundaries of
670 quartz and feldspar in gneisses and granulites. *Neues Jahrb. Mineral.*
671 *Monatsh*, **4**, 165–183.

- 672 Miyashiro, A., 1961. Evolution of metamorphic belts. *Journal of Petrology*, **2**(3), 277–311.
- 673 Montel, J.-M. & Vielzeuf, D., 1997. Partial melting of metagreywackes, Part II.
- 674 Compositions of minerals and melts. *Contributions to Mineralogy and*
- 675 *Petrology*, **128**(2-3), 176–196.
- 676 Norton, D. & Knapp, R., 1977. Transport phenomena in hydrothermal systems; the nature
- 677 of porosity. *American Journal of Science*, **277**, 937–981.
- 678 Patino-Douce, A. E., 1997. Generation of metaluminous A-type granites by low-pressure
- 679 melting of calc-alkaline granitoids. *Geology*, **25**(8), 743–746.
- 680 Patino-Douce, A. E., 2004. Vapor-absent melting of tonalite at 15–32 kbar. *Journal of*
- 681 *Petrology*, **46**(2), 275–290.
- 682 Patino-Douce, A. E. & Beard, J. S., 1994. H₂O loss from hydrous melts during fluid-absent
- 683 piston cylinder experiments. *American Mineralogist*, **79**, 585–588.
- 684 Patino-Douce, A. E. & Beard, J. S., 1996. Effects of P, f(O₂) and Mg/Fe ratio on
- 685 dehydration melting of model metagreywackes. *Journal of Petrology*, **37**(5), 999–1024.
- 686 Peterson, J. W., Chacko, T. & Kuehner, S. M., 1991. The effects of fluorine on the
- 687 vapor-absent melting of phlogopite + quartz; implications for deep-crustal processes.
- 688 *American Mineralogist*, **76**(3-4), 470–476.
- 689 Pickering, J. M. & Johnston, D. A., 1998. Fluid-absent melting behavior of a two-mica
- 690 metapelite: experimental constraints on the origin of Black Hills granite. *Journal of*
- 691 *Petrology*, **39**(10), 1787–1804.

- 692 Powell, R. & Holland, T. J. B., 1988. An internally consistent dataset with uncertainties
693 and correlations: 3. Applications to geobarometry, worked examples and a computer
694 program. *Journal of Metamorphic Geology*, **6**, 173–204.
- 695 Powell, R., Guiraud, M. & White, R. W., 2005. Truth and beauty in metamorphic phase
696 equilibria: Conjugate variables and phase diagrams. *Canadian Mineralogist*, **43**, 21–33.
- 697 Rosenberg, C. L. & Handy, M. R., 2005. Experimental deformation of partially melted
698 granite revisited: implications for the continental crust. *Journal of Metamorphic
699 Geology*, **23**, 19–28.
- 700 Rudnick, R. L. & Gao, S., 2003. Composition of the continental crust. In: *Treatise on
701 Geochemistry*, (eds Rudnick, R. L., Holland, H. D. & Turekian, K. K.), pp. 1–64. Elsevier.
- 702 Rushmer, T., 1991. Partial melting of two amphibolites: Contrasting experimental results
703 under fluid-absent conditions. *Contributions to Mineralogy and Petrology*, **107**(1), 41–59.
- 704 Rutter, M. J. & Wyllie, P. J., 1988. Melting of vapour-absent tonalite at 10 kbar to
705 simulate dehydration–melting in the deep crust. *Nature*, **331**(6152), 159–160.
- 706 Sawyer, E. W., 1994. Melt segregation in the continental crust. *Geology*, **22**(11), 1019.
- 707 Sawyer, E. W., 1998. Formation and evolution of granite magmas during crustal reworking:
708 The significance of diatexites. *Journal of Petrology*, **39**(6), 1147–1167.
- 709 Sawyer, E. W., 2010. Migmatites formed by water-fluxed partial melting of a
710 leucogranodiorite protolith: Microstructures in the residual rocks and source of the fluid.
711 *Lithos*, **116**(3–4), 273–286.

- 712 Skjerlie, K. P. & Johnston, A. D., 1993. Fluid-absent melting behavior of an F-rich tonalitic
713 gneiss at mid-crustal pressures: Implications for the generation of anorogenic granites.
714 *Journal of Petrology*, **34**(4), 785–815.
- 715 Thompson, A. B., 1983. Fluid-absent metamorphism. *Journal of the Geological*
716 *Society*, **140**(4), 533–547.
- 717 Thompson, A. B., 1996. Fertility of crustal rocks during anatexis. *Transactions of the Royal*
718 *Society of Edinburgh: Earth Sciences*, **87**(1-2), 1–10.
- 719 Thompson, A. B., 1999. Some time-space relationships for crustal melting and granitic
720 intrusion at various depths. *Geological Society, London, Special*
721 *Publications*, **168**(1), 7–25.
- 722 Thompson, A. B. & Connolly, J. A. D., 1990. Metamorphic fluids and anomalous porosities
723 in the lower crust. *Tectonophysics*, **182**(1-2), 47–55.
- 724 Thompson, A. B. & Connolly, J. A. D., 1995. Melting of the continental crust: Some
725 thermal and petrological constraints on anatexis in continental collision zones and other
726 tectonic settings. *Journal of Geophysical Research: Solid Earth*, **100**(B8), 15565–15579.
- 727 Tuttle, O. F. & Bowen, N. L., 1958. Origin of Granite in the light of experimental studies in
728 the system NaAlSi₃O₈–KAlSi₃O₈–SiO₂–H₂O, *GSA Memoirs*, pp. 1–146. Geological
729 Society of America.
- 730 Vielzeuf, D. & Holloway, J. R., 1988. Experimental determination of the fluid-absent
731 melting relations in the pelitic system. *Contributions to Mineralogy and*
732 *Petrology*, **98**, 257–276.

- 733 Vielzeuf, D. & Montel, J.-M., 1994. Partial melting of metagreywackes. Part I. Fluid-absent
734 experiments and phase relationships. *Contributions to Mineralogy and*
735 *Petrology*, **117**(4), 375–393.
- 736 Vigneresse, J. L., Barbey, P. & Cuney, M., 1996. Rheological transitions during partial
737 melting and crystallization with application to felsic magma segregation and transfer.
738 *Journal of Petrology*, **37**(6), 1579–1600.
- 739 White, R. W. & Powell, R., 2002. Melt loss and the preservation of granulite facies mineral
740 assemblages. *Journal of Metamorphic Geology*, **20**(7), 621–632.
- 741 White, R. W. & Powell, R., 2010. Retrograde melt-residue interaction and the formation of
742 near-anhydrous leucosomes in migmatites. *Journal of Metamorphic*
743 *Geology*, **28**(6), 579–597.
- 744 White, R. W., Pomroy, N. E. & Powell, R., 2005. An in situ metatexite-diatexite transition
745 in upper amphibolite facies rocks from Broken Hill, Australia. *Journal of Metamorphic*
746 *Geology*, **23**(7), 579–602.
- 747 White, R. W., Powell, R. & Holland, T. J. B., 2001. Calculation of partial melting
748 equilibria in the system Na₂O–CaO–K₂O–FeO–MgO–Al₂O₃–SiO₂–H₂O (NCKFMASH).
749 *Journal of Metamorphic Geology*, **19**(2), 139–153.
- 750 White, R. W., Powell, R. & Holland, T. J. B., 2007. Progress relating to calculation of
751 partial melting equilibria for metapelites. *Journal of Metamorphic*
752 *Geology*, **25**(5), 511–527.
- 753 White, R. W., Powell, R., Holland, T. J. B. & Worley, B. A., 2000. The effect of TiO₂ and
754 Fe₂O₃ on metapelitic assemblages at greenschist and amphibolite facies conditions:

- 755 mineral equilibria calculations in the system
756 K₂O–FeO–MgO–Al₂O₃–SiO₂–H₂O–TiO₂–Fe₂O₃. *Journal of Metamorphic*
757 *Geology*, **18**(5), 497–511.
- 758 White, R. W., Stevens, G. & Johnson, T. E., 2011. Is the crucible reproducible? Reconciling
759 melting experiments with thermodynamic calculations. *Elements*, **7**(4), 241–246.
- 760 Wolf, M. B. & Wyllie, P. J., 1994. Dehydration-melting of amphibolite at 10 kbar: the
761 effects of temperature and time. *Contributions to Mineralogy and*
762 *Petrology*, **115**(4), 369–383.
- 763 Yardley, B. W. D., 2009. The role of water in the evolution of the continental crust. *Journal*
764 *of the Geological Society*, **166**(4), 585–600.

765 TABLES

Table 1. Starting compositions used in this study in wt.% and equivalent NCKFMASHTO mol.% values. The compositions are and the corresponding experimental studies are as follows: VH88P - high-Al metapelite from Vielzeuf & Holloway (1988), G95P - low-Al metapelite from Gardien *et al.* (1995), VM94G - metagreywacke from Vielzeuf & Montel (1994), and PD97T - metatonalite from Patino-Douce (1997).

Sample	VH88P		G95P		VM94G		PD97T	
Units	wt. %	mol. %	wt. %	mol. %	wt. %	mol. %	wt. %	mol. %
SiO ₂	64.35	66.64	70.04	72.14	69.99	72.27	61.48	66.56
Al ₂ O ₃	18.13	11.06	14.92	9.06	12.96	7.89	17.07	10.87
Fe ₂ O ₃	—	—	—	—	0.45	—	—	—
FeO	6.26	5.42	3.65	3.14	4.42	4.16	5.92	5.35
MgO	2.44	3.77	1.39	2.13	2.36	3.64	2.68	4.32
MnO	0.09	—	0.09	—	0.06	—	0.11	—
CaO	1.52	1.69	1.79	1.98	1.67	1.85	5.39	6.24
Na ₂ O	1.66	1.67	2.65	2.65	2.95	2.95	3.88	4.07
K ₂ O	2.56	1.69	3.18	2.09	2.41	1.59	2.57	1.77
TiO ₂	0.82	0.64	0.38	0.29	0.7	0.54	0.9	0.73
P ₂ O ₅	—	—	—	—	0.2	—	—	—
CO ₂	—	—	—	—	0.1	—	—	—
water	2.15	7.42	1.9	6.52	1.43	4.92	—	—
F	—	—	—	—	0.09	—	—	—
S	—	—	—	—	0.067	—	—	—
B(ppm)	—	—	—	—	20	—	—	—
Total	99.98	100.00	99.99	100.00	99.857	99.81	99.89	100.00

Table 2. Compositions used to construct pseudosections (in mol.%).

Figure	Composition	Diagram	H ₂ O	SiO ₂	Al ₂ O ₃	CaO	MgO	FeO	K ₂ O	Na ₂ O	TiO ₂	O
Figures 2 & 3												
(a)	VH88P	$T-M_{\text{H}_2\text{O}}$ lhs	0.11	71.90	11.93	1.82	4.07	5.85	1.82	1.80	0.69	0.01
		$T-M_{\text{H}_2\text{O}}$ rhs	14.02	61.88	10.27	1.57	3.50	5.03	1.57	1.55	0.59	0.01
(b)	G95P	$T-M_{\text{H}_2\text{O}}$ lhs	0.11	76.94	9.67	2.11	3.35	2.27	2.23	2.82	0.31	0.20
		$T-M_{\text{H}_2\text{O}}$ rhs	13.80	66.39	8.34	1.82	2.89	1.96	1.92	2.44	0.27	0.17
(c)	VM94G	$T-M_{\text{H}_2\text{O}}$ lhs	0.52	75.62	8.25	1.93	3.81	4.35	1.66	3.09	0.56	0.20
		$T-M_{\text{H}_2\text{O}}$ rhs	12.41	66.58	7.26	1.70	3.36	3.83	1.46	2.72	0.50	0.17
(d)	PD97T	$T-M_{\text{H}_2\text{O}}$ lhs	0.50	66.22	10.83	6.22	4.30	5.33	1.76	4.06	0.73	0.05
		$T-M_{\text{H}_2\text{O}}$ rhs	9.92	59.95	9.81	5.63	3.90	4.83	1.60	3.67	0.66	0.05
Figure 4												
(a)	VH88P - CSW	mode(X)-T	7.42	66.63	11.06	1.69	3.77	5.42	1.69	1.67	0.64	0.01
(b)	VH88P - SWL	mode(X)-T	3.45	69.50	11.53	1.76	3.93	5.65	1.76	1.74	0.67	0.01
(c)	G95P - CSW	mode(X)-T	6.52	72.00	9.05	1.97	3.13	2.12	2.08	2.64	0.29	0.19
(d)	G95P - SWL	mode(X)-T	4.08	73.88	9.28	2.03	3.22	2.18	2.14	2.71	0.30	0.19
(e)	VM94G - CSW	mode(X)-T	5.28	72.01	7.85	1.84	3.63	4.14	1.58	2.94	0.54	0.19
(f)	VM94G - SWL	mode(X)-T	2.78	73.91	8.06	1.89	3.72	4.25	1.62	3.02	0.55	0.19
(g)	PD97T - CSW	mode(X)-T	3.19	64.43	10.54	6.05	4.19	5.19	1.71	3.95	0.71	0.05
(h)	PD97T - SWL	mode(X)-T	2.57	64.84	10.61	6.09	4.21	5.22	1.72	3.97	0.71	0.05
Figure 5												
(a)	VH88P - SWL	mode(X)-T	3.47	69.48	11.53	1.76	3.93	5.65	1.76	1.74	0.67	0.01
(b)	G95P - SWL	mode(X)-T	3.55	74.29	9.34	2.04	3.23	2.19	2.15	2.73	0.30	0.18
(c)	VM94G - SWL	mode(X)-T	2.88	73.84	8.05	1.88	3.72	4.25	1.62	3.02	0.55	0.19
(d)	PD97T - SWL	mode(X)-T	2.48	64.90	10.62	6.10	4.22	5.22	1.73	3.98	0.72	0.05

FIGURE CAPTIONS

Fig. 1 Harker plots of mol.% oxides, molar K_2O / Na_2O and ASI (molar $Al_2O_3 / (CaO + Na_2O + K_2O)$) for the starting compositions: VH88P - metapelite, G95P - metapelite, VM94G - metagreywacke and PD97T - metatonalite. L, M and U represent the estimated average compositions of the lower-, middle- and upper-crust, respectively, of Rudnick & Gao (2003). All values are plotted on an anhydrous basis.

Fig. 2 Calculated NCKFMASHTO T - M_{H_2O} pseudosections for the compositions: (a) VH88P - metapelite, (b) G95P - metapelite and (c) VM94G - metagreywacke at 10 kbar and (d) PD97T - metatonalite at 8 kbar (see Table 2). The dashed line indicates the solidus for each diagram. The white dot indicates the locus of the minimum saturation point for each diagram.

Fig. 3 Calculated contours for the modal abundances of melt (solid lines) and free H_2O (dashed lines) for the compositions: (a) VH88P - metapelite, (b) G95P - metapelite, (c) VM94G - metagreywacke and (d) PD97T - metatonalite (see Table 2). The dark grey areas represent the regions over which free-water is stable. The grey arrows represent the path followed for heating in a closed system with the H_2O contents fixed at the value reported for the experimental starting composition. The H_2O contents for the tonalitic bulk composition, (d), is unknown so the H_2O contents were determined so that 30% melt is generated at 950 °C as reported in Patino-Douce (1997). As with Fig. 2 the white dot indicates the locus of the minimum saturation point for each diagram. The black arrows represent heating combined with subsolidus water loss. The stars represent melt fraction estimates with increasing temperature derived from the corresponding melting experiments (assuming

uncertainty in the H₂O contents only; see text for discussion).

Fig 4 Calculated molecular proportions of phases (X) with increasing temperature for the four compositions: VH88P, G95P and VM94G at 10 kbar and PD97T at 8 kbar under conditions of conservation of water () and sub-solidus water loss (SWL). The dashed vertical line indicates the solidus for each diagram.

Fig 5 Calculated molecular proportions of phases (X) with increasing temperature for the four compositions: VH88P, G95P and VM94G at 5 kbar and PD97T at 4 kbar under conditions of sub-solidus water loss (SWL). The solidus is at the left edge for each diagram.

Fig 6 Comparison of the CSW and SWL melt fraction curves (shown in figure 5) with the available experimental estimates for the four compositions VH88P, G95P and VM94G at 10 kbar and PD97T at 8 kbar . The horizontal dashed lines indicate the melt fractions corresponding to the solid-liquid transition (SLT) and melt connectivity threshold (MCT; both after Rosenberg & Handy, 2005) and the melt escape threshold (MET; after Vigneresse *et al.*, 1996). The arrows indicate the difference in the temperature of the MCT between the CSW and SWL scenarios (see text for discussion).

Fig 7 Comparison of the melt fraction curves with increasing temperature at low P (light grey curve) and high P (dark grey curve) for the four compositions VH88P, G95P and VM94G (5 kbar and 10 kbar) and PD97T at (4 kbar and 8 kbar). The respective solidi at each pressure are indicated by the corresponding grey arrows. The position of the MCT is shown by the horizontal dashed line.

811 Fig 8 Schematic diagram showing the relationships between various melting paths (dark
812 grey dotted lines) and crystallisation paths (light grey dotted lines) as a function of
813 water contents. Thin grey lines represent melt contours (increasing to the top-right of
814 the diagram). The striped area is inaccessible because H_2O contents less than this
815 value are unobtainable without invoking changes in other compositional vectors (e.g.
816 via melt and or solid segregation; see text for discussion).

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure 8.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Webb, G;Powell, R;McLaren, S

Title:

Phase equilibria constraints on the melt fertility of crustal rocks: the effect of subsolidus water loss

Date:

2015-02-01

Citation:

Webb, G., Powell, R. & McLaren, S. (2015). Phase equilibria constraints on the melt fertility of crustal rocks: the effect of subsolidus water loss. JOURNAL OF METAMORPHIC GEOLOGY, 33 (2), pp.147-165. <https://doi.org/10.1111/jmg.12114>.

Persistent Link:

<http://hdl.handle.net/11343/112394>