Abstract

Currently there is a lack of evidence on whether women and men respond differently to ~~trauma-focused evidence-based~~ psychological treatments for PTSD. This study was a systematic review and meta-analysis to examine whether gender is associated with response to trauma-focused psychological interventions for PTSD~~gender modifies the effect of trauma-focused psychological interventions for PTSD~~. Randomized controlled trials comparing trauma-focused interventions for PTSD with comparison conditions were identified in a literature review. Fifty-six randomized controlled trials were included in the meta-analysis: 31 had a mixed gender sample, 19 were female only and six were male only. There was evidence that gender modified the effect of trauma-focused interventions, with larger effects for females compared with any condition based on the primary outcome of clinician-rated PTSD symptoms at post-intervention and short-term follow-up. This finding was supported by direct effects meta-analyses of studies that provided data on both females and males. There was also a larger effect for females in an analysis restricted to trials that evaluated an intervention likely to be provided in routine clinical practice. These findings support a gender difference in outcomes following trauma-focused psychological interventions for PTSD. Future research should seek to identify specific factors related to gender that facilitate or inhibit response to these interventions.

Key words: gender, systematic review, meta-analysis, posttraumatic stress disorder, treatment, trauma-focused, outcomes

According to DSM-5 (American Psychiatric Association, 2013), Posttraumatic Stress Disorder (PTSD) is a mental disorder characterized by severe and persistent stress reactions following exposure to a potentially traumatic event (PTE), including re-experiencing of the event, avoidance of memories and reminders of the event, negative cognitions and mood, and increased arousal symptoms. Gender differences in both exposure to PTEs and the development of PTSD following exposure have been well-documented. Women are more likely to report partner abuse, sexual assault, rape and childhood sexual abuse, whereas men are more likely to experience non-sexual physical assault, combat, accidental injury and witnessing someone being badly injured or killed (Mills et al., 2011; Tolin & Foa, 2006). Epidemiological studies from several countries have shown indicated that, while women are less likely than men to be exposed to traumatic events, the prevalence of PTSD within the community is significantly higher among women (Breslau et al., 1998; Chapman et al., 2012; de Vries & Olf, 2009; R. Kessler, Sonnega, Bromet, Hughes, & Nelson, 1995; R. C. Kessler, Sonnega, Hughes, & Nelson, 1995; Norris et al., 2003; Pietrzak, Goldstein, Southwick, & Grant, 2011; Stein, Walker, Hazen, & Forde, 1997). For example, Chapman and colleagues (2012) found that Australian women had a higher lifetime (9.7% versus 4.7%) and 12-month (5.9% versus 2.8%) rate of PTSD compared with men. PTEs involving interpersonal violence (that are more likely to be experienced by women) are associated with a high probability of PTSD (R. Kessler et al., 1995), although the higher prevalence of PTSD in women may not simply be the result of increased exposure to certain types of trauma (Tolin & Foa, 2006). Other factors to consider include gender differences in severity of trauma, type and severity of symptoms, comorbid mental and physical disorders, appraisal and processing of trauma, emotional expression, social support, and physiological reactivity (Cason, Grubaugh, & Resick, 2002; Christiansen & Elklit, 2013; Chung & Breslau, 2008; R. Kessler et al., 1995).

Currently, there is a lack of evidence to determine whether there is a gender difference in response to psychological and other interventions for PTSD. If firm evidence is found, then this would necessitate efforts to identify gender-related factors that influence this differential

Comment [DJW1]: Reference repeated for some reason

response in order to develop optimal treatment for both women and men with PTSD.

Previous systematic reviews and meta-analyses have found that the effects of different types of psychotherapies were larger in studies with a higher proportion of women (Watts et al., 2013), and that the effects of trauma-focused cognitive-behavioural therapies were larger in women-only studies (all sexual assault or assault survivors) compared with all studies (Bisson et al., 2007). However, both studies used 'indirect' analyses (based on female only, male only and mixed gender studies) to assess for a gender difference in outcomes, that is, regression analysis or sub-group analysis to assess whether gender predicted ~~or modified~~ the effect size. In contrast, direct effects analyses include only studies that have data from both women and men and calculate the difference between the treatment response for each in each study, then pools these differences across studies. This has the benefit of limiting clinical and methodological heterogeneity and potential confounders in included studies.

Comment [DJW2]: Need this reference

Comment [DJW3]: Change this to Cochrane 2013 reference

The aim of the current systematic review and meta-analysis was to undertake indirect and direct effects analyses to examine whether gender is associated with response to ~~gender modifies the effect of evidence-based~~ trauma-focused psychological interventions for PTSD, ~~that is, interventions that involve confronting trauma-related memories and situations and addressing unhelpful beliefs about the traumatic experience (Australian Centre for Posttraumatic Mental Health, 2013).~~

Method

The Cochrane Collaboration systematic review methodology (Higgins & Green, 2011) was used as a guide for this study.

Search Strategy

Relevant randomized controlled trials (RCTs) that were published up until July 11, 2012, were identified via MEDLINE, PsychINFO, Embase and Cochrane Central Register of Controlled Studies (CENTRAL). The keywords or equivalent terms used to search the databases were: 'posttraumatic stress disorder', 'post-traumatic stress disorder', 'stress disorder' and 'traumatic stress'. In addition, hand-searches were undertaken of the abstracts of the International Society for Traumatic Stress Studies (ISTSS) conferences for the years

2008 – 2012, as well as the reference list of the revised Australian PTSD guidelines (Australian Centre for Posttraumatic Mental Health, 2013).

Study selection

Any published or unpublished English language RCT that tested the efficacy of a trauma-focused psychological intervention for PTSD was considered for inclusion. A trauma-focused intervention was defined as one that encourages the person to confront their traumatic memory or situations avoided since the traumatic event, or that challenged the person's unhelpful thoughts or beliefs about their traumatic experience (Australian Centre for Posttraumatic Mental Health, 2013). There were no restrictions in terms of the mode of delivering the intervention, such that individual, group, face-to-face, self-help, internet and virtual reality interventions were all eligible for inclusion. Both single (i.e. female only or male only) and mixed (i.e. both females and males) gender studies were eligible. Studies with multiple intervention arms (e.g., psychological and pharmacotherapy) were included if at least one intervention arm was a trauma-focused intervention. Studies that compared two trauma-focused interventions only (without any other comparison group) were excluded.

For participants, the following inclusion criteria applied: mean age at least 15 years old; at least 70% had a primary diagnosis of PTSD or subclinical PTSD as defined as having at least one symptom in each of recognized three PTSD symptom clusters (e.g. re-experiencing, avoidance and arousal), or any acceptable and equivalent definition adopted by the authors; the diagnosis was based on a structured diagnostic or clinical interview using DSM or ICD criteria, or a self-rated measure of PTSD using a cut-off score pre-defined by the authors; and the duration of PTSD symptoms was at least one month.

Comparison conditions were categorized as follows: wait list or no intervention, usual care, attention control, pharmacotherapy, and non-trauma-focused psychological interventions. Individual comparison conditions were combined into 'any comparison' which was the primary comparison condition. Wait list or no intervention included placement on a wait list for intervention or no intervention at all. Usual care included interventions or services that would normally be offered in the absence of the study. Attention control included non-

directive interventions such as non-specific supportive counselling that was judged to primarily provide an equivalent period of contact time as the trauma-focussed intervention. Non-trauma-focused interventions included skills-based interventions that did not explicitly encourage the individual to confront their traumatic memory or situations, or challenge the person's unhelpful thoughts or beliefs about their traumatic experience, such as non-trauma-focussed cognitive and/or behavioural therapy, problem solving therapy and relaxation training.

~~Non-trauma-focused interventions included skills-based interventions that did not explicitly encourage the individual to confront their traumatic memory or situations, or challenge the person's unhelpful thoughts or beliefs about their traumatic experience.~~

The primary outcome measure was severity of PTSD symptoms using a standardized clinician-rated measure. The secondary outcome measure was severity of PTSD symptoms using a standardized self-rated measure.

Where a full-text publication was not available for a study which passed the screening stage, authors were contacted directly to request a manuscript. Two reviewers (D.K. and T.V.) independently assessed all potentially relevant articles for inclusion. Any disagreements about inclusion or exclusion of studies were resolved by discussion with other authors (S.H. and D.W.).

Assessment of risk of bias

Studies were assessed for risk of bias with regard to random sequence generation, allocation concealment, and blinded assessment of outcomes, respectively (Higgins & Green, 2011). Studies were further rated with regard to missing data ($\geq 15\%$ missing data, and methods of imputation). Two reviewers (D.K. and D.W.) independently carried out the assessment of risk of bias, with any disagreement resolved by discussion.

Data extraction

Data were extracted from all included studies following the full-text assessment. One reviewer (D.K.) extracted the data and two others (T.V. and D.W.) checked the data, with any disagreement resolved by discussion.

Statistical analysis

Common assessment time-points for RCTs include post-treatment, and 3 months and 6 months following intervention. For this reason, we sought outcome data for the following time-points: post-intervention, short-term follow-up (i.e. one to five months following intervention), and long-term follow-up (i.e. six or more months following intervention).

Outcome data were sought for the following time-points: post-intervention, short-term follow-up (i.e. one to five months following intervention), and long-term follow-up (i.e. six or more months following intervention). When a study provided two sets of outcome data for a single follow-up time-point, preference was given to the earlier ~~the data collected~~ ~~data at the~~ ~~earlier time-point~~.

When necessary, study authors were contacted directly to request gender-specific outcome data including the number of participants and the means and standard deviations of PTSD outcome measures.

For studies that included two trauma-focused interventions, the data from each arm were combined. For studies with more than one comparison condition, data from only one condition was included as 'any comparison' selected according to the following prioritization: wait list or no intervention, usual care, attention control, pharmacotherapy, and non-trauma-focused psychological intervention.

The Review Manager statistical software program (Review Manager (RevMan), 2011) was used for all analyses. The outcome measures were continuous and were analyzed as standardized mean differences (SMDs) with 95% confidence intervals to allow for ease of comparison across studies that used different measures of the same outcomes. For all meta-analyses, intention to treat (ITT) data were combined with completer data and the random effects model was used.

Statistical heterogeneity was assessed based on Cochrane recommendations: I^2 values of 0-40% may not be important; 30% to 60% may represent moderate heterogeneity; 50% to 90% may represent substantial heterogeneity; and 75% to 100% represents considerable heterogeneity (Higgins & Green, 2011).

Planned subgroup analyses were undertaken to investigate the impact of the comparison group on the treatment outcome. In order to examine whether gender modified intervention effects, subgroup analyses were undertaken on females versus males.

Direct effects meta-analyses were performed using studies that included and reported data on both females and males. This enabled us to investigate the impact of indirect effects from female only and male only studies on treatment outcome. The mean difference between the intervention effect of females and males and the standard error of this difference were calculated. The pooled mean difference was calculated using inverse variance weighting. DerSimonian and Laird's method of moments estimator was used to estimate between-study variance (Der Simonian & Laird, 1986). A second direct effects meta-analysis included only those mixed gender studies where participants experienced the same trauma type to avoid the potential confound of trauma type.

Sensitivity analyses were undertaken based on the primary comparison and outcome at the post-intervention time-point. Firstly, we investigated the impact of restricting the analysis to only those trials that tested an intervention most likely to be provided in routine practice, that is, the intervention was not exclusively group-based, self-help or delivered online, and the prescribed number of sessions was between 5 and 20. Secondly, we investigated the impact of excluding trials judged to be at high risk of bias due to the following not clearly stated to have been undertaken: random sequence generation, allocation concealment, and blinded assessment of outcomes. Additional analyses also examined the impact of excluding those trials with more than 15% of data unavailable for analysis, and those with only completer data available.

Results

Description of studies

We were unable to obtain data for 30 (34.9%) of the 86 primary studies that met the inclusion criteria (see flowchart of the search for studies in Figure 1). As a result, 56 studies provided data for inclusion in the meta-analysis: 31 mixed gender, 19 female only, and six male only. Usable data were obtained for females only in two mixed gender studies (Asukai

et al., 2010; Başoğlu et al., 2005), and males only in five mixed gender studies (Difede et al., 2007; Forbes et al., 2012; Litz et al., 2007; McLay et al., 2011; Nacasch et al., 2011). Details of studies included in the meta-analysis are shown in Table 1 in the online supplement.

The majority of trauma-focused interventions evaluated in these studies were cognitive and/or behavioral interventions. There were nine studies that evaluated two trauma-focused interventions (in addition to a comparison condition) for which the data were combined. The duration of interventions lasted between one day and 30 weeks.

Twenty-three studies included a wait list or no intervention comparison condition, nine included usual care, four included attention control, one included pharmacotherapy, and eight studies included a non-trauma-focused intervention.

Study participants experienced a wide range of trauma types, often within a single sample of participants. ~~Study participants experienced a wide range of trauma types, often within a single study.~~ Of note, participants experienced military-related combat or other trauma in nine studies, rape or sexual assault in nine studies, and child sexual or physical abuse in seven studies. Four studies did not report trauma type.

Risk of bias

For included studies, a high or unclear rating of bias was made for 28 (50.0%) studies regarding random sequence generation, 42 (75.0%) studies regarding allocation concealment, and 7 (12.5%) studies regarding blinded assessment of outcomes (see Table 2 in the online supplement). Forty studies (71.0%) provided only completer data, while 16 (29.0%) studies provided ITT data or reported no missing data. Twenty (35.7%) studies had more than 15% of data unavailable for analysis.

Gender differences in treatment effects of trauma-focused psychological interventions for PTSD

There were medium to large statistically significant reductions in clinician- and self-rated PTSD symptoms favoring trauma-focused interventions for females and males combined at all available time-points (see Table 3 in the online supplement). Heterogeneity was substantial for most of these analyses.

For the 'any comparison' condition based on clinician-rated PTSD symptoms, there was evidence that gender modified the intervention effect, with a larger effect for females compared with males at post-intervention (Standard Mean Difference (95% CI): -1.02 (-1.27, -0.76) versus -0.60 (-0.86, -0.33)) and short-term follow-up (-0.54 (-0.81, -0.26) versus -0.18 (-0.35, -0.00)). The forest plot of clinician-rated PTSD symptoms at post-intervention is shown in Figure 2. There was no evidence that gender modified the intervention effect at long-term follow-up, although it should be noted there were relatively few data for males. For self-rated PTSD symptoms, there was some evidence for a larger intervention effect for females at post-intervention (SMD (95% CI): -0.93 (-1.17, -0.69) for females versus -0.62 (-0.90, -0.34) for males), and for males at long-term follow-up (-0.50 (-0.81, -0.20) for females versus -1.02 (-1.46, -0.59) for males), although both results did not reach statistical significance and there were relatively few data available for males in the latter comparison. There was no evidence that gender modified the intervention effect at short-term follow-up.

Further analyses were conducted to examine whether gender modified the intervention effect for trauma-focused interventions compared with wait list or no intervention, usual care, attention control, pharmacotherapy, and non-trauma-focused interventions separately (detailed results provided in online supplement Table 3). For comparisons with wait list or no intervention, there was some evidence for a non-significant larger effect for females on clinician-rated PTSD symptoms at post-intervention (SMD (95% CI): -1.26 (-1.53, -0.98) versus -0.78 (-1.21, -0.35)), and evidence for a larger effect for females on self-rated PTSD symptoms at post-intervention (SMD (95% CI): -1.22 (-1.45, -0.99) vs -0.76 (-1.14, -0.38)). For comparisons with usual care, there was some evidence for a non-significant larger effect for females on clinician-rated PTSD symptoms at short-term follow-up (SMD (95% CI): -0.77 (-1.16, -0.38) vs -0.15 (-0.63, 0.33)). For comparisons with attention control there was some evidence for a non-significant larger effect for females compared with males on self-rated PTSD symptoms at post-intervention (-0.43 (-0.68, -0.17) vs -0.12 (-0.34, 0.09)). There were few data available for comparisons with pharmacotherapy and non-trauma-focused interventions, particularly for males in the latter analyses.

Direct effects meta-analyses

A direct effects meta-analysis that included only those studies that provided data on both females and males (Beck et al., 2009; Ehlers et al., 2003; Ehlers et al., 2005; T. Jensen et al., 2014; Monson et al., 2012; Monson et al., 2006; Mueser et al., 2008; Pacella et al., 2012; Schnyder et al., 2011; Shalev et al., 2012) demonstrated a larger effect for females compared with males (mean difference (95% CI): 10.00 (1.21, 18.79)). Heterogeneity appeared to be relatively unimportant (chi-square = 12.18, df=9, p=0.20; $I^2 = 26\%$). A second direct effects analysis was restricted to those studies that recruited participants with the same type of trauma, namely, two studies of motor vehicle accident survivors (Beck et al., 2009; Ehlers et al., 2003). The findings showed a non-significant larger effect for females compared with males (mean difference (mean difference (95% CI): 5.89 (-17.42, 29.21) with heterogeneity unlikely to be important ($I^2 = 0\%$)).

Sensitivity analyses

Seven of 38 studies (Başoğlu et al., 2005; Beck et al., 2009; Bohus et al., 2013; J. A. Jensen, 1994; Litz et al., 2007; Rothbaum, 1997; Schnurr et al., 2003) were excluded from an analysis restricted to trials that evaluated an intervention likely to be provided in routine practice. For two studies that tested two trauma-focused interventions, we used the 'unmerged' data for the intervention that did not include a non-trauma component; that is, prolonged exposure (and not prolonged exposure plus stress inoculation) in the Foa and colleagues (1999) study, and affect and interpersonal regulation plus prolonged exposure (and not supportive counselling plus prolonged exposure) in the Cloitre and colleagues (2010) study. The results showed that gender modified the intervention effect, with a larger effect for females compared with males (chi-square=4.33, df =1, p=0.04; heterogeneity for females and males combined: $I^2 = 78\%$).

Further analyses showed a smaller intervention effect for studies with low ratings of bias for random sequence generation (chi-square =7.36, df=1, p=0.007) and randomization allocation concealment (chi-square =6.54, df=1, p=0.01). There was also some evidence for a smaller intervention effect for studies with ITT data compared with completer data,

although this finding was not significant (chi-square =3.01, df = 1, p=0.08). The intervention effect was not modified by blinding of outcome assessors (chi-square =0.27, df=1, p=0.60) or the percentage of participants with more than 15% missing data (chi-square =0.45, df=1, p=0.50).

Discussion

The current study identified 56 randomized controlled trials that were included in the meta-analysis. The current study supports the available research evidence and practice guidelines that trauma-focused interventions are effective for both women and men with PTSD (Australian Centre for Posttraumatic Mental Health, 2013). The main finding was that women had greater reductions than men in the primary outcome measure of clinician-rated PTSD symptoms when trauma-focused psychological interventions were compared with 'any comparison' at both post-intervention and short-term follow-up. Women were also found to have greater reductions in self-rated PTSD symptoms at post-intervention when trauma-focused interventions were compared with wait list or no intervention. The main finding was supported by a direct effects meta-analysis of data from studies that included both women and men. Finally, there was a larger effect for women in an analysis restricted to trials that evaluated an intervention likely to be provided in routine clinical practice.

Methodological factors that are largely unrelated to gender, such as treatment quality and fidelity, the type of control condition, and the level of general functioning of patients, may help to explain the finding that women appear to respond better to psychological treatments for PTSD Some researchers have suggested that methodological factors that are largely independent of gender, such as treatment quality and fidelity, the type of control condition, and the level of general functioning of patients, may help to explain the finding that women appear to respond better to psychological treatments for PTSD (Cason et al., 2002). For example, it may be the case that there are substantive differences in methodology and quality between studies of (female) rape victims and (male) military veterans. In the current study, the results of the direct effects meta-analysis of data from studies that recruited both women and men suggests that the larger intervention effect for women is unlikely to be

related to such factors. Moreover, the current study provides some preliminary evidence of a gender difference in treatment response irrespective of trauma type. While not statistically significant, the results of the direct effects analysis of data from two treatment studies where participants were included on the basis of a single trauma type (motor vehicle accident) suggest that women had greater reductions in PTSD symptoms compared with men. It should be noted, however, that controlling for trauma type does not exclude the possibility that a difference between women and men in the *severity* of trauma exposure is related to a gender difference in treatment response. It is also possible that such a difference in treatment response may be related to other gender-related differences, including expression of emotions, coping strategies, interpersonal and social support, civilian versus military status, and the chronicity and severity of PTSD and comorbid mental health conditions (Cason et al., 2002). The present study was unable to determine whether such variables were implicated in gender differences in PTSD outcomes.

Some caution should be taken when interpreting the current findings. ~~First, First, one third of eligible studies were not included due to a lack of gender-specific data~~ ~~approximately one third of the studies deemed eligible for inclusion were not able to be included because of a lack of available or usable gender-specific data.~~ Most of the studies that were not included were mixed gender studies that could have contributed data to both indirect and direct effects meta-analyses, and the inclusion of these additional studies could potentially have changed the current findings. Second and relatedly, for some comparisons there were relatively few studies and data which resulted in inadequate power to assess whether gender modified the intervention effect. Third, there was significant statistical heterogeneity in many of the analyses undertaken. There was considerable clinical diversity within the trauma-focused interventions, which included cognitive and/or behavioral therapies, EMDR, narrative exposure therapy, and other types of interventions, and they varied considerably in terms of duration and planned number of sessions. The planned use of random effects models that take statistical heterogeneity into account was appropriate, and in particular the direct effects meta-analyses had low heterogeneity. Overall, however, it is unclear whether

heterogeneity may have influenced gender-related effects. Fourth, there was often inadequate reporting of aspects of the conduct of studies known to introduce risk of bias and potentially inflate intervention effects. For example, the risk of bias was assessed to be unclear for random sequence generation in almost half of the included studies, and unclear for randomization allocation concealment in three-quarters of the studies. Sensitivity analyses undertaken showed that studies with lower ratings of bias regarding some, but not all, aspects of the conduct of studies tended to have smaller intervention effects. Again, however, it is not clear if the quality of studies had a differential impact on the outcomes of females versus males.

The current study has a number of strengths. First, to our knowledge, this is the first study to undertake a systematic review and meta-analysis that included direct effects analyses to examine a potential gender difference in outcomes following psychological interventions for PTSD. Second, the current review was conducted using explicit and systematic methods that adhered closely to the Cochrane guidelines to minimize bias in the review process. Third, we attempted to find unpublished studies by hand-searching conference proceedings of the ISTSS, as well as searching the Cochrane Central Register of Controlled Studies which is a database that includes the results of grey literature searches undertaken by different review groups internationally.

The current findings highlight the need to more closely investigate potential gender differences in treatment response to psychological interventions for PTSD. Further well-designed controlled trials are required that report on outcomes for women and men separately. [Trials that recruit both women and men who have been exposed to the same discrete trauma type would be particularly useful to assess the influence of gender on treatment outcomes.](#) [More broadly,](#) future research efforts should seek to identify specific gender-related factors that facilitate or inhibit response to trauma-focused psychological interventions to inform enhancements or adaptations of those interventions for women and men with PTSD.

Acknowledgements

This study was supported by a Department of Veterans' Affairs (DVA) Applied Research Program Grant (1131). The study was undertaken and reported without editorial or other influence by DVA. The authors have no conflicts of interest to report.

References

- *Ahrens, J., & Rexford, L. (2002). Cognitive Processing Therapy for incarcerated adolescents with PTSD. *Journal of Aggression, Maltreatment & Trauma*, 6(1), 201-216.
- American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders* (5th ed.). Washington DC: American Psychiatric Association.
- Asukai, N., Saito, A., Tsuruta, N., Kishimoto, J., & Nishikawa, T. (2010). Efficacy of exposure therapy for Japanese patients with posttraumatic stress disorder due to mixed traumatic events: A randomized controlled study. *Journal of Traumatic Stress*, 23(6), 744-750. doi: 10.1002/jts.20589
- Australian Centre for Posttraumatic Mental Health. (2013). *Australian Guidelines for the Treatment of Acute Stress Disorder and Posttraumatic Stress Disorder*. Melbourne, Victoria: ACPMH.
- Başoğlu, M., Salcioğlu, E., Livanou, M., Kalender, D., & Acar, G. (2005). Single-session behavioral treatment of earthquake-related posttraumatic stress disorder: a randomized waiting list controlled trial. *Journal of Traumatic Stress*, 18(1), 1-11.
- Beck, J. G., Coffey, S. F., Foy, D. W., Keane, T. M., & Blanchard, E. B. (2009). Group Cognitive Behavior Therapy for chronic Posttraumatic Stress Disorder: An initial randomized pilot study. *Behavior Therapy*, 40(1), 82-92. doi: 10.1016/j.beth.2008.01.003
- Bohus, M., Dyer, A., Priebe, K., Kruger, A., Kleindienst, N., Schmahl, C., . . . Steil, R. (2013). DBT for PTSD after childhood sexual Abuse in patients with and without Borderline Personality Disorder: A randomised controlled trial. *Psychotherapy and Psychosomatics*, 82(4), 221-233. doi: 10.1159/000348451

- Breslau, N., Kessler, R. C., Chilcoat, H. D., Schultz, L. R., Davis, G. C., & Andreski, P. (1998). Trauma and posttraumatic stress disorder in the community. *Archives of General Psychiatry*, 55(7), 626-632.
- Carlson, J. G., Chemtob, C. M., Rusnak, K., Hedlund, N. L., & Muraoka, M. Y. (1998). Eye movement desensitization and reprocessing (EDMR) treatment for combat-related posttraumatic stress disorder. *Journal of Traumatic Stress*, 11(1), 3-24. doi: 10.1023/a:1024448814268
- Cason, D., Grubaugh, A., & Resick, P. (2002). Gender and PTSD treatment: Efficacy and effectiveness. In R. Kimerling, P. Ouimette & J. Wolfe (Eds.), *Gender and PTSD* (pp. 305-334). New York: Guilford Press.
- Chapman, C., Mills, K., Slade, T., McFarlane, A. C., Bryant, R. A., Creamer, M., . . . Teesson, M. (2012). Remission from post-traumatic stress disorder in the general population. *Psychological Medicine*, 42(8), 1695-1703. doi: 10.1017/S0033291711002856
- Chard, K. M. (2005). An evaluation of cognitive processing therapy for the treatment of posttraumatic stress disorder related to childhood sexual abuse. *Journal of Consulting and Clinical Psychology*, 73(5), 965-971. doi: 10.1037/0022-006x.73.5.965
- Christiansen, D., & Elklit, A. (2013). Sex differences in PTSD. In E. Ovuga (Ed.), *Posttraumatic Stress Disorders in a Global Context* (pp. 113-142): InTech.
- Chung, H., & Breslau, N. (2008). The latent structure of post-traumatic stress disorder: tests of invariance by gender and trauma type. *Psychological Medicine*, 38(4), 563-573. doi: 10.1017/s0033291707002589
- Cloitre, M., Koenen, K. C., Cohen, L. R., & Han, H. (2002). Skills training in affective and interpersonal regulation followed by exposure: A phase-based treatment for PTSD

- related to childhood abuse. *Journal of Consulting and Clinical Psychology*, 70(5), 1067-1074. doi: 10.1037/0022-006x.70.5.1067
- Cloitre, M., Stovall-McClough, K. C., Nooner, K., Zorbas, P., Cherry, S., Jackson, C. L., . . . Petkova, E. (2010). Treatment for PTSD related to childhood abuse: A randomized controlled trial. *American Journal of Psychiatry*, 167(8), 915-924. doi: 10.1176/appi.ajp.2010.09081247
- de Vries, G. J., & Olf, M. (2009). The lifetime prevalence of traumatic events and posttraumatic stress disorder in the Netherlands. *Journal of Traumatic Stress*, 22(4), 259-267.
- Der Simonian, R., & Laird, N. (1986). Meta-analysis in clinical trials. *Controlled Clinical Trials*, 7, 177-188.
- Difede, J., Malta, L. S., Best, S., Henn-Haase, C., Metzler, T., Bryant, R., & Marmar, C. (2007). A randomized controlled clinical treatment trial for World Trade Center attack-related PTSD in disaster workers. *Journal of Nervous and Mental Disease*, 195(10), 861-865. doi: 10.1097/NMD.0b013e3181568612
- DuHamel, K. N., Mosher, C. E., Winkel, G., Labay, L. E., Rini, C., Meschian, Y. M., . . . Redd, W. H. (2010). Randomized clinical trial of telephone-administered cognitive-behavioral therapy to reduce post-traumatic stress disorder and distress symptoms after hematopoietic stem-cell transplantation. *Journal of Clinical Oncology*, 28(23), 3754-3761. doi: 10.1200/jco.2009.26.8722
- Dunne, R. L., Kenardy, J., & Sterling, M. (2012). A randomized controlled trial of cognitive behavioral therapy for treatment of PTSD in the context of chronic Whiplash. *Clinical Journal of Pain*, 28(9), 755-765. doi: 10.1097/AJP.0b013e318243e16b

- Echeburua, E., De Corral, P., Sarasua, B., & Zubizarreta, I. (1996). Treatment of Acute Posttraumatic Stress Disorder in rape victims: An experimental study. *Journal of Anxiety Disorders, 10*(3), 185-200. doi: 10.1016/0887-6185(96)89842-2
- Ehlers, A., Clark, D. M., Hackmann, A., McManus, F., Fennell, M., Herbert, C., & Mayou, R. (2003). A randomized controlled trial of cognitive therapy, a self-help booklet, and repeated assessments as early interventions for posttraumatic stress disorder. *Archives of General Psychiatry, 60*(10), 1024-1032. doi: 10.1001/archpsyc.60.10.1024
- Ehlers, A., Clark, M., Hackmann, A., McManus, F., & Fennell, M. (2005). Cognitive therapy for post-traumatic stress disorder: development and evaluation. *Behavior Research and Therapy, 43*, 413-431. doi: 10.1016/j.brat.2004.03.006
- Ertl, V., Pfeiffer, A., Schauer, E., Elbert, T., & Neuner, F. (2011). Community-implemented trauma therapy for former child soldiers in Northern Uganda: A randomized controlled trial. *JAMA, 306*(5), 503-512. doi: 10.1001/jama.2011.1060
- Falsetti, S. A., Resnick, H. S., & Davis, J. L. (2008). Multiple Channel Exposure Therapy for women with PTSD and comorbid panic attacks. *Cognitive Behaviour Therapy, 37*(2), 117-130. doi: 10.1080/16506070801969088
- Feske, U. (2008). Treating low-income and minority women with posttraumatic stress disorder: A pilot study comparing prolonged exposure and treatment as usual conducted by community therapists. *Journal of Interpersonal Violence, 23*(8), 1027-1040. doi: 10.1177/0886260507313967
- Foa, E. B., Dancu, C. V., Hembree, E. A., Jaycox, L. H., Meadows, E. A., & Street, G. P. (1999). A comparison of exposure therapy, stress inoculation training, and their combination for reducing posttraumatic stress disorder in female assault victims. *Journal of Consulting and Clinical Psychology, 67*(2), 194-200. doi: 10.1037/0022-006x.67.2.194

- Foa, E. B., Hembree, E. A., Cahill, S. P., Rauch, S. A. M., Riggs, D. S., Feeny, N. C., & Yadin, E. (2005). Randomized trial of prolonged exposure for posttraumatic stress disorder with and without cognitive restructuring: Outcome at academic and community clinics. *Journal of Consulting and Clinical Psychology*, 73(5), 953-964. doi: 10.1037/0022-006x.73.5.953
- Foa, E. B., Rothbaum, B. O., Riggs, D. S., & Murdock, T. B. (1991). Treatment of posttraumatic stress disorder in rape victims: a comparison between cognitive-behavioral procedures and counseling. *Journal of consulting and clinical psychology*, 59(5), 715-723. doi: 10.1037//0022-006x.59.5.715
- Forbes, D., Lloyd, D., Nixon, R. D. V., Elliott, P., Varker, T., Perry, D., . . . Creamer, M. (2012). A multisite randomized controlled effectiveness trial of cognitive processing therapy for military-related posttraumatic stress disorder. *Journal of Anxiety Disorders*, 26(3), 442-452. doi: 10.1016/j.janxdis.2012.01.006
- Higgins, J. P. T., & Green, S. (2011). *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0*.
- Jensen, J. A. (1994). An investigation of eye movement desensitization and reprocessing (EMD/R) as a treatment for posttraumatic stress disorder (PTSD) symptoms of Vietnam combat veterans. *Behavior Therapy*, 25(2), 311-325. doi: 10.1016/S0005-7894(05)80290-4
- Jensen, T., Holt, T., Ormhaug, S., Granly, L., Egeland, K., Hoaas, L. C., . . . Wenzel-Larsen, T. (2014). A randomized effectiveness study comparing Trauma-Focused Cognitive Behavioral Therapy with therapy as usual for youth. *Journal of Child and Adolescent Psychology*, 43(3), 356-369. doi: 10.1080/15374416.2013.822307
- Katz, L., Douglas, S., Zaleski, K., Williams, J., Huffman, C., & Cojucar, G. (2014). Comparing Holographic Reprocessing and Prolonged Exposure for women veterans

- with sexual trauma: A randomized trial. *Journal of Contemporary Psychotherapy*, 44(1), 9-19. doi: 10.1037/0033-3204.45.2.186
- Kessler, R., Sonnega, A., Bromet, E., Hughes, M., & Nelson, C. B. (1995). Posttraumatic stress disorder in the National Comorbidity Survey. *Arch General Psychiatry*, 52(12), 1048 - 1060.
- Kessler, R. C., Sonnega, A., Hughes, M., & Nelson, C. B. (1995). Posttraumatic stress disorder in the National Comorbidity Survey. *Archives of General Psychiatry*, 52(12), 1048-1060.
- Knaevelsrud, C., & Maercker, A. (2007). Internet-based treatment for PTSD reduces distress and facilitates the development of a strong therapeutic alliance: a randomized controlled clinical trial. *BMC Psychiatry*, 7(13). doi: 10.1186/1471-244X-7-13
- Kubany, E. S., Hill, E. E., & Owens, J. A. (2003). Cognitive trauma therapy for battered women with PTSD: preliminary findings. *Journal of Traumatic Stress*, 16(1), 81-91. doi: 10.1023/A:1022019629803
- Kubany, E. S., Hill, E. E., Owens, J. A., Iannce-Spencer, C., McCaig, M. A., Tremayne, K. J., & Williams, P. L. (2004). Cognitive trauma therapy for battered women with PTSD (CTT-BW). *Journal of Consulting and Clinical Psychology*, 72(1), 3-18. doi: 10.1037/0022-006X.72.1.3
- Litz, B. T., Engel, C. C., Bryant, R. A., & Papa, A. (2007). A randomized, controlled proof-of-concept trial of an Internet-based, therapist-assisted self-management treatment for posttraumatic stress disorder. *The American Journal of Psychiatry*, 164(11), 1676-1683. doi: 10.1016/0005-7967(96)00033-2.; 10.1037/1082-989x.7.2.147; 10.1176/appi.ajp.2007.06122057
- McDonagh, A., McHugo, G., Sengupta, A., Demment, C. C., Schnurr, P. P., Friedman, M., . . . Descamps, M. (2005). Randomized trial of cognitive-behavioral therapy for chronic

posttraumatic stress disorder in adult female survivors of childhood sexual abuse.

Journal of Consulting and Clinical Psychology, 73(3), 515-524. doi: 10.1037/0022-006X.73.3.515

McGovern, M. P., Lambert-Harris, C., Alterman, A. I., Xie, H., & Meier, A. (2011). A randomized controlled trial comparing integrated cognitive behavioral therapy versus individual addiction counseling for co-occurring substance use and posttraumatic stress disorders. *Journal of Dual Diagnosis*, 7(4), 207-227.

McLay, R. N., Wood, D. P., Webb-Murphy, J. A., Spira, J. L., Wiederhold, M. D., Pyne, J. M., & Wiederhold, B. K. (2011). A randomized, controlled trial of virtual reality-graded exposure therapy for post-traumatic stress disorder in active duty service members with combat-related post-traumatic stress disorder. *Cyberpsychology, Behavior, and Social Networking*, 14(4), 223-229. doi: 10.1089/cyber.2011.0003

Mills, K. L., McFarlane, A. C., Slade, T., Creamer, M., Silove, D., Teesson, M., & Bryant, R. (2011). Assessing the prevalence of trauma exposure in epidemiological surveys. *Australian and New Zealand Journal of Psychiatry*, 45(5), 407-415. doi: 10.3109/00048674.2010.543654

Monson, C. M., Fredman, S., Macdonald, A., Pukay, N. D., Resick, P., & Schnurr, P. (2012). Effect of cognitive-behavioral couple therapy for PTSD: A randomized controlled trial. *JAMA*, 308(7), 700-709. doi: 10.1001/jama.2012.9307

Monson, C. M., Schnurr, P. P., Resick, P. A., Friedman, M. J., Young-Xu, Y., & Stevens, S. P. (2006). Cognitive processing therapy for veterans with military-related posttraumatic stress disorder. *Journal of Consulting and Clinical Psychology*, 74(5), 898-907. doi: 10.1037/0022-006x.74.5.898

Mueser, K., Rosenberg, S. D., Xie, H., Jankowski, M. K., Bolton, E. E., Lu, W., . . . Wolfe, R. (2008). A randomized controlled trial of cognitive-behavioral treatment for

- posttraumatic stress disorder in severe mental illness. *Journal of Consulting and Clinical Psychology*, 76(2), 259-271. doi: 10.1037/0022-006x.76.2.259
- Nacasch, N., Foa, E. B., Huppert, J. D., Tzur, D., Fostick, L., Dinstein, Y., . . . Zohar, J. (2011). Prolonged exposure therapy for combat- and terror-related posttraumatic stress disorder: a randomized control comparison with treatment as usual. *Journal of Clinical Psychiatry*, 72(9), 1174-1180. doi: 10.4088/JCP.09m05682blu
- Neuner, F., Kurreck, S., Ruf, M., Odenwald, M., Elbert, T., & Schauer, M. (2010). Can asylum-seekers with posttraumatic stress disorder be successfully treated? A randomized controlled pilot study. *Cognitive Behaviour Therapy*, 39(2), 81-91. doi: 10.1080/16506070903121042
- Neuner, F., Onyut, P. L., Ertl, V., Odenwald, M., Schauer, E., & Elbert, T. (2008). Treatment of posttraumatic stress disorder by trained lay counselors in an African refugee settlement: A randomized controlled trial. *Journal of Consulting and Clinical Psychology*, 76(4), 686-694. doi: 10.1037/0022-006x.76.4.686
- Neuner, F., Schauer, M., Klaschik, C., Karunakara, U., & Elbert, T. (2004). A comparison of narrative exposure therapy, supportive counseling, and psychoeducation for treating posttraumatic stress disorder in an African refugee settlement. *Journal of Consulting and Clinical Psychology*, 72(4), 579-587. doi: 10.1037/0022-006x.72.4.579
- Norris, F. H., Murphy, A. D., Baker, C. K., Perilla, J. L., Rodriguez, F. G., & Rodriguez, J. D. (2003). Epidemiology of trauma and posttraumatic stress disorder in Mexico. *Journal of Abnormal Psychology*, 112(4), 646-656.
- Pacella, M. L., Armelie, A., Boarts, J., Wagner, G., Jones, T., Feeny, N., & Delahanty, D. L. (2012). The impact of prolonged exposure on PTSD symptoms and associated psychopathology in people living with HIV: A randomized test of concept. *AIDS and Behavior*, 16(5), 1327-1340. doi: 10.1007/s10461-011-0076-y

- Pietrzak, R. H., Goldstein, R. B., Southwick, S. M., & Grant, B. F. (2011). Prevalence and Axis I comorbidity of full and partial posttraumatic stress disorder in the United States: Results from Wave 2 of the National Epidemiologic Survey on Alcohol and Related Conditions. *Journal of Anxiety Disorders*, 25(3), 456-465. doi: 10.1016/j.janxdis.2010.11.010
- Power, K., McGoldrick, T., Brown, K., Buchanan, R., Sharp, D., Swanson, V., & Karatzias, A. (2002). A controlled comparison of eye movement desensitization and reprocessing versus exposure plus cognitive restructuring versus waiting list in the treatment of post-traumatic stress disorder. *Clinical Psychology and Psychotherapy*, 9(5), 299-318. doi: 10.1002/cpp.341
- Ready, D. J., Gerardi, R. J., Backscheider, A. G., Mascaro, N., & Rothbaum, B. O. (2010). Comparing virtual reality exposure therapy to present-centered therapy with 11 U.S. Vietnam veterans with PTSD. *Cyberpsychology, Behavior and Social Networking*, 13(1), 49-54. doi: 10.1089/cyber.2009.0239
- Resick, P. A., Nishith, P., Weaver, T. L., Astin, M. C., & Feuer, C. A. (2002). A comparison of cognitive-processing therapy with prolonged exposure and a waiting condition for the treatment of chronic posttraumatic stress disorder in female rape victims. *Journal of Consulting and Clinical Psychology*, 70(4), 867-879. doi: 10.1037//0022-006x.70.4.867
- Review Manager (RevMan). (2011). Review Manager (Version 5.1.). Copenhagen: The Nordic Cochrane Centre: The Cochrane Collaboration.
- Rothbaum, B. O. (1997). A controlled study of eye movement desensitization and reprocessing in the treatment of posttraumatic stress disorder sexual assault victims. *Bulletin of the Menninger Clinic*, 61(3), 317-334.

- Rothbaum, B. O., Astin, M. C., & Marsteller, F. (2005). Prolonged Exposure versus Eye Movement Desensitization and Reprocessing (EMDR) for PTSD rape victims. *Journal of Traumatic Stress, 18*(6), 607-616. doi: 10.1002/jts.20069
- Schaal, S., Elbert, T., & Neuner, F. (2009). Narrative exposure therapy versus interpersonal psychotherapy: A pilot randomized controlled trial with Rwandan genocide orphans. *Psychotherapy and Psychosomatics, 78*(5), 298-306. doi: 10.1002/jts.20104; 10.1159/000229768
- Scheck, M. M., Schaeffer, J. A., & Gillette, C. (1998). Brief psychological intervention with traumatized young women: the efficacy of eye movement desensitization and reprocessing. *Journal of Traumatic Stress, 11*(1), 25-44. doi: 10.1023/a:1024400931106
- Schnurr, P. P., Friedman, M. J., Engel, C. C., Foa, E. B., Shea, M. T., Chow, B. K., . . . Bernardy, N. (2007). Cognitive behavioral therapy for posttraumatic stress disorder in women - A randomized controlled trial. *JAMA, 297*(8), 820-830. doi: 10.1001/jama.297.8.820
- Schnurr, P. P., Friedman, M. J., Foy, D. W., Shea, M. T., Hsieh, F. Y., Lavori, P. W., . . . Bernardy, N. C. (2003). Randomized trial of trauma-focused group therapy for posttraumatic stress disorder: results from a department of veterans affairs cooperative study. *Archives of General Psychiatry, 60*(5), 481-489. doi: 10.1001/archpsyc.60.5.481
- Schnyder, U., Müller, J., Maercker, A., & Wittmann, L. (2011). Brief eclectic psychotherapy for PTSD: a randomized controlled trial. *The Journal of Clinical Psychiatry, 72*(4), 564-566.
- Shalev, A. Y., Ankri, Y., Israeli-Shalev, Y., Peleg, T., Adessky, R., & Freedman, S. (2012). Prevention of Posttraumatic Stress Disorder by early treatment: Results from the

- Jerusalem Trauma Outreach and Prevention Study. *Archives of General Psychiatry*, 69(2), 166-176. doi: 10.1001/archgenpsychiatry.2011.127
- Sloan, D. M., Marx, B. P., Epstein, E. M., & Lexington, J. M. (2007). Does altering the writing instructions influence outcome associated with written disclosure? *Behavior Therapy*, 38(2), 155-168. doi: 10.1016/j.beth.2006.06.005
- Sloan, D. M., Marx, B. P., & Greenberg, E. M. (2011). A test of written emotional disclosure as an intervention for posttraumatic stress disorder. *Behaviour Research and Therapy*, 49(4), 299-304. doi: 10.1016/j.brat.2011.02.001
- Spence, J., Titov, N., Dear, B. F., Johnston, L., Solley, K., Lorian, C., . . . Schwenke, G. (2011). Randomized controlled trial of internet-delivered cognitive behavioral therapy for posttraumatic stress disorder. *Depression and Anxiety*, 28(7), 541-550. doi: 10.1002/da.20835
- Stein, M. B., Walker, J. R., Hazen, A. L., & Forde, D. R. (1997). Full and partial posttraumatic stress disorder: Findings from a community survey. *American Journal of Psychiatry*, 154, 1114-1119.
- Tolin, D. F., & Foa, E. B. (2006). Sex differences in trauma and posttraumatic stress disorder: A quantitative review of 25 years of research. *Psychological Bulletin*, 132(6), 959-992.
- van Emmerik, A. A., Kamphuis, J. H., & Emmelkamp, P. M. (2008). Treating acute stress disorder and posttraumatic stress disorder with cognitive behavioral therapy or structured writing therapy: a randomized controlled trial. *Psychotherapy and Psychosomatics*, 77(2), 93-100. doi: 10.1159/000112886
- Vera, M., Reyes-Rabanillo, M., Juarbe, D., Pérez-Pedrogo, C., Olmo, A., Kichic, R., & Chaplin, W. (2011). Prolonged exposure for the treatment of Spanish-speaking Puerto

Figure 2. Forest plot of clinician-rated PTSD symptoms post-intervention for trauma-focused interventions versus any comparison condition

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Wade, D;Varker, T;Kartal, D;Hetrick, S;O'Donnell, M;Forbes, D

Title:

Gender Difference in Outcomes Following Trauma-Focused Interventions for Posttraumatic Stress Disorder: Systematic Review and Meta-Analysis

Date:

2016-05-01

Citation:

Wade, D., Varker, T., Kartal, D., Hetrick, S., O'Donnell, M. & Forbes, D. (2016). Gender Difference in Outcomes Following Trauma-Focused Interventions for Posttraumatic Stress Disorder: Systematic Review and Meta-Analysis. PSYCHOLOGICAL TRAUMA-THEORY RESEARCH PRACTICE AND POLICY, 8 (3), pp.356-364. <https://doi.org/10.1037/tra0000110>.

Persistent Link:

<http://hdl.handle.net/11343/115318>