ECOLOGICAL SOCIETY OF AMERICA

Ecology/Ecological Monographs/Ecological Applications

PREPRINT

This preprint is a PDF of a manuscript that has been accepted for publication in an ESA journal. It is the final version that was uploaded and approved by the author(s). While the paper has been through the usual rigorous peer review process of ESA journals, it has not been copy-edited, nor have the graphics and tables been modified for final publication. Also note that the paper may refer to online Appendices and/or Supplements that are not yet available. We have posted this preliminary version of the manuscript online in the interest of making the scientific findings available for distribution and citation as quickly as possible following acceptance. However, readers should be aware that the final, published version will look different from this version and may also have some differences in content.

The doi for this manuscript and the correct format for citing the paper are given at the top of the online (html) abstract.

Once the final published version of this paper is posted online, it will replace the preliminary version at the specified doi.

1 Running title: Optimal fire management for biodiversity

2

3 **Choice of biodiversity index drives optimal fire management decisions**

4 Giljohann, K.M.¹, McCarthy, M.A.¹, Kelly, L.T.¹ and Regan, T.J.¹

5

6 1. Centre of Excellence for Environmental Decisions, The School of Botany, University of
7 Melbourne, Parkville, Victoria, Australia, 3010

8 Corresponding author address: K. Giljohann, Centre of Excellence for Environmental

9 Decisions, School of Botany, University of Melbourne, Parkville, 3070, Vic, Australia.

10 **Abstract**

11 Preservation of biodiversity is a central goal of conservation management, yet the conditions that
 12 promote persistence may differ for the species in the community. For systems subject to
 13 stochastic disturbances such as fire, understanding which management practices promote
 14 persistence for all species in a community is complex. Before deciding on the best course of
 15 action an objective must be specified. Yet an overarching goal of species persistence can be
 16 specified into a measureable objective many different ways. We investigated four alternative
 17 management objectives for maximizing species persistence that use common biodiversity
 18 indices: i) attaining the minimally acceptable mix of successional vegetation states to support
 19 species' relative abundances, ii) maximizing the arithmetic mean abundance of species, iii)
 20 maximizing the geometric mean abundance of species and iv) minimizing the average extinction
 21 risk of species. We used stochastic dynamic programming to model successional changes in
 22 vegetation in the presence of both planned and unplanned fires, and utilize an extensive data set
 23 on the occurrence of birds, reptiles and small mammals in different successional states in semi-
 24 arid Australia. We investigated the influence the choice of objective function and taxonomic
 25 focus has on the optimal fire management recommendations. We also evaluated a recent hazard
 26 reduction policy to annually burn a fixed amount of the landscape and compare results to the
 27 optimal solution. The optimal management strategy to maximize species persistence over a 100-
 28 year period is predominantly to minimize wildfires. This is because the majority of species are
 29 more likely to occur in intermediate and late-successional vegetation. However the optimal
 30 solution showed sensitivity to the objective and the species included in the analysis. These
 31 results highlight the need for careful consideration when specifying an objective to represent
 32 overarching conservation goals. Using the extinction risk objective, we show that a policy to

33 annually burn 5% of the landscape could increase the average probability of extinction for the
34 modelled species by 7% over the next 100 years compared to the optimal management scenario.

35

36 Key words: biodiversity index, fauna, management, objectives, policy, prescribed burning,
37 stochastic dynamic programming, successional vegetation, wildfire

essa
preprint

38 **Introduction**

39 Conservation management aims to preserve biological diversity, yet the conditions that promote
 40 persistence may differ among the species in the community (Margules and Pressey 2000) and
 41 management objectives may be specified in many different ways. Thus, the best management
 42 action to take now to ensure the preservation of species into the future can be complex, possibly
 43 unintuitive and difficult to identify. This is especially pertinent for systems subject to large
 44 stochastic disturbances such as unplanned fires that can profoundly change the state of the
 45 landscape.

46 Inappropriate fire regimes are a key threatening process for species persistence (Driscoll
 47 et al. 2010). A dominant paradigm for maintaining biological diversity in fire-prone
 48 environments is ‘pyrodiversity begets biodiversity’, where the broad objective is to create a
 49 dynamic mosaic of vegetation patches throughout the landscape. The underlying assumption is
 50 that increasing variability in fire regime characteristics will support the needs of multiple species
 51 with diverse habitat requirements (Martin and Sapsis 1992, Parr and Andersen 2006). This
 52 assumes that the structure and availability of resources is affected by the post-fire successional
 53 state of the vegetation. Although a uniform fire regime will clearly not suit all species, the
 54 benefit of arbitrarily adding variability into a landscape is questionable without a clear
 55 understanding of the effects on the biota.

56 The degree to which fire is used as a management tool to support community diversity
 57 ought to be intrinsically connected to species’ requirements. If we assume that species will most
 58 commonly be found in areas that provide suitable habitat and resources, the probability that a
 59 species persists should increase with the area of suitable habitat. This is a reasonable assumption

when species have a relatively low probability of occurrence. Then relating species abundance in successional vegetation ages to the proportion of area covered by the different vegetation ages presents a means of defining the most suitable age structure of the vegetation, in relation to time since last fire, for the species in question (McCarthy 2012, Kelly et al. *in press*). When this information is summarized over multiple species, the optimal vegetation age structure for the community can be identified (McCarthy 2012). Despite the ability to manipulate the vegetation age structure using standard fire management strategies, such as lighting fires (prescribed burning) or minimizing (fighting) wildfire extent (Keeley and Fotheringham 2001a), the question of how to attain and then retain an optimal age structure for different species is complex and unintuitive. This is especially true when accounting for the dynamics of vegetation succession and stochastic wildfire regimes.

Richards et al. (1999) used stochastic dynamic programming (SDP) to determine optimal fire management in the face of succession and unplanned fires. SDP, a formal mathematical approach to decision theory, can assist decision-makers by requiring explicitly-stated objectives and by accounting for uncertainty in system dynamics when identifying the best strategy from a set of competing strategies (Mangel and Clark 1988). The method requires: representing the system as a series of mutually exclusive alternative states and identifying the processes that govern the transition of the system between the states; a decision model of alternative management strategies that alter the rate in which the system changes state; a management objective formalized as an objective function; and a solution technique, such as backwards iteration, to identify the optimal state-dependent management strategies over a finite time horizon (Mangel and Clark 1988, Possingham et al. 2001, Mareshot et al. 2013). Richards et al. (1999) specified a management objective to attain a landscape vegetation age structure with at

least 20% of the area in each of three successional states. However, this objective does not link directly to the persistence of species in the community. Further, the objective to equally represent all successional states implies that vegetation in each state is equally important. The objective is essentially based on the pyrodiversity begets biodiversity paradigm, which has been questioned (Parr and Andersen 2006) because some successional states are more important for particular taxa (Kelly et al. 2012, Taylor et al. 2012).

A benefit of the SDP approach is that the objective needs to be stated explicitly and needs to be measureable; however, an overarching goal of species persistence can be specified as a measureable management objective in many different ways. For example, an objective may measure species persistence in terms of available habitat area, abundance, or probability of extinction. The importance of how management objectives are specified and their underlying biases has been addressed in conservation planning (Hof and Raphael 1993, Williams and Araujo 2000, Nicholson and Possingham 2006) and species conservation (McCarthy et al. 2008), yet despite the integral role of management objectives in fire planning they have not been systematically compared.

Data on multiple species in a community is commonly unavailable because it is too difficult or expensive to collect. Instead management decisions are often based on indicator species or taxa (Landres et al. 1988) which have been widely adopted as surrogates for the state of the system for other species or taxa (Landres et al. 1988). Birds are commonly used as an indicator for ecosystem health as they are well studied and have detailed data (Brooks et al. 2001, Rodrigues and Brooks 2007). Yet management planning that uses data for only one taxon may have negative repercussions for community persistence if other species and taxa require different habitat.

Multiple objectives commonly underpin fire management decisions, so contentions can arise due to competing priorities to mitigate risks to human life and assets versus meeting ecological objectives (Hesseln 2000, Gibbons et al. 2012). In the aftermath of large wildfires, more prescribed burning is often advocated to reduce the impact of future fires (Penman et al. 2011). However, prescribed burning has generated considerable debate within the public and scientific domains due to uncertainty surrounding its ability to effectively reduce the size, rate of spread and intensity of future fires (Keeley and Fotheringham 2001b, Fernandes and Botelho 2003, Moritz et al. 2004, Gibbons et al. 2012). Also, the potential risks posed to biodiversity from increased prescribed burning programs have seldom been quantified (but see Bradstock et al. 1998).

Kelly et al. (*in press*) presented a framework for fire management decision-making that links species distribution models, biodiversity indices and decision models. In this study we build on this framework by asking if the goal of maximizing the persistence of species, using the optimal vegetation age class structure for the community, is influenced by the way the management objective is specified. We adapt the model of Richards et al. (1999) and investigate four alternate specifications of a management objective to maximize community abundance i) attaining a minimally acceptable mix of successional vegetation states to support species' relative abundances (*sensu* Richards et al. 1999), ii) maximizing the arithmetic mean abundance of species, iii) maximizing the geometric mean abundance of species and iv) minimizing extinction risk using an index based on the geometric mean abundance of species (McCarthy et al. 2014). We test both the arithmetic mean and geometric mean as these are standard measures of central tendency, yet with different properties. The arithmetic mean combines values on an additive scale thereby rendering it relatively insensitive to small values, such as the low

probability of occurrence of the less common species, yet highly sensitive to large outlying values, such that one very common species can exert undue influence upon the mean. Whereas, as the geometric mean combines values on a multiplicative scale, the mean is less susceptible to extreme values and therefore more sensitive to all species requirements.

The objectives use biodiversity indices to calculate the optimal vegetation age structure (McCarthy 2012) based on data for 22 vertebrates (birds, reptiles and small mammals) within the Murray Sunset National Park, a semi-arid mallee conservation reserve in southeastern Australia. The four objectives are evaluated in terms of their influence upon the resultant optimal fire management strategies.

To better understand whether strategies based upon only one taxon reflect priorities for other taxa we also investigate the influence specific taxonomic groups have upon the optimal fire management strategy. We separately model data for birds and reptiles. We also explore the effect of annual prescribed burning targets on species persistence to assess a recent policy to implement fuel reduction burns to 5% per annum of the public estate in southeastern Australia (Teague et al. 2010). The policy was devised in response to severe wildfires in 2009 that were associated with significant loss of human life, but was not directly related to the requirements of species within the different ecosystems across the region.

Methods

Case Study: Vertebrate diversity in semi-arid Australia

Murray Sunset National Park (MSNP) is a large conservation reserve (633, 000 ha) in the semi-arid mallee region of southern Australia. The presence of many threatened and endangered small mammal and bird species makes this an important region for biodiversity (Garnett and Crowley

2000, Brown et al. 2009) and a model system for exploring fire management strategies due to its naturally occurring fire and species with diverse habitat requirements. The reserve is composed of predominantly mallee vegetation; open woodland characterised by a canopy of resprouting shrubby multi-stemmed *Eucalyptus* trees (<10 m) and an understory dominated by the perennial hummock grass *Triodia scariosa*, by multiple species of the Chenopodiaceae, or heathy shrubs. The focus of this analysis is the *Triodia* mallee which covers approximately half the reserve (Haslem et al. 2011) and provides important habitat for many native animals (birds, reptiles and small mammals) that utilize resources and structural elements such as *T. scariosa* hummocks, tree hollows, litter and decorticated bark whose development is influenced by time since last fire (Bradstock and Cohn 2002, Kelly et al. 2011, Watson et al. 2012, Nimmo et al. 2012). The direct impact of fire upon the local fauna varies. For fire-sensitive species, the lack of specific resources in the early post-fire environment can induce population crashes, or movement out of the area, with recolonization rates dependent on resources that are often only available once the system has recovered to a particular successional vegetation state (Driscoll and Henderson 2008, Watson et al. 2012).

Wildfire from lightning strikes is a natural disturbance in the mallee. Although fires have a propensity to be small (< 5% of the reserve), large fires occur infrequently and can profoundly alter the state of the reserve. In the last forty years the MSNP has experienced two fires greater than 100 000 ha (>15% of the reserve; Appendix A). As fire removes both canopy and understorey of *Triodia* mallee the vegetation structure varies markedly with time since fire (Bradstock and Cohn 2002, Haslem et al. 2011).

System & decision models

We represent the system as a Markov chain of vegetation succession, whereby transition to a future state depends only on the current state of the system. Based on the availability of key structural habitat features we allocate the vegetation of the system into three age classes which correspond to distinct successional states: early (0-10 yrs), characterised by bare ground with small coppicing *Eucalytus* spp. and post-fire ephemerals; intermediate (11-35 yrs) by high cover of *T. scariosa*, and late (36-110 yrs) by the presence of larger trees with decortivating bark and the progressive development of tree hollows. The system dynamics are the same as described by Richards et al. (1999) but with parameters values matching the MSNP. To characterize the state of the vegetation within the reserve, we modelled the reserve as a collection of 20 equal sized sites (site $\approx 17\ 000$ ha), with each site classified as either early (E), intermediate (M) or late (L) successional. The state of the reserve is then described by the number of sites in each successional state, which can take any combination of the three mutually exclusive vegetation states giving a total of $S = 231$ states. For a reserve in state j , the number of sites present in each of the three vegetation states is denoted E_j , M_j and L_j ; where $E_j + M_j + L_j$ are constrained to sum to 20. The processes of succession and fire govern changes in the successional composition of the reserve (Fig. 1). In the absence of fire, the succession transition probabilities ($s_{j,k}$) from early to intermediate, and intermediate to late, are calculated as the reciprocal of the average number of years spent in the younger successional age. A change of state is conditional on the number of sites in each successional age in the previous year (state k) and the following year (state j). Wildfire is modelled as the probability of burning $x\%$ of the reserve in each annual time-step. The spatially implicit nature of the model means that fires are assumed to occur randomly across the landscape. A fire matrix was constructed where sites are considered either entirely burnt or unburnt and all burnt sites revert to early successional vegetation regardless of their successional

state prior to the fire event. To determine the annual probability of wildfire we created a frequency distribution of *Triodia* mallee fire scars from 1972-2011 (Appendix B, Fig. B1) using data converted from satellite imagery by Avitabile et al. (2013) (Appendix A, Fig. A1). The wildfire frequency distribution was modified to approximate the situation where wildfires are left to burn unhindered (no management) and wildfires are actively minimized (fight); details are provided in Appendix B. The probability of wildfire at a site was modelled as constant through time, thus all successional states were assigned equal fire hazard. This implies that wildfire is just as likely to occur in early successional vegetation as it is in intermediate or late successional vegetation. As the model is spatially implicit, the occurrence of fire in one site does not influence the probability of fire occurring in surrounding sites. The elements ($f_{j,k}$) of the fire matrix are the probabilities the reserve moves from state k to state j during the year, conditional upon the number of sites in intermediate and late successional states in the previous year and following year multiplied by the annual probability of wildfire under either of the two scenarios. The succession and fire matrices are then multiplied to form a transition matrix which represents the probability the reserve will be in state j next year given it was in state k the previous year, $T_{j,k} = F_{j,k} \cdot S_{j,k}$.

We defined a set of six management strategies, $a \in A$, that influence how the state of the system changes: fight all wildfires to minimize the amount of area burnt and maintain the existing vegetation structure, let wildfires burn unhindered (no management) to utilize naturally occurring disturbance, or prescribed burn either 10% or 20% of intermediate or late successional vegetation to create areas of early successional vegetation and ensure the long-term supply of intermediate successional vegetation. At each time-step the six strategies are evaluated over the set of states; these are then ranked on their ability to satisfy the objective function, with one

strategy being optimal. Prescribed burns were modelled such that, the state k of the reserve after burning either n intermediate or n late successional sites is denoted as $B_m(k, n)$ or $B_l(k, n)$. For example, when burning intermediate successional sites the transition matrix is modified by replacing the $T_{j,k}$ with $T_{j,B_m(k,n)}$. This management strategy was only an option when the state of the reserve has n sites in the target successional state. Prescribed burns were implemented at the start of the model year, and wildfires can occur after the prescribed burns have been implemented. Prescribed burning of early successional *Triodia* mallee was not a proposed strategy as the vegetation cannot effectively carry fire at this age under conditions suitable for prescribed burning (Noble and Vines 1993). We assume that management actions are 100% effective, e.g. prescribed burns are not patchy; therefore the post-fire successional trajectory after a prescribed burn was modelled as described above for wildfire. It is likely that burn severity would be lesser under conditions suitable for prescribed fires, however as there is a lack of data to substantiate this we took a conservative approach and assumed the vegetation response to be similar (Haslem et al. 2011).

Species data

To characterise species response to the age class structure of the vegetation, we use empirically derived estimates of the probability of occurrence, for each of 22 species of birds, reptiles and small mammals, in each of three age-classes (E , M , L). The data were collected as part of a large project investigating the influence of fire history on the occurrence of small mammals, reptiles, birds in *Triodia* mallee (The Mallee Fire and Biodiversity Project: Kelly et al 2011, Nimmo et al 2012, Watson et al 2012). Of the fifty-plus species that were recorded, 22 showed strong differential usage of vegetation age classes. The responses (presence/absence) of these 22 species

were modelled by Kelly et al (*in press*) and estimates of probability of occurrence were calculated for each of the three vegetation age classes. Of the 10 bird species modelled six reached their highest probability of occurrence in late successional vegetation, three in intermediate successional vegetation and one in early successional vegetation. For the reptiles associated with successional vegetation, four of the 11 species reached their highest probability of occurrence in early successional vegetation, four in intermediate successional vegetation, and three in late successional vegetation. Only one small mammal showed a strong fire response, attaining its highest probability of occurrence in intermediate and late successional vegetation (Table 1; see Kelly et al. *in press* for details). We interpret this probability as a measure of the species' relative abundance. The species whose occurrence was invariant to successional habitat change were not included in the analysis as their requirements should be met by provision of habitat in any successional state.

The average probability of occurrence (P) of species i in any state depends upon the proportion of area in each vegetation age class. For each species i and state j this probability was calculated as

$$P_{ij} = (y_{Ei}p_{Ej} + y_{Mi}p_{Mj} + y_{Li}p_{Lj}) \quad (1)$$

where y is the probability of species i occurrence and p is the proportion of area in either E , M or L successional vegetation.

Objective functions

We specify four alternate management objectives explicitly linked to the resources required for species preservation. The objective functions identify the vertebrate community's response to the vegetation age structure of the reserve via a biodiversity index that synthesizes the individual species responses. The objectives were calculated for all possible states of the reserve, for the 22 species combined and for two taxonomic subsets of the data: birds (10 species) and reptiles (11 species).

1. *Binary objective*

A relatively simple approach is to use a binary function. This classifies each state of the reserve as either nominally 'desirable' (value 1) or 'undesirable' (value 0) for species conservation. 'Desirable' and 'undesirable' are defined by whether the state contains a pre-determined minimum proportion of the reserve in each successional age class (*sensu* Richards et al. 1999). For the objective to have relevance to the species in the community, the target minimum area in each successional state (%) was determined by numerically optimising the geometric mean probability of species' occurrence across the three vegetation age classes (Kelly et al. *in press*). This method identifies the state of the reserve that jointly maximizes each species' probability of occurrence. By iteratively cycling through the range of states (proportion of the reserve in each successional age class, *E*, *M* & *L*) the state that simultaneously maximizes all species' probability of occurrence is identified. The optimised state values were 45% intermediate and 55% late for all species, 100% late for birds, and 75% intermediate and 25% late for reptiles. However, these values are approximate given error in the data and cannot easily be achieved precisely in practice given the occurrence of unplanned fires. Therefore, we specified ranges around the optimal values that were close to the optimal vegetation age structure and which we regarded as

acceptable age structures for all species ($\geq 30\%$ intermediate and $\geq 40\%$ late), for birds ($\geq 70\%$ late), and for reptiles ($\geq 50\%$ intermediate and $\geq 20\%$ late). The proportion of early vegetation was never a limiting condition, so did not influence the specification of the desirable state.

The objective function (Ψ) for each state j of the reserve was calculated such that

$$\Psi(j) = \begin{cases} 1, & \text{if } p_{Mj} \geq \text{opt}_M \text{ and } p_{Lj} \geq \text{opt}_L \\ 0, & \text{otherwise} \end{cases} \quad (2)$$

Where opt is the minimum area target (%) in either intermediate (M) or late (L) successional vegetation that is regarded as acceptable.

2. Arithmetic mean probability of species' occurrence

The arithmetic mean of the probability of species' occurrence is the sum of each species average probability of occurrence across the reserve divided by the total number of species (N). This is a standard measure for calculating measures of central tendency and represents the expected species richness.

For each state j of the reserve the second objective function is calculated as

$$\Psi(j) = \frac{1}{N} \sum_{i=1}^N P_{ij} \quad (3)$$

3. Geometric mean probability of species' occurrence

The geometric mean of the probability of species occurrence is calculated by taking the n th root of the product of each species probability of occurrence. This metric is sensitive to changes in the number of species, species' abundances and community evenness; three key criteria for monitoring changes in biodiversity (Buckland et al. 2011).

299 For each state j of the reserve the third objective is calculated as

$$\Psi(j) = \left(\prod_{i=1}^N P_{ij} \right)^{\frac{1}{N}} \quad (4)$$

300 4. *Index of extinction risk*

301 This index relates the geometric mean of species' abundances to the proportion of species going
 302 extinct within time t . This is based on the logarithm of the geometric mean of species abundance
 303 being proportional to the mean time to extinction for species undergoing exponential decline
 304 (McCarthy et al. 2014). The index is correlated with the extinction risk of plant and animal
 305 populations (McCarthy et al. 2014). Thus, it can be used to quantify how changes in abundance
 306 affect persistence and infer future risks to species populations when absolute abundances are
 307 known. To calculate this index we used the modelled occupancy data as a proxy for abundance
 308 data.

309 The objective function for each state j of the reserve is calculated as

$$\Psi(j) = \left(\frac{1}{\ln \left(\left(\prod_{i=1}^N P_{ij} \right)^{\frac{1}{N}} \right)} \right) \quad (5)$$

310 *Solution method*

311 Using Bellman's (Bellman 1966) dynamic programming equation, the expected value of the
 312 reserve (V) for objectives 1 to 3 is calculated for each state of the system by maximizing the
 313 expected state value at each time step such that:

$$V(j, t, T_f) = \max_{a \in A} \left[\Psi(j) + \sum_{k=1}^S T_{k,j}(a) V(k, t + 1, T_f) \right] \quad (6)$$

Where j and k reference the state of the reserve from 1 to S , t is the current time, T_f the terminal future year, and a of A the set of strategies to be evaluated. As objective 4 aims to minimize the extinction risk of species, the expected value of the reserve is calculated by minimizing (rather than maximizing) Eqn 6. To find the SDP solution using backward iteration requires initializing the expected value vector for all possible states of the reserve at the final time-step; this was set as $V(j, T_f, T_f) = \Psi(j)$ for $j = 1$ to S .

The optimal fire management strategy was modelled over a 100-year period for each objective, for all species combined and for two taxonomic subsets: birds and reptiles. By comparing results at the final time-step we sought to identify how sensitive the optimal management strategy and the expected value of each reserve state are to the choice of objective function. We also investigate the optimal state of the system by modifying the fire frequency distribution to simulate an average inter-fire interval of 200 yrs; a scenario of very infrequent fire. Models were written in R version 3.1.0 (R Development Core Team 2013). To compare between objectives the terminal values of V were rescaled to lie between 0-1 (with 1 being best).

Performance of the alternate objective functions

Different objective functions can result in a different sequence of optimal management strategies through time. Therefore we used forward value-iteration to compare the values for each objective function when it followed the optimal management strategy for one of the other objective functions. In the event that each of the objectives appear equally relevant, identifying the optimal strategy that induces the least reduction in state value for all objectives can assist in determining

which strategy to implement. The starting state for simulations at t_1 was the current vegetation state of the MSNP *Triodia* vegetation (5% early, 55% intermediate and 40% late successional vegetation). At each yearly time-step ($t=1$ to $T_f=100$) the optimal strategy a under objective q for the current state k at time t was followed. The strategy taken at time t influences the probability the reserve will transition to any randomly selected state j at $t+1$. At each time-step the value of state j under each objective Ψ_q was recorded. State values were scaled to lie between 0-1, with 1 being optimal. Each objective-strategy combination was simulated 1000 times and the average of these taken. The values are expressed as a percentage of the value it would be if the optimal strategy identified for that objective was followed.

Incorporating a fuel reduction target

Annual fuel reduction targets of up to 5% of the reserve, in increments of 0.05%, were implemented by limiting the set of management strategies when the state of the reserve did not meet the target. For example, when there is a target to burn 5% of the reserve per year, this equates to a goal of having 50% of the vegetation in the early successional state (≤ 10 years). The 50% is calculated on the basis that if ‘early’ successional vegetation takes at least 10 years to transition to ‘intermediate’ successional vegetation then a prescribed burning target of 5% per year requires that over the long term 50% of the vegetation will be present as ‘early’ successional. Therefore, in states where less than 50% of the vegetation is early successional, management actions were limited to one of the prescribed burning options to shift the reserve towards a state that attains the target. In contrast, when the current reserve state meets the management target the selected strategy is the one that maximizes the expected biodiversity value for the reserve. Due to the spatially implicit nature of the model the location of fires was

again assumed to be random. We assess the implications for species persistence of annual targets to prescribed burn between 0-5% of the reserve.

Results

The optimal strategy for three of the four objectives, over a 100-year period and when prioritising all species, was fight wildfires (Fig. 2). The threshold objective (Fig. 2a) was different from the other three, with burning late successional vegetation being optimal when ~60% of the reserve was composed of late successional vegetation. Burning late successional vegetation was optimal in this case to allow development of the intermediate vegetation. However, the objectives that were continuously based did not strictly limit the required area of intermediate vegetation. Indeed, burning was never optimal because older vegetation was likely to burn eventually from wildfire, and the benefit of promoting younger vegetation was outweighed both by the uncertain prospect of it developing and by the loss of older vegetation that was particularly important for many species. When prioritising only reptiles the optimal solutions diverged considerably, revealing sensitivity to the choice of objective function.

To investigate the optimal state of the system - regardless of ignition source, we modified the fire frequency distribution to simulate an average inter-fire interval of 200 yrs (10 times the average inter-fire interval from the MSNP data). Under such a low fire frequency the likelihood of wildfire burning the vegetation and promoting young and intermediate vegetation is highly uncertain, so lighting fires is the optimal management strategy when the reserve is dominated by late vegetation (Fig 3). However in the majority of states fighting fires is optimal under such a scenario due to the need to protect young vegetation because of its relative scarcity and enable the development of intermediate successional vegetation. Indicating that the optimal fire regime

for the species modelled is more frequent than a 200 yr interval but less frequent than currently occurs.

Sensitivity of expected value of the reserve to objective function

The expected value of the reserve (V) was sensitive to the objective. When prioritising all species, the solution of the threshold (BN) objective was most dissimilar to the continuous objectives (the arithmetic mean (AM), geometric mean (GM) and extinction risk index (ER)). This is due to BN classifying states as either strictly ‘desirable’ or ‘undesirable’; therefore, states that fall within the limits of the area target (within the inner triangle) are allocated a value and states that do not meet the target are not (Fig. 2a). The optimal strategy is the one most likely to shift the reserve into a valued state. Thus under the BN objective it is optimal to prescribed burn when late successional vegetation predominates; the highest value states under the continuous objectives. For the continuous objectives, as the state of the system moves away from the most optimal state (darkest shading) the expected values decline steeply for the AM and more gradually for GM and ER objective functions. The gradual decline under the GM and ER objectives is a result of averaging on the log scale, thus reducing the influence of the more prevalent species.

Sensitivity of the optimal fire management strategy to taxonomic focus

For the two taxonomic subsets, results are only presented for the GM objective; results for the other objectives are in Appendices C & D. As birds attain maximum probability of occurrence in late successional vegetation, the optimal strategy over all states is to fight wildfires. Despite differences between the expected state values under the four objectives, the optimal strategies are

identical. Due to the recurrent risk of wildfire a fight fire strategy is the only way to maximize the likelihood of obtaining a reserve composed of entirely late successional vegetation (Fig. 3b & Appendix C, Fig. C1). However for reptiles, the optimal strategy was sensitive to choice of objective function. As reptile probability of occurrence is greatest in intermediate successional vegetation, the optimal strategy is to burn late successional vegetation when it dominates the reserve and to fight wildfires when early or intermediate successional vegetation predominates (Fig. 3c & Appendix D, Fig. D1). Yet the amount of prescribed burning considered optimal is contingent on the objective function. As the AM objective had very few states of high value, it was optimal to burn late successional vegetation when it comprised greater than 10% of the reserve. In contrast, it was only optimal to burn late successional vegetation when it comprised >30% of the reserve for BN, or >40% for GM and ER.

Performance of the objective functions

The value of each objective function was compared using the optimal strategies for all species and those for reptiles - as these strategies were the most divergent. When prioritising for all species, the AM, GM and ER objectives performed equally well under all strategies (i.e. percent reduction in value relative to that achieved if using the optimal strategy) (Appendix E, Fig. E1a, c & d). The BN objective despite performing equally well on average, varied up to 20% better and worse under strategies other than its' own (Appendix E, Fig. E1b). When prioritising for reptiles, the BN strategy gave the best results for all objectives, the value of the GM and ER objectives was reduced on average only 2% and the AM 17% (Fig. 4b). The GM and ER strategies were very similar, with both objectives performing optimally under the other. Under these strategies the AM objective performed on average 26% worse and the BN objective 35%

worse (Fig. 4c & d). The strategy of the AM objective resulted in the greatest divergence between the objectives, with the GM and ER performing on average 20% worse, yet the BN 97% worse.

Annual fuel reduction targets

Implementing a fuel reduction target reduced the expected value of the reserve by minimizing time spent in the intermediate and late successional states. The optimal strategy reflects the policy constraint by recommending burning of intermediate or late successional vegetation for any state of the reserve with less than the target amount of early successional vegetation. For the majority of states where prescribed burning was recommended it was preferable to burn intermediate successional vegetation as opposed to late successional vegetation (Appendix F, Fig. F1). This is due to species preference for late successional vegetation and the length of time taken for early successional vegetation to transition to a late successional state. The influence of the fuel reduction burning policy upon the optimal strategy was similar for all objectives and so results are only presented for the ER objective.

Assuming the ER objective is proportional to the extinction risk, a 3% target could increase the probability of extinction for the 22 species by 1.9% over the next 100 yrs. This is a relative increase in extinction risk when compared to the optimal management scenario (Fig. 5). Under a 3% scenario the relative change in the ER objective is almost equivalent to that resulting from a scenario where management takes a hands-off approach (no management). Burning <2% has a better outcome than undertaking no management. If a 5% target were implemented the probability of extinction could increase by 7% over the next 100 years.

Discussion

In fire prone environments, determining the management strategy most likely to achieve an overarching objective is complicated. Objectives may be specified in a number of ways, each capturing a different aspect of community structure. This work has demonstrated that the choice of biodiversity index used to define the management objective, in this instance the optimal age class structure of the vegetation, influences the best way to manage fires.

Our results confirm the expectation that management decisions based on one group of species will not necessarily promote conditions suitable to preserve all species. This is consistent with other research that has found single-taxon prioritization to be inefficient and inadequate for protecting non-target species (Prendergast et al. 1993, Lindenmayer and Margules 2000). Thus, multi-taxon prioritization can offer a better outcome in community conservation planning. Although surrogate or partial measures of biodiversity must be used in the absence of complete knowledge (Margules and Pressey 2000) these must be carefully chosen. Our analyses included extensive data on birds, reptiles and small mammals, nonetheless including other taxa, such as plants and invertebrates, would further improve the applicability of the results.

When all 22 species are modelled together, the predominant optimal management strategy is to fight wildfires, irrespective of the objective function. This result will not necessarily be applicable to other systems, although the methods definitely are. A strategy to fight wildfires may seem a surprising result for species that have evolved with fire. However, this strategy does not recommend the complete exclusion of fire from the environment, but aims to minimize the extent burnt by wildfire. This strategy is optimal for three reasons. The first is the overall preference of the modelled species for intermediate and late successional vegetation. If other species and taxa were included in the analysis (e.g. threatened species, invertebrates and

plants) then this may alter the result. However the strategy is congruent with what is known of the habitat requirements for threatened species such as the Mallee Emu-wren, Black-eared Miner and Malleefowl; species which are understood to require large areas of long unburnt vegetation for habitat (Brown et al. 2009). As the structure of the vegetation continues to develop throughout the length of our late successional class (35-100 yrs) (Haslem et al. 2011), fighting wildfires should enable significant stands of late successional vegetation to develop, thereby supporting species that respond to these continuing structural changes (Clarke 2008, Kelly et al. 2011, Watson et al. 2012). It is also plausible that a strategy to minimize wildfire will be suitable for the plant community, as many species in the mallee exhibit great longevity (e.g. *Callitris*), coupled with an ability to resprout following fire (e.g. *Eucalypts*, *Triodia*) or are ephemeral with a persistent soil seedbank (Kenny 2012). However quantifying these traits and including them in the analysis would reduce uncertainty in the results.

The second reason a fight wildfire strategy is optimal is due to the likelihood of recurrent wildfire altering the distribution of successional states. If wildfire was expected to occur much less frequently (e.g. average inter-fire interval of 200 yrs) then prescribed burning would be required as the optimal fire return interval for the species would be exceeded. This change of management strategy implies that in the current environment there is a risk of too much fire. The fire history data we used only dates back to the 1970s and so it is possible that by chance, this was a period of increased fire activity. Fire history records over a longer period may tell a different story. As decades elapse before intermediate and late successional vegetation recur the expected fire frequency is therefore a critical parameter for management to know when identifying management actions to optimize the state of the system for the benefit of the local species.

The third factor influencing the optimal strategy is related to the efficacy of fighting wildfires. In the study we assumed that fighting fires was most effective at minimizing small to medium sized fires, but had minimal effect on the moderating the size of large fires. If the assumed efficacy was sufficiently large to reduce the risk of late vegetation burning, or spatial interactions between areas of reduced fuels and wildfire movement had been explicitly accounted for, then prescribed burning in some system states might be desirable to help conserve the species. An assumption we made was that fire-fighting efficacy was unrelated to the successional state of the vegetation (e.g. the three successional vegetation classes have equal probability of burning) as it was shown by Richards et al. (1999) that the optimal management strategy was similar even when the successional states had varying probabilities of fire. Although not addressed in this study, prescribed burning is widely used as a wildfire mitigation tool (Keeley and Fotheringham 2001a). Such as the linear strip burns (approximately 14, 000 ha) implemented in the MSNP (Sandell et al 2006); the explicit aim being to minimize the risk of landscape-scale fires. A spatially explicit analysis would be useful to see how it impacts upon the results.

Time-since-fire thresholds have been advocated for defining the habitat requirements of species and thus their conservation (Keith et al. 2001, Sandell et al 2006). The thresholds are used to calculate the proportion of area within the tolerable limits (desirable states) and then management is implemented to attain a reserve in this state. In this study, setting thresholds based on the minimum desirable proportions of the three successional states (the BN objective) did not accurately represent the species response to habitat. Despite the thresholds of the BN objective being based on acceptable (tolerable) ranges around the optimized geometric mean probability of species occurrences the objective is not sensitive to incremental changes. The three

continuous objective functions examined (AM, GM and ER) provide a more realistic measure of the expected habitat value of the reserve as they contain considerably more information on the relative value of each state for the community. This difference is equivalent to comparisons of continuous and discrete indices of habitat suitability (Williams and Araujo 2000).

As the biodiversity indices upon which the objectives were based each uniquely summarise the component information, the expected state values and optimal fire management strategies differ. The change in management strategy can be seen most clearly for the reptile-only prioritisation. As these species are most prevalent in intermediate successional vegetation the optimal fire management strategy is not obvious. For example, the number of states recommended for prescribed burning increased substantially under the AM versus the GM or ER objectives. The AM is highly influenced by common species (Buckland et al. 2005) and therefore shows a steeper decline in value away from the optimum, whereas the multiplicative scale of the GM and ER objectives weight proportional changes in species equally. The similar output from the latter two objectives is due to their common basis in the geometric mean; in this case study the GM and ER could be considered interchangeable; although they are slightly different when averaged over time.

In evaluating objectives for conservation decision-making selection will depend on what aspect of biodiversity is of primary interest and the goals of management (Nicholson and Possingham 2006, Magurran and McGill 2011). If the management goal is to conserve all species, then an objective that is sensitive to the less common species, such as the GM or ER, may be most relevant. A benefit of the ER objective is that through correlation with the extinction risk, the index provides information on the likely future state of biodiversity

(McCarthy et al. 2014). Extinction risk is also an easily communicated and tangible concept with relevance to management (Williams and Araujo 2000).

In the event that multiple objectives appear equally relevant, the strategy of the BN presents the most suitable option for the semi-arid mallee vertebrates. This is because under the BN strategy each objective exhibited the least reduction in value from its possible optimum (the state values obtained when following the optimal strategy for that objective) (Fig. 4b & Appendix E). This method of selecting a strategy, from a set of competing strategies, presents a trade-off between obtaining the best value, for the species modelled, under any single objective and identifying the strategy under which all objectives perform best. The GM and ER objectives were the least sensitive to the strategies of the other objectives. However as it is unlikely any single objective will be suitable for all situations we believe the choice of objective should be guided by the management question. The most robust strategy will nonetheless be one that promotes large areas of late successional vegetation, such as the GM or ER. This is not only because the species in the analysis prefer intermediate-late successional vegetation, but also because of the likelihood of wildfires changing the state of the reserve and because of the length of time, without fire, required for late vegetation to develop. Although early successional vegetation is necessary to provide a supply of intermediate successional vegetation, this is easily gained through wildfire or prescribed burning.

Incorporating a fuel reduction policy into a decision-theoretic model can help predict potential impacts to the species modelled, and provides a means to discriminate between alternate policies in an explicit and transparent manner (Failing and Gregory 2003). The policy we simulated, to annually burn 5% of the public estate, was implemented in southeastern Australia as a response to devastating wildfires in 2009 that were associated with a significant

loss of human life. However, the policy was not evaluated for its impact on the ecosystems it would affect. By simulating a range of targets for fuel reduction, we identified resulting differences in risk associated with the alternate targets. Hence, our technique can help identify fuel reduction targets that present a better outcome for biodiversity conservation while also meeting fire management objectives. The finding that burning less than 2% of the reserve has a better outcome than undertaking no management indicates that a ‘hands-off’ approach to conservation is not the best strategy, and that trade-offs can be found when managing for risk mitigation and biodiversity persistence. As such, a blanket state-wide fuel reduction target may not be an appropriate policy initiative for the persistence of species in the mallee. As this work has shown, if a 5% annual fuel reduction target were implemented in the *Triodia* mallee, the landscape wide reduction in preferred habitat for the species modelled could increase their risk of extinction by an average of 7% over 100 years. Our formulation of the fuel reduction policy is more akin to a risk reduction strategy, as prescribed burns are only enforced when intermediate and late successional vegetation exceeds a predetermined level. This is a more conservative approach than annually burning regardless of the state of the reserve and results should be interpreted as such. When interpreting changes in extinction risk under the alternate scenarios, it is important to remember that the projected differences in risk assume species abundance depends predictably on the successional vegetation states over the modelled time frame. Thus, the projections do not account for other pressures (e.g. due to climate change) that may induce changes in abundances and which could alter the result.

The best way to maintain a mosaic of successional vegetation by fire management that meets the needs of the resident community is typically subject to considerable uncertainty (Parr and Andersen 2006). Our research has demonstrated a novel extension to the Richards’ et al.

(1999) decision-support model by using empirical data on the resident species to identify the optimal vegetation age structure (mosaic) appropriate for the community and thus the fire management strategies most likely to retain this desired mosaic. These results will be sensitive to the definition and number of successional age classes used to describe system dynamics. In this study we used three age classes that are consistent with the understanding of vegetation change in the system (Haslem et al. 2011), are correlated with faunal distributions (Kelly et al. 2012, Nimmo et al. 2013, Taylor et al. 2012) and are able to be mapped throughout the reserve (Avitabile et al. 2013). The age classes therefore reflect the requirements of the vertebrate species rather than varying fuel loadings.

Additionally, the model of system dynamics does not account for differential patchiness, intensity and fuel removal between and within prescribed burns and wildfires. As wildfires and prescribed burns are likely to generate different patterns in vegetation and therefore faunal response, the recommended fire management actions are likely to be sensitive to this assumption of equivalence. Similarly the results will be sensitive to the way area is aggregated into homogenous patches, such as the 20 sites the reserve was apportioned into for this study.

Surprisingly, given its integral role in fire planning, the influence of the specific objective on the optimal management strategy has received little attention. Although we focused on species' responses to time since last fire, we do not to imply that other parts of the fire regime, such as severity and inter-fire interval, are not important; they could be included if relevant data were available.

The objectives we formulated utilize data typically collected as part of a monitoring program, thereby providing a clear link with species presence and the management goal. The optimal management strategy to promote species persistence showed sensitivity to the objective

function and the species included in the analysis. This indicates that management objectives and the focal species must be specified carefully. By implementing a prescribed burning policy we demonstrate the applicability of this approach for improving conservation outcomes through a structured approach to management scenario planning.

Acknowledgements

Many thanks to the Mallee Fire and Biodiversity Project for providing animal data and fire history mapping, and to Shane Richards for access to code from the original model. This research was supported by funding from the Australian Research Council (ARC) Centre of Excellence for Environmental Decisions. LK was also supported by funding from the Department of Environment and Primary Industries, and MM by an ARC Future Fellowship. We also thank two anonymous reviewers for comments that improved the manuscript.

614 **References**

- 615 Avitabile, S. C., K. E. Callister, L. T. Kelly, A. Haslem, L. Fraser, D. G. Nimmo, S. J. Watson,
 616 S. A. Kenny, R. S. Taylor, L. M. Spence-Bailey, A. F. Bennett, and M. F. Clarke. 2013.
 617 Landscape and Urban Planning. *Landscape and Urban Planning* 117:81–91.
- 618 Bellman, R. 1966. Dynamic programming. *Science* 153:34–37.
- 619 Bradstock, R. A., and J. S. Cohn. 2002. Demographic characteristics of mallee pine (*Callitris*
 620 *verrucosa*) in fire-prone mallee communities of central New South Wales. *Australian Journal*
 621 *of Botany* 50:653.
- 622 Bradstock, R. A., M. Bedward, B. J. Kenny, and J. Scott. 1998. Spatially-explicit simulation of
 623 the effect of prescribed burning on fire regimes and plant extinctions in shrublands typical of
 624 south-eastern Australia. *Biological Conservation* 86:83–95.
- 625 Brooks, T., A. Balmford, N. Burgess, L. A. Hansen, J. Moore, C. Rahbek, P. Williams, L. A.
 626 Bennun, A. Byaruhanga, and P. Kasoma. 2001. Conservation priorities for birds and
 627 biodiversity: do East African Important Bird Areas represent species diversity in other
 628 terrestrial vertebrate groups. *Ostrich* 15:3–12.
- 629 Brown, S., M. Clarke, and R. Clarke. 2009. Fire is a key element in the landscape-scale habitat
 630 requirements and global population status of a threatened bird: The Mallee Emu-wren
 631 (*Stipiturus mallee*). *Biological Conservation* 142:432–445.
- 632 Buckland, S. T., A. C. Studeny, A. E. Magurran, J. B. Illian, and S. E. Newson. 2011. The
 633 geometric mean of relative abundance indices: a biodiversity measure with a difference.
 634 *Ecosphere* 2:art100.

- 635 Buckland, S. T., A. E. Magurran, R. E. Green, and R. M. Fewster. 2005. Monitoring change in
636 biodiversity through composite indices. *Philosophical Transactions of the Royal Society B:*
637 *Biological Sciences* 360:243–254.
- 638 Cheal, D. 2010. Growth stages and tolerable fire intervals for Victoria’s native vegetation data
639 sets. Pages 1–42. Department of Sustainability and Environment.
- 640 Clarke, M. F. 2008. Catering for the needs of fauna in fire management: science or just wishful
641 thinking? *Wildlife Research* 35:385.
- 642 Driscoll, D. A., and M. K. Henderson. 2008. How many common reptile species are fire
643 specialists? A replicated natural experiment highlights the predictive weakness of a fire
644 succession model. *Biological Conservation* 141:460–471.
- 645 Driscoll, D. A., D. B. Lindenmayer, A. F. Bennett, M. Bode, R. A. Bradstock, G. J. Cary, M. F.
646 Clarke, N. Dexter, R. Fensham, G. Friend, M. Gill, S. James, G. Kay, D. A. Keith, C.
647 MacGregor, J. Russell-Smith, D. Salt, J. E. M. Watson, R. J. Williams, and A. York. 2010.
648 Fire management for biodiversity conservation: Key research questions and our capacity to
649 answer them. *Biological Conservation* 143:1928–1939.
- 650 Failing, L., and R. Gregory. 2003. Ten common mistakes in designing biodiversity indicators for
651 forest policy. *Journal of Environmental Management* 68:121–132.
- 652 Fernandes, P. M., and H. S. Botelho. 2003. A review of prescribed burning effectiveness in fire
653 hazard reduction. *International Journal of Wildland Fire* 12:117.
- 654 Garnett, S. T., and G. M. Crowley. 2000. The action plan for Australian birds 2000. Environment
655 Australia.
- 656 Gibbons, P., L. van Bommel, A. M. Gill, G. J. Cary, D. A. Driscoll, R. A. Bradstock, E. Knight,
657 M. A. Moritz, S. L. Stephens, and D. B. Lindenmayer. 2012. Land management practices

- 658 associated with house loss in wildfires. PLoS ONE 7:e29212.
- 659 Hargreaves, B. 2007. Forests Act 1958. Pages 1–142. State Government of Victoria, Australia.
- 660 Haslem, A., L. T. Kelly, D. G. Nimmo, S. J. Watson, S. A. Kenny, R. S. Taylor, S. C. Avitabile,
 661 K. E. Callister, L. M. Spence-Bailey, M. F. Clarke, and A. F. Bennett. 2011. Habitat or fuel?
 662 Implications of long-term, post-fire dynamics for the development of key resources for fauna
 663 and fire. *Journal of Applied Ecology* 48:247–256.
- 664 Hessel, H. 2000. The economics of prescribed burning: a research review. *Forest Science*
 665 46:322–334.
- 666 Hof, J. G., and M. G. Raphael. 1993. Some mathematical programming approaches for
 667 optimizing timber age-class distributions to meet multispecies wildlife population objectives.
 668 *Canadian Journal of Forest Research* 23:828–834.
- 669 Keeley, J. E., and C. J. Fotheringham. 2001a. History and management of crown-fire
 670 ecosystems: a summary and response. *Conservation Biology* 15:1561–1567.
- 671 Keeley, J. E., and C. J. Fotheringham. 2001b. Historic fire regime in southern California
 672 shrublands. *Conservation Biology*:1–14.
- 673 Keith, D. A., J. E. Williams, and J. C. Z. Woinarski. 2001. Fire management and biodiversity
 674 conservation: key approaches and principles. Pages 401–425 in R. A. Bradstock, J. E.
 675 Williams, and A. M. Gill, editors. *Flammable Australia: the fire regimes and biodiversity of*
 676 *a continent*. Cambridge University Press, USA.
- 677 Kelly, L. T., A. F. Bennett, M. F. Clarke, and M. A. McCarthy. (*in press*). Optimal fire histories
 678 for biodiversity conservation. *Conservation Biology*.
- 679 Kelly, L. T., D. G. Nimmo, L. M. Spence-Bailey, A. Haslem, S. J. Watson, M. F. Clarke, and A.
 680 F. Bennett. 2011. Influence of fire history on small mammal distributions: insights from a

- 681 100-year post-fire chronosequence. *Diversity and Distributions* 17:462–473.
- 682 Kelly, L. T., D. G. Nimmo, L. M. Spence-Bailey, R. S. Taylor, S. J. Watson, M. F. Clarke, and
683 A. F. Bennett. 2012. Managing fire mosaics for small mammal conservation: a landscape
684 perspective. *Journal of Applied Ecology* 49:412–421.
- 685 Kenny, S. A. 2012. The relative influence of time-since-fire and environment on vegetation
686 composition and structure in mallee shrublands, south-eastern Australia. PhD thesis, La
687 Trobe University. Melbourne, Australia.
- 688 Landres, P. B., J. Verner, and J. W. Thomas. 1988. Ecological uses of vertebrate indicator
689 species: a critique. *Conservation Biology* 2:316–328.
- 690 Lindenmayer, D. B., and C. R. Margules. 2000. Indicators of biodiversity for ecologically
691 sustainable forest management. *Conservation Biology* 14:941–950.
- 692 Magurran, A. E., and B. J. McGill. 2011. *Biological diversity: frontiers in measurement and*
693 *assessment*. Oxford University Press, UK.
- 694 Mangel, M., and C. W. Clark. 1988. *Dynamic modeling in behavioral ecology*. Princeton
695 University Press, Princeton.
- 696 Marescot, L., G. Chapron, I. Chadès, P. L. Fackler, C. Duchamp, E. Marboutin, and O. Gimenez.
697 2013. Complex decisions made simple: a primer on stochastic dynamic programming.
698 *Methods in Ecology and Evolution* 4:872–884.
- 699 Margules, C. R., and R. L. Pressey. 2000. Systematic conservation planning. *Nature* 405:243–
700 253.
- 701 Martin, R. E., and D. B. Sapsis. 1992. Fires as agents of biodiversity: pyrodiversity promotes
702 biodiversity. *Proceedings of the symposium on biodiversity in Northwestern California*
703 Berkeley: Wildland Resources Centre, University of California.

- 704 McCarthy, M. A. 2012. Review of resilience concepts and their measurement for fire
705 management. Pages 1–38. Department of Sustainability and Environment, Victorian State
706 Government, Melbourne.
- 707 McCarthy, M. A., A. L. Moore, J. Krauss, J. W. Morgan, and C. F. Clements. 2014. Linking
708 indices for biodiversity monitoring to extinction risk. *Conservation Biology*:dx.doi.org–
709 10.1101–000760.
- 710 McCarthy, M. A., C. J. Thompson, and S. T. Garnett. 2008. Optimal investment in conservation
711 of species. *Journal of Applied Ecology* 45:1428–1435.
- 712 Moritz, M. A., J. E. Keeley, E. A. Johnson, and A. A. Schaffner. 2004. Testing a basic
713 assumption of shrubland fire management: how important is fuel age? *Frontiers in Ecology*
714 *and the Environment* 2:67–72.
- 715 Nicholson, E., and H. P. Possingham. 2006. Objectives for multiple-species conservation
716 planning. *Conservation Biology* 20:871–881.
- 717 Nimmo, D. G., L. T. Kelly, L. M. Spence-Bailey, S. J. Watson, A. Haslem, J. G. White, M. F.
718 Clarke, and A. F. Bennett. 2012. Predicting the century-long post-fire responses of reptiles.
719 *Global Ecology and Biogeography* 21:1062–1073.
- 720 Noble, J. C., and R. G. Vines. 1993. Fire studies in mallee (*Eucalyptus* spp.) communities of
721 western New South Wales: grass fuel dynamics and associated weather patterns. *The*
722 *Rangeland Journal* 15:270–297.
- 723 Parr, C. L., and A. N. Andersen. 2006. Patch mosaic burning for biodiversity conservation: a
724 critique of the pyrodiversity paradigm. *Conservation Biology* 20:1610–1619.
- 725 Penman, T. D., F. J. Christie, A. N. Andersen, R. A. Bradstock, G. J. Cary, M. K. Henderson, O.
726 Price, C. Tran, G. M. Wardle, R. J. Williams, and A. York. 2011. Prescribed burning: how

- 727 can it work to conserve the things we value? *International Journal of Wildland Fire* 20:721.
- 728 Possingham, H. P., S. J. Andelman, B. R. Noon, S. Trombulak, and H. R. Pulliam. 2001. Making
729 smart conservation decisions. Pages 225–244 *in* M. E. Soulé and G. Orians, editors.
730 *Conservation Biology: Research Priorities For The Next Decade*. Island Press, USA.
- 731 Prendergast, J. R., R. M. Quinn, J. H. Lawton, B. C. Eversham, and D. W. Gibbons. 1993. Rare
732 species, the coincidence of diversity hotspots and conservation strategies. *Nature* 365:335–
733 337.
- 734 R Development Core Team. 2013. R: A language and environment for statistical computing. R
735 Foundation for Statistical Computing, Vienna.
- 736 Richards, S. A., H. P. Possingham, and J. Tizard. 1999. Optimal fire management for
737 maintaining community diversity. *Ecological Applications* 9:880–892.
- 738 Rodrigues, A. S. L., and T. M. Brooks. 2007. Shortcuts for biodiversity conservation planning:
739 the effectiveness of surrogates. *Annual Review of Ecology, Evolution, and Systematics*
740 38:713–737.
- 741 Sandell, P., K. G. Tolhurst, J. Dalton, B. Scott and M. Smith. 2006. Fire management
742 prescriptions for the Victorian Mallee Parks. *Proceedings of the Royal Society of Victoria*
743 118:395–412.
- 744 Taylor, R. S., S. J. Watson, D. G. Nimmo, L. T. Kelly, A. F. Bennett, and M. F. Clarke. 2012.
745 Landscape-scale effects of fire on bird assemblages: does pyrodiversity beget biodiversity?
746 *Diversity and Distributions* 18:519–529.
- 747 Teague, B., R. McLeod, and S. Pascoe. 2010. 2009 Victorian Bushfires Royal Commission final
748 report, Vol 2. Fire preparation, response and recovery. Parliament of Victoria, Melbourne.
- 749 Watson, S. J., R. S. Taylor, D. G. Nimmo, L. T. Kelly, A. Haslem, M. F. Clarke, and A. F.

750 Bennett. 2012. Effects of time since fire on birds: how informative are generalized fire
751 response curves for conservation management? *Ecological Applications* 22:685–696.
752 Williams, P. H., and M. B. Araujo. 2000. Using probability of persistence to identify important
753 areas for biodiversity conservation. *Proceedings of the Royal Society B: Biological Sciences*
754 267:1959–1966.
755

essa
preprint

756 **Ecological Archives**

757 Appendices

- 758 A. Map of wildfire scars from 1972-2011 within the Murray Sunset National Park.
- 759 B. Annual probability of wildfire in *T. scariosa* mallee vegetation, within the Murray Sunset
- 760 National Park.
- 761 C. Optimal strategy and expected value of the reserve for the bird-only analysis.
- 762 D. Optimal strategy and expected value of the reserve for the reptile-only analysis.
- 763 E. Relative performance of the objective functions, through time, when adhering to the
- 764 optimal strategy for each of the alternate objective functions. Data is for all 22 species
- 765 combined.
- 766 F. Optimal strategy and expected value of the reserve when constraining the solution to
- 767 achieve a 3% annual fuel reduction target.

768

769 Table 1. Habitat preferences of the 22 vertebrate species in three vegetation successional age
 770 classes; Early 0-10 yrs, Intermediate 11-35 yrs, Late 36-110 yrs. Species are grouped by the age
 771 class in which they have the highest probability of occurrence.

Age class (successional state)	Birds	Reptiles	All species (birds, reptiles & 1 small mammal)
Early (0-10 yrs)	1	4	5
Intermediate (11-35 yrs)	3	4	7
Late (35+ yrs)	6	3	10

772

Fig. 1. Schematic representation of the state and transition model of system dynamics. System states are represented by circles (early, intermediate and late successional vegetation), solid arrows denote successional transitions between the vegetation states and dashed arrows denote transitions occurring with fire.

Fig. 2. Optimal strategy and expected value of the reserve for the four objectives that aim to maximize the abundance of the 22 species throughout the reserve over a time frame of 100 years. The relative value of each state for biodiversity is represented by the grey scale; with 1 being the highest value. The objective functions are a) Binary: constrain to have 30% intermediate and 40% late; b) maximize the arithmetic mean probability of species occurrence; c) maximize the geometric mean probability of species occurrence; d) minimize extinction risk. The way in which each objective function combines the individual species requirements modifies the expected state-specific value of the reserve. The optimal strategy for each state of the reserve is denoted by a letter: *L* burn 20% late; *I* burn 10% late; fight wildfires has no letter. Taking no action and burning intermediate successional vegetation was never optimal in these cases.

Fig. 3. Optimal strategy and expected value of the reserve under an average fire-interval of 200 yrs. The four objectives aim to maximize the abundance of the 22 species throughout the reserve over a time frame of 100 years. The relative value of each state for biodiversity is represented by the grey scale; with 1 being the highest value. The objective functions are a) Binary: constrain to have 30% intermediate and 40% late; b) maximize the arithmetic mean probability of species occurrence; c) maximize the geometric mean probability of species occurrence; d) minimize extinction risk. The way in which each objective function combines the individual species

requirements modifies the expected state-specific value of the reserve. The optimal strategy for each state of the reserve is denoted by a letter: *L* burn 20% late; *I* burn 10% late; fight wildfires has no letter. Taking no action and burning intermediate successional vegetation was never optimal in these cases.

Fig. 4. Results for the objective function based on the geometric mean comparing the optimal strategy and value of the reserve, over a 100 year period, when a) all 22 species are modelled, b) only birds are modelled, and c) only reptiles are modelled. The value of each state for biodiversity is represented by the grey scale; with 1 being the highest value. The optimal strategy for each state of the reserve is denoted by a letter: *L* burn 20% late; *I* burn 10% late; fight wildfires has no letter. Taking no action and burning intermediate successional vegetation was never optimal in these cases.

Fig. 5. Relative performance of the objective functions, through time, when adhering to the optimal strategy for each of the alternate objective functions. Results are the average of 1000 simulations for the 100-year period. The values are expressed as a percentage of the value it would be if the optimal strategy identified for that objective was followed. Strategies presented are the a) arithmetic mean (AM), b) binary (BN), c) geometric mean (GM) and d) extinction risk (ER) with the objectives denoted by the legend. Data is for the 11 reptile species.

Fig. 6. Relative change in the index proportional to the probability of extinction (ER) for all 22 species when constraining the optimal solution to account for different levels of annual

818 prescribed burning. The horizontal black line represents the optimal management scenario. Time
819 frame is 100 years. Values are the arithmetic mean of all reserve states +/- standard deviation.

820
821

esa
preprint

822

823 Fig. 1.

Fig. 2.

Fig. 3.

829

830 Fig. 4.

preprint

831

832 Fig. 5.

833

834 Fig. 6.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Giljohann, KM;Mccarthy, MA;Kelly, LT;Regan, TJ

Title:

Choice of biodiversity index drives optimal fire management decisions

Date:

2015-01-01

Citation:

Giljohann, K. M., Mccarthy, M. A., Kelly, L. T. & Regan, T. J. (2015). Choice of biodiversity index drives optimal fire management decisions. ECOLOGICAL APPLICATIONS, 25 (1), pp.264-277. <https://doi.org/10.1890/14-0257.1>.

Persistent Link:

<http://hdl.handle.net/11343/116180>