Title: Predicting Novel Riparian Ecosystems in a Changing Climate

Running head: Novel riparian ecosystems in future climates

Authors: Jane A. Catford^{1*}, Robert J. Naiman^{2,3}, Lynda E. Chambers⁴, Jane Roberts^{5,6}, Michael Douglas⁷, and Peter Davies³

5 **Addresses:**

¹School of Botany, The University of Melbourne, Victoria 3010, Australia

²School of Aquatic and Fishery Sciences - 355020, University of Washington, Seattle, Washington 98195, USA

10 ³Centre of Excellence in Natural Resource Management, University of Western Australia, Albany, Western Australia 6330, Australia

⁴Centre for Australian Weather and Climate Research, Bureau of Meteorology, GPO Box 1289, Victoria 3001, Australia

⁵Institute of Land, Water and Society, Charles Sturt University, Albury-Wodonga, New South Wales 2640

15 ⁶PO Box 6191, O'Connor, Australian Capital Territory 2602, Australia

⁷Research Institute for the Environment and Livelihoods, Charles Darwin University, Darwin, Northern Territory 0909, Australia

***Corresponding author:** E. catfordj@unimelb.edu.au T. +613 8344 5422, F. +613 9347 5460

RECEIVED 15 DECEMBER 2011; ACCEPTED 25 MAY 2012

20 **Author contributions:** JAC, RJN, LEC and JR conceived and designed the study; all authors performed the research and wrote the paper under the leadership of JAC.

Abstract:

Rapid changes in global climate are likely to alter species assemblages and environmental characteristics resulting in novel ecosystems. The ability to predict characteristics of future ecosystems is crucial for environmental planning and the development of effective climate change adaptation strategies. This paper presents an approach for envisioning novel ecosystems in future climates. Focusing on riparian ecosystems, we use qualitative process models to predict likely abiotic and biotic changes in four case study systems: tropical coastal floodplains, temperate streams, high mountain streams and urban riparian zones. We concentrate on functional groups rather than individual species and consider dispersal constraints and the capacity for genetic adaptation. Our scenarios suggest that climatic changes will reduce indigenous diversity, facilitate non-indigenous invasion (especially C4 graminoids), increase fragmentation and result in simplified and less distinctive riparian ecosystems. Compared to models based on biota-environment correlations, process models built on mechanistic understanding (like Bayesian belief networks) are more likely to remain valid under novel climatic conditions. We posit that predictions based on species' functional traits will facilitate regional comparisons and can highlight effects of climate change on ecosystem structure and function. Ecosystems that have experienced similar modification to that expected under climate change (for example, altered flow regimes of regulated rivers) can be used to help inform and evaluate predictions. By manipulating attributes of these system models (for example, magnitude of climatic changes or adaptation strategies used), implications of various scenarios can be assessed and optimal management strategies identified.

Keywords: Non-indigenous species invasions; community composition; environmental planning; functional traits; emerging ecosystems; process models

Introduction

Novel (or emerging) ecosystems are characterized by new species assemblages that result from human-induced changes to the environment and biota (Hobbs and others 2006; Walther and others 2009). Species occur in combinations and abundances that have not occurred previously, which can prompt changes in ecosystem structure and function (Hobbs and others 2006). Humans intentionally and unintentionally facilitate the development of novel ecosystems through the extinction of indigenous taxa or depletion of their propagule pool, introduction of non-indigenous taxa, creation of dispersal barriers (for example, cities, roads, dams) and environmental modifications (for example, agriculture, fire suppression, river regulation, overgrazing; Hobbs and others 2006). Hybrid ecosystems have a combination of original and novel components that are outside the historical range of ecological variability (Hobbs and others 2009). There are no definitive boundaries between hybrid and novel ecosystems, so – for simplicity – we refer to both as novel ecosystems in this paper.

Biological invasions and climate change are among the most widespread and dominant causes of ecological change in all types of environments (Pressey and others 2007); their global reach and increasing rates of change suggest that many of the world's ecosystems will transition to novel states, including riparian zones. Human-induced climate change and the associated increase in greenhouse gas concentrations are predicted to affect precipitation, temperature, storminess and precipitate a rise in sea level (Hulme 2005). The type, extent and intensity of these changes will vary among ecosystems and their geographic location, so myriad scenarios may unfold. Being the interface between terrestrial and freshwater ecosystems (Naiman and Décamps 1997; Naiman and others 2005), riparian ecosystems are affected by changes that occur to watercourses, catchments and riparian zones themselves (Davies 2010). Of particular concern are global alterations in river discharge; in future climates, many river basins will experience water stress either from extended periods of waterlogging or from drought (Palmer and others 2008; Vörösmarty and others 2010). Rapid changes in climate may reduce or eliminate habitable space for many species (Pressey and others 2007), alter species distributions, prompt changes in biotic interactions (Morin and Thuiller

2009) and will potentially alter ecosystem processes and services (S.J. Capon, S. Williams, L. Chambers and others, this issue).

In addition to changes prompted by various types of environmental modification (Catford and others 2011), human-mediated introduction of non-indigenous (non-native, alien, exotic) species is altering community composition at an ever-increasing rate (Hulme and others 2009). Predictions suggest that, under climate change, non-indigenous species will expand and reproduce in previously inhospitable areas (Walther and others 2009) and that higher temperatures will favor traits that are usually possessed by non-indigenous species, facilitating their dominance (Sandel and Dangremond 2012). Growth rates of invasive plant species may increase more than their indigenous counterparts from CO₂ enrichment (Dukes and others 2011). Combined with their greater plasticity (Davidson and others 2011) and broader environmental ranges (Canning-Clode and others 2011; Davey and others 2012), CO₂ enrichment may further increase the dominance of invasive species. Dispersal will become more important as environmental conditions shift geographically thus giving a further advantage to non-indigenous species, which are likely to experience greater dispersal opportunities than indigenous species because of their close association with humans (Catford and others 2012). Unlike indigenous species that are adapted to historical conditions, non-indigenous species may also be pre-adapted (or adapt rapidly) to the new environmental conditions (Catford and others 2012). Riparian ecosystems are among the most invaded types of ecosystem (Richardson and others 2007) and it is likely that they will continue to have a high non-indigenous component in the future.

Natural resource management, maintenance of ecosystem services and conservation of biodiversity require information about patterns of biodiversity and ecosystem processes and condition (Pressey and others 2007). The ability to predict the ecological character and function of future ecosystems is therefore crucial for environmental planning, like reserve design, and the development of effective mitigation and adaptation strategies (Morin and Thuiller 2009). Because, by definition, novel ecosystems have not previously existed, it is very difficult to predict their characteristics and few attempts have been made to do so. Controlled experiments cannot be carried out at appropriate

spatial or temporal scales, so modeling approaches are required.

In this paper, we propose a qualitative approach for envisioning future riparian ecosystems. Based around four recommendations, we present the approach in the first part of the paper. We then use four case studies from contrasting environments to illustrate the approach and to determine:

- 5 - Whether certain characteristics make some ecosystems more susceptible to climate-induced shifts in community structure than others; and
- Which aspect of climate change seems to have the greatest effect on community structure and therefore should be a research priority.

10 Focusing on changes in community structure, we investigate likely consequences of climate change for tropical floodplains, temperate streams, high mountain streams and urban riparian zones. The paper centers on Australian riparian zones experiencing climate change, but we note that the approach can be applied to ecosystems in other biomes that are subject to environmental change. Although human adaptations to climate change can lead to novel ecosystems, we do not address these changes. Instead we concentrate on unintentional physical and biogeochemical changes that, 15 although the result of human activities, are outside of direct and immediate human control and manipulation.

Ways to predict novel ecosystems: laying out an approach

1. *Use process models to predict characteristics of future ecosystems, but integrate their predictions with those from other models.*

20 We advocate the use of process (or mechanistic or causal) models, rather than correlative models, when envisioning characteristics of novel ecosystems. Process models are based on ecological mechanisms, dynamics and feedbacks; they consider the various factors that influence ecosystem characteristics and indicate how a change in one factor can lead to changes in others (Pearl 2009). Using forecasts about environmental change combined with current understanding of ecological

relationships and processes, process models can be used to predict ecological responses. Because they are based on mechanistic understanding, such predictions are more general than those based purely on correlative relationships and are more likely to remain valid under novel climatic conditions (Koca and others 2006). However, like any predictive models, process models assume that current knowledge can predict future relationships (Hulme 2005). However, if relationships are non-linear and move to a level and type not witnessed before, the capacity to predict cause-effect relationships will diminish. Updating cause-effect relationships as new information becomes available is crucial for the efficacy of process models.

There are a number of forms that predictive models can take (Ferrier and Guisan 2006; Koca and others 2006; Morin and Thuiller 2009; Kearney and others 2010). To help counter deficiencies in individual modelling approaches, it is important to develop several different types of models and compare their predictions (Morin and Thuiller 2009; Kearney and others 2010). Accordingly, a number of existing modelling approaches could be adapted to help predict novel ecosystems, for example, models based on responses of specific species (Morin and Thuiller 2009), vegetation types (Koca and others 2006) and communities (Ferrier and Guisan 2006).

Here we devise qualitative process models based on expert knowledge. A pre-requisite of quantitative models, qualitative models enable rapid appraisal of the type of ecological changes that can be expected under climate change. Qualitative models can be used to gauge the relative importance of different drivers, though the magnitude of changes cannot be accurately assessed.

2. *Use functional groups, as well as species, to predict the types of communities that will develop in the future.*

We recommend predictions based on functional groups to complement predictions about individual species and vegetation types. Functional groups are collections of taxa grouped according to shared traits, life history characteristics or ways they interact with the environment (Keith and others 2007).

Classifications can be based on species' morphological, physiological, phenological and behavioral

characteristics. Functional traits affect growth, reproduction and survival of organisms (Violle and others 2007) and can indicate early responses to environmental change (Merritt and others 2010), including climate. In this paper, we are specifically interested in shifts in community structure that relate to climate change so we concentrate on the biological form of animal and plant taxa and, when germane, use these descriptions in combination with the environmental preferences of the taxa (Table 1). However, for illustrative purposes, we occasionally refer to culturally significant species or ones that may be considered “keystone” or “umbrella” species.

To date, most predictions about community composition under future climates have relied upon species-based forecasts (Hijmans and Graham 2006). Correlative species distribution models and mechanistic niche models can indicate the likely future distribution and abundance of individual species (Elith and others 2010), and these approaches have merit when interested in the fate of particular species (especially when used in combination; Morin and Thuiller 2009). However, the large number of species (including those in Australia’s biodiverse riparian zones) makes broad-scale application of this approach untenable; there are insufficient data and resources to model the distributions of thousands of species. Plus, individual species respond differently in different places making it hard to generalize from one region to the next. Even in mechanistic niche models that incorporate species’ ecophysiological requirements, community interactions cannot be accurately predicted. Although lack of information about community dynamics limits the ability of any method (including those based on functional groups) to accurately predict the composition of future biological communities (compare Ferrier and Guisan 2006), more general approaches may be less sensitive to this uncertainty. For instance, individual species from one functional group may buffer the response of others.

Use of functional groups overcomes some of the disadvantages of using a species-based approach (Neilson and others 2005). Based on traits rather than the identity of individual taxa, functional groups enable findings to be generalized and compared across ecosystems and biomes (Westoby 1998). Depending on the type of functional classification used, they enable implications of climate

change for ecosystem structure and function to be more apparent (Violle and others 2007) and often can integrate information about constituent taxa.

The influence of history on community composition may be less important when considering functional groups as opposed to individual species. Variation in the order of species immigration results in diverse and often unpredictable community assemblages, but community assembly based on trait-based functional groups may be less stochastic and more deterministic (Fukami and others 2005; though evidence for this remains mixed: Trowbridge 2007). It should, however, be acknowledged that not all species will follow the trends of their group (Keith and others 2007), and it can be harder to evaluate model predictions when using groups of taxa (Neilson and others 2005).

3. *Use of analogue systems to help envisage the sorts of ecosystems that might develop in the future.*

Although novel ecosystems are – by definition – “new”, it may be possible to use ecosystems with similar attributes (or those that have experienced similar conditions at some point in time) to help inform predictions. Flow regulation and diversion of water from rivers may result in similar effects as the reduction in precipitation associated with climate change. Similarly, urbanization and an increase in the proportion of impervious surfaces may be similar to an increase in extreme flow events: both result in a “flashy” hydrograph with increased river discharge over a short period of time. In this sense, information about ecological changes in regulated and urban rivers could be used to inform predictions about climate change effects in unregulated and rural rivers. As well as modified systems, ecosystems in different biogeographic regions may serve as suitable comparisons. For example, many lowland rivers in Australia’s temperate south-east may become more like dryland rivers in arid and semi-arid Australia. At a broader level, these systems could even start to resemble those in southern states of the US, like Arizona and Nevada.

The use of functional groups will facilitate comparisons across biogeographic regions. The world’s biota is becoming increasingly similar (Stohlgren and others 2011) as the rate of species transfer and

biological invasions continues to rise (Hulme and others 2009), so regional differences in biota are declining (Villéger and others 2011). Knowledge of similar ecosystems could be used to help inform the development of process-based models for predicting novel ecosystems, but care must be taken as differences may still be great.

5 4. *Incorporate information about taxon migration rates, dispersal patterns and genetic adaptation.*

The majority of models that predict biological responses to climate change do not explicitly consider species migration or dispersal (Neilson and others 2005; but refer to work on invasive species, for example, Elith and others 2010). However, it cannot be assumed that, just because a species can tolerate the environmental conditions of a site, it will occur there now or in the future. Whether
10 based on individual species or functional groups, modelling approaches need to include information about migration and succession processes (Neilson and others 2005), dispersal pathways (including the role of human as dispersal vectors) and proximity of source populations.

Similarly, the ability of a species or population to adapt to altered conditions must be considered. It is likely that species with higher phenotypic or genotypic plasticity will tolerate changes in climate
15 better than species with low genetic diversity or plasticity. This is thought to be one reason why the abundance of invasive non-indigenous species will increase under climate change (Davidson and others 2011).

Scenarios of change

Using the four recommendations presented above, we develop process models to examine the
20 potential consequences of climate change in four riparian ecosystems in Australia. The patterns and consequences of climate change for riparian zones will vary according to local conditions. These conditions include not only the rate and magnitude of climate alterations but also how they interact with topography, land use, human legacies (for example, past land uses), flow regimes for water and sediment, and indigenous and non-indigenous species. We illustrate this complexity by examining
25 expected changes for riparian zones in contrasting environmental settings and with different

histories of human activities. The sites represent a broad range in environmental characteristics and thus have the potential to indicate general trends as well as regional differences. They are also ecosystems with which we are most familiar. We consider climate change impacts on temperature, precipitation, sea level rise and carbon dioxide concentrations. Addressing the aims of the paper, we use the models to assess whether some ecosystems are more vulnerable to change than others, and which aspect of climate change has the greatest effect on community structure. We first provide an overview of climate change impacts predicted in Australia.

Climate change impacts

As global climate change escalates, it is expected that average temperatures in Australia will rise by 0.6-1.5 °C above 1990 temperatures by 2030 and 2.2 °C and 5 °C by 2070 (Cleugh and others 2011). Rainfall projections are less certain than temperature projections, but most models suggest that rainfall will decline in southern Australia, particularly during winter (Davies 2010), and spring rainfall will markedly decline in southern and eastern Australia (Cleugh and others 2011). The magnitude of extreme rainfall events is expected to increase throughout the country (Cleugh and others 2011). In south-eastern Australia, reductions in snow depth and season length and an earlier timing of maximum snow depth are predicted (CSIRO and Bureau of Meteorology 2007). The El Niño-Southern Oscillation, a major driver of climate variability in eastern Australia, will become more extreme, and droughts become more severe (DECCW 2010). Changes in the seasonality and amount of precipitation will have important implications for stream flow and soil moisture. Warmer and drier conditions in southern Australia are likely to result in an increased fire-weather risk and a lengthening of the fire season (Lucas and others 2007). There is some uncertainty as to the magnitude of future sea level rise, but it is anticipated that sea level rise in the Australian region will increase the frequency of coastal flooding. For instance, coastal flooding events that used to occur once in every 100 years may occur more than once a year by 2100 (Cleugh and others 2011). CO₂ levels are expected to continue to rise until at least 2040 (Nakicenovic and others 2000), and further increases in CO₂ levels are likely to amplify long-term average warming (IPCC 2007).

Tropical floodplains in northern Australia

Coastal floodplain wetlands that flank macro-tidal river systems are conspicuous riparian habitats in northern Australia (Bayliss and Yeomans 1990). Having formed only 6,000-1,500 years ago, these seasonally inundated floodplains are recent landscape features yet they are highly productive and rich in biodiversity (Finlayson and others 2006). Internationally recognized for significant natural and cultural values they include, among others, the Ramsar and World Heritage listed wetlands of Kakadu National Park (Finlayson and others 2006). Environmental alterations caused by climate change will be large and complex in these tropical floodplain ecosystems.

The floodplains are still largely vegetated by indigenous species, but they are currently being modified by non-indigenous plants and animals and by altered fire regimes (Pusey and others 2011). They are also susceptible to rising sea levels (Bartolo and others 2008) because of their low elevation (most are <2 m above sea level). The floodplains were formed as recently as 1,500-6,000 years ago after a long period of sediment deposition by rivers when sea-levels retreated (Mulrennan and Woodroffe 1998). The landward margins of the floodplains are topographically constrained and cannot migrate inland as sea levels rise. Projections for sea level rise during the next century range from 0.4–1.0 m (Horton and others 2008). Although it is difficult to reliably quantify the extent of seawater intrusion in northern Australia due to a lack of high resolution elevation models, it is likely that more than half of the floodplain wetlands will be inundated with seawater over the next 50 years (Bartolo and others 2008) and nearly all within 300-400 years (Traill and others 2010).

The likely effects of seawater intrusion are based on previous experiences of seawater intrusion in the Northern Territory, on recent modelling of the potential impacts on key fauna and on recent research into the ecological role of tropical floodplains (Figure 1). Seawater intrusion has already occurred in some areas due to human activity, extreme weather events and the effects of non-indigenous water buffalo (*Bubalus bubalis*). These provide a valuable analogue for the likely effects of climate-induced changes. Elevated sea levels will alter floodplain hydrology and water quality.

Seawater intrusion will dramatically change the pattern of floodplain inundation as it will be driven by tidal movement and storm surge rather than rainfall alone. Saline water makes floodplain soils more dispersive, which will in turn extend seawater channels, facilitate the ingress of seawater (Winn and others 2006) and increase the drainage of freshwater from floodplains (Petty and others 2007). Increased salinity can kill salt-intolerant vegetation and facilitate a transition to salt-tolerant trees, forbs and graminoids, such as mangroves. For example, the East Alligator River has already experienced tidal intrusion up to 4 km inland, resulting in a nine-fold increase in the area of saline mudflats and a loss of salt-intolerant semi-aquatic vegetation, including 64% of the original *Melaleuca* forest (Winn and others 2006). Seawater intrusion has also prompted upstream colonization by mangroves in the Wildman and South Alligator Rivers (Cobb and others 2007). The most dramatic example of seawater intrusion in northern Australia is on the Mary River. Since tidal creeks began extending into Mary River wetlands in the 1930s and 1940s, more than 17,000 ha of freshwater floodplain have been affected by seawater, which has resulted in vast areas of salt-intolerant vegetation being replaced by a saline swamp and a transition to salt-tolerant vegetation (Mulrennan and Woodroffe 1998).

The loss of salt-intolerant vegetation has serious consequences for floodplain fauna including semi-aquatic birds such as magpie geese (*Anseranas semipalmata*, Figure 1), which rely on freshwater vegetation for forage and nesting. Magpie geese, once widespread across Australia but now restricted to the north, have a population of less than 3 million (Delaney and others 2009). They are a tourist icon, a popular species for recreational hunting, an important food source for Aboriginal people and are ecologically important because of their abundance (Delaney and others 2009).

Modelling has shown that a loss of just 5% of the wetland habitat currently available to magpie geese could reduce populations to just a few thousand individuals and possibly initiate an ecological bottleneck or “point-of-no-return” (Traill and others 2010). Other waterbirds, reptiles, such as the saltwater crocodile (*Crocodylus porosus*), and mammals that use the floodplains in the dry season will also be affected by reductions in the availability of suitable food and habitat (Figure 1; BMT

WBM 2010). For other animals though, effects will be more direct: it will simply be impossible for some to persist in the saline water, for example, salt-intolerant fish and reptiles such as freshwater turtles.

Saline water and the loss of freshwater floodplain vegetation will likely alter the food supply for many species through a reduction in freshwater epiphytic algae, which grows on plants and is a primary carbon source on these floodplains (Figure 1; Douglas and others 2005). Change in salinity and primary productivity may affect other aquatic habitats as many fish species feed, grow and reproduce on the floodplains and transport this energy to upstream riverine or downstream estuarine and coastal ecosystems (Jardine and others 2012). The floodplains are critical to the life history of many species including economically important fish species such as barramundi (*Lates calcarifer*).

Floodplains unaffected by seawater intrusion are also likely to be altered by direct climatic changes (Figure 1). Elevated CO₂ (Williams and Baruch 2000) and temperature (Hamilton 2010) favor non-indigenous, fire-tolerant semi-aquatic C4 graminoids (Williams and Baruch 2000) such as para grass (*Brachiaria mutica*), which could replace the diverse indigenous vegetation with a non-indigenous monoculture (Douglas and O'Connor 2003) and increase dry season fuel loads (Douglas and O'Connor 2004) and fire intensities. Increased temperatures are also expected to result in more frequent and intense dry season fires, which could further reduce the abundance of salt-intolerant semi-aquatic trees (for example, *Melaleuca* sp.) as well as fire-intolerant terrestrial trees in riparian vine forests. It is also highly likely that non-indigenous pigs (*Sus scrofa*), which disturb the soil, alter local hydrology and compete with indigenous fauna for plants and invertebrates, will concentrate in the remaining floodplain areas putting further pressure on these refugial habitats.

Riparian zones in temperate south-western Australia

South-western Australia is a global biodiversity hotspot and the only one located on the Australian continent (Myers and others 2000). The Gondwanic origin of the biota, combined with a long period

of geographic isolation, has resulted in distinctive riparian habitats and a high level of endemism in both freshwater and terrestrial species (Bunn and Davies 1990). Streams and rivers are characterized by low rates of primary production and aquatic food webs rely heavily on the input of riparian carbon (Davies 2010).

5 Riparian zones in this region are highly degraded as a result of flow regulation, intensive agriculture (Horwitz and others 2008) and uncontrolled livestock access (Bunn and others 1999). The clearing of indigenous vegetation commenced with European settlers in 1829 and escalated in the early part of the 20th century (Bolton 1972) resulting in a fragmented distribution of remnant vegetation.

Riparian restoration efforts have been patchy and success limited. Increasingly harsh climatic
10 conditions (typically warming and drying) will interact with the already fragmented landscape of south-western Australia (Horwitz and others 2008) leading to further degradation and the creation of novel riparian ecosystems.

Many ecological processes in river and riparian habitats are regulated by the magnitude, frequency and seasonality of flows (Poff and others 1997). As a consequence, substantial and complex changes
15 are expected in riparian areas from climate change (Figure 2). Altering the timing of rainfall will affect seed-set and the subsequent recruitment of riparian plants, particularly indigenous trees, which may in turn disrupt host-pollinator relationships (for example, a mismatch between insects emerging and plants flowering: Davies 2010). A reduction in the magnitude of winter flows will limit overbank flows and riparian inundation. Flow regulation in rivers in south-western Australia narrows
20 the riparian zone and reduces overall plant species richness (Pettit and others 2001). Such effects will likely be compounded under climate change, though an increase in invasive non-indigenous species in this region may partially compensate for declines in indigenous plant richness (O'Donnell and others 2012).

Drying and warming of the riparian zone will also lead to the removal of stenothermic animals and
25 may indirectly facilitate the establishment of invasive animal species, which often tolerate a broader

thermal range (in marine systems at least; Canning-Clode and others 2011). The predicted increase in the severity of fires will open the riparian canopy increasing light into the understory, thus providing more suitable conditions for non-indigenous plant species (Davies 2010). In lower river reaches, sea level rise will increase the extent of floodplain inundation and will inhibit salt-intolerant species. Many river and riparian systems in south-western Australia have undergone secondary salinization as a consequence of overclearing (Horwitz and others 2008). The community structure of macroinvertebrates in these rivers has shifted from insect forms to a dominance of crustaceans, common in nearby still-water habitats (Bunn and Davies 1992). Increased sea levels and intrusion into lower river floodplains is expected to result in a similar response with a shift in community structure favoring crustaceans.

Overall, we foresee climate change leading to increased fragmentation, narrowing of riparian zones and simplification (and homogenization) of the biota due to the loss of stenothermic species and increased abundance of non-indigenous flora that are dispersed widely by humans (Figure 2). In particular, an increase in non-indigenous plants in these novel ecosystems, particularly C4 graminoids, will reduce the current strong subsidy between the riparian zone and in-stream food webs (Bunn and others 1997). A more open canopy will encourage algal growth potentially leading to a shift from heterotrophy to autotrophy in streams. Increased autotrophy in association with low water levels is expected to increase the likelihood of anoxia. This could hasten the decline of indigenous species, especially fish and macroinvertebrates, and lead to an increase in more tolerant non-indigenous species.

Many riverine plant and animal species in south-western Australia are endemic with small range sizes and restricted geographic distributions (Bunn and Davies 1990). A result of range compression about 18,000 years ago, these endemic species have little genetic diversity (De Deckker 1986) and many are now restricted to national parks and nature reserves in the cool, wet south coast of south-

western Australia. As conditions become warmer and drier, these species are not be able to move southward nor is it likely that they can genetically adapt to the changes.

Riparian zones in the high country

In south-eastern Australia, land over 1000 m in elevation occurs as topographic islands with rounded
5 tops and steep flanks, the largest and highest being the northern part of the Snowy Mountains. The high country is significant for its glacial landscapes, endemic plant and animal alpine species, and as the water source for economically important rivers such as the Snowy, Murray and Murrumbidgee. Although now largely reserved in national parks, the high country has a history of human use.

Alluvial gold mining and timber harvesting have ceased, but tourism and hydro-electricity generation
10 continue. Plant communities are mostly intact except where disturbed by roads and engineering structures, or by the 100-year legacy of cattle grazing, particularly in bogs and fens. Non-indigenous plant species are relatively few, with low cover (McDougall and others 2005), though some highly invasive species have established recently and are spreading rapidly (*for example, Hieracium aurantiacum*, Orange hawkweed, Williams and others 2008). Non-indigenous animals (for example,
15 horses, deer and pigs) damage vegetation, especially along stream edges and in wetlands (McDougall and Walsh 2002).

The riparian plant communities of the high country are few, small in area and are floristically and structurally distinct from each other. They form a discontinuous riparian zone, arranged in altitudinal sequence from short alpine herbfields (forbs and graminoids) above the tree-line, to bogs
20 in the alpine and subalpine zones (typically a mix of semi-aquatic mosses, namely *Sphagnum cristatum*, forbs, graminoids and shrubs), then a fringe of medium-tall semi-aquatic and terrestrial shrubs at lower forested elevations (Costin 1954).

A prolonged drought (2001-2010) and the 2003 wildfires indicate some of the potential effects of a warmer, drier climate in the high country. Observations from this period thus serve as a suitable
25 analogue for climate change. After a few years of reduced rainfall and higher temperatures, bogs

and terrestrial herb communities will become drier (Good and others 2010) resulting in mortality and areal contraction and the development of bare patches (Figure 3). These bare patches may be colonized by regenerating terrestrial graminoids and shrubs, invaded by indigenous and non-indigenous species, or eroded by summer storms or frost-heave (McDougall and others 2005). Once

5 dry, bogs are more susceptible to wildfires, further shifting the balance from semi-aquatic herbs to terrestrial plants better adapted to fire (McDougall 2007). Compounding this trend, the forbs and graminoids (including endemic species) dependent on snow patches may be replaced by other plants as the cover and persistence of snow declines (Green and Pickering 2009).

If the exposed patches of soils are eroded, the material is likely to accumulate in headwater streams, especially if perennial streams become intermittent because of reduced rainfall. Like the desiccated and burnt bogs, in-stream accumulations of sediment and organic materials may be colonized by non-indigenous, as well as indigenous, species (McDougall and others 2011). Most non-indigenous species are expected to be eurythermic generalists from lower altitudes that can move to higher elevations via wind dispersal, non-indigenous herbivores and humans (McDougall and others 2011).

15 The variety of sites and the type of invasion pathways available suggest that species establishing in stream channels could include semi-aquatic herbs, shrubs and trees such as willow (*Salix* spp.), recognized as a serious environmental weed in Australia and one that would dramatically alter the vegetation structure and in-stream functioning. The combined effects of sedimentation and vegetation change on high country fauna are potentially dramatic: a shift from perennial to intermittent streams and loss of bogs will cause localized and possibly even total extinctions of endemic fish, amphibians and small mammals, such as galaxids, the Northern Corroboree frog *Pseudophryne pengilleyi* and the broad-toothed rat *Mastacomys fuscus*; in-stream sedimentation is likely to change invertebrate abundance and composition (Figure 3; Harrison and others 2008).

Under climate change, these riparian systems are likely to comprise three elements with respect to vegetation: fragments of the current riparian vegetation; replacement of semi-aquatic moss and shrub communities by terrestrial graminoids and shrubs; and the presence of opportunistic semi-

aquatic herbs, shrubs and possibly even trees in patches of in-stream sediment, analogous to those that develop on fine and coarse sediments of regulated rivers at lower altitudes. Although smaller in extent and more species-rich than contemporary systems, future riparian ecosystems are likely to become more variable in structure, less distinct and their altitudinal sequence more diffuse.

5 ***Urban Melbourne***

Urban areas represent one of the most dramatic ways humans can alter ecosystems, especially riparian zones (Groffman and others 2003; Meyer and others 2005). Urbanization impacts on riparian zones may exceed those associated with climate change, but it is likely that ecosystem alterations are compounded when they occur simultaneously.

- 10 Over half of the world's human population lives in urban areas (United Nations 2010) and this proportion will increase in coming decades (Meyer and others 2005). Australia is one of the more urbanized countries in the world, with about 90% of its population residing in cities (United Nations 2010). In south-eastern Australia, over four million people currently reside in Melbourne (Australian Bureau of Statistics 2011). Over the period 2001 to 2011, this city has seen the largest annual growth
- 15 of all Australian capital cities with an average increase of over 1,300 people per week (Australian Bureau of Statistics 2011).

Founded in the 1830s, Melbourne experienced rapid growth following the discovery of gold in 1851.

- This led to wholesale ecosystem changes. River hydrology, hydraulics and geomorphology were altered through a number of processes, including: clearance of streamside vegetation, flattening of
- 20 hills, draining of wetlands, diversion of streams and rivers (sometimes underground), dam construction and water extraction, conversion of retentive small upland drainage lines into a vast network of hydraulically efficient stormwater pipes, and an increase in impervious (non-porous) surfaces (Leahy and others 2005; Otto 2005). Although many measures were put in place to reduce flood risk, they actually increased it: removal of vegetation and an increase in hard surfaces
- 25 accelerated runoff leading to increased flooding (Otto 2005; Lake 2008). Melbourne's metropolitan

area is now almost entirely served by stormwater drains that bypass riparian zones and thus do not make use of their capacity to ameliorate diffuse runoff pollution (Walsh and others 2001). The hyperconnection of impervious surfaces throughout the catchment to the stream by pipes, the disconnection of rivers from floodplains, and alteration of riparian vegetation has severely reduced the ecological quality of Melbourne's streams and rivers (Walsh and others 2001).

Most indigenous riparian plant communities in Melbourne have been eliminated, degraded or reduced to small remnants (though there are exceptions; Walsh and others 2007). Together with the loss of many wetlands and direct urban impacts, this has resulted in significant faunal changes (Leahy and others 2005). Substantial changes in riparian flora and fauna also occurred through planned (including willow *Salix* spp., blackberry *Rubus* spp., rabbit *Oryctolagus cuniculus* and fox *Vulpes vulpes*) and accidental introductions of non-indigenous species (Otto 2005). Collectively, these changes have resulted in novel riparian ecosystems that are expected to be further modified by climate change.

In many ways, potential effects of climate change on riparian systems have a number of similarities with effects of urbanization (Table 2, Figure 3): channel form can be altered through engineering works or channel incision, which limit over-bank flows and isolate floodplains for long periods; floodplains are fragmented in ways that impede dispersal of terrestrial species along the river corridor; the balance of erosion and sedimentation is outside historic experiences; non-indigenous species dominate; efficiency of nutrient cycling on floodplains and in channels is low with urban riparian zones acting as sources rather than sinks for nitrate; populations of indigenous plants and animals fail to recruit sufficiently to maintain numbers across range of age classes; and the diversity of components and processes declines. With a growing human population size, it is likely that Melbourne's urban riparian zones will continue to experience high levels of disturbance and there will be increased demand for water interception and extraction, particularly in response to the drying effects of climate change (Seavy and others 2009). Although reductions in rainfall in pervious catchments will lead to disproportionately greater reductions in runoff, proportional reductions in

rainfall and runoff in impervious catchments will be similar. As a consequence, by re-engineering stormwater drainage systems to artificially maintain more natural wetting and drying cycles of riparian soils, there is a greater opportunity to buffer some effects of climate change in urban areas (Walsh and others 2005).

- 5 Climate change and urbanization act together to amplify system decline. However, provided suitable remnant vegetation occurs and riparian soils experience wetting events, it is likely that urban riparian zones will continue to act as local micro-climate refugia from the warmer, higher density urban areas. As such, riparian ecosystems in urban Melbourne, novel as they may be, have the potential to act as local biodiversity hotspots and wildlife corridors and offer some relief from
- 10 climate change if managed “creatively” (Seavy and others 2009).

Discussion

- The four scenarios highlight the complexity and potential magnitude of climate change impacts on riparian ecosystems in Australia. Each suggests an increase in non-indigenous species, especially C4 graminoids, and a reduction in indigenous diversity. Even if overall diversity changes little, loss of
- 15 specialization through the decline of specialist species and range expansion of generalist species will lead to homogenization and loss of ecological distinctiveness, as observed elsewhere (Davey and others 2012). The models suggest that changes in rainfall (that is, stream discharge) are likely to have the largest direct influence on riparian ecosystems overall, with the exception of coastal floodplains that will be affected by sea level rise. We suggest that improving predictions about
- 20 future rainfall patterns and rainfall-discharge relationships is a key research priority for predicting the magnitude of ecological change in riparian zones. Currently, predictions about rainfall in Australia are generally less certain than temperature projections (Cleugh and others 2011). For coastal floodplains, improved digital elevation models are needed to underpin hydrodynamic models so that more accurate predictions can be made about the extent of seawater intrusion.
- 25 Changes in flow and increases in temperature were predicted to affect all four ecosystems, but

community structures were not predicted to change in the same way or to the same extent. Because they are geographically restricted, riparian ecosystems in the high country and in south-western Australia will be especially susceptible to species extinctions from increases in temperature. Changes in these ecosystems will likely be compounded by their geographic isolation and high levels of

5 endemism as well: without dispersal of similar taxa from similar types of ecosystems and with limited capacity for genetic adaptation, the loss of endemic species will likely herald a marked shift in community structure. In contrast, the recent age of tropical floodplains and the highly dynamic hydrology means that there are few endemic species and much of the fauna undertakes seasonal migration and is highly adapted to dispersal. However, their uniformly low elevation means that

10 massive losses of freshwater habitat will occur rapidly once sea levels exceed a certain threshold. Although human-mediated dispersal of non-indigenous species may partially compensate for loss of indigenous species in the high country and south-western Australia when focusing on total species richness, it is likely that the “new arrivals” will be functionally different. In contrast to the other ecosystems, riparian zones in Melbourne have already experienced a high degree of modification

15 through urbanization, so changes resulting from climate change may be relatively negligible. Although riparian zones, as an ecosystem type, will persist under climate change, we expect that their structure, processes and ecosystem services will change, at least in part. For instance, fragmentation in southern temperate ecosystems will reduce the role of riparian zones as dispersal corridors, and loss of freshwater wetlands in northern Australia will reduce recreational and

20 economic opportunities associated with hunting and fishing. These changes create complex challenges for management especially if fundamental ecosystem properties vary significantly among the types of novel riparian zones. Anticipating the type of changes likely to occur in the future is essential for the efficacy of long-term environmental management and planning. Knowledge of the key factors that shape ecosystem characteristics – and the extent to which those factors will change

25 in the future – will be instructive in predicting the properties of the emerging ecosystems.

Value and extension of approach

Despite encompassing considerable uncertainty (for example, nature of climate change impacts and ecological responses to these impacts; Hulme 2005; Koca and others 2006), our predictions illustrate that altering several key abiotic factors may lead to the development of a range of novel ecosystems. By using process models and functional groups, we were able to envisage the potential implications of climate change on the abiotic characteristics and community structure of these riparian ecosystems. Considering the dispersal and genetic constraints of taxa was fundamental to this goal. Comparison with systems that have similar conditions or have experienced similar changes enabled us to assess the likelihood of our predictions. Such comparisons will become more important if quantitative predictions are desired.

To gauge the magnitude of predicted changes and the effectiveness of different management approaches, the models presented here could be further refined by including quantitative relationships. This would facilitate the development of quantitative predictions. Based on analyses of similar issues (Murray and others 2012; Rumpff and others 2011), Bayesian belief networks appear to be particularly suitable for including both qualitative and quantitative understanding (Figures 1-4).

Bayesian networks are a type of process model that are increasingly being used to predict ecological responses to environmental change and management actions (Rumpff and others 2011). They take the form of flow charts where the relationships among, and values of, factors are specified. They can include uncertainty about both the understanding and variability of ecosystem responses, which is instructive when making management decisions (McCann and others 2006). Data from previous

studies can be used to parameterize the models, as can information elicited from experts. By manipulating the value of variables included in the models (for example, altering the magnitude of climate change impacts or altering the adaptation strategies used), the implications of a number of different scenarios on ecosystem structure and function can be rapidly assessed. The models can be updated as more information becomes available and understanding increases, and the way they operate can be easily communicated to non-experts (McCann and others 2006). These attributes make Bayesian networks particularly attractive for use in adaptive management (Rumpff and others

2011), which would be a suitable approach for climate change adaptation.

Most conservation activities have focused on conserving particular suites of species in particular locations (Pressey and others 2007; Hobbs and others 2009). Such a goal, however, may be

unrealistic as ecosystems and their species assemblages change. Instead, it may be more pragmatic

5 to focus on maximizing genetic, species and functional diversity, maintaining the biogeochemical configuration of a system (Seastedt and others 2008) or increasing habitat complexity. The ability to accurately predict the characteristics of future ecosystems is crucial for setting realistic management objectives and represents an exciting area of research.

10 **Acknowledgments:**

Ideas for this paper were conceived at a “Riparian Ecosystems under Climate Change” workshop

hosted by the National Climate Change Adaptation Research Facility – Water Resources and

Freshwater Biodiversity Network and Terrestrial Biodiversity Network. Thanks to other workshop

participants and especially Ugo Arbieu for comments in the early stages of this paper. Thanks also to

15 Brendan Wintle for insightful comments on a draft manuscript, Chris Walsh for reviewing the

Melbourne scenario and an anonymous reviewer for constructive feedback. P.D. and M.D.

acknowledge the Tropical Rivers and Coastal Knowledge (TRaCK) research consortium and M.D. also

acknowledges support from the NERP North Australian Hub; JC was partially supported by the ARC

Centre of Excellence for Environmental Decisions.

20 **References**

Australian Bureau of Statistics. 2011. Regional Population Growth, Australia, 2010-11.

<http://www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/3218.0Main+Features12010-11?OpenDocument> (last accessed 23/4/2012)

- Bartolo R, Wasson R, Valentine E, Cleland S, Bayliss P, Winderlich S. 2008. Climate change – The status of climate change research in the Kakadu landscape context. Kakadu National Park Landscape Symposia Series 2007-2009. Symposium 1: Landscape Change Overview. Darwin. p84-96.
- Bayliss P, Yeomans KM. 1990. Seasonal distribution and abundance of magpie geese, *Anseranas semipalmata* in the Northern Territory, and their relationship to habitat, 1983-86. Australian Wildlife Research 17: 15-38.
- Bolton GC. 1972. A Fine Country to Starve In. Perth: University of WA Press. 278p.
- Bunn SE, Davies PM. 1990. Why is the stream fauna of south-western Australia so impoverished? Hydrobiologia 194: 169-176.
- 10 Bunn SE, Davies PM. 1992. Community structure of the macroinvertebrate fauna and water chemistry of a salinised river system in south-western Australia. Hydrobiologia. 248: 143-160.
- Bunn SE, Davies PM, Kellaway D. 1997. Contributions of sugar cane and invasive pasture grass to the aquatic food web of a tropical lowland stream. Marine and Freshwater Research 48: 173-179.
- Bunn SE, Davies PM, Mosisch TD. 1999. Ecosystem measures of river health and their response to
15 riparian and catchment degradation. Freshwater Biology 41: 333-345.
- Canning-Clode J, Fowler AE, Byers JE, Carlton JT, Ruiz GM. 2011. 'Caribbean Creep' chills out: climate change and marine invasive species. PLoS ONE 6: e29657.
- Catford JA, Downes BJ, Gippel CJ, Vesk PA. 2011. Flow regulation reduces native plant cover and facilitates exotic invasion in riparian wetlands. Journal of Applied Ecology 48: 432-442.
- 20 Catford JA, Daehler CC, Murphy HT, Sheppard AW, Hardesty BD, Westcott DA, Rejmánek M, Bellingham PJ, Pergl J, Horvitz CC, Hulme PE. 2012. The intermediate disturbance hypothesis and plant invasions: Implications for species richness and management. Perspectives in Plant Ecology, Evolution and Systematics, in press.
- Cleugh H, Stafford Smith M, Battaglia M, Graham P. 2011. Climate change: science and solutions for

Australia. Collingwood: CSIRO Publishing.

Cobb SM, Saynor MJ, Eliot M, Eliot I, Hall R. 2007. Saltwater intrusion and mangrove encroachment of coastal wetlands in the Alligator Rivers Region, Northern Territory, Australia. Darwin: Environmental Research Institute of the Supervising Scientist.

- 5 Costin AB. 1954. Ecosystems of the Monaro region of NSW, with special reference to soil erosion. Sydney: Government Printer.

CSIRO, Bureau of Meteorology. 2007. Climate Change in Australia. Canberra: Department of Climate Change and Energy Efficiency http://www.climatechangeinaustralia.gov.au/technical_report.php

- Davey CM, Chamberlain DE, Newson SE, Noble DG, Johnston A. 2012. Rise of the generalists: evidence for climate driven homogenization in avian communities. *Global Ecology and Biogeography* 21: 568-578.

Davidson AM, Jennions M, Nicotra AB. 2011. Do invasive species show higher phenotypic plasticity than native species and, if so, is it adaptive? A meta-analysis. *Ecology Letters* 14: 419-431.

- Davies PM. 2010. Climate change implications for river restoration in global biodiversity hotspots. *Restoration Ecology* 18: 261-268.

DECCW. 2010. NSW Climate Impact Profile. The impacts of climate change on the biophysical environment of New South Wales. Sydney: Department of Environment, Climate Change and Water NSW.

- De Deckker P. 1986. What happened to the Australian aquatic biota 18 000 years ago? De Deckker P, Williams WD, editors. *Limnology in Australia*. Melbourne: CSIRO Publishing. p487-496.

Delaney R, Fukuda Y, Saalfeld K. 2009. Management Program for the Magpie Goose (*Anseranas semipalmata*) in the Northern Territory of Australia, 2009 – 2014. Darwin: Northern Territory Department of Natural Resources, Environment, the Arts and Sport.

Douglas MM, Bunn SE, Davies PM. 2005. River and wetland food webs in Australia's wet-dry tropics:

general principles and implications for management. *Marine and Freshwater Research* 56: 329-342.

Douglas MM, O'Connor RA. 2003. Effects of the exotic macrophyte, para grass (*Urochloa mutica*), on benthic and epiphytic macroinvertebrates of a tropical floodplain. *Freshwater Biology* 48: 962-971.

Douglas MM, O'Connor RA. 2004. Weed invasion changes fuel characteristics: Para Grass (*Urochloa mutica* (Forssk.) T.Q. Nguyen) on a tropical floodplain. *Ecological Management & Restoration* 5: 143-145.

Dukes JS, Chiariello NR, Loarie SR, Field CB. 2011. Strong response of an invasive plant species (*Centaurea solstitialis* L.) to global environmental changes. *Ecological Applications* 21: 1887-1894.

Elith J, Kearney M, Phillips S. 2010. The art of modelling range-shifting species. *Methods in Ecology and Evolution* 1: 330-342.

Ferrier S, Guisan A. 2006. Spatial modelling of biodiversity at the community level. *Journal of Applied Ecology* 43: 393-404.

Finlayson C, Lowry J, Bellio M, Nou S, Pidgeon R, Walden D, Humphrey C, Fox G. 2006. Biodiversity of the wetlands of the Kakadu Region, northern Australia. *Aquatic Sciences - Research Across Boundaries* 68: 374-399.

Fukami T, Martijn Bezemer T, Mortimer SR, van der Putten WH. 2005. Species divergence and trait convergence in experimental plant community assembly. *Ecology Letters* 8: 1283-1290.

Good R, Wright G, Whinam J, Hope G. 2010. Restoration of mires of the Australian Alps following the 2003 wildfires. Haberle S, Stevenson J, Prebble M, editors. *Altered ecologies: fire, climate and human influences on terrestrial landscapes*. Canberra: ANU Press. p353-362.

Green K, Pickering CM. 2009. The decline of snowpatches in the Snowy Mountains of Australia: importance of climate warming, variable snow and wind. *Arctic, Antarctic and Alpine Research* 41: 212-218.

Groffman PM, Bain DJ, Band LE, Belt KT, Brush GS, Grove JM, Pouyat RV, Yesilonis IC, Zipperer WC. 2003. Down by the riverside: urban riparian ecology. *Frontiers in Ecology and the Environment* 1:

315-321.

Hamilton S. 2010. Biogeochemical implications of climate change for tropical rivers and floodplains. *Hydrobiologia* 657: 19-35.

5 Harrison ET, Norris RH, Wilkinson SN. 2008. Can an indicator of river health be related to assessments from a catchment scale sediment model? *Hydrobiologia* 600: 48-64.

Hijmans RJ, Graham CH. 2006. The ability of climate envelope models to predict the effect of climate change on species distributions. *Global Change Biology* 12: 2272-2281.

10 Hobbs RJ, Arico S, Aronson J, Baron JS, Bridgewater P, Cramer VA, Epstein PR, Ewel JJ, Klink CA, Lugo AE, Norton D, Ojima D, Richardson DM, Sanderson EW, Valladares F, Vilà M, Zamora R, Zobel M. 2006. Novel ecosystems: theoretical and management aspects of the new ecological world order. *Global Ecology and Biogeography* 15: 1-7.

Hobbs RJ, Higgs E, Harris JA. 2009. Novel ecosystems: implications for conservation and restoration. *Trends in Ecology & Evolution* 24: 599-605.

15 Horton R, Herweijer C, Rosenzweig C, Liu J, Gornitz V, Ruane AC. 2008. Sea level rise projections for current generation CGCMs based on the semi-empirical method. *Geophysical Research Letters* 35: L02715.

Horwitz P, Bradshaw D, Hopper S, Davies P, Froend P, Bradshaw F. 2008. Hydrological change escalates risk of ecosystem stress in Australia's threatened biodiversity hotspot. *Journal of the Royal Society of Western Australia* 91: 1-11.

20 Hulme PE. 2005. Adapting to climate change: is there scope for ecological management in the face of a global threat? *Journal of Applied Ecology* 42: 784-794.

Hulme PE, Pyšek P, Nentwig W, Vilà M. 2009. Will threat of biological invasions unite the European Union? *Science* 324: 40-41.

IPCC. 2007. Climate Change 2007: Synthesis Report. In: Change IPoC (Ed.). Cambridge: Cambridge

University Press.

Jardine TD, Pusey BJ, Hamilton SK, Pettit NE, Davies PM, Douglas MM, Sinnamon V, Halliday IA, Bunn SE. 2012. Fish mediate high food web connectivity in the lower reaches of a tropical floodplain river. *Oecologia* 168: 829-838.

- 5 Kearney MR, Wintle BA, Porter WP. 2010. Correlative and mechanistic models of species distribution provide congruent forecasts under climate change. *Conservation Letters* 3: 203-213.

Keith DA, Holman L, Rodoreda S, Lemmon J, Bedward M. 2007. Plant functional types can predict decade-scale changes in fire-prone vegetation. *Journal of Ecology* 95: 1324-1337.

- Koca D, Smith B, Sykes M. 2006. Modelling regional climate change effects on potential natural
10 ecosystems in Sweden. *Climatic Change* 78: 381-406.

Lake PS. 2008. Drought, the “creeping disaster”: effects on aquatic ecosystems. Canberra: Land & Water Australia.

- Leahy PJ, Tibby J, Kershaw AP, Heijnis H, Kershaw JS. 2005. The impact of European settlement on Bolin Billabong, a Yarra River floodplain lake, Melbourne, Australia. *River Research and Applications*
15 21: 131-149.

Lu P, Yu Q, Liu J, Lee X. 2006. Advance of tree-flowering dates in response to urban climate change. *Agricultural and Forest Meteorology* 138: 120-131.

Lucas C, Hennessy K, Mills G, Bathols J. 2007. Bushfire weather in southeast Australia: recent trends and projected climate change impacts. Melbourne: Bushfire Cooperative Research Centre.

- 20 McCann RK, Marcot BG, Ellis R. 2006. Bayesian belief networks: applications in ecology and natural resource management. *Canadian Journal of Forest Research* 36: 3053-3062.

McDougall KL, Alexander JM, Haider S, Pauchard A, Walsh NG, Kueffer C. 2011. Alien flora of mountains: global comparisons for the development of local preventive measures against plant invasions. *Diversity and Distributions* 17: 103-111.

- McDougall KL, Morgan JW, Walsh NG, Williams RJ. 2005. Plant invasions in treeless vegetation of the Australian Alps. *Perspectives in Plant Ecology, Evolution and Systematics* 7: 159-171.
- McDougall KL, Walsh NG. 2002. The flora of Nungar Plain, a treeless sub-alpine frost hollow in Kosciuszko National Park. *Cunninghamia* 7: 601-610.
- 5 McDougall KL. 2007. Grazing and fire in two subalpine peatlands. *Australian Journal of Botany* 55: 42-57.
- Merritt DM, Scott ML, Poff NL, Auble GT, Lytle DA. 2010. Theory, methods and tools for determining environmental flows for riparian vegetation: riparian vegetation-flow response guilds. *Freshwater Biology* 55: 206-225.
- 10 Meyer JL, Paul MJ, Taulbee WK. 2005. Stream ecosystem function in urbanizing landscapes. *Journal of the North American Benthological Society* 24: 602-612.
- Morin X, Thuiller W. 2009. Comparing niche- and process-based models to reduce prediction uncertainty in species range shifts under climate change. *Ecology* 90: 1301-1313.
- Mulrennan ME, Woodroffe CD. 1998. Saltwater intrusion into the coastal plains of the Lower Mary River, Northern Territory, Australia. *Journal of Environmental Management* 54: 169-188.
- 15 Murray JV, Stokes KE, van Klinken RD. 2012. Predicting the potential distribution of a riparian invasive plant: the effects of changing climate, flood regimes and land-use patterns. *Global Change Biology* 18: 1738-1753.
- Myers N, Mittermeier RA, Mittermeier CG, de Fonseca GAB, Kent J. 2000. Biodiversity hotspots for conservation priorities. *Nature (Lond.)* 403: 853-858.
- 20 Naiman RJ, Décamps, H. 1997. The ecology of interfaces -- riparian zones. *Annual Review of Ecology and Systematics* 28:621-658.
- Naiman RJ, Décamps H, McClain ME. 2005. *Riparia: Ecology, Conservation and Management of Streamside Communities*. San Diego: Elsevier/Academic Press.

Nakicenovic N, Alcamo J, Davis G, Vries HJMd, Fenhann J, Gaffin S, Gregory K, Grubler A, Jung TY, Kram T, Michaelis ELLRL, Mori S, Morita T, Papper W, Pitcher H, Price L, Riahi K, Roehrl A, Rogner H, Sankovski A, Schlesinger M, Shukla P, Smith S, Swart R, Rooijen Sv, Victor N, Dadi Z. 2000. IPCC Special Report on Emissions Scenarios. Cambridge: Cambridge University Press.

- 5 Neilson RP, Pitelka LF, Solomon AM, Nathan RAN, Midgley GF, Fragoso JSMV, Lischke H, Thompson KEN. 2005. Forecasting regional to global plant migration in response to climate change. *BioScience* 55: 749-759.

O'Donnell J, Gallagher RV, Wilson PD, Downey PO, Hughes L, Leishman MR. 2012. Invasion hotspots for non-native plants in Australia under current and future climates. *Global Change Biology* 18: 617-
10 629.

Otto K. 2005. *Yarra a diverting history of Melbourne's Murky River*. Melbourne: The Text Publishing Company.

Palmer MA, Liermann CAR, Nilsson C, Flörke M, Alcamo J, Lake PS, Bond N. 2008. Climate change and the world's river basins: anticipating management options. *Frontiers in Ecology and the Environment*
15 6: 81-89.

Pearl J. 2009. *Causality: Models, Reasoning, and Inference*. New York, USA: Cambridge University Press.

Pettit NE, Froend RH, Davies PM. 2001. Identifying the natural flow regime and the relationship with riparian vegetation for two contrasting western Australian rivers. *Regulated Rivers: Research &*
20 *Management* 17: 201-215.

Petty AM, Werner PA, Lehmann CER, Riley JE, Banfai DS, Elliott LP. 2007. Savanna responses to feral buffalo in Kakadu National Park, Australia. *Ecological Monographs* 77: 441-463.

Poff NL, Allan JD, Bain MB, Karr JR, Prestegard KL, Richter BD, Sparks RE, Stromberg JC. 1997. The natural flow regime: a paradigm for river conservation and restoration. *Bioscience* 47: 769-784.

- Pressey RL, Cabeza M, Watts ME, Cowling RM, Wilson KA. 2007. Conservation planning in a changing world. *Trends in Ecology & Evolution* 22: 583-592.
- Pusey BJ, Warfe DM, Townsend SA, Douglas MM, Burrows D, Kennard MK, Close P. 2011. Condition, impacts and threats to aquatic biodiversity. Pusey BJ, editor. *Aquatic biodiversity in northern Australia: patterns, threats and future*. Darwin: Charles Darwin University Press.
- Richardson DM, Holmes PM, Esler KJ, Galatowitsch SM, Stromberg JC, Kirkman SP, Pyšek P, Hobbs RJ. 2007. Riparian vegetation: degradation, alien plant invasions, and restoration prospects. *Diversity and Distributions* 13: 126-139.
- Rumpff L, Duncan DH, Vesik PA, Keith DA, Wintle BA. 2011. State-and-transition modelling for Adaptive Management of native woodlands. *Biological Conservation* 144: 1224-1236.
- Sandel B, Dangremond EM. 2012. Climate change and the invasion of California by grasses. *Global Change Biology* 18: 277-289.
- Seastedt TR, Hobbs RJ, Suding KN. 2008. Management of novel ecosystems: are novel approaches required? *Frontiers in Ecology and the Environment* 6: 547-553.
- Seavy NE, Gardali T, Golet GH, Griggs FT, Howell CA, Kelsey R, Small SL, Viers JH, Weigand JF. 2009. Why climate change makes riparian restoration more important than ever: recommendations for practice and research. *Ecological Restoration* 27: 330-338.
- Stohlgren T, Pyšek P, Kartesz J, Nishino M, Pauchard A, Winter M, Pino J, Richardson D, Wilson J, Murray B, Phillips M, Ming-yang L, Celesti-Grapo L, Font X. 2011. Widespread plant species: natives versus aliens in our changing world. *Biological Invasions* 13: 1931-1944.
- Traill LW, Bradshaw CJA, Delean S, Brook BW. 2010. Wetland conservation and sustainable use under global change: a tropical Australian case study using magpie geese. *Ecography* 33: 818-825.
- Trowbridge WB. 2007. The role of stochasticity and priority effects in floodplain restoration. *Ecological Applications* 17: 1312-1324.

United Nations. 2010. World Urbanization Prospects: The 2009 Revision Population Database.

United Nations Population Division.

Villéger S, Blanchet S, Beauchard O, Oberdorff T, Brosse S. 2011. Homogenization patterns of the world's freshwater fish faunas. *Proceedings of the National Academy of Sciences* 108: 18003-18008.

- 5 Violle C, Navas M-L, Vile D, Kazakou E, Fortunel C, Hummel I, Garnier E. 2007. Let the concept of trait be functional! *Oikos* 116: 882-892.

Vörösmarty CJ, McIntyre PB, Gessner MO, Dudgeon D, Prusevich A, Green P, Glidden S, Bunn SE, Sullivan CA, Liermann CR, Davies PM. 2010. Global threats to human water security and river biodiversity. *Nature* 467: 555-561.

- 10 Walsh CJ, Roy AH, Feminella JW, Cottingham PD, Groffman PM, Morgan RP. 2005. The urban stream syndrome: current knowledge and the search for a cure. *Journal of the North American Benthological Society* 24: 706-723.

Walsh CJ, Sharpe AK, Breen PF, Sonneman JA. 2001. Effects of urbanization on streams of the Melbourne region, Victoria, Australia. I. Benthic macroinvertebrate communities. *Freshwater Biology* 46: 535-551.

Walsh CJ, Waller KA, Gehling J, Nally RM. 2007. Riverine invertebrate assemblages are degraded more by catchment urbanisation than by riparian deforestation. *Freshwater Biology* 52: 574-587.

Walther G-R, Roques A, Hulme PE, Sykes MT, Pyšek P, Kühn I, Zobel M, Bacher S, Botta-Dukát Z, Bugmann H, Czúcz B, Dauber J, Hickler T, Jarosík V, Kenis M, Klotz S, Minchin D, Moora M, Nentwig W, Ott J, Panov VE, Reineking B, Robinet C, Semchenko V, Solarz W, Thuiller W, Vilà M, Vohland K, Settele J. 2009. Alien species in a warmer world: risks and opportunities. *Trends in Ecology & Evolution* 24: 686-693.

BMT WBM. 2010. Kakadu-Vulnerability to climate change impacts. Canberra: Department of Climate Change and Energy Efficiency.

Westoby M. 1998. A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and Soil* 199: 213-227.

Williams D, Baruch Z. 2000. African grass invasion in the Americas: ecosystem consequences and the role of ecophysiology. *Biological Invasions* 2: 123-140.

- 5 Williams NSG, Hahs AK, Morgan JW. 2008. A dispersal-constrained habitat suitability model for predicting invasion of alpine vegetation. *Ecological Applications* 18: 347-359.

Winn KO, Saynor MJ, Eliot MJ, Elio I. 2006. Saltwater intrusion and morphological change at the mouth of the East Alligator River, Northern Territory. *Journal of Coastal Research*: 137-149.

Figures

Figure 1: Conceptual model of ecological responses to projected changes in sea level, storm intensity and carbon dioxide levels in tropical coastal floodplains in northern Australia.

Figure 2: Conceptual model of ecological responses to projected changes in sea level, storm

5 intensity, temperature and rainfall on riparian ecosystems in temperate south-western Australia.

Based on evidence from Davies (2010), CSIRO and Bureau of Meteorology (2007) and Nakicenovic and others (2000). Legend as in Figure 1.

Figure 3: Conceptual model of ecological responses to projected changes in temperature and rainfall

in high country riparian systems. Climate change projections for the high country include warmer

10 temperatures, less annual rainfall (but with an increase in summer) and less rainfall as snow,

reduced streamflow and loss of spring snowmelt signal, increased drought intensity and increased

fire intensity (DECCW 2010). The levels of uncertainty assume that no mitigation is in place. Legend

as in Figure 1.

Figure 4: Conceptual model of ecological responses to projected changes in temperature and rainfall

15 in urban (Melbourne) riparian systems. Climate change projections for Melbourne include warmer

temperatures (particularly in summer), decreased annual rainfall (with the strongest signal in winter

and spring), reduced streamflow, decreased relative humidity, and increased potential evaporation

and solar radiation (CSIRO and Bureau of Meteorology 2007). The levels of uncertainty assume that

no additional mitigation strategies are used, that is, system management follows the status quo.

20 Legend as in Figure 1.

Table 1 . Details of the Functional Classification Scheme Used in the Paper

Descriptors		Definition
Response to climate change		
Temperature	Stenothermic	Organisms that can only function in a narrow range of temperatures
	Eurythermic	Organisms that can function over a broad range of temperatures
Hydrology	Aquatic	Organisms that live, grow and reproduce underwater
	Semi-aquatic	Organisms that inhabit wet-dry ecotone
	Terrestrial	Organisms that inhabit dry land. Includes organisms that live on saturated soil or where the water table is below soil surface
Fire	Fire adapted	Organisms that are able to withstand fire through physiological adaptation (resistance or tolerance in plants) or behavioural responses (avoidance in animals)
	Fire sensitive	Organisms that are killed by fire
Salinity	Salt-tolerant	Organisms that can function in saline and brackish (and potentially freshwater) conditions
	Salt-intolerant	Organisms that can only function in freshwater and are very sensitive to increases in salinity
Carbon dioxide	CO ₂ sensitive	Organisms whose growth rate and physiological processes are sensitive to CO ₂ concentrations
	CO ₂ insensitive	Organisms whose growth rate and physiological processes vary little with changes in CO ₂ concentrations
Origin of biota		
Indigenous		Species that are locally native to the region of interest
Non-indigenous		Species that are not locally native; includes alien/exotic/non-native species and species that are native to Australia but have not historically occurred in the region of interest
Structural form of biota		
Plants	Trees	Perennial woody plants that usually have many secondary branches that are clear of the ground and are on a single main stem or trunk with strong apical dominance
	Shrubs	A woody plant that branches below or near the ground into several main stems, so has no clear trunk. Includes scramblers (for example, <i>Rubus</i> species) and subshrubs (for example, <i>Erica</i> species)
	Climbers	Plants that have trailing or scandent stems or runners. Includes vines

	Forbs*		Herbaceous flowering plants that are not graminoids
	Graminoids*	C3	Graminoids are monocotyledonous plants with narrow leaves growing from the base. They include grasses, sedges, rushes. C3 graminoids use a C3 pathway to capture carbon from photosynthesis and produce a 3-carbon molecule; C3 plants are adapted to cool season establishment and growth in either wet or dry environments
		C4	C4 graminoids use their additional C4 pathway to produce a 4-carbon molecule; C4 plants are better adapted to warm or hot seasonal conditions under moist or dry environments; compared to C3 plants, C4 plants have higher light requirements, higher productivity, lower moisture requirements and are more sensitive to frost
Algae			Eukaryotic, typically autotrophic, organisms that lack many of distinct the organs found in land plants (for example, roots, stems, leaves)
Animals	Invertebrates	Microinvertebrates	Organisms without a backbone that are generally less than 1-5 m in size
		Macroinvertebrates	Organisms without a backbone that are generally 5 mm or larger in size; macroinvertebrates are typically more tolerant of environmental fluctuations and are longer lived than microinvertebrates
	Vertebrates	Mammals	Air-breathing endothermic organisms characterised by the possession of hair, three middle ear bones and mammary glands that are functional in mothers with young
		Birds	Air-breathing endothermic organisms that lay eggs, are feathered, winged and bipedal
		Reptiles	Air-breathing, ectothermic tetrapods characterized by skin covered in scales and/or scutes
		Amphibians	Air-breathing ectothermic tetrapods with eggs that are not surrounded by membranes
		Fish	Gill-bearing aquatic craniate organsims that lack limbs with digits

*With some exceptions (for example, *Bambusa* species), forbs and graminoids are collectively referred to as herbs; herbs lack woody stems.

Table 2. Comparison of the Effects of Urbanization and Climate Change on Riparian Systems

Issue	Urbanization effect	Climate change effect
Disconnection of river from floodplain ecosystems (reduced lateral connectivity)	Reduced frequency of connection in urban environments through building over streams, raised river banks and walls, incised river channels and construction of stormwater drains	Reduction in rainfall and runoff reduces frequency, duration and magnitude of overbank flow
Water quality	Degradation of water quality, for example, increased pollution and contaminant concentrations due to reduced base flows, through rubbish and runoff from roads, industrial land and stormwater	Reduction in runoff and discharge reduces dilution effect
Hydrology changes – altered frequency, duration, seasonality and magnitude of river flow	<p>Increased proportion of surfaces in a catchment that are impervious increases the volume of surface runoff and the rate at which it is transferred to streams after a rainfall event (flashy hydrograph)</p> <p>Stormwater drains that bypass riparian zones increase the speed at which surface runoff reaches streams</p> <p>High runoff volumes over short periods of time (that is, from rainfall events) can incise urban stream channels which, together with impervious uplands, can lead to lower groundwater levels in riparian zone.</p>	<p>Reduced rainfall and increased evaporation reducing magnitude and altering seasonality of run-off</p> <p>Increased frequency and severity of extreme weather events, including severe flooding</p> <p>Altered surface flows</p> <p>Increased frequency and magnitude of extreme rainfall events results in increased channel erosion and altered ecosystem processes</p>
Drought	Land clearance and subsequent drying, reductions in soil carbon, increased surface albedo and compaction of soils may give rise to catchments that have increased surface runoff, but less groundwater recharge, reducing the capacity of groundwater to maintain streams during drought.	More frequent and intense droughts
Species composition and habitat	Increased fragmentation of remnant vegetation patches – reduction in	Alteration of connections between sites and species'

connectivity	<p>connectivity and impediment to dispersal of terrestrial species along river corridors</p> <p>Increased proportion of exotic species</p> <p>Habitat degradation or simplification (for example, altered patterns of nutrient and energy cycling, altered production: respiration ratio)</p>	<p>dispersal abilities, for example, faster drying of ponds due to warmer and drier summers reduces connections between wetland systems</p> <p>Prolonged dry periods lead to shift away from biota requiring wetter conditions</p> <p>Smaller patch sizes leads to loss of species richness</p> <p>Disturbances, including climate, beyond an ecosystem's natural range likely to lead to proliferation of disturbance-adapted invasive exotic species</p>
Warming	<p>Loss of vegetation (shading and evapotranspiration) and heat storage capacity of concrete/brick (heat island effect) leading to warmer local temperatures, including in stream.</p>	<p>Increased air and surface water temperatures</p>
Phenology and distribution	<p>Urbanization can facilitate an increase in the distribution and abundance of disturbance-tolerant species</p> <p>Heat island effect can influence phenology, typically by advancing spring events (Lu and others 2006)</p>	<p>Warmer and drier conditions, together with altered rainfall seasonality, influencing reproductive timing and species distributions – the direction of the effect dependent on the species (IPCC 2007)</p>

Information sourced from: Groffman and others (2003); Walsh and others (2005); Lake (2008); Seavy and others (2009); Davies (2010); also see Figure 4.