Brief Report: Children with ADHD without co-morbid autism do not have impaired motor proficiency on the Movement Assessment Battery for Children.

Authors, names and addresses:

Nicole Papadopoulos BBNS, Grad Dip (Psych): Centre for Developmental Psychiatry & Psychology. School of Psychology and Psychiatry, Monash University. Building 1, 270 Ferntree Gully Road Notting hill, Victoria, Australia, 3168

Nicole Rinehart BA(Hons), MClin. Psych., PhD, MAPS, Centre for Developmental Psychiatry & Psychology. School of Psychology and Psychiatry, Monash University. Building 1, 270 Ferntree Gully Road Notting hill, Victoria, Australia, 3168

John Bradshaw MA, PhD, DSc, Centre for Developmental Psychiatry & Psychology. School of Psychology and Psychiatry, Monash University. Building 1, 270 Ferntree Gully Road Notting hill, Victoria, Australia, 3168

Jennifer McGinley B.App.Sci (PT), Grad Dip (Neurosci), PhD. School of Physiotherapy, The University of Melbourne, and the Clinical Research Centre for Movement Disorders & Gait, Cheltenham, Parkville 3052, Australia.

This research was funded by the National Health and Medical Research Council (NHMRC) [Project grant 436609; Project grant APP1004387].

Abstract

Motor proficiency was investigated in a sample of children with Attention Deficit Hyperactivity Disorder-Combined type (ADHD-CT) without autism. Two groups of children aged between 7-14 years were recruited; children with ADHD-CT (n=16; mean age 10y, 7m [SD= 1y, 10m]) and a typically developing (n=16; mean age 10y, 6m [SD= 2y, 6m]) group. Motor proficiency was measured using the Movement Assessment Battery for Children- 2nd Edition and ADHD symptoms were measured using the Conner's Parent Rating Scale. Children with ADHD-CT without autism did not display motor difficulties on the MABC-2. Higher levels of Inattention, but not hyperactivity or impulsivity was associated with poorer motor performance. These findings provide indirect evidence that co-morbid autism may contribute to motor problems in children with ADHD.

Correspondence/reprint requests to: Nicole Papadopoulos, Centre for Developmental Psychiatry & Psychology, School of Psychology and Psychiatry, Monash University Building 1, 270 Ferntree Gully Road Notting hill, Victoria, Australia, 3168 Ph: 9902 4559

Email. Nicole.Papadopoulos@monash.edu

Key words: autism, ADHD, co-morbidity, motor performance

Authors Note

The authors would like to thank all the children and families who participated in this research study and Dr Anne-Marie Turner, Dr Katie Heathershaw, & Dr Harriet Hiscock from the Melbourne Children's Clinic.

Declaration of conflicting interests

Nicole Rinehart is an associate professor at Monash University and is also a clinical psychologist who consults at the Melbourne Children's Clinic where children with ADHD were recruited from. Assoc. Professor Rinehart is also a member of the Australian National Health and Medical Research Council Expert Working Group on ADHD Clinical Practice.

Funding

This research was funded by the National Health and Medical Research Council (NHMRC) [Project grant 436609; Project grant APP1004387].

Brief Report: Children with ADHD without co-morbid autism do not have impaired motor proficiency on the Movement Assessment Battery for Children.

Abstract

Motor proficiency was investigated in a sample of children with Attention Deficit Hyperactivity Disorder-Combined type (ADHD-CT) without autism. Accounting for the influence of co-morbid autistic symptoms in ADHD motor studies is vital given that motor impairment has been linked to social –communication symptoms in children who have co-morbid ADHD and autistic-like symptoms. Two groups of children aged between 7-14 years were recruited; children with ADHD-CT (n=16; mean age 10y, 7m [SD= 1y, 10m]) and a typically developing (n=16; mean age 10y, 6m [SD= 2y, 6m]) group. Motor proficiency was measured using the Movement Assessment Battery for Children- 2nd Edition, ADHD symptoms were measured using the Conner's Parent Rating Scale. Children with ADHD-CT who had been screened for co-morbid autism did not display motor difficulties on the MABC-2. Higher levels of inattention, but not hyperactivity or impulsivity were associated with poorer motor performance. These findings provide indirect evidence that the motor problems that children with ADHD experience may be related to co-occurring social responsiveness impairments.

Attention deficit hyperactivity disorder (ADHD) is one of the most common neurodevelopmental disorders, affecting approximately 3-7 % of school aged children (APA 2000). Neuroimaging studies reveal abnormalities in brain structure and function for ADHD, although the precise pathogenesis of this disorder still remains unclear (Konrad & Eickhoff, 2010). The DSM-IV-TR (APA, 2000) describes three subtypes of ADHD; predominantly Hyperactive-Impulsive subtype, predominantly Inattentive subtype (ADHD-PI), and a Combined type (ADHD-CT) which includes a combination of inattentive, impulsive, and hyperactive symptoms. In addition to psychosocial complications, up to 50% of children with ADHD are reported to show motor coordination ability below age expected performance (Piek et al. 1999).

Investigations using standardized motor batteries commonly report motor disturbances in individuals with ADHD (Pan et al. 2009; Brossard- Racine et al. 2011, Piek et al. 1999, Reiersen et al. 2008). Additionally, the type and degree of motor impairment has been shown to differ between subtypes, with the ADHD- PI showing significantly poorer fine motor skills, while children with ADHD-CT show more difficulty with gross motor skills (Piek et al. 1999). Piek et al. (1999) also investigated the relationship between core clinical ADHD symptoms and motor problems indicating that the severity of children's inattentive symptomatology was found to predict motor coordination difficulties. Additionally, inattention in ADHD-CT has been shown to be associated with the presence of motor problems on the Children Behaviour Checklist- Parent Report (Reiersen et al. 2008).

The association between motor problems and inattention has also been demonstrated in motor studies that have investigated the effect of medication known to enhance attention/cognitive abilities on motor performance in ADHD. A recent study conducted by Brossard-Racine et al. (2012) investigated the effect of medication (methylphenidate) on motor proficiency using the Movement Assessment Battery for Children (MABC). A subset of children (approximately 20%) with ADHD (n=9) were found to perform better on the MABC after the administration of methylphenidate over a three month period. This provides preliminary evidence that motor problems may in part be associated with core inattentive symptoms in ADHD.

The combination of autistic and ADHD symptoms present in a given individual may also underlie some of the motor problems commonly reported in children diagnosed with ADHD. It has been reported that nearly one-third of boys with population-defined ADHD-CT meet clinical cut-offs for autistic symptomatology however, this is not traditionally considered in studies of motor functioning (Reiersen et al. 2008). Of particular relevance, a large cohort study (n = 851), found that children with a combination of ADHD and parent-reported autistic symptoms exhibited greater movement difficulties measured on the Child Behaviour Checklist (Reiersen et al. 2008).

A meta-analysis by Fournier et al. (2010) has suggested that motor features are a persistent clinical feature of Autism Spectrum Disorders (ASD). Neuroimaging studies suggest an association between brain structure, core ASD symptoms and motor disturbance, for example, Qiu et al. (2009) reported

that abnormalities in the basal ganglia correlated with core ASD symptoms as measured by the Autism Diagnostic Observation Scale (ADOS) and motor impairment as measured by the Physical and Neurological Examination of Subtle Signs (PANESS). Our own work has shown that children with ASD who experience greater difficulties with social responsiveness also experience greater motor problems on the MABC (Papadopoulos et al. in press).

The question as to whether autism and ADHD should be co-morbidly diagnosed has been a topic of great interest recently (see Gargaro et al. 2011 for a review). The current diagnostic classification system precludes a dual diagnosis of autism and ADHD (APA 2000). Despite this, it is well documented that children with ASD show **ADHD behaviours** such as hyperactivity, impulsivity and attentional deficits (Goldstein & Schewebach 2004), and similarly young people with ADHD often experience social-communicative difficulties (Clark et al. 1999; Geurts et al. 2008).

A recent study conducted by Grzadzinski et al. (2010) indicated that the co-morbid autistic traits evident in a subgroup of individuals with ADHD could not be accounted for by the primary impulsive, hyperactive, and inattentive behaviours of ADHD. Thus, children with ADHD with comorbid autistic symptoms may form a subgroup with a common genetic pathway, consistent with the reported genetic overlap in autism and ADHD (Nijmeijer et al. 2010).

The aim of this study is to investigate motor difficulties, as measured by motor

proficiency scores on the Movement Assessment Battery for Children (MABC-2) in children with ADHD without autism.

It was hypothesised that there would be no difference in motor proficiency between a sample of individuals diagnosed with ADHD-CT **without autism** and a sample of Typically Developing (TD) controls. **Additionally, based on Piek et al. (1999) and Reiersen et al. (2008) findings that inattention is associated with motor problems in children with ADHD, it was further predicted that inattentive symptoms measured on the Conner's Rating Scale (CRS) would be associated with greater deficits in motor proficiency.**

Method

Participants

Sixteen boys diagnosed with ADHD-CT aged between 7 and 14 years were recruited from a private Paediatrician clinic in Melbourne. **Paediatricians referred children with a diagnosis of ADHD-CT who had been screened for autistic disorder, Asperger's disorder or pervasive developmental disorder not otherwise specified. Exclusion criteria included a co-morbid medical (e.g. tuberous sclerosis), hearing or visual, or genetic (e.g. Fragile X syndrome) disorder other than the primary diagnosis of ADHD-CT as described in Langmaid et al. (in press). No participants were excluded for having these conditions.** Additionally, a doctoral level trained graduate student interviewed parents to review the child's medical, developmental and behavioural history, to confirm that the child met criteria for ADHD-CT according to the DSM-IV-TR.

Participants were also administered the Conner's Rating Scale Parent report (Conners 2001), and further screened for a possible autism diagnosis using the Autistic Diagnostic Observation Scale (ADOS) by a researcher trained in administration, author (NP). No participants met the criteria for autistic disorder on the ADOS and thus no clinical participants were excluded from the study.

The majority of participants in the study (13/16) were on stimulant medication such as methylphenidate (Ritalin). Participants discontinued medication at least 24 hours prior to testing (Langleben et al. 2006; Leitner et al. 2007).

Sixteen typically developing boys aged between 7 and 12 years with no prior history of psychological, neurological or psychiatric diagnoses were recruited from local schools and the local community.

Intellectual functioning was measured for all participants in both groups using the Weschler Intelligence Scale for Children 4th Edition (WISC-IV) (Wechsler 2005) with the exception of two clinical participants who did not complete a cognitive assessment. **A series of Independent samples t-tests were conducted to investigate the effect of age and IQ between the two groups. There was no significant difference in age($t(30)= 0.172, p= 0.865$), Full Scale Intelligence Quotient (FSIQ)($t(28)= 1.37, p=0.183$), Verbal Comprehension Index (VCI) ($t(28)= 1.78, p = 0.087$)and Perceptual Reasoning Index (PRI) ($t(28)= 0.565, p = 0.580$)between the groups (see table 1).**

“ Insert Table 1 here”

Measures

Motor Proficiency

Motor proficiency was **measured** using an age appropriate measure of motor performance: the Movement Assessment Battery for Children (MABC-2) Henderson et al. (2007). **The MABC-2 has been found to have good reliability with Pearson's coefficients of .77, .84, .75 and .80 for manual dexterity, aiming and catching, balance and total MABC score respectively (Henderson et al., 2007). Inter-rater reliability coefficients ranging from .51-.71 have been reported for the first age band (3-6 years) (Ellinoudis et al., 2011), however, no study has yet reported inter-rater reliability for the older age bands used in this study.**

The MABC has been previously used in children with ADHD (e.g. Piek and Pitcher, 2004). It comprises of a total of eight tasks grouped into three subtests; manual dexterity, aiming and catching and balance. Percentile ranks and age-adjusted standard scores are provided for subtest scores and total impairment scores. Higher percentile ranks/scores on the MABC-2 indicate better motor proficiency. The MABC-2 was administered by a graduate student who had undergone training in the administration and scoring of the MABC-2, from a Physiotherapist. The MABC-2 was conducted in a standardized test setting at the Clinical Research Centre for Movement Disorders and Gait, Kingston Centre, Melbourne.

Severity of total ADHD symptoms (CRS-total score), severity of Inattention (CRS- Inattention Score) and severity of hyperactivity (CRS- hyperactivity score)

symptoms were measured using the Conners' Rating Scale (CRS)- Parent report. Raw scores were converted to t-scores, where a higher t-score indicates greater severity of ADHD symptoms experienced.

Data Analysis

A series of Independent sample's t-tests were conducted to assess group differences in Total MABC and its three component subtest scores. Spearman correlations were conducted on the ADHD group data ($n=16$) to investigate the association between movement impairment and Inattention, Hyperactivity and Impulsivity measured on the Conner's Rating Scale.

Results

Motor Proficiency

The motor proficiency total and subtest percentile scores of the MABC-2 are illustrated in Figure 1. There were no significant differences between the ADHD ($M=80.8$, $SD=12.5$) and TD ($M=82.0$, $SD=12.7$) groups when compared on total mean MABC-2 percentile scores ($t(30) = 0.27$, $p = .792$). Additionally there were no significant differences between the mean percentile scores for the ADHD and TD groups across the three subtests: manual dexterity ($t(30) = .44$, $p = .662$; ADHD; $M=36.8$, $SD=28.7$, TD; $M=32.8$, $SD=23.8$), throwing and catching ($t(30) = .220$, $p = .827$; ADHD; $M=64.2$, $SD=26.0$, TD; $M=66.3$, $SD=27.8$) and balance ($t(30) = .449$, $p = .657$, ADHD; $M=61.4$, $SD=28.7$, TD; $M=66.4$, $SD=34.1$).

“Insert Figure 1 here”

Clinical Correlations

There was a significant negative correlation between Inattention scores and Total MABC-2 ($n(16)$, $r_s = -0.58$, $p = 0.02$) (Higher scores on the Inattention- CRS reflect greater inattentive symptoms, while the lower percentile scores on the MABC reflect poorer motor performance). Inattention was also significantly correlated with the ball skills subtest ($n(16)$, $r_s = -0.69$, $p = 0.003$) but not the manual dexterity and balance subtest ($p > .05$, see Figures 2a & 2b). There was no significant correlation between Hyperactive- Impulsive symptoms (**$n(16)$, $r_s = -0.046$, $p = 0.867$**) and total MABC-2 score or subtest scores.

“Insert Figures 2a and 2b”

Discussion

This study aimed to investigate motor proficiency in a well defined sample of individuals diagnosed with ADHD-CT without autism. In contrast to past studies, the current study excluded autism using DSM-IV criteria in addition to the “gold standard” autistic diagnostic measure: the Autism Diagnostic Observation Scale (ADOS-G) (Lord et al. 2000). Our hypothesis, that children with ADHD-CT without co-morbid autistic symptoms would not display motor difficulties was supported. The findings that children with “ADHD only” in this sample did not have motor difficulties contrasts with previous research that commonly report motor deficits in children with

ADHD (e.g. Piek et al. 1999). These findings suggest that it may be that the co-morbid autistic symptoms in children with ADHD contribute to more impaired motor proficiency difficulties.

As predicted, inattention, but not hyperactivity or impulsivity, correlated with degree of motor proficiency deficits in children with ADHD. Higher levels of inattention were associated with greater motor impairment in ADHD. This could be interpreted in the context of a recent neuropsychological review by O'Halloran et al. (2012) who commented on parallel development of brain regions such as the cerebellum responsible for both motor and cognitive (attentional) processes.

There are several limitations to this study. The use of a standardized motor measure may have limited our ability to detect subtle motor anomalies commonly reported in individuals with ADHD. Additionally, it may be that this sample comprised of the “milder type” of ADHD, and that motor deficits in “ADHD-CT only” only become apparent in individuals who exhibit greater severity of ADHD symptoms. **Two clinical participants did not complete a cognitive assessment, however there was no evidence to suggest that these children had impaired intellectual function. It is possible that this study was underpowered and we acknowledge that these findings are preliminary and exploratory and require further replication with a larger sample size.** Future research using motor tasks which can detect the more subtle aspects of motor impairment which have previously been suggested by the literature, e.g. motor timing deficits (Buderath et al., 2008) for samples of individuals diagnosed with ADHD only, autism only and a subgroup of

individuals with autism and ADHD, will enable a closer investigation of the interplay between autistic symptoms, ADHD symptoms and motor functioning in autism and ADHD.

Conclusions

This study is the first to our knowledge to describe the motor profile of a well-defined group of individuals diagnosed with ADHD-CT without autism. Preliminary findings indicate that children with ADHD-CT who do not exhibit co-morbid autism do not have impaired motor proficiency. Inattentive symptoms were shown to correlate with overall movement impairment. These findings provide indirect evidence that co-morbid autistic symptoms that may be present in ADHD samples may contribute to motor difficulties commonly reported in ADHD (e.g. Reiersen et al. (2008) study). These findings have important clinical implications as motor symptoms in ADHD may serve as a useful “red flag” in identifying autistic symptoms.

Table 1- Participant Characteristics

	ADHD	Typically Developing Controls (TD)
No of participants	16	16
Age Mean (SD)	10yrs 7months (1 yr, 10 months)	10yrs, 6 months (2yrs 6 months)
FSIQ Mean (SD)*	97.50 (12.48)	104.88 (16.50)
VCI Mean (SD)*	94.79 (13.44)	104.88 (17.13)
PRI Mean (SD)*	102.50 (14.53)	105.69 (16.38)
CRS Total ADHD Score	62.1(9.9)	44.4(3.3)
CRS Inattention Score	59.9 (10.2)	45.4 (5.1)
CRS Hyperactivity Score	65.2 (14.1)	47.6 (5.9)

*Cognitive assessment was not completed for two of the ADHD participants

Figure 1- Movement proficiency (mean percentile ranks) measured on the Movement Assessment Battery for Children-2.

Percentile rank represents the proportion of same aged peers with a lower score (e.g. 25th percentile is interpreted as 25% of same age peers with an equivalent or lower score).

Figure 2a - A significant negative correlation between Inattention and mean total movement impairment percentile score, $n(16)$, $r_s = -0.581$, $p = 0.02$.

Figure 2b- A significant negative correlation ($n(16)$, $r_s = -0.686$, $p = 0.003$) between the ball skills subtest of the MABC-2 and Inattention.

References

- American Psychiatric Association. (2000). *Diagnostic and Statistical Manual of Mental Disorders IV-TR* (4th ed.). Washington, DC: American Psychiatric Association Press.
- Brossard-Racine, M., Shevell, M., Snider, L., Belanger, S.A., & Majnemer, A. (2012). Motor skills in children newly diagnosed with Attention Deficit Hyperactivity Disorder prior to and following treatment with stimulant medication. *Research in Developmental Disabilities*, 33, 2080-2087.
- Brossard-Racine, M., Majnemer, A., & Shevell, M. I. (2011). Exploring the neural mechanisms that underlie motor difficulties in children with attention deficit hyperactivity disorder. *Developmental Neurorehabilitation*, 14, 101-111.
- Buderath, P., Gartner, K., Frings, M., Christiansen, H., Schoch, B., Konczak, J., Timmann, D. (2009). Postural and gait performance in children with attention-deficit/hyperactivity disorder. *Gait and Posture*, 29(2), 249-254.
- Clark, T., Feehan, C., Tinline, C., & Vostanis, P. (1999). Autistic symptoms in children with attention deficit-hyperactivity disorder. *European Child and Adolescent Psychiatry*, 8, 50-55.
- Conners, C. (2001). *Conners' Rating Scales - revised (CRS-R): technical manual*. North Tonawanda, N.Y.: Multi-Health Systems Inc.
- Ellinoudis, T., Evaggelinou, C., Kourtessis, T., Konstantinidou, Z. (2011). Reliability and validity of age band 1 of the *Movement Assessment*. *Research in Developmental Disability*, 32(1), 1046-1051.
- Gargaro, B. A., Rinehart, N. J., Bradshaw, J., Tonge, B. J., & Sheppard, D. M. (2011). Autism and ADHD: How far have we come in the comorbidity debate? *Neuroscience and Biobehavioral Reviews*, 35, 1081-1088.
- Geurts, H., Grasman, R., Verte, S., Oosterlaan, J., Roeyers, H., Kammen, S., et al. (2008). Intra-individual variability in ADHD, autism spectrum disorders and Tourette's syndrome. *Neuropsychologia*, 46, 3030-3041.
- Goldstein, S., & Schwebach, A. (2004). The comorbidity of pervasive developmental disorder and attention deficit hyperactivity disorder: Results of a retrospective chart review. *Journal of Autism and Developmental Disorders*, 34, 329-339.
- Grzadzinski, R., Di Martino, A., Brandy, E., Mairene, M. A., O'Neale, M., Petkova, E., et al. (2010). Examining autistic traits in children with ADHD: Does the autism spectrum extend to ADHD? *Journal of Autism and Developmental Disorders*, 41, 1178-1191.
- Henderson, S. E., Sugden, D. A., & Barnett, A. L. (2007). *Movement Assessment Battery for Children - 2* (2 ed.). London: Pearson Assessment.
- Johnson, K., I, R., S, K., Silk, T., Barry, E., Daibhis, A., et al. (2007). Dissociation in performance of children with ADHD and high-functioning autism on a task of sustained attention. *Neuropsychologia*, 45, 2234-2245.
- Konrad, K., & Eickhoff, S. (2010). Is the ADHD brain wired differently? A review on structural and functional connectivity in Attention deficit hyperactivity disorder. *Human Brain Mapping*, 31, 904-916.
- Langleben, D. D., Monterosso, J., Elman, I., Ash, B., Krikorian, G., & Austin, G. (2006). Effect of methylphenidate on Stroop Color-Word task performance in

- children with attention deficit hyperactivity disorder. *Psychiatry Research*, 141(3), 315-320.
- Langmaid, R.A., Papadopoulos, N., Johnson, B.P., Phillips, J.G., & Rinehart, N.J. (In press 12/2011). Handwriting in children with Attention Deficit Hyperactivity Disorder. *Journal of Attention Disorders*.
- Leitner, Y., Barak, R., Giladi, N., Peretz, C., Eshel, R., Gruendlinger, L., et al. (2007). Gait in attention deficit hyperactivity disorder: Effects of methylphenidate and dual tasking. *Journal of Neurology*, 254 (10), 1330-1338.
- Lord, C., Risi, S., Lambrecht, L., Cook, E. H., Jr., Leventhal, B. L., DiLavore, P. C., et al. (2000). The autism diagnostic observation schedule-generic: a standard measure of social and communication deficits associated with the spectrum of autism. *Journal of Autism and Developmental Disorders*, 30(3), 205-223.
- Nijmeijer, J.S., Arias-Vasquez, A., Rommelse, N.J., Altink, A.E., Anney, R.J., Asherson, P et al. (2010). Identifying loci for the overlap between attention-deficit /hyperactivity disorder and autism spectrum disorder using a genome-wide QLT linkage approach. *Journal of the American Academy of Child and Adolescent Psychiatry*, 49 (7), 675-685.
- O'Halloran, C.J., Kinsella, G.J., & Storey, E. (2012). The cerebellum and neuropsychological functioning: A critical review. *Journal of clinical and experimental neuropsychology*, 34(1), 35-56.
- Pan, C., Tsai, C., & Chu, C. (2009). Fundamental movement skills in children diagnosed with Autism Spectrum Disorders and Attention Deficit Hyperactivity Disorder. *Journal of Autism and Developmental Disorders*, 39, 1694-1705.
- Piek, J., Pitcher, T., & Hay, D. (1999). Motor coordination and kinaesthesia in boys with attention deficit-hyperactivity disorder. *Developmental Medicine and Child Neurology*, 41, 195-165.
- Qiu, A., Adler, M., Crocetti, D., Miller, M., & Mostofsky, S. (2010). Basal ganglia shapes predict social, communication, and motor dysfunctions in boys with autism spectrum disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*, 49(6), 539-551.
- Reiersen, A., Constantino, M., & Todd, R. (2008). Co-occurrence of motor problems and autistic symptoms in Attention-Deficit/Hyperactivity Disorder. *Journal of American Academy of Child Adolescent Psychiatry*, 47 (6), 662-672.

Figure 1- Movement proficiency (mean percentile ranks) measured on the Movement Assessment Battery for Children-2.

Percentile rank represents the proportion of same aged peers with a lower score (e.g. 25th percentile is interpreted as 25% of same age peers with an equivalent or lower score).

Figure 2a - A significant negative correlation between Inattention and mean total movement impairment percentile score, $n(16)$, $r_s = -0.581$, $p = 0.02$.

Figure 2b- A significant negative correlation ($n(16)$, $r_s = -0.686$, $p = 0.003$) between the ball skills subtest of the MABC-2 and Inattention.

Table 1- Participant Characteristics

	ADHD	Typically Developing Controls (TD)
No of participants	16	16
Age Mean (SD)	10yrs 7months (1 yr, 10 months)	10yrs, 6 months (2yrs 6 months)
FSIQ Mean (SD)*	97.50 (12.48)	104.88 (16.50)
VCI Mean (SD)*	94.79 (13.44)	104.88 (17.13)
PRI Mean (SD)*	102.50 (14.53)	105.69 (16.38)
CRS Total ADHD Score	62.1(9.9)	44.4(3.3)
CRS Inattention Score	59.9 (10.2)	45.4 (5.1)

CRS Hyperactivity Score	65.2 (14.1)	47.6 (5.9)
--------------------------------	-------------	------------

*Cognitive assessment was not completed for two of the ADHD participants