1 **Growth races in The Mallee: Height growth in woody plants examined**
2 **with a trait-based model**

3

4 Freya M Thomas^{1*} and Peter A Vesk¹

5

6

7 ¹School of BioSciences, ARC Centre of Excellence for Environmental Decisions,
8 The University of Melbourne, Victoria 3010, Australia

9

10 * Corresponding author: FM. Thomas, e: freyamthomas@gmail.com,

11

12

13

14

15

16

17

18

19

20

21

22

Abstract

1. Plant height and growth are fundamental to the understanding of species ecological strategies, to the description and prediction of ecosystem dynamics and to vegetation management, such as plant species' fire responses. However, a convenient way to characterise the height growth strategies for multiple species has been elusive.
 2. We examine the height growth trajectories in 18 woody plant species in a light-saturated, fire-prone, semi-arid environment as well as the influence of functional traits on those trajectories. We test trait-growth relationships by examining the influence of specific leaf area, woody density, seed size and leaf nitrogen content on three aspects of plant growth; maximum relative growth rate, age at maximum growth and asymptotic height.
 3. Woody plant species in the semi-arid mallee exhibit fast growth trajectories. Small seeded species were likely to be the fastest to reach maximum height, while large-seeded species with high leaf nitrogen were likely the slowest. Tall species had low stem densities and tended to have low specific leaf area.
- Synthesis:* We modeled plant growth using a hierarchical multi-species model that formally incorporates plant functional traits as species-level predictors of growth, which provides a method for predicting species height growth strategies as a function of their traits. We extend this approach by using the modeled relationships from our trait-growth model to predict: growth trajectories of species with limited data; real species with only trait data and; hypothetical species based only on trait coordination. We hope this highlights the potential to use trait

46 information for ecological inference and to generate predictions that could be
47 used for management.

48

49 Keywords: height-growth, functional traits, mallee, prediction, woody plants

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

Introduction

Plants compete for sunlight. The tallest plant in a community will have the greatest access to light and consequently will experience a more productive environment relative to shorter plants (Givnish 1988; Monsi & Saeki 2005; Craine & Dybzinski 2013).

Competition for light can be thought of as the partitioning of height dominance and is divided along two gradients. The vertical gradient reflects direct competition for light between individuals, whereby plants can gain dominance of available light by growing the fastest and tallest. By contrast, temporal gradients reflect light competition through time, whereby individuals may dominate time in the sun for some period before being overtopped (Kobe 1999; Falster & Westoby 2003; 2005b). Light competition can be studied by measuring plant height through time, which can also provide measures of vegetation structure and change. Plant height is also a commonly used plant functional trait associated with competitive vigour, whole plant fecundity, and is an important indicator of community disturbance intervals and environmental tolerances (Westoby *et al.* 2002; Cornellissen *et al.* 2003). In fire-prone ecosystems, measurements of plant height are often used to reflect vegetation structure and habitat redevelopment after disturbance. Height can be used as an indicator of habitat provision and tolerable fire intervals (Haslem *et al.* 2010; Muir, Vesk & Hepworth 2014) although quantitative estimates of growth trajectories for this purpose remain scarce (but see Muir, Vesk & Hepworth 2014).

Despite the importance of plant height to these various branches of ecology, a general model of cross species variation in height growth is lacking. Developing functional trait-

92 based models of growth has become an important research goal for plant ecologists in an
93 effort to generalize trait-growth relationships. Traits have been linked to species growth
94 rates in several field studies (Kobe 1999; Reich *et al.* 1999; Falster & Westoby 2005a;
95 Falster & Westoby 2005b; King *et al.* 2005; Poorter *et al.* 2008; Adler *et al.* 2014), but
96 the influence traits have on aspects of plant growth can seem inconsistent or ambiguous.
97 Seed mass has previously been variously reported as a positive predictor of seedling
98 growth (Poorter *et al.* 2008), thought unlikely to directly affect growth of large plants
99 (Ruger *et al.* 2012), and been negatively related to individual growth and fecundity
100 (Adler *et al.* 2014). Wood density has previously been highlighted as one of the most
101 important traits influencing adult woody plant growth (King *et al.* 2005; Falster &
102 Westoby 2005a; Poorter *et al.* 2008; Ruger *et al.* 2012; Reich *et al.* 2014) but also found
103 to be a poor predictor of inherent growth rates (Herault *et al.* 2010). Leaf traits appear to
104 have particularly unclear implications for growth; some studies reporting strong support
105 (Kitajima 1994; Falster & Westoby 2005a; Sterk, Poorter & Schieving 2006; Reich *et al.*
106 2014; Kunstler *et al.* 2016), whilst others report poor or ambiguous relationships (Poorter
107 *et al.* 2008; Paine *et al.* 2015) and assert no justification for using leaf economics traits to
108 describe growth rates (Ruger *et al.* 2012). Leaf nitrogen has previously been related to
109 growth, where species with fast growth rates tending to have high leaf nitrogen (Poorter,
110 Remkes & Lambers 1990).

111

112 Trait-based growth research typically takes advantage of long term monitoring data sets
113 from well-sampled, dense and productive forests, such as temperate and tropical forests,
114 where access to light represents a major resource gradient (Thomas 1996; Kobe 1999;

King *et al.* 2005; Condit *et al.* 2006; Sterck *et al.* 2006; Poorter *et al.* 2008; Herault *et al.* 2011; Ruger *et al.* 2012; Prior & Bowman 2014). The bias of trait-growth rate studies to mesic systems is a bias towards systems with high light competition and frequently dominated by tall, non- or weakly-resprouting species. Vegetation without a strong vertical light profile may show different growth strategies in response to contrasting resource gradients. Prior & Bowman (2014) found that asymmetric-size competition for light was greater at high productivity sites. Areas of relatively scarce resources, resulting in low productivity, or frequent disturbance are often understudied from a functional trait-based perspective. This impedes a truly general understanding of height growth strategies.

In the Mallee region of south-eastern Australia, fire responses of broad vegetation associations and growth forms have been well described (Bradstock & Gill 1993; White 2006; Giljohann *et al.* 2015), and despite many papers on mallee eucalypt demography (Wellington 1984; Wellington & Noble 1985a; Noble 1985; Wellington & Noble 1985b; Clarke *et al.* 2010), and studies focused on *Callitris* sp. (Bradstock, Bedward & Cohn 2006), and *Triodia* sp. (Rice & Westoby 1999, Giljohann *et al.* 2017), quantitative data on individual growth responses for many of the common plant species remain limited (Pausas *et al.* 2004). We know little about the coexistence and height-growth trajectories of species in the Mallee region.

We use data from this high-light, low precipitation and low productivity region of South Eastern Australia which is on the border of temperate and semi-arid fire-regime (Murphy

et al. 2013) to complement the growing literature on trait-based growth models. This ecosystem differs substantially from tropical forest systems, where previous trait-growth work has been published, with an estimated net primary productivity of Mallee ecosystems in Australia of $0.9 \text{ t C ha}^{-1} \text{ year}^{-1}$ within a gradient of productivity in Australia ranging from Tussock Grasslands ($0.3 \text{ t C ha}^{-1} \text{ year}^{-1}$) to Eucalypt Woodlands ($2.2 \text{ t C ha}^{-1} \text{ year}^{-1}$) and rainforest ($7.9 \text{ t C ha}^{-1} \text{ year}^{-1}$) (Murphy *et al.* 2013). We model plant height using a hierarchical approach, where traits are incorporated into our model as linear predictors of growth parameters. Relating growth characteristics to functional traits in a flexible yet formal manner such as this not only increases our understanding of trait-growth rate relationships but also aids our ability to draw predictive inferences from them, particularly for species for which we may have few data. We believe this is an important research endeavor intrinsically, but could also have consequences for decisions about many plant species but where managers only have limited data for few species. An example of this is plant growth and life history trajectories in fire-prone environments.

We test trait-growth relationships by examining the influence of specific leaf area, woody density, seed size and leaf nitrogen content on three aspects of plant growth; maximum relative growth rate, age at maximum growth and asymptotic height. We do this in a light-saturated and disturbance-prone environment with woody semi-arid plant species, to test the theory of an ‘arms race for light’ outside of low-light, mesic forests. Finally we demonstrate how formally incorporating functional traits into models can aid in predicting growth strategies of plant species.

Methods

The study area is in Murray Sunset National Park, Victoria (34.7683° S, 141.8542° E), a large conservation area within the semi-arid Murray Mallee region of South-Eastern Australia. This region of Victoria has hot dry summers and cold winters. Mean annual rainfall is 290 mm, falling mostly in the cooler months from May-November, and mean maximum daily temperatures ranging from 15.4 °C in July to 32.3 °C in January (BOM). Two main soil formations occur in this area, The Lowan Sands and The Woorinen Formation, which differ in colour, sand grain size and grain surface features driven largely by differing clay content (Pell, Chivas and Williams 2001). The landscape is dominated by dune/swale systems, which reflect subtle but important variation in soil characteristics and moisture availabilities (White 2006).

The vegetation and fire histories of the Murray Mallee region of South-Eastern Australia have previously been comprehensively mapped (Haslem *et al.* 2012; Aitavale *et al.* 2013). The most common vegetation type of the region is ‘tree mallee’ which is characterised by small multi-stemmed eucalypts (‘mallees’) with a shrubby and grassy understorey (Haslem *et al.* 2012). Within this broad vegetation type, three distinct classes occur; *Triodia* mallee, Chenopod/Shrubby mallee and Heathy Mallee (White 2006; Haslem *et al.* 2012). Large wildfires are a prominent disturbance in this region with 40% of tree mallee vegetation in the region being burnt in the last 40 years, however, only a small percentage (<3%) has burnt more than once during this period (Aitavale *et al.* 2013).

184 This study took advantage of fire histories within Murray Sunset National Park (Avitable
185 *et al.* 2013) to conduct a chronosequence study of plant species growth over time (Falster
186 & Westoby 2005a; Muir, Vesk & Hepworth 2014). Eleven areas varying in time-since-
187 fire (1,2,4,8,13,15, 26, 33, 36,46,86 years), were selected within a relatively small region
188 of Murray Sunset National Park, sites encompassed a range of post-fire ages whilst
189 restricting the amount of edaphic variation, and all areas were within *Triodia* Mallee.
190 Between three and five different patches within a ‘time-since-fire’ area were surveyed,
191 totaling 50 surveys across 11 areas. Eighteen woody plant species were selected that had
192 at least three individuals across each of the 50 survey sites, a total of 1250 individual
193 plants were sampled.

194

195 *Size Measurements*

196

197 We measured heights of at least five (and up to 10) individuals per species in each time-
198 since-fire area. Many species in this region resprout after fire, but for species with post-
199 fire seed regeneration, separate measurements were made on seedling and resprouting
200 individuals. It is easy to determine seeded individuals from resprouting individuals when
201 plants are young, but as they age it becomes increasingly difficult to have confidence of
202 whether an individual seeded or resprouted. Additionally, we thought it likely that
203 resprouters would generally outpace reseeders due to their underground resource stores,
204 and so modeling both functional types together might conflate growth rates based on
205 functional traits and inherently different growth mechanisms (Falster & Westoby, 2005a).
206 Due to this, *Eucalyptus oleosa*, *Eucalyptus gracilis* and *Melaleuca lanceolata* (the

resprouting species) were left out of the trait-based multi-species model, and their growth was modeled separately to represent the resprouting species. Canopy length, canopy width and basal diameter of the biggest stem were also measured for each individual of each species; we focussed our analysis on height-growth to complement the growing literature of height-growth, mainly from more temperate ecosystems. Regressions showing the relationships between these alternative size variables to canopy height are in the appendix (Appendix Figs S2, S3 and S4).

Measuring plant traits

Guidance on sample sizes for measuring plant traits were followed from Cornelissen *et al.* (2003), and we took five samples (ie five leaves or five stem sections) from each of five individual plants for each species. Specific leaf area (SLA; mg mm^{-2}), the one-sided leaf surface area of fresh leaves divided by the mass of leaves oven-dried at 65°C for 48 h was measured for at least five leaves from each of five individuals for each species. Leaf nitrogen concentrations (N_{mass} ; %) were calculated on five fully expanded leaves taken from well-lit positions on each of five individuals per species. Leaves from each species were then pooled and finely ground for nitrogen analysis. Total nitrogen concentration based on mass (%) was measured using complete combustion gas chromatography performed by The Surface and Chemical Analysis Network at The University of Melbourne. Stem tissue density (dry mass per unit fresh volume; mg mm^{-3}) was measured using 40-60 mm long, stem segments from at least five individuals per species. Samples were collected in the field and refrigerated before processing as soon as feasible.

Stem tissue density was then determined following Archimedes principle using the protocol from Cornellissen *et al.* (2003). Seed mass (mg) was measured for as many species as possible with field-collected seed, oven-dried and weighed. Seed mass data was also supplemented from the literature and a global seed mass database (Moles & Westoby 2003). See Table 1 in Appendix for list of all species and their mean trait values.

A hierarchical, trait-based nonlinear growth model

We modeled the growth of species using a hierarchical multi-species non-linear growth model, which formally incorporates plant traits as species-level predictors of growth. Our chosen model has biologically interpretable parameters; the maximum height achieved, the maximum relative growth rate (maximum RGR), which is measured in units of cm/cm/yr (Atwell 1999) and the time at which this maximum growth occurs, measured in years.

In deciding which functional traits to include in our models we drew on literature to generate hypotheses of trait effects on growth. Specifically that wood density, seed mass and leaf traits would be influential traits affecting growth trajectories (Falster & Westoby 2005a; Reich *et al.* 2014), whereby seed mass should be most influential on initial growth (Moles & Westoby 2006), stem density of achievable height (Falster & Westoby 2005a; Falster & Westoby 2005b) and leaf traits throughout the whole growth process (Sterck, Poorter & Schieving 2006; Reich *et al.* 2014), leaf nitrogen being important for the pace and the timing of maximum growth (Poorter, Remkes & Lambers 1990). Our final model

had stem density and SLA influencing maximum height, nitrogen leaf content and seed mass on age at maximum growth and seed mass on maximum relative growth rate. These enter into our multi-species model in a way similar to that of Pollock *et al.* (2011).

We chose to use the three-parameter sigmoidal model form due to the biological interpretable parameters and its use in previous studies (Thomas 1996; Falster & Westoby 2005a) and the increased information due to a parameter specifying age at maximum relative growth rate. As well as biologically interpretable parameters, our model choice sought to adequately represent the growth of all species modeled. One consequence of modeling multiple species and growth forms together is that some species are better fit than others. We display some model fits to raw data of our chosen model (Fig. 1a-f). We compared our three-parameter sigmoidal model to four other possible growth models; two two-parameter models (Power and Exponential) and one other three-parameter model (Logistic), plus our chosen Hillslope model with log time (Appendix Table S2). We used Bayesian inference employing Markov chain Monte Carlo (MCMC) methods with the open-source software package JAGS version 3.3.0 (Plummer 2003) run via the statistical software environment R version 2.15.2 (R Development Core Team, 2015) with the package R2jags (Su & Yajima 2015). Please see the Supporting Information for a more detailed model description.

Predictions

We used modeled relationships between growth parameter and traits to make predictions for real species for which we have no growth data but we do have trait data and also hypothetical species based on trait information only. We did this by including our real

species and our ‘hypothetical species’ into our dataset, with missing height values, but with trait values included for them in the dataset. In a Bayesian framework, these missing data values are estimated with predictive distributions. For our real species we used their measured trait data collected from the field. For our hypothetical species, we use extreme trait values that lay with two standard deviations of those observed in our trait data set. We present a tall, fast and early growing species (low stem density 0.65 mg mm^{-3} , low SLA 2.55 mg mm^{-2} , small seed mass 0.430 mg and low leaf nitrogen content 0.97%); a tall, late and slower growing species (low stem density 0.65 mg mm^{-3} , low SLA 2.55 mg mm^{-2} , heavy seed mass 100 mg , high leaf nitrogen 4.67%); a short, late and slow growing species (high stem density 0.87 mg mm^{-3} , high SLA 6.53 mg mm^{-2} , heavy seed mass 21.86 mg , high leaf nitrogen 2.91%); and a short, early and fast growing species (high stem density 0.87 mg mm^{-3} , high SLA 6.53 mg mm^{-2} , small seed mass 0.350 mg and low leaf nitrogen 0.97%).

Results

Growth trajectories

Maximum height of woody species in the Mallee ranged between short woody shrubs such as *Halgania cyanea* (Boraginaceae), reaching an average maximum height of approximately 40 cm and the tallest species *Codonocarpus cotinifolius*, which reached average maximum heights of approximately 5 m . The four next tallest woody species, which dominate the overstorey canopy are *Callitris verrucosa* (321 cm), *Eucalyptus*

gracilis (258 cm), *Eucalyptus oleosa* (268 cm) and *Melaleuca lanceolata* (209 cm) (Fig. 2). The seed-derived individuals of the resprouter species have much slower relative growth rates, achieve these later and have lower heights compared to their resprouting individuals (Fig. 2a-b). The species that achieved the maximum growth earliest had the fastest maximum relative growth rates and that reached the greatest eventual heights were *C. verrucosa* and *C. cotinifolius* and then the resprouting species (Fig. 2). Of the seed-derived species data, one clear trade-off existed whereby high relative growth rates (parameter b) were negatively correlated (Pearson's correlation coefficient of -0.73) with age at maximum growth rate (parameter c), so species that grew relatively quickly always reached their maximum growth rate early (Appendix Fig. S3).

Growth variation among species

All species modeled had height-growth characterized by maximum relative growth rates of between 3 cm/cm/yr and 7 cm/cm/yr, onset of maximum growth occurring between one and two years following disturbance, and asymptotic heights of between 40 cm and 500 cm (Figs 2 and 3). The fastest growing species were the two resprouting eucalypts, which both had maximum relative growth rates of 6.7 cm/cm/yr, and these species also reached their maximum relative growth rate the earliest, at 1.2 years of age. Resprouting plants in the Mallee accelerated faster to achieve higher maximum growth and higher maximum heights compared to seed-derived conspecifics and to other non-resprouting species (Fig. 3). There was a clear difference between the growth rates and the age at maximum growth between the seed-derived plants of resprouting eucalypt species - *E.*

gracilis and *E. oleosa* - and their resprouting counterparts. Resprouting eucalypt individuals had nearly two-fold higher maximum growth rates (6.98 and 6.91 cm/cm/yr respectively for resprouters and 3.56 and 3.66 cm/cm/yr for seed derived cohorts) and achieved these by half the age of the seed derived individuals (2.21 and 2.01 years respectively for resprouters and 1.11 and 1.11 years for seed derived cohorts).

The tallest species, *C. cotinifolius*, had one of the fastest growth rates, reaching its maximum height of 468 cm at 1.8 yrs at a rate of 4.5 cm/cm/yr (Fig. 3). This species unusual relative to the other species in our dataset as it is known as a polycarpic fire ephemeral and characteristically has relatively short life spans and very rapid growth (Pate et al 1985). The second tallest species, *Callitris verrucosa* (Cupressaceae) had a below average growth rate. All other species had mean asymptotic heights between 40 cm and 3.2 m. The next tallest suite of species were all resprouters, the two eucalypt species and *Melaleuca lanceolata* (Myrtaceae). The slowest growing species was *Beyeria opaca* (Euphorbiaceae) with a maximum growth rate of 3.2 cm/cm/yr. In these data, there appears to be little evidence for a clear succession of leaders. For example, both *E. gracilis* (Myrtaceae) and *E. oleosa* (Myrtaceae) were always the tallest species at a site, apart from *C. cotinifolius* at two years. In the Mallee, seed-derived individuals of short species could reach their maximum growth either early or late. Species that grew fast could achieve both tall and short heights. There were, however, no tall species that had low maximum growth rates (Fig. 3, Fig. S1 Appendix).

The greatest variation between species was in the parameter asymptotic height (Standard Deviation = 0.494). There was relatively little variation between species in age at

maximum relative growth rate (SD =0.13) and the least variation between species was for maximum relative growth rate (SD =0.02).

Trait effects on growth parameters

Leaf, seed and stem traits all explained some variation in sub-models for the three growth parameters (Fig. 4a, see Table S2 Appendix for further details). Effect sizes were largest for asymptotic height, which is the parameter that varied most between species. Smaller effect sizes were found for the other two parameters, reflecting the smaller variation to explain. The largest single effect size was for stem density on maximum height. Seed mass had smaller effects and those of leaf traits were of intermediate size, but highly uncertain.

Species with lighter seeds achieved faster relative growth rates and did so earlier (Fig 4a, b, d). Species with high leaf Nitrogen content and heavy seeds were likely to be the slowest to reach maximum growth (Fig 4c, d). In general, species with low stem densities and low specific leaf area reach greater asymptotic heights (Fig 4 e, f).

Codonocarpus cotinifolius appeared an outlier, having high SLA and yet achieved a tall height, but it has an unusually low stem density, nearly 4 standard deviations below the mean value of this community. Species traits explained similar amounts of variance in the parameters; maximum growth rate has $R^2 = 0.65$, age at maximum growth $R^2 = 0.66$ and maximum asymptotic height $R^2 = 0.59$. Stem density and specific leaf area varied the most, with stem density having the largest and strongest effect size. Seed mass and

nitrogen content both had little variation around the mean parameter value but had relatively small effect sizes (Fig. 4a).

Predictions based on traits

We used the modeled relationships from our hierarchical trait-growth model to predict the growth trajectory of two real species (*Dodonaea viscosa* and *Myoporum platycarpum*) for which we only have field collected trait data as well as four hypothetical species, which represent different regions of ‘trait space’ from within our dataset (Fig. 5). These scenarios mostly differ in maximum height within the first three years and these height values commonly range between 50 cm and 250 cm.

Discussion

Mallee growth trajectories

The theoretical arms race for light suggests that two prominent gradients of competition for light exist; a vertical gradient where species compete at one point in time and a temporal gradient, where over time, each species will have some dominating time in the sun (Falster & Westoby 2003). Falster & Westoby (2005) and Muir (pers. comms) found distinct periods of prominence across species in forest systems where shorter species temporarily out-pace taller species, but are subsequently overgrown, allowing a temporal partitioning of dominance between species after disturbance. Our results suggest that this does not occur in Mallee, and all species in this system have relatively fast growth rates and trajectories compared to growth of species in other ecosystems such as open

coastal woodland (Falster & Westoby 2005) and damp forest (Muir *et al.* 2014) (Figs 2 and 3). All species in the Mallee reached an average asymptotic height after 5 years and there were no distinct prolonged overlapping series of species within the canopy (Fig. 2). Light is abundant in this ecosystem and direct competition for light appears to be less pronounced in the temporal species dynamics. In fact, two of the tallest species, *Eucalyptus gracilis* (Myrtaceae) and *Eucalyptus oleosa* (Myrtaceae), reach their period of maximum growth the earliest, at 1.2 yrs and have the fastest maximum relative growth rates (Fig. 3), which allows these species to share competitive dominance in the upperstorey. Mallee eucalypts resprout from lignotubers and have multi-stemmed growth forms (Noble 1985; White 2006). Our results suggest these attributes contribute to these species' persistence in this environment, as they resprout readily after fire and maintain their height dominance over all other species apart from the short lived *Codonocarpus cotinifolius* (Gyrostemonaceae) and the slower growing *Callitris verrucosa* (Cupressaceae) (Figs 2 and 3). When growth trajectories of the seed-derived individuals of the resprouting species were compared to their resprouting counterparts, the resprouting individuals achieved taller heights, younger ages of maximum growth and faster relative growth (Fig. 3), reflecting well appreciated principles of resprouting species being strong competitors due to large underground energy reserves stored below ground in lignotubers (Bellingham & Sparrow 2000; Pausas & Keeley 2014). These resprouting cohorts also appear to fit the description of 'super-species', which grow fast, achieve tall heights and live a long time (Huston & Smith 1987). However, seeding species also demonstrate fast growth rates, which may reflect an adequate share, or less competition for post fire resources, light and space, for both resprouters and non-

410 resprouters in this system. But the question remains, if light is not limiting, then why
411 grow so fast?
412
413 Falster & Westoby (2005) suggest there may be strong selection for traits that maximise
414 height growth rate, and thereby light acquisition, in vegetation undergoing frequent
415 disturbance. When light is not a limiting resource, fast growth may not be a race for light
416 but instead reflect an ability to start either reproducing or storing resources quickly in the
417 face of unpredictable disturbance or environmental fluctuations such as drought
418 (Bellingham & Sparrow 2000). Our data contains both resprouters and non-resprouters
419 and both syndromes have the ability for rapid height growth; resprouters have persistent
420 underground resources that facilitate high competitive ability in fertile environments and
421 non-resprouters have rapid seedling growth and early emergence postfire (Pausas &
422 Keeley 2014). The trait composition in our study site may reflect an ‘intermediate’
423 position in regards to fire frequency, productivity and rainfall, as resprouters dominate
424 the canopy and non-resprouters dominate the understory woody vegetation and all
425 species demonstrate rapid post-fire height-growth (Fig. 2). Co-existence of these two
426 strategies is expected when there has been long-term variation in environmental
427 conditions or patchy disturbance that may differentially benefit the survival of seedlings
428 or adults, for example long terms trends in fluctuating water availability (Bellingham &
429 Sparrow 2000; Pausas & Keeley 2014). One of the most important considerations for an
430 individual plant is that it is able to re-occupy a site and take advantage of abundant but
431 short pulses of resources before the next disturbance event (Bellingham & Sparrow 2000)
432 and if there is a time constraint on growth, rates of growth are often fast (Arendt 1997).

Our results suggest that both the resprouters and the non-resprouters in the Mallee region are capable of achieving rapid height-growth.

Trait relationships with growth parameters

In theory, the rate of individual plant growth and when this growth occurs directly influences species position and length of time at a competitively dominant position or ‘at the top’, where it has the access to the most light (Huston & Smith 1987; Loehle 2000; Falster & Westoby 2003). We found trends suggesting that smaller seeded species are likely to grow faster per year and reach their period of maximum growth rate earlier than larger seeded species. Species with low leaf nitrogen are also likely to reach a period of fastest annual growth before others. Species with low stem densities and low specific leaf area will generally grow taller than species with high stem densities and high SLA. Our results seem consistent with some literature, for example Adler *et al.* (2014) found that species with low wood density, small seeds, high SLA and rapid leaf turnover rates (indicative of resource exploitation as a high priority) are expected to grow faster. This suite of traits leading to fast growth is indicative of plants prioritising short-term gains over long-term benefits and are thought to possess a higher capacity to change their growth rates to changing resource conditions (Adler *et al.* 2014). This may be particular relevant to species in resource-poor, disturbance-prone or areas with pronounced environmental variation, such as mallee vegetation, where resources (ie rainfall) and disturbance (ie fire) fluctuate.

Predictions based on trait-relationships

The ability to move from descriptive to predictive science is a goal of much trait-based research (Ruger *et al.* 2012; Adler *et al.* 2014). The formal incorporation of traits into our hierarchical and biologically interpretable multi-species model, allows us to make predictions of growth trajectories based on trait values and hence estimate the growth parameters for species we have few data for but can access some easily measurable traits; either from existing trait databases, or with a targeted field collection. For example, we can predict a growth trajectory for a species with low stem density, low SLA and small seeds and demonstrate that this species is likely to achieve a greater height compared to other species and it will reach its maximum growth rate earlier. We can also use only trait information to predict the growth curves of species which may occur locally, but for which we do not have growth data (Fig. 5). We were able to conduct this type of prediction because we chose to use a semi-mechanistic and hierarchical model that allows overall trends in growth to be consistent between species with modeled variation, but also specific trait effects to have a variable impact on each of our growth parameters. An important future research goal is to determine how robust trait based predictions are across different species, between different sites. Additionally, understanding the differences in growth strategies between different light environments and how these change through time across ecosystems would be a fascinating area of comparative ecology.

This study is one of a handful to examine how plant functional traits influence growth

trajectories and complements existing literature by focusing on a high light, disturbance-prone and low productivity region. We built a hierarchical multi-species model of plant growth with biologically interpretable parameters; maximum growth rate, age at maximum growth rate and asymptotic height achieved. Including functional traits as predictors of these growth parameters allows us to formally capture the role of seed size, stem density, leaf nitrogen content and specific leaf area in species space and timing of growth and their competitive dominance. There are multiple ways this approach may be extended, such as specifically incorporating environmental variables and intra-specific trait information. We believe this is a good first start in developing relatively simple models that can be used to describe multi-species growth trajectories, but also produce testable predictions about growth trajectories based on easy to collect functional data.

Acknowledgements

We thank Brett Murphy, James Camac and Chris Jones for thoughtful comments. We also thank three anonymous reviewers for thoughtful and constructive reviews. We thank Jian Yen for analytical discussions. FMT is supported by an Australian Postgraduate Award, a Holsworth Wildlife Research Scholarship and top up from Australian Research Council Centre of Excellence in Environmental Decisions (CEED). PV is supported by CEED. Author contributions: FMT and PAV formulated idea and developed methodology. FMT performed fieldwork, statistical analysis and wrote the first draft of the manuscript with both contributing to its development.

References

Adler, P. B., Salguero-Gomez, R., Compagnoni, A., Hsu, J.S., Ray-Mukherjee, J., Mbeau-Ache, C. & Franco, M. (2014) Functional traits explain variation in plant life history strategies. *Proceedings of the National Academy of Science*, **111**(27), 740 – 745.

Arendt, J. D. (1997) Adaptive intrinsic growth rates: an integration across taxa. *The Quarterly Review of Biology*, **72**(2), 149 – 177.

Atwell, B. (1999) Growth analysis: a quantitative approach. *Plants in action*. (eds. B. Atwell, P. Kriedemann, & C. Turnbull), pp. 188. Macmillan Education, Australia.

Avitabile, S. C., Callister, K.E., Kelly, L.T., Haslem, A., Fraser, L., Nimmo, D.G., *et al.* (2013) Systematic fire mapping is critical for fire ecology, planning and management: A case study in the semi-arid Murray Mallee, south-eastern Australia. *Landscape and Urban Planning*, **117**, 81–91.

Bellingham, P. J. & Sparrow, A. D. (2000) Resprouting as a life history strategy in woody plant communities. *Oikos*, **89**(2), 409–416.

Bolker, B. M., Brooks, M.E., Clark, C.J., Geange, S.W., Poulsen, J.R., Stevens, M.H.H., & White, J.S. (2009) Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in Ecology and Evolution*, **24**(3), 127–135.

Bradstock, R.A., & Gill, A.M. (1993) Fire in semi-arid, Mallee shrublands: size of flames from discrete fuel arrays and their role in the spread of fire. *International Journal of Wildland Fire*, **3**(1), 3 – 12.

Bradstock, R.A., Bedward, M. & Cohn, J.S. (2006) The modeled effects of differing fire management strategies on the conifer *Callitris verrucosa* within semi-arid mallee vegetation in Australia. *Journal of Applied Ecology*, **43**, 281-292.

Clarke, J.S. (2003) Uncertainty and variability in demography and population growth: a hierarchical approach. *Ecology*, **84**(6), 1370 – 1381.

Clark, J. S. (2005) Why environmental scientists are becoming Bayesians. *Ecological Letters*, **8**(1), 2–14.

Clarke M.F., Avitabile, S.C., Brown, L., Callister, K.E., Haslem, A., Holland, G.J., Kelly, L.T., *et al.* (2010) Ageing mallee eucalypt vegetation after fire: insights for successional trajectories in semi-arid mallee ecosystems. *Australian Journal of Botany*, **58**(5), 363–372.

Condit, R., Ashton, P., Bunyavejchewin, S., Dattaraja, H.S., & Davies, S. (2006) The Importance of demographic niches to tree diversity. *Science*, **313**, 98 – 101.

Cornelissen, J.H.C., Lavorel, S., Garnier, E., Diaz, S., Buchmann, N., Gurvich, D.E., *et al.* (2003) A handbook of protocols for standardised and easy measurement of plant functional traits worldwide. *Australian Journal of Botany*, **51**(4), 335 – 380.

Craine, J.M. & Dybzinski, R. (2013) Mechanisms of plant competition for nutrients, water and light. *Functional Ecology*, **27**(4), 833 – 840.

Falster, D. S. & Westoby, M. (2003) Plant height and evolutionary games. *Trends in Ecology and Evolution*, **18**(7), 337–343.

Falster, D. S. & Westoby, M. (2005a) Tradeoffs between height growth rate, stem persistence and maximum height among plant species in a post-fire succession. *Oikos*, **111**(1), 57–66.

Falster, D. S. & Westoby, M. (2005b) Alternative height strategies among 45 dicot rain forest species from tropical Queensland, Australia. *Journal of Ecology*, **93**, 521 – 535.

563 Giljohann, K. M., McCarthy, M.A., Kelly, L.T., & Regan, T.J. (2015) Choice of
564 biodiversity index drives optimal fire management decisions. *Ecological Applications*,
565 **25**(1), 264–277.
566

567 Giljohann, K. M., McCarthy, M.A., Keith, D.A., Kelly, L.T., Tozer, M.G., & Regan, T.J.
568 (2017) Interactions between rainfall, fire and herbivory drive resprouter vital rates in a
569 semi-arid ecosystem. *Journal of Ecology*,
570

571 Givnish, T.J. (1988) Adaption to sun and shade: a whole-plant perspective. *Australian*
572 *Journal of Plant Physiology*, 15, 63 – 92.
573

574 Haslem, A., Kelly, L.T., Nimmo, D.G., Watson, S.J., Jenny, S.A., Taylor, R.S. *et al.*
575 (2010) Habitat or fuel? Implications of long-term, post-fire dynamics for the development
576 of key resources for fauna and fire. *Journal of Applied Ecology*, **48**(1), 247–256
577

578 Haslem, A., Avitable, S.C., Taylor, R.S., Kelly, L.T., Watson, S.J., Nimmo, D.G. *et al.*
579 (2012) Time-since-fire and inter-fire interval influence hollow availability for fauna in a
580 fire-prone system. *Biological Conservation*, **152**I, 212–221.
581

582 Herault, B., Ouallet, J., Blanc, L., Wagner, F. & Baraloto, C. (2010) Growth responses of
583 neotropical trees to logging gaps. *Journal of Applied Ecology*, **47**(4), 821–831.
584

585 Hérault, B., Bachelot, B., Poorter, L., Rossi, V., Bongers, F., Chave, J., *et al.* (2011)
586 Functional traits shape ontogenetic growth trajectories of rain forest tree species. *Journal*
587 *of Ecology*, **99**, 1431–1440.
588

589 Huston, M. & Smith, T. (1987) Plant succession: life history and competition. *The*
590 *American Naturalist*, **130**(2), 168 – 198.
591

- King, D. A., Davies, S.J., Supardi, M.N.N., & Tan, S. (2005) Tree growth is related to light interception and wood density in two mixed dipterocarp forests of Malaysia. *Functional Ecology*, **19**(3), 445–453.
- Kitajima, K. (1994) Relative Importance of Photosynthetic Traits and allocation Patterns as Correlates of Seedling Shade Tolerance of 13 Tropical Trees. *Oecologia*, **98**, 419 – 428.
- Kobe, R.K. (1999) Light Gradient Partitioning among Tropical Tree Species through Differential Seedling Mortality and Growth. *Ecology*, **80**(1), 187 – 201.
- Kunstler, G., Falster, D., Coomes, D.A., Hui, F., Kooyman, R., Laughlin, D., *et al.* (2016) Plant functional traits have globally consistent effects on competition. *Nature*, 529, 204 – 218.
- Loehle, C. (2000) Strategy Space and the Disturbance Spectrum: A Life-History Model for Tree Species Coexistence. *The American Naturalist*, **156**(1), 14–33.
- Moles, A. T. & Westoby, M. (2003) Latitude, seed predation and seed mass. *Journal of Biogeography*, **30**, 105 - 128.
- Moles, A. T. & Westoby, M. (2004) Seedling survival and seed size: a synthesis of the literature. *Journal of Ecology*, **92**, 372 – 383.
- Moles, A. T. & Westoby, M. (2006) Seed size and plant strategy across the whole life cycle. *Oikos*, **113**(1), 91 – 105.
- Monsi, M. & Saeki, T. (2005) On the Factor Light in Plant Communities and its Importance for Matter Production. *Annals of Botany*, **95**(3), 549 – 567.

- 622 Muir, A. M., Vesk, P.A., & Heopworth, G. (2014) Reproductive trajectories over decadal
623 time-spans after fire for eight obligate-seeder shrub species in south-eastern Australia.
624 *Australian Journal of Botany*, **62**(5), 369 – 378.
625
- 626 Murphy, B. P., Bradstock, R.A., Boer, M.M., Carter, J., Cary, G.J., Cochrane, M.A., *et al.*
627 (2013) Fire regimes of Australia: a pyrogeographic model system. *Journal of*
628 *Biogeography*, **40**(6), 1048–1058.
629
- 630 Noble, J.C. (1985) Fires and Emus: the population ecology of some woody plants in arid
631 Australia. *Studies on plant demography: a festschrift for John L. Harper*. (eds Harper,
632 J.L. and White, J.) pp. 33 – 49. London Academic Press.
633
- 634 Paine, C.E.T., Marthews, T.R., Vogt, D.R., Purves, D., Rees, M. *et al.* (2012) How to fit
635 nonlinear plant growth models and calculate growth rates: an update for ecologists.
636 *Methods in Ecology and Evolution*, **3**(2), 245–256.
637
- 638 Paine, C.E.T., Amissah, L., Auge, H., Baraloto, C., Baruffol, M., Bourland, N., *et al.*
639 (2015) Globally, functional traits are weak predictors of juvenile tree growth, and we do
640 not know why. *Journal of Ecology*, **103**(4), 978–989.
641
- 642 Pausas, J. G., & Keeley, J. E. (2014) Evolutionary ecology of resprouting and seeding in
643 fire-prone ecosystems. *New Phytologist*, **204**(1), 55–65.
644
- 645 Plummer, M. 2013. JAGS Version 3.4.0 user manual.
646 http://www.stats.ox.ac.uk/~nicholls/MScMCMC14/jags_user_manual.pdf
647
- 648 Pollock, L. J., Morris, W.K., & Vesk, P.A. (2011) The role of functional traits in species
649 distributions revealed through a hierarchical model. *Ecography*, **35**(8), 716–725.
650
- 651 Poorter, H., Remkes, C., & Lambers, H. (1990) Carbon and nitrogen economy of 24 wild
652 species differing in relative growth rate. *Plant Physiology*, **94**(2), 621 – 627.

- 653
654 Poorter, L., Bongers, L., & Bongers, F. (2006) Architecture of 54 moist-forest tree
655 species: traits, trade-offs, and functional groups. *Ecology*, **87**(5), 1289–1301.
656
657 Poorter, L., Wright, S.J., Paz, H., Ackerly, D.D., Condit, R., Ibarra-Manriques, G., (2008)
658 Are functional traits good predictors of demographic rates? Evidence from five
659 Neotropical forests. *Ecology*, **89**(7), 1908 – 1920.
660
661 Poorter, L., McDonald, I., Alarcon, A., Fichtler, E., Licona, J., Pena-Claros, M., (2009)
662 The importance of wood traits and hydraulic conductance for the performance and life
663 history strategies of 42 rainforest tree species. *New Phytologist*, 185(2), 481–492.
664
665 Prior, L.D. & Bowman, D.M.J.S. (2014) Across a macro-ecological gradient forest
666 competition is strongest at the most productive sites. *Frontiers in Plant Science*, **5**, 1 –
667 12.
668
669 Reich, P. B., Ellsworth, D.S., Walters, M.B., Vose, J.M., Gresham, C., Volin, J.C., &
670 Bowman, W.D. (1999) Generality of leaf trait relationships: a test across six biomes.
671 *Ecology*, **80**(6), 1955 – 1969.
672
673 Reich, P. B. (2014) The world-wide “fast-slow” plant economics spectrum: a traits
674 manifesto. *Journal of Ecology*, **102**(2), 275–301
675
676 Rice, B., & Westoby, M. (1999) Regeneration after fire in *Triodia* R. Br. *Austral*
677 *Ecology*, 24 (5), 563 – 572.
678
679 Ruger, N., Berger, U., Hubbell, S.P., Vieilledent, G., & Condit, R. (2011) Growth
680 Strategies of Tropical Tree Species: Disentangling Light and Size Effects. *PLoS ONE*,
681 **6**(9), e25330.
682
683 Ruger N., Wirth, C., Wright, S.J., & Condit, R. (2012) Functional traits explain light and

- size response of growth rates in tropical tree species. *Ecology*, **93**(12), 2626 – 2636.
- Sterck, F. J., Poorter, L., & Schieving, F. (2006) Leaf traits determine the growth-survival trade-off across rain forest tree species. *The American Naturalist*, **167**(5), 758–765.
- Su, Y. S. & Yajima, M. (2015) R2jags: a package for running jags from R. R package version 0.5-7. [http://CRAN.R-project.org/ package=R2jags](http://CRAN.R-project.org/package=R2jags).
- Thomas, S.C. (1996) Asymptotic height as a predictor of growth and allometric characteristics in Malaysian rain forest trees. *The American Journal of Botany*, **83**(5), 556 – 566.
- Tjørve, E. (2003) Shapes and functions of species–area curves: a review of possible models. *Journal of Biogeography*, **30**(6), 827–835.
- Tjørve, E. (2009) Shapes and functions of species-area curves (II): a review of new models and parameterizations. *Journal of Biogeography*, **36**(8), 1435–1445.
- Tjørve, E., & Tjørve, K. M. C. (2010) A unified approach to the Richards-model family for use in growth analyses Why we need only two model forms. *Journal of Theoretical Biology*, **267**(3), 417–425.
- Vesk, P. A. (2006) Plant size and resprouting ability: trading tolerance and avoidance of damage? *Journal of Ecology*, **94**(5), 1027–1034.
- Violle, C., Navas, M., Vile, D., Kazakou, E., Fortunel, C., Hummel, R., & Garnier, E. (2007) Let the concept of trait be functional! *Oikos*, **116**(5), 882–892.
- Wellington, B.A. (1984) Leaf water potentials, fire and refeneration of Mallee Eucalypts in Semi-Arid, South-Eastern Australia. *Oecologia*, **64**(3), 360-362.

Wellington, B.A. & Noble I.R. (1985a) Seed dynamics and factors limiting recruitment of the mallee *Eucalyptus incrassata* in semi-arid, south-eastern Australia. *Journal of Ecology*, **73**, 657 – 666.

Wellington, B.A. & Noble, I.R. (1985b) Post-fire recruitment and mortality in a population of the mallee *Eucalyptus incrassata* in semi-arid, south-eastern Australia. *Journal of Ecology*, **73**, 645-656.

Westoby, M. (1998) A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and Soil*, **199**, 213 – 227.

White, M. D. (2006) The Mallee vegetation of north western Victoria. *Proceedings of the Royal Society of Victoria*, **118**, 229 – 243.

Wright, I. J., Leishman, M.R., Read, C., & Westoby, M. (2006) Gradients of light availability and leaf traits with leaf age and canopy position in 28 Australian shrubs and trees. *Functional Plant Biology*, **33**(5), 407 – 419.

Figure 1. Plant height over times since fire and hierarchical model fits showing reseedling species a) *Acacia bractybotrya* (Mimosaceae), b) *Westringia rigida* (Lamiaceae), c) *Halgania cyanea* (Boraginaceae), d) *Beyeria opaca* (Euphorbiaceae), e) *Acacia montana* (Mimosaceae), and f) *Eremophila glabra* (Myoporaceae). The black dots represent observations.

Figure 2. Modeled height growth trajectories for 18 semi-arid woody plant species after fire. Resprouting species and their seed-derived growth comparison in (a), whereby blue dashed lines are modeled growth based on resprouting individuals, dashed black lines are seed derived plants from resprouting species. Non-sprouting species in (b), and species were modeled to 40 years, but all species reached asymptotic height by ten years, so we only display these values and include a close up of the growth trajectories from 0.5 to 3.5 years; the period where the most rapid growth occurs for all species in (c).

Figure 3. Species estimates for the three separate growth parameters in rank order; a) maximum growth rate, b) age at maximum growth and c) maximum height. The shaded panels at the base of each figure show the cross-species mean. The black dots and lines represent non-resprouting species, the blue dots and lines represent the separately modeled re-sprouting species. Error bars are 95% credible intervals. Species names are abbreviated to the first letter of the genus and species: *Beyeria opaca* (Bo), *Westringia rigida* (Wr), *Eremophila glabra* (Eg), *Phebalium squamulosum* (Ps), *Olearia muelleri* (Om), *Acacia montana* (Am), *Callitris verrucosa* (Cv), *Acacia wilhelmiana* (Aw), *Senna artemisioides* subsp *zygophylla* (Sa), *Dodonaea bursarifolia* (Db), *Acacia brachybotrya*

792 (Ab), *Olearia pimeleiodes* (Op), *Olearia subspicata* (Os), *Halgania cyanea* (Hc),
793 *Codonocarpus cotinifolius* (Cc), *Melaleuca lanceolata* (Ml), *Eucalyptus oleosa* (Euco),
794 *Eucalyptus gracilis* (Eucg).

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

Figure 4. Coefficients estimates (effect sizes) for linear predictors in the three submodels, which describe how traits influence the growth parameter. (a) Positive coefficients (numbers larger than 0) indicate a positive relationship between the trait values and the growth parameters. Bars represent 95% credible intervals. We also show 50% credible intervals with the light grey bars. (b-f) Partial dependency plots representing the effect of one trait on one growth parameter with all other parameters and coefficients held at their mean. (b) Seed mass had a negative influence on maximum growth rate, (c,d) leaf nitrogen content and seed mass positively influenced age of maximum growth, (e,f) stem

density and SLA negatively influenced asymptotic height. Along the x-axis are the scaled and centered trait values and along the y-axis are the parameter values for each species. . This trait-based growth model was constructed using only non-resprouting species. See Table S4 (appendix) for further details on effect sizes.

Figure 5. Example predicted height growth trajectories based on traits. (a) is a prediction of the growth trajectory of *Acacia brachybotrya* (Mimosaceae) which we have limited data for in our dataset. The black dots represent the height data for this species, the black line represents the fitted growth curve in our hierarchical model, the gray line represents using *A. brachybotrya* traits only to predict its growth. (b) represents two real species, *Dodonaea viscosa* (Sapindaceae) and *Myoporum platycarpum* (Myoporaceae), for which we have trait data but no height data and their predicted growth curves based on their traits. (c) two hypothetical species, where we have used trait values representing fast/slow and tall/short strategies. We present a tall, fast and early growing species (low stem density, low SLA, small seed mass, and low leaf nitrogen); a tall, late and slower growing species (low stem density, low SLA, heavy seed mass, high leaf nitrogen); a short, late and slow growing species (high stem density, high SLA, heavy seed mass, high leaf nitrogen); and a short, early and fast growing species (high stem density, high SLA, small seed mass and low leaf nitrogen). See methods for quantitative trait values. All predicted relationships are based on median predicted parameter values.

Appendix

Table S1. Mean trait values for 18 species from Murray Sunset National Park, the numbers in parentheses next to species names are the number of individuals sampled for trait measurements.

Family	Species	Specific leaf area (mg mm ⁻²)	Stem density (mg mm ⁻³)	Seed mass (mg)	Nitrogen content (%)
Mimosaceae	<i>Acacia bractybotrya</i> (16)	4.11	0.81	21.86	1.46
Mimosaceae	<i>Acacia montana</i> (15)	3.59	0.91	10.4	1.56
Mimosaceae	<i>Acacia wilhelmiana</i> (20)	3.44	0.83	5.9	1.65
Euphorbiaceae	<i>Beyeria opaca</i> (15)	3.9	0.76	20.32	1.57
Cupressaceae	<i>Callitris verrucosa</i> (15)	3.3	0.65	8.93	0.97
Gyrostemnaceae	<i>Codonocarpus cotinifolius</i> (5)	9.4	0.32	2.5	4.67
Sapindaceae	<i>Dodonaea bursarifolia</i> (20)	4.12	0.87	8	1.34
Myoporaceae	<i>Eremophila glabra</i> (20)	4.17	0.76	83	1.82
Myrtaceae	<i>Eucalyptus oleosa</i> (20)	2.55	0.84	0.58	0.97
Myrtaceae	<i>Eucalyptus gracilis</i> (20)	2.24	0.84	0.43	1.16
Boraginaceae	<i>Halgania cyanea</i> (20)	4.49	0.71	5.79	1.87
Myrtaceae	<i>Melaleuca lanceolata</i> (15)	2.26	0.77	0.35	1.34
Asteraceae	<i>Olearia muelleri</i> (16)	4.36	0.87	0.86	1.67
Asteraceae	<i>Olearia pimeleiodes</i> (25)	6.53	0.85	0.77	2.91
Asteraceae	<i>Olearia subspicata</i> (15)	3.5	0.82	0.8	2.09
Rutaceae	<i>Phebalium squamulosum</i> (15)	3.09	0.83	1.143	1.76
Mimosaceae	<i>Senna artemisioides</i> subsp. <i>zygophylla</i> (10)	1.68	0.87	12.15	1.59
Lamiaceae	<i>Westringia rigida</i> (20)	3.01	0.87	1.25	1.68

Figure S1. Scatter plots of model parameters and functional traits, the numbers in the top right corner of each panel are the Pearson's correlation coefficient.

Figure S2. Regressions showing relationships and R squared values between species structural attributes – canopy height (cm), canopy width (cm), canopy length (cm) and species basal diameter (mm) measured for all species.

899

900 Figure S3 – The relationships between species height (cm) and canopy length (cm) for
901 each species.

902

903

904

905

906

907

Figure S4 – The relationships between height (cm) and basal diameter (mm) measured for each species.

920 Table S2. Alternative non-linear model forms, convergence, Deviance Information
921 Criteria (DIC), the effective number of parameters (pD), mean R squared values of
922 observed versus predicted heights across all species (Naïve Rsq) and mean R squared
923 values of observed versus predicted heights based on internal cross validation, where
924 each species was left out of the training dataset and it's heights predicted (CV Rsq) and
925 key references for model forms.
926

Model	Equation	Convergence	DIC	Naïve Rsq	CV Rsq	Reference
Power	aT^b	50000	9759.1	0.792	0.001	Tjørve 2003
Exponential	$a + b \log(T)$	50000	8687.7	0.815	0.000	Tjørve 2003
Logistic	$a/(1 + \exp(-bT + c))$	100000	9276.1	0.780	0.011	Tjørve 2003
Hillslope log(T)	$a/\{1 + \exp[-b(\log(T) - c)]\}$	90000	8772.7	0.784	0.518	Tjørve 2009
Hillslope	$a/\{1 + \exp[-b(T - c)]\}$	90000	8723.0	0.779	0.514	Tjørve 2009

927
928
929
930
931
932
933

934
 935 Figure S5. Plant height data over times since fire and hierarchical model fits showing
 936 resprouting species a) *Eucalyptus oleosa*, b) *Eucalyptus gracilis* and c) *Melaleuca*
 937 *lanceolata*. The black dots represent field-collected data; the blue stars for the resprouting
 938 species represent heights of seedlings; the blue dots represent randomly selected height data
 939 points from individuals of indeterminate origin to use as asymptotic height estimates for the
 940 seeders of the same species.

941

942

943

944

945

946

947

948

949

950

951

952

953

954

955
956 Table S3 : Linear trait model statistics - Gelman’s R squared values represent the
957 proportion of variance in the response explained by the predictors for each level of a
958 multi-level model, values approaching 1 are better. Pooling factors assess the
959 contribution of within group estimates (species level data) versus population estimates
960 (all data combined), a pooling factor of less than 0.5 indicates the model is using more
961 within group information than population mean information. As models improve, both
962 Gelman’s R squared and the pooling factor should approach 1.
963

Growth parameter	Sub-model	Gelman’s R squared	Pooling factor
Asymptotic height (a)	$\log(a_j) = \exp[\beta_0 + \beta_1 \times SD_j + \beta_2 \times SLA_j + \varepsilon_j]$	0.57	0.19
Maximum relative growth rate (b)	$\log(b_j) = \exp[\beta_0 + \beta_1 \times SM_j + \varepsilon_j]$	0.77	0.64
Age at maximum growth rate (c)	$\log(c_j) = \exp[\beta_0 + \beta_1 \times N_j + \beta_2 \times SM_j + \varepsilon_j]$	0.83	0.46

964
965 Table S4. Effect sizes of Figure 4 displaying the growth parameter, traits that influence
966 that parameter as corresponding for Figure 4 and the mean effect size with lower and
967 upper bounds of 95% credible interval.

968

Growth parameter	Trait	Mean standardised effect size (95% credible intervals)
Asymptotic height (a)	Stem density	- 0.58 [- 0.99, - 0.20]
Asymptotic height (a)	Specific leaf area	- 0.19 [- 0.57, 0.19]
Age at maximum growth rate (c)	Seed mass	0.08 [- 0.02, 0.19]
Age at maximum growth rate (c)	Leaf nitrogen	0.05 [- 0.05, 0.15]
Maximum relative growth rate (b)	Seed mass	- 0.06 [- 0.14, 0.03]

975

976

977

1 **Appendix**

2 **Detailed statistical modeling methods**

3

4 While a multi-species hierarchical approach is not as simple as single species regressions,
5 we believe the added complexity is worthwhile, as explained and unexplained variance
6 can be partitioned between the observation and the species level, and species with few
7 data can still be included. Furthermore this approach is relatively simple compared to
8 fully mechanistic models, which can be hard to parameterise with empirical data. Similar
9 approaches have been used for hierarchical models of plant resprouting (Vesk 2006),
10 quantifying distributions of demographic rates in tropical forests (Condit *et al.* 2006) and
11 species distributions (Pollock *et al.* 2011). We believe this represents a practical approach
12 to describing the growth of multiple plant species.

13

14 We were interested in the height growth of plant species; our observations were heights
15 (cm) of individual plants for each species across time (yr). Because plant heights are
16 finite, height growth must be nonlinear, and as such so must our model. Our model had a
17 hierarchical structure corresponding to two levels of variation: between species and
18 between individuals. The strength of hierarchical modeling is that it allows explained and
19 unexplained variation to be partitioned within and between multiple levels (in this case,
20 species and individual) of a dataset. It does this by allowing parameters to vary between
21 levels by assuming they are drawn from a common distribution (Condit *et al.* 2006,
22 Gelman & Hill 2007, Ruger *et al.* 2012). The advantage of this is that rare or data-scarce
23 species can also be modeled from these common distributions. When field data are
24 scarce for less common species, the ability to retain species with few observations is
25 attractive. A hierarchical model structure also allows specifying sub-models for model
26 parameters (Pollock *et al.* 2011). In our case, functional traits are formally included as
27 species-specific linear predictors of the growth parameters in a Hillslope growth model
28 (Tjörve 2009). This approach reveals not just plant growth through time for individual
29 species, but also how functional traits contribute to inter and intra-species variation in
30 growth and represents a more, but not fully, mechanistic understanding of how traits
31 affect the function of plants.

32

33 *A hierarchical, nonlinear growth model*

34

35 We modeled the growth of species using a hierarchical multi-species non-linear growth
36 model, which formally incorporates plant traits as species-level predictors of growth.

37 We have an **observation model** (Eq. 1.1), in which the heights of individual plants are
38 drawn from a lognormal distribution. The lognormal distribution reflects the natural
39 constraints of height data, which are only positive numbers, with relatively few extreme
40 height values (Limpert, Stahel & Abbt 2001):

41

$$H_i \sim \text{lnorm}(\mu_i, \sigma^{-2}) \qquad \text{Equation 1.1}$$

Where H is the observed height (cm) for each individual i , which is drawn from a lognormal distribution with mean μ_i precision σ^{-2} .

We have a **process model**, where each of our modeled height observations is estimated from a growth model. A range of possible growth models exist (Tjørve 2003; Tjørve & Tjørve 2010; Paine *et al.* 2012) and our aim was not to systematically compare every growth model, but instead to choose one that has biologically relevant parameters (Thomas 1996; Falster & Westoby 2005a). We chose a sigmoidal model referred to as the Hillslope equation (Tjørve 2009), which is re-paramaterisation of a logistic equation with three parameters (Eq 1.2):

$$\mu_{i,j} = \frac{\alpha_{i,j}}{(1 + \exp[-\beta_{i,j} (T_i - c_{i,j})])} \quad \text{Equation 1.2}$$

Where μ is the observed height of individual plant i of species j (cm) and T_i is the observed time since fire (yr) of each individual i , and α , β , and c are species-specific constants. These are biologically interpretable and will be referred to from here as the maximum height achieved (α), the slope (β) represents the maximum relative growth rate (maximum RGR), which is measured in units of cm/cm/yr (Atwell 1999) and the time at which this maximum growth occurs (c , the location parameter) measured in years.

Each of the three parameters in our process model of growth are estimated via **parameter sub-models**, equations 1.3 – 1.5. Mean parameter values are estimated by species-specific linear models, with species trait values as covariates to influence each of the parameters and with intercepts varying by species. In deciding which functional traits to include in our models we relied on previously published literature to generate hypothesis of trait effects on growth. Specifically that wood density, seed mass and leaf traits would be influential traits affecting growth trajectories (Falster & Westoby 2005a; Reich *et al.* 2014), whereby seed mass should be most influential on initial growth (Moles & Westoby 2006), stem density of achievable height (Falster & Westoby 2005a; Falster & Westoby 2005b) and leaf traits throughout the whole growth process (Sterck, Poorter & Schieving 2006; Reich *et al.* 2014). Initially we let all traits influence each parameter in our growth model, and used a process of backwards selection to test different trait-parameter combinations. Our final model had stem density and SLA influencing maximum height, nitrogen leaf content and seed mass on age at maximum growth and seed mass on maximum relative growth rate, (see equations 3 - 5).

$$\log(\alpha_j) = \exp[\beta_0 + \beta_1 \times SD_j + \beta_2 \times SLA_j + \varepsilon_j] \quad \text{Equation 1.3}$$

$$\log(\beta_j) = \exp[\beta_0 + \beta_1 \times SM_j + \varepsilon_j] \quad \text{Equation 1.4}$$

$$\log(c_j) = \exp[\beta_0 + \beta_1 \times N_j + \beta_2 \times SM_j + \varepsilon_j] \quad \text{Equation 1.5}$$

Where the ε terms represent separate species-level random effects and describe the residual departure of species from the fixed effects.

We used Bayesian inference employing Markov chain Monte Carlo (MCMC) methods with the open-source software package JAGS version 3.3.0 (Plummer 2003) run via the statistical software environment R version 2.15.2 (R Development Core Team, 2015) with the package R2jags (Su & Yajima 2015). Priors for parameters were normally distributed, centered on zero with a standard deviation of 0.0001. We used positive half Cauchy distributions truncated at 0, with prior mean 0 and prior scale 25 for the observation error and random effects (Gelman 2006). We initialized our model by using mean parameter values taken from single species non-linear models. We logged trait values and then centered and scaled our trait data by one standard deviation before modeling.

Model choice

As well as biologically interpretable parameters, our model choice sought to adequately represent the growth of all species modeled. One consequence of modeling multiple species and growth forms together is that some species are better fit than others. We compared our three-parameter sigmoidal model to four other possible growth models; two two-parameter models (Power and Exponential) and one other three-parameter model (Logistic), plus our chosen Hillslope model with log time. To compare these models, we looked at DIC and R squared values between predicted and observed heights for each species. Additionally we performed internal cross validation for each model, where each species was left out of the training dataset and then it's height over time predicted and observed and predicted heights assessed by R squared values (Appendix Table S2). Overall, the Hillslope model had the lowest DIC and performed well in internal cross validation (Appendix Table S2). We did not consider four, five or six parameter models, because we were cautious of over-parameterisation with our relatively small empirical dataset. We chose to use the three-parameter sigmoidal model form due to the biological interpretable parameters and its use in previous studies (Thomas 1996; Falster & Westoby 2005a) and the increased information due to an extra parameter specifying age at maximum relative growth rate. We do not consider these alternative models further. We display some model fits to raw data of our chosen model (Fig. 1a-f).

For the resprouting species, heights of seedlings were measured when the seedling origins were obvious in the field. However, past approximately ten years this became difficult to distinguish and measurements were made on resprouters only. For this reason, and also because we were unsure about using a multi-species model of growth which included both seed derived individuals and resprouters we chose to model the resprouters separately; we present the resprouter data in the Appendix Fig. S2. In order to incorporate the seedlings of resprouter species for comparison against the resprouter growth trajectories, we randomly sampled some top heights from the resprouting individuals (grey circles, Appendix Fig. S2) to use in our model. This approach assumes

that resprouting and seed derived individuals of resprouting species achieve similar top heights. The sigmoidal model for resprouter species tended to under predict the asymptotic height of *Eucalyptus oleosa* and *Eucalyptus gracilis* (Appendix Fig. S2).

Model checking

Three chains were monitored to ensure convergence, which was assessed both visually and via the Brooks-Gelman-Rubin convergence diagnostic (Brooks & Gelman 1998). We evaluated model fit through visual checks of predicted versus observed heights and by comparing Deviance Information Criteria (DIC, Spiegelhalter *et al.* 2002) between alternative models. We evaluated the fit of our species-specific linear sub-models by calculating Gelman's R squared statistic (this statistic can be interpreted like a classical R squared statistic, whereby it tells us the proportion of variance in the response explained by the predictors for each level of a multi-level model) and pooling factors (which assesses the contribution of mean model group estimates versus data) for each submodel (Gelman & Pardoe 2006), these statistics are reported in the Appendix Table S3. We plot coefficient estimates with 95% and 50% credible intervals of the linear predictors in our growth parameter sub-models. While the effect sizes are sometimes small (effect sizes can be seen in Fig 4a and Appendix Table S4) we discuss the trends apparent in the data, particularly as we chose our predictor variables based on a priori information based on previous trait-based work (see methods).

References

- Atwell, B. (1999) Growth analysis: a quantitative approach. *Plants in action*. (eds. B. Atwell, P. Kriedemann, & C. Turnbull), pp. 188. Macmillan Education, Australia.
- Brooks, S.P. & Gelman, A. (1998) General methods for monitoring convergence of iterative simulations. *Journal of Computation and Graphical Statistics*, **7**(4), 434-455.
- Condit, R., Ashton, P., Bunyavejchewin, S., Dattaraja, H.S., & Davies, S. (2006) The Importance of demographic niches to tree diversity. *Science*, **313**, 98 – 101.
- Falster, D. S. & Westoby, M. (2005a) Tradeoffs between height growth rate, stem persistence and maximum height among plant species in a post-fire succession. *Oikos*, **111**(1), 57–66.
- Falster, D. S. & Westoby, M. (2005b) Alternative height strategies among 45 dicot rain forest species from tropical Queensland, Australia. *Journal of Ecology*, **93**, 521 – 535.
- Gelman, A. (2006) Prior distributions for variance parameters in hierarchical models (comment on article by Browne and Draper). *Bayesian Analysis*, **1**, 515 – 534.
- Gelman, A. & Hill, J. (2007) Data Analysis using regression and multilevel/hierarchical models. Cambridge University Press.

- 165 Gelman, A. & Pardoe, I. (2006) Bayesian Measures of Explained Variance and Pooling
166 in Multilevel (Hierarchical) Models. *Technometrics*, **48**(2), 241–251.
167
- 168 Limpert, E., Stahel, W.A., & Abbt, M. (2001) Lognormal distributions across the
169 sciences: keys and clues. *Bioscience*, **51**(5), 341 – 352.
170
- 171 Moles, A. T. & Westoby, M. (2006) Seed size and plant strategy across the whole life
172 cycle. *Oikos*, **113**(1), 91 – 105.
173
- 174 Paine, C.E.T., Marthews, T.R., Vogt, D.R., Purves, D., Rees, M. *et al.* (2012) How to fit
175 nonlinear plant growth models and calculate growth rates: an update for ecologists.
176 *Methods in Ecology and Evolution*, **3**(2), 245–256.
177
- 178 Pollock, L. J., Morris, W.K., & Veski, P.A. (2011) The role of functional traits in species
179 distributions revealed through a hierarchical model. *Ecography*, **35**(8), 716–725.
180
- 181 Reich, P. B. (2014) The world-wide “fast-slow” plant economics spectrum: a traits
182 manifesto. *Journal of Ecology*, **102**(2), 275–301
183
- 184 Ruger N., Wirth, C., Wright, S.J., & Condit, R. (2012) Functional traits explain light and
185 size response of growth rates in tropical tree species. *Ecology*, **93**(12), 2626 – 2636.
186
- 187 Sterck, F. J., Poorter, L., & Schieving, F. (2006) Leaf traits determine the growth-survival
188 trade-off across rain forest tree species. *The American Naturalist*, **167**(5), 758–765.
189
- 190 Spiegelhalter, D.J., Best, N.G., Carlin, B.P., & van der Linde, A. (2002) Bayesian
191 measures of model complexity and fit. *The Journal of the Royal Statistical Society B*,
192 **64**(4), 583–639.
- 193 Su, Y. S. & Yajima, M. (2015) R2jags: a package for running jags from R. R package
194 version 0.5-7. <http://CRAN.R-project.org/package=R2jags>.
- 195 Thomas, S.C. (1996) Asymptotic height as a predictor of growth and allometric
196 characteristics in Malaysian rain forest trees. *The American Journal of Botany*, **83**(5), 556
197 – 566.
198
- 199 Tjørve, E. (2003) Shapes and functions of species–area curves: a review of possible
200 models. *Journal of Biogeography*, **30**(6), 827–835.
201
- 202 Tjørve, E. (2009) Shapes and functions of species-area curves (II): a review of new
203 models and parameterizations. *Journal of Biogeography*, **36**(8), 1435–1445.
204
- 205 Tjørve, E., & Tjørve, K. M. C. (2010) A unified approach to the Richards-model family
206 for use in growth analyses Why we need only two model forms. *Journal of Theoretical*
207 *Biology*, **267**(3), 417–425.

208

209 Vesk, P. A. (2006) Plant size and resprouting ability: trading tolerance and avoidance of
210 damage? *Journal of Ecology*, **94**(5), 1027–1034.

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227 Table S1. Mean trait values for 18 species from Murray Sunset National Park, the
228 numbers in parentheses next to species names are the number of individuals sampled for
229 trait measurements.

230

Family	Species	Specific leaf area (mg mm ⁻²)	Stem density (mg mm ⁻³)	Seed mass (mg)	Nitrogen content (%)
Mimosaceae	<i>Acacia bractybotrya</i> (16)	4.11	0.81	21.86	1.46
Mimosaceae	<i>Acacia montana</i> (15)	3.59	0.91	10.4	1.56
Mimosaceae	<i>Acacia wilhelmiana</i> (20)	3.44	0.83	5.9	1.65
Euphorbiaceae	<i>Beyeria opaca</i> (15)	3.9	0.76	20.32	1.57
Cupressaceae	<i>Callitris verrucosa</i> (15)	3.3	0.65	8.93	0.97
Gyrostemnaceae	<i>Codonocarpus cotinifolius</i> (5)	9.4	0.32	2.5	4.67
Sapindaceae	<i>Dodonaea bursarifolia</i> (20)	4.12	0.87	8	1.34
Myoporaceae	<i>Eremophila glabra</i> (20)	4.17	0.76	83	1.82
Myrtaceae	<i>Eucalyptus oleosa</i> (20)	2.55	0.84	0.58	0.97
Myrtaceae	<i>Eucalyptus gracilis</i> (20)	2.24	0.84	0.43	1.16
Boraginaceae	<i>Halgania cyanea</i> (20)	4.49	0.71	5.79	1.87
Myrtaceae	<i>Melaleuca lanceolata</i> (15)	2.26	0.77	0.35	1.34
Asteraceae	<i>Olearia muelleri</i> (16)	4.36	0.87	0.86	1.67
Asteraceae	<i>Olearia pimeleiodes</i> (25)	6.53	0.85	0.77	2.91
Asteraceae	<i>Olearia subspicata</i> (15)	3.5	0.82	0.8	2.09
Rutaceae	<i>Phebalium squamulosum</i> (15)	3.09	0.83	1.143	1.76
Mimosaceae	<i>Senna artemisioides</i> subsp. <i>zygophylla</i> (10)	1.68	0.87	12.15	1.59
Lamiaceae	<i>Westringia rigida</i> (20)	3.01	0.87	1.25	1.68

231
232
233
234
235
236
237
238
239

Figure S1. Scatter plots of model parameters and functional traits, the numbers in the top right corner of each panel are the Pearson's correlation coefficient.

251

252

253 Figure S2. Regressions showing relationships and R squared values between species

254 structural attributes – canopy height (cm), canopy width (cm), canopy length (cm) and

255 species basal diameter (mm) measured for all species.

256

257

258

259

260

261

Figure S3 – The relationships between species height (cm) and canopy length (cm) for each species.

Figure S4 – The relationships between height (cm) and basal diameter (mm) measured for each species.

Table S2. Alternative non-linear model forms, convergence, Deviance Information Criteria (DIC), the effective number of parameters (pD), mean R squared values of observed versus predicted heights across all species (Naïve Rsq) and mean R squared values of observed versus predicted heights based on internal cross validation, where each species was left out of the training dataset and it's heights predicted (CV Rsq) and key references for model forms.

Model	Equation	Convergence	DIC	Naïve Rsq	CV Rsq	Reference
Power	aT^b	50000	9759.1	0.792	0.001	Tjørve 2003
Exponential	$a + b \log(T)$	50000	8687.7	0.815	0.000	Tjørve 2003
Logistic	$a/(1 + \exp(-bT + c))$	100000	9276.1	0.780	0.011	Tjørve 2003
Hillslope log(T)	$a/\{1 + \exp[-b(\log(T) - c)]\}$	90000	8772.7	0.784	0.518	Tjørve 2009
Hillslope	$a/\{1 + \exp[-b(T - c)]\}$	90000	8723.0	0.779	0.514	Tjørve 2009

298
299 Figure S5. Plant height data over times since fire and hierarchical model fits showing
300 resprouting species a) *Eucalyptus oleosa*, b) *Eucalyptus gracilis* and c) *Melaleuca*
301 *lanceolata*. The black dots represent field-collected data; the blue stars for the resprouting
302 species represent heights of seedlings; the blue dots represent randomly selected height data
303 points from individuals of indeterminate origin to use as asymptotic height estimates for the
304 seeders of the same species.

319

320 Table S3 : Linear trait model statistics - Gelman's R squared values represent the
 321 proportion of variance in the response explained by the predictors for each level of a
 322 multi-level model, values approaching 1 are better. Pooling factors assess the
 323 contribution of within group estimates (species level data) versus population estimates
 324 (all data combined), a pooling factor of less than 0.5 indicates the model is using more
 325 within group information than population mean information. As models improve, both
 326 Gelman's R squared and the pooling factor should approach 1.
 327

Growth parameter	Sub-model	Gelman's R squared	Pooling factor
Asymptotic height (a)	$\log(a_j) = \exp[\beta_0 + \beta_1 \times SD_j + \beta_2 \times SLA_j + \varepsilon_j]$	0.57	0.19
Maximum relative growth rate (b)	$\log(b_j) = \exp[\beta_0 + \beta_1 \times SM_j + \varepsilon_j]$	0.77	0.64
Age at maximum growth rate (c)	$\log(c_j) = \exp[\beta_0 + \beta_1 \times N_j + \beta_2 \times SM_j + \varepsilon_j]$	0.83	0.46

328

329 Table S4. Effect sizes of Figure 4 displaying the growth parameter, traits that influence
 330 that parameter as corresponding for Figure 4 and the mean effect size with lower and
 331 upper bounds of 95% credible interval.

332

Growth parameter	Trait	Mean standardised effect size (95% credible intervals)
Asymptotic height (a)	Stem density	- 0.58 [- 0.99, - 0.20]
Asymptotic height (a)	Specific leaf area	- 0.19 [- 0.57, 0.19]
Age at maximum growth rate (c)	Seed mass	0.08 [- 0.02, 0.19]
Age at maximum growth rate (c)	Leaf nitrogen	0.05 [- 0.05, 0.15]
Maximum relative growth rate (b)	Seed mass	- 0.06 [- 0.14, 0.03]

338

339

340

301x177mm (72 x 72 DPI)

376x161mm (72 x 72 DPI)

341x229mm (72 x 72 DPI)

266x229mm (72 x 72 DPI)

469x229mm (72 x 72 DPI)