Knowns and Unknowns of Influenza B Viruses

Marios Koutsakos¹, Thi HO Nguyen¹, Wendy S Barclay² and Katherine Kedzierska^{1*}

¹Department of Microbiology and Immunology, University of Melbourne, at the Peter Doherty Institute for Infection and Immunity, Parkville VIC 3010, Australia; ²Section of Virology, Faculty of Medicine, Wright Fleming Institute, Imperial College London, Norfolk Place, London W2 1PG, UK

Running title: Flu B virology and immunology

Keywords: influenza B, virology, epidemiology and immunology

*Correspondence should be addressed to:

Katherine Kedzierska, PhD

Department of Microbiology and Immunology

University of Melbourne

At the Peter Doherty Institute for Infection and Immunity

Vic 3010, Australia

Ph: +61 03 8344 7962

Fax: +61 03 9347 1540

kkedz@unimelb.edu.au

26 ABSTRACT (Words: 103)

27

28 Influenza B (FluB) viruses circulate annually along with influenza A (FluA)
29 strains during seasonal epidemics. FluB can dominate influenza seasons and
30 cause severe disease, particularly in children and adolescents. Research has
31 revealed interesting aspects of FluB and highlighted the importance of these
32 viruses in clinical settings. Yet, many important questions remain
33 unanswered. In this review, the clinical relevance of FluB is emphasized,
34 unique features are highlighted and gaps in knowledge pinpointed. Multiple
35 aspects of FluB epidemiology, evolution, virology and immunology are
36 discussed. Future research into FluB is needed to understand how we can
37 prevent severe disease in high-risk groups, especially children and elderly.

In 1940, an outbreak of acute respiratory illness in children with clinical presentations identical to the contemporary influenza A epidemics led to the isolation of a new virus that was serologically distinct from previous influenza A isolates. This novel influenza virus was named Influenza B [1].

It is now well established that influenza A (FluA) and Influenza B (FluB) viruses co-circulate annually within the human population and cause respiratory illness with varying levels of severity. Unlike FluA, FluB does not have an established animal reservoir and thus does not pose the same pandemic risk as FluA [2]. Consequently, FluB has received considerably less attention. Nonetheless, FluB presents a substantial challenge during annual epidemics. Although current knowledge of FluB viruses highlights their unique features, a plethora of questions on the virology, immunology, epidemiology and evolution of FluB remains to be elucidated.

FluB viruses belong to the family of Orthomyxoviruses and have a segmented negative-sense single-stranded RNA genome. FluB has 8 segments and encodes for 3 polymerase proteins (PA, PB1 and PB2), the nucleoprotein (NP), the non-structural protein (NS1), the nuclear export protein (NEP or sometimes called BNS2), the matrix protein (BM1), the BM2 ion channel and three surface glycoproteins (HA, NA and NB) [3] (Fig 1A-B). FluA and FluB proteins generally show substantial differences in length, amino acid composition and function [2]. FluB viruses are currently divided into two genetically and antigenically distinct lineages, B/Victoria/2/1987-like and B/Yamagata/16/1988-like viruses [4-6].

Epidemiology and Clinical Presentation of FluB

FluB viruses from one or both lineages co-circulate annually with the seasonal H1N1 and H3N2, with FluB being the main cause of the seasonal epidemic every 2-4 years [7,8]. Multiple lines of evidence suggest that FluB can cause substantial disease [9-11]. In an analysis of influenza-related mortality between 1976 and 1999 in the US, highest morbidity was associated with H3N2, followed by FluB and then by H1N1 [12]. This highlights that FluB infections can cause a significant degree of morbidity, although the analysis was conducted in 2009 and the above hierarchy might have changed with the establishment of the H1N1pdm strain in seasonal epidemics. Regardless, various reports highlight a strong presence of FluB in seasonal epidemics across the world. An epidemiological analysis of influenza-attributable mortality in the US between 1997 and 2009, reported that 29% of total influenza associated deaths were attributable to a FluB infection. Importantly, in 4 out of 12 seasons FluB infection was the predominant cause of seasonal influenza associated mortality [13]. In the 2008 season in Australia, 54% of influenza notifications were due to FluB infections, and the majority of these infections were in children [14]. In England, in the first half of the 2012/2013 season, FluB was the predominant cause of influenza infections, with the highest activity occurring in children aged 5-14 [15]. Additionally, in Thailand between 2010 and 2014 the majority of FluB infections occurred in children aged 5-19 [16]. These are only a few studies indicating that FluB can cause severe disease, especially in the young. The factors determining the predominance of FluB over FluA in a given season are largely unknown, but most likely depend on specific viral characteristics and interactions from all three circulating viruses.

The symptoms of FluB infection include a sore throat, coughing, nasal discharge and fever [17]. Overall, disease caused by FluB is indistinguishable from that of FluA [18], although complications can differ. Complications of FluB can include neurological and muscular manifestations [19,20], cardiologic complications [21-23] and secondary bacterial infections [22,24,25]. Paddock *et al* (2012) observed that secondary bacterial pneumonia was more frequently observed in adult fatal cases of FluB than pediatric fatal cases, and the reverse was observed for cardiac complications. The same study also determined that in fatal cases of influenza, the time from disease onset to death was shorter for FluB than that determined for the 1918, 1957, 1968, 2009 pandemic FluA or seasonal H3N2 [22]. It is however important to assess the validity of this observation in other FluB cohorts. It also appears that lineage does not impact disease outcome [26]. As mentioned above, although FluB can infect adults and the elderly, disease manifestations are more prevalent in children [27-29]. Moreover, FluB viruses of the Victoria lineage are more prevalent in younger populations than Yamagata viruses [6,30,31]. It has been proposed that this could be explained by higher expression of α -2,3 linked sialic acid glycans in children than in adults and the differential α -2,3 and α -2,6 sialic acid glycan binding preferences of viruses from the two lineages [6,32]. Overall, FluB can cause significant disease, particularly in children. The viral and host determinants of FluB pathogenicity and the reasons of increased susceptibility of young children remain to be elucidated.

Evolution and Phylodynamics of FluB

113 In the early decades of the 20th century FluB viruses belonged to a
114 single lineage. Two antigenically distinct lineages emerged prior to 1980 [4-6]
115 and now co-circulate globally, with one lineage usually predominating in any
116 given year, although equal circulation of the two lineages in certain areas has
117 been reported. For instance in the 2011-2012 season in Canada, 24% of all
118 influenza cases were caused by B/Victoria-like viruses and 27% were of
119 B/Yamagata-like origin [33]. In general, after the appearance of the two
120 lineages, Victoria-like viruses dominated in the late 1980s, while viruses of the
121 Yamagata lineage prevailed in the 1990s, with the exception of an outbreak of
122 Victoria-like viruses in Asia in 1996/1997. The Victoria lineage re-emerged
123 globally in 2001, and the two lineages have co-circulated ever since [34].
124 Interestingly, while new H3N2 viruses seem to originate from East and
125 Southeast Asia in every season, FluB strains can persist during multiple
126 seasons and re-seed subsequent epidemics [35,36]. This appears to be
127 linked to less frequent global movements of FluB compared to H3N2 FluA
128 [35].

129 Evolution of FluB involves a constant turnover of antigenically-distinct
130 viruses, in a process similar to the antigenic drift of FluA, as well as
131 reassortment. The evolution rate of FluB is slower than that of FluA, as
132 calculated by the rate of nucleotide substitutions for each segment. (0.14×10^{-3} –
133 3.32×10^{-3} substitutions/site/year for FluB, 2.68×10^{-3} – 12.5×10^{-3}
134 substitutions/site/year for FluA H3N2) [34,37]. HA exhibits the highest rate of
135 adaptive evolution, suggesting a degree of positive selection, depicting the
136 effect of immune evolutionary pressure [5,34] Additionally, the NA and NS1
137 segments exhibit high levels of adaptive evolution, possibly due to immune

pressure [34]. In addition to these changes, reassortment also contributes to FluB genetic diversity. Reassortment can occur between viruses of the same lineage or viruses of the two distinct lineages [5,6,34,38]. Inter-lineage reassortment, in which Victoria viruses acquire genes from the Yamagata lineage have been more frequent than the reverse [6]. Interestingly, different segments can exhibit different patterns during reassortment events. Two bioinformatics studies revealed that only the PB1, PB2 and HA segments have maintained separate Victoria or Yamagata lineages. In contrast, the NS1/NS2 segment of current strains is derived from the Victoria lineage, while the rest of the segments (PA, NP, NA/NB, M1/BM2) come from the Yamagata lineage, suggesting substantial exchange of genetic information between the two lineages [6,38]. The observation that novel reassortant viruses can dominate the viral population implies that reassortment can generate novel strains with increased fitness.

Intriguingly, the two lineages exhibit different phylodynamics. Distinct clades of Victoria viruses generally circulate for 1-3 years before being replaced by new clades, while multiple clades of Yamagata viruses can co-circulate for longer periods of time. Additionally, Victoria viruses exhibit great genetic diversity between seasons and a stronger antigenic drift compared to Yamagata viruses. Furthermore, viruses of the Victoria lineage contain more amino acid substitutions in the HA and these are located near the receptor binding site of the HA, while changes in Yamagata viruses are less frequent and tend to appear in sites more distant to the receptor binding site. Overall, the Yamagata viruses are more conserved and show weaker antigenic drift and selection than the faster evolving Victoria viruses [6].

In addition to these evolutionary forces, interactions seem to occur between FluA and FluB. For instance, FluA (H1N1pdm09) infection can prevent FluB infection within a short interval of time but the opposite is not true, in the ferret model [39]. Additionally, analysis of FluA and FluB viruses over 30 years revealed that in certain years, strong genetic bottlenecks have been imposed on FluB viruses and these were associated with high prevalence of FluA. Such bottlenecks have led to changes in the dominant FluB antigenic cluster[34].

In summary, FluB evolution is a complex function of interactions between FluB viruses of the different lineages, interactions between co-circulating FluB and FluA viruses as well as the selective pressures from the host's immune mechanisms.

Virology of FluB

The Haemagglutinin (HA)

The HA of FluB mediates attachment and entry of the virus into cells via interaction with sialic acids. FluB HA binds to sialic acids with lower affinity than FluA [40,41]. Also, the FluB HA can interact with both α -2,6- and α -2,3-linked sialic acids. Despite their low sequence homology (~20-30%), the overall structures of FluB and FluA HAs are similar. Similarly to FluA, the FluB HA has an elongated membrane-proximal domain and a globular membrane-distal domain. The functional form of the FluB HA is a homotrimer. The Receptor Binding Site (RBS) of FluB HA has some significant differences to that of FluA HA. One important difference is located at amino acid position 95

(FluB HA numbering), which in FluB is a phenylalanine, while in FluA is a highly conserved tyrosine [40,42]. The Phe95 residue has been shown to be involved in the ability of FluB HA to interact with both human-like and avian-like receptor analogues and with the lower affinity of FluB HA. Replacement of Phe95 with a tyrosine residue increased receptor binding to levels comparable to FluA and reduced binding to avian-like receptor analogues [43]. The role of the HA protein in determining pathogenicity of FluA is well established [44], however this has not been studied for FluB.

The Neuraminidase (NA)

The overall structure of FluB NA is similar to that of FluA NA, despite only 30% sequence homology. The active form of the FluB NA is a tetramer and each monomer contains two glycosylation sites (Asn283 and Asn143) and a calcium-ion binding site. Asn283 is specific for FluB, while Asn143 is conserved among Influenza viruses [45]. Nineteen highly conserved residues form the NA active site of influenza A and B viruses. Residues in the catalytic site (R118, D151, R152, R224, E276, R292, R371, and Y406) directly interact with sialic acids, while framework residues structurally support the enzymatic active site (E119, R156, W178, S179, D198, I222, E227, H274, E277, N294, and E425). Neuraminidase inhibitors (NAIs) licensed in the clinic for treatment of influenza infection appear to be less effective against FluB [46].

The NB Protein

The NB protein is uniquely encoded by FluB in segment 6, overlapping with the NA coding sequence and is not present in FluA. The start codon of

213 NB is located 4 nucleotides upstream of the NA start codon [47]. NB is an
214 integral membrane glycoprotein of 100 amino acids that is incorporated into
215 the virion [48]. The N-terminus contains two N-glycosylation sites, which are
216 further modified by the addition of multiple N-acetyl-glucosamine residues
217 [49]. Palmitoylation of C-terminal residues is important for trafficking of NB to
218 the cell surface [50]. NB is dispensable for virus rescue and *in vitro* replication
219 [51] (unpublished observations Elderfield, Koutsakos & Barclay). Hatta and
220 Kawaoka observed attenuation in absence of NB in BALB/c mice [51],
221 however, this could be due to an altered NA activity as the mutations
222 introduced have been previously reported to affect the NA expression [47]. In
223 contrast, our data suggest that a NB-knockout virus with unaffected NA
224 activity replicates equivalently to the WT virus in C57B/6 mice. This mutant
225 virus showed no fitness cost in ferrets and also transmitted to a new host,
226 although overall transmission in that study was low (unpublished observations
227 Elderfield, Koutsakos & Barclay). This apparent redundancy of NB is
228 surprising and could be due to a host-specific role of NB. Additionally, the *in*
229 *vivo* role of NB has only been assessed in the early stages of infection (up to
230 day 4). It is possible that NB plays a role in the later stages of FluB infection.

231 In various experimental settings, NB has displayed the ability to
232 conduct ions. In *Xenopus* oocytes, NB could stimulate the conductance of Cl⁻
233 ions [52], while in *E.coli*-derived membranes NB was permeable to protons
234 [53]. Additionally, purified protein and peptides corresponding to the
235 transmembrane region of NB incorporated into artificial lipid bilayers could
236 form cation-selective channels [54,55]. A review of these studies questioned
237 the observed ion channel activity due to the limitations of the experimental

systems utilized, especially as previous observations showed that membrane proteins with no known ion channel activity were able to form multimers and conduct ions in similar experimental settings [56]. Subsequently, it was reported that a single amino acid substitution (S20A) in the NB transmembrane domain resulted in altered gating and proton permeability, supporting the idea of a NB ion channel. However, unlike the AM2, BM2, and CM2 ion channels of FluA, FluB and Influenza C viruses, NB has no pH-modulating activity in the *trans*-Golgi network (TGN), which the AM2, BM2 and CM2 ion channels do [57]. It is therefore unclear whether NB operates as an ion channel. Overall, more than 30 years after the discovery of NB, its role and function remain enigmatic. Understanding the role of NB could benefit the design of novel live-attenuated vaccines [51].

The Matrix Protein (BM1)

Very little is known about the M1 protein of FluB, which is encoded in segment 7. M1 plays an important role in FluB adaptation in mice and attenuation of the master donor vaccine strain (B/Ann Arbor/1/66) of the Live Attenuated Influenza Vaccine (LAIV) [58]. The functional role of M1 in the life cycle of FluB is not entirely known. M1 contains a Nuclear Localization Signal (NLS), two Nuclear Export Signals (NES) and can act as a shuttling protein between the nucleus and the cytoplasm in infected cells. All of these features were necessary for the function of M1, as they were required for the production of viable virus [59]. Thus, the exact role of BM1, and whether it is of a similar function to that of FluA M1, remains to be determined.

The BM2 Ion Channel

Like in FluA, segment 7 encodes for a second protein, namely BM2 [60]. The BM2 protein is a small hydrophobic, integral membrane protein [61] that is synthesized in the late stages of infection [62]. The protein is transported through the TGN to the cell surface [63] and is incorporated in the virions [62]. The N-terminus of the protein appears to be important for its transport within infected cells [63]. An interaction of BM2 with M1 has been proposed [64], supporting previous observations that BM2 is involved in the incorporation of the vRNP in the virions [65]. Additionally, BM2 is necessary for the production of viable virus [65-67] and decreased incorporation of BM2 in virions reduces infectivity, indicating a role for BM2 similar to that of AM2 [66]. Indeed, BM2 contains a HXXXW motif like AM2 and thus can conduct ions. This activity is similar to the one of AM2 in acidifying the endosome and modulating pH in the Golgi [68]. The crystal structure of BM2 reveals similarities to AM2 but also unique features. BM2 forms coiled-coil tetramers, with polar residues lining the pore that is formed, a feature important for the inactivity of adamantanes against BM2 [69]. Zhang *et al* , recently suggested a novel role for BM2 in inhibiting p53-mediated transcription and apoptosis [70].

A unique feature of BM2 is its expression mechanism. The BM2 coding sequence lies directly downstream of the M1 coding sequence and the two are translated from the same mRNA molecule. The translation termination codon of M1 overlaps with the start codon of BM2 in the following pentanucleotide sequence: UAAUG (M1 stop-codon underlined, BM2 start codon in bold) [71]. The 45 nucleotides in the M1 coding sequence upstream of penta-

nucleotide appear essential for BM2 expression [72]. Additionally, the sequence upstream of the penta-nucleotide exhibits complementarity to the 18S component of the small ribosomal subunit [73]. An interaction between the mRNA and the rRNA as well as the involvement of the translation initiation factor eIF3 are important for BM2 expression [74]. This termination:reinitiation mechanism was the first one of its kind when initially identified, but since then similar mechanisms have been identified in other viruses, like caliciviruses [73].

The Nucleocapsid Protein (NP)

The NP of FluB (BNP) is encoded by segment 5. It differs greatly from the NP of FluA (ANP). The most striking differences are the lack of the two NLSs of ANP and the extended N-terminus of the protein in FluB. Specifically, BNP contains 50 amino acids in its N-terminus that are not found in ANP [75].

Structural studies revealed that BNP consists of a head, a body and a flexible tail loop. The groove between the head and the body contains an extended charged loop (residues 125-149), within which there are two lysine clusters critical for RNA binding. The flexible tail loop of BNP enables the assembly of homo-oligomers. Comparisons with the structure of ANP revealed an interesting similarity. In both proteins, a highly conserved salt bridge (R472-E395 in BNP, R416-E339 in ANP) seems to be involved in the stability of the loop and thus the assembly of homo-oligomers [76]. Pharmacological inhibition of this salt bridge in ANP can inhibit FluA [77], and thus might be a possible drug target for FluB.

The N-terminus of BNP has been of particular interest, as it is considerably different from ANP. Wanitchang *et al* showed that the N-terminus of BNP is required for nuclear localization and suggested that the K₄₄RTR₄₇ motif in BNP might be acting as a NLS [78]. Reverse genetics were later employed to generate FluB lacking the N-terminus of BNP, in order to investigate the role of the N-terminus of BNP in the context of infection and viral replication. Different mutations in the N-terminus affected the ability to rescue virus or viral replication and transcription. Moreover, the entire N-terminus appears to be essential for optimal localization in the nucleus during infection and residues 51-81 were suggested to contain a bipartite NLS [79]. The NP protein of FluA is known to have multiple functions that are crucial for effective virus replication, however, there are no similar studies for BNP. Current knowledge suggests that BNP has features similar to ANP but also possesses some unique domains, suggesting that at least some functions of the two proteins are unique.

The Polymerases (PA, PB1 & PB2)

Although there is a vast amount of knowledge about the polymerases of FluA, minimal data exist on the polymerase proteins of FluB, with the general view being that they work similarly to those of FluA. However it is pertinent that the FluB heterotrimeric polymerase complex structure was solved before that of a human or avian FluA virus [80]. Indeed, similarities exist, although recent studies suggest key differences. Localization studies showed that all three FluB proteins have inherent nuclear import function and encode NLSs. PB2 was found to predominantly localize in the nucleus, while

PB1 and PA were localized both in the nucleus and the cytoplasm. Pairwise interactions for all three proteins were also observed and localized in the nucleus. These observations are similar to what has been observed for FluA polymerase proteins [81].

A major difference between FluA and FluB polymerases is their cap-binding mechanism. Specifically FluB polymerases can bind to and cleave both m⁷G-capped and unmethylated G-capped mRNAs. This is in contrast to FluA polymerases that exclusively bind to and cleave m⁷G-capped mRNA [82]. Structural studies suggested that the overall structure and binding of the polymerases to the vRNA is similar between FluA and FluB [80], but also elucidate the details of the unique cap-binding mechanism of FluB [83]. The polymerases of FluA have central roles in host-adaptation and pathogenicity [44,84]. Conversely, the contributions of FluB polymerases to pathogenicity and host-restriction are unknown.

The Non-Structural 1 Protein (NS1)

The NS1 protein of FluB (NS1B) is a 281 amino acid long protein encoded by segment 8. NS1B is important for virus replication as knock-out viruses display impaired replicative capacity both *in vitro* [85] and *in vivo* [86]. It consists of an N-terminus (residues 1-90), a C-terminus (residues 120-281) and a linker domain (residues 91-119) [87]. NS1B localizes in the nucleus of infected cells in the early stages of infection, but mostly in the cytoplasm in the later stages. Residues 46-56 in the N-terminus function as a monopartite NLS and NS1B interacts with importin α 3. Within the nucleus, NS1B localizes in nuclear speckles containing splicing factor SC35 [88]. Although NS1B can

bind RNA via its N terminus it cannot not inhibit export of host mRNA like NS1A does, as it does not interact with CPSF30 via its C terminus [89].

NS1B can antagonize multiple aspects of the interferon antiviral response [85,90]. Firstly, it can inhibit the IFN $\alpha\beta$ response by preventing activation and nuclear translocation of IRF3. Particularly, FluB lacking NS1B induces stronger IFN β expression and activation of the IFN β promoter than WT FluB [85]. Additionally, ectopic overexpression of NS1B can inhibit IFN β promoter activity [90]. Interestingly, NS1B can complement the attenuated growth of Δ NS1A FluA [90]. All these studies implicate NS1B as a potent inhibitor of the IFN response. Both the N- and C- termini are involved in this function. The N-terminus of NS1B sequesters dsRNA to prevent activation of the antiviral PKR kinase, while the C-terminus is more involved in IFN β promoter inhibition [91], although either termini could independently prevent activation of IRF-3 [90]. Interestingly, NS1B knock-out viruses are attenuated in IFN-deficient Vero cells [85], suggesting that NS1B has additional effects separate of counteracting the IFN response.

NS1B also can directly bind to and block the interferon-stimulated Ubiquitin-like ISG15 protein, which is naturally conjugated to newly synthesized proteins [92]. By binding to ISG15, NS1B prevents activation of ISG15 by the UBE1L enzyme, and this activity is mediated by the N-terminus of the NS1B protein [87,93]. Published evidence suggests that NS1B re-localizes the bound ISG15 to the nuclear speckles [94]. NS1B specifically interacts with human and non-human primate ISG15, but not with other mammalian ISG15-like mouse or canine, implicating the ISG15-NS1B interaction in the host-restriction of FluB [94].

Overall, NS1B has a key role in innate immune response evasion, like NS1A, although the mechanism seems to be different for the two viruses. The unique targeting of ISG15 suggests that this interferon stimulated gene would otherwise play an important antiviral role against influenza B virus.

The Nuclear Export Protein (or Non-Structural 2) Protein (NEP)

The NEP protein of FluB interacts with the nuclear export machinery, contains a Nuclear Export Signal (NES) sequence in its N-terminus and possesses nuclear export activity [95]. Furthermore, it is produced in the late stages of infection, initially accumulating in the nucleus, then being transported to the cytoplasm and finally, in plasma membrane proximal regions, the sites of virion budding. NEP can also be detected in virions as part of the vRNP complex. Interestingly, in FluB NEP seems to interact directly with the vRNP and BM1 as opposed to FluA where M1 acts as a bridge between NEP and the vRNP [96]. Overall, FluB NEP appears to have a similar function to its FluA counterpart, although its precise function is yet to be determined.

Non-coding Sequences

The coding sequences of Flu viruses are flanked by stretches of non-coding sequences in both termini. These sequences form the promoter that drives viral replication and transcription and also contain packaging signals [3]. In the specific context of FluB, they have shown to be important for vRNA replication [97]. The sequences at the extreme of each terminus are highly conserved. Specifically, the first 9 nucleotides of the 3' vRNA end (5'-

AGCAGAAGC-3' in cDNA) and the first 10 nucleotides of the 5' vRNA end (with the exception of position 6) (3'- TCATC_xTTGT-5' in cDNA) are highly conserved among segments and strains (Fig 1C). The sequences in the two termini show a high degree of complementarity and thus form secondary RNA structures. Between these extreme terminal sequences and the coding sequences, lie non-coding sequences that are conserved for each segment but differ between segments [98]. Understanding the non-coding sequences of influenza viruses has been pivotal in the generation of reverse genetics systems[2,98-100] as well as the design of novel vaccination strategies [101].

Immunology of FluB

Innate Immune Recognition and Response

As with other aspects of FluB immunology, the innate immune response has been largely unexplored. The recognition receptors for FluB have not been identified but RIG-I seems to be essential in the response to FluB. MDA-5 might play an additional, although not essential, role in initiating an innate response against FluB [102]. PKR can also detect viral RNAs and inhibit replication of FluB both in cell culture and in mice, as highlighted by the inhibition of PKR by NS1B. Activation of PKR was temporally linked to the appearance of viral RNPs in the cytoplasm and purified vRNPs could activate PKR activity *in vitro* [103]. FluB induces IFN-dependent and independent responses through activation of IRF3 [104]. The MAP Kinase and NFκB pathways can be activated. Interestingly, the innate immune recognition of FluB occurs earlier in infection of dendritic cells than FluA (peak IFN-λ mRNA at 2-4 hours for FluB and at 8-16 hours for FluA) and is independent of viral

replication or transcription [105]. The mechanism behind this early recognition is not yet understood.

In response to *in vitro* FluB infection, epithelial and innate cells can produce type I and III interferons, specifically IFN α , IFN β , IFN λ 1, and other cytokines/chemokines, including CXCL10, CXCL8, CCL5, CCL2, IL1- β , IL-18 and IL-6, although the specific levels and roles remain to be investigated [105-107]. Similarly, although $\gamma\delta$ -T cell hybridomas can respond to both FluA- and FluB-infected cells [108], the roles and functions of $\gamma\delta$ -T cells or other innate immune cells in controlling FluB are unknown.

Antibody Responses

Similarly to FluA, the vast majority of antibody responses against FluB are directed at the HA. The four main antigenic sites are in the globular head of the HA1 and include the 120 loop (residues 116-137), the 150 loop (residues 141-150), the 160 loop (residues 162-167) and the 190 loop (residues 194-202) [42]. These sites of the HA are under positive selection, indicating the effect of the antibody response to the evolution of the virus [109]. Antibodies against the NA can also neutralize FluB and protect mice from sub-lethal infections. Interestingly, immunization with B/Yamagata/16/88 NA can confer protection against B/Yamagata/16/87 as well as B/Victoria/2/87 and B/Malaysia/2506/04, both of which belong to the Victoria lineage [110]. Although the two lineages are considered antigenically distinct, HA-directed antibody cross-reactivity between the two lineages has been observed in various contexts [111-115] potentially due to original antigenic sin[116].

Broadly neutralizing monoclonal antibodies have also been identified for FluB, firstly in a seminal paper by Dreyfus et al [117]. Antibody CR8033 binds to the RBS region of the HA, while antibody CR8071 binds to the vestigial esterase domain at the base of the HA head. Both can neutralize viruses from the two FluB lineages *in vivo*, even at low doses. Interestingly, they can inhibit the release of viral progeny rather than the entry of viruses into cells. Another antibody, CR9114, binds to the HA stem, on an epitope conserved across FluB and FluA viruses. Structural studies showed that this antibody binds to the HA of H1, H3, H5, H7, H9 and FluB. CR9114 blocks the pH-induced conformational changes in HA and subsequent membrane fusion. CR9114 could protect mice against lethal infection. Interestingly, sequence analysis of the V_H region of CR9114 showed a small number of somatic mutations, suggesting that this broadly neutralizing antibody can be generated from genomic sequences without extensive hypermutation [117]. Yasugi et al [118] reported another broadly neutralizing antibody against FluB (5A7), which targets the C-terminal region of HA1. This antibody showed therapeutic efficacy even when administered after challenge. Broadly neutralizing antibodies against the NA have also been reported. An antibody recognizing residues 222-230 (N2 numbering), an epitope highly conserved across FluB and FluA, can inhibit growth of FluB in cells but was not assessed *in vivo* [119,120]. Vaccines that would prime antibody responses against such highly conserved and cross-reactive epitopes would be of vital importance in generating universal immunity against influenza viruses.

T cells Responses

The involvement of CD8⁺ T cells in the immune response to FluA has been extensively studied in mice and human PBMCs [121-126]. However, little is known about T cell responses to FluB. Mice lacking any antibody responses are protected from lethal FluB challenge in a CD8⁺-T cell-dependent manner. Primary infection can protect from secondary FluB infection and this is also dependent on CD8⁺ as well as CD4⁺ T cells [127]. Thus, similarly to FluA, T cells have an important role in promoting recovery from FluB infection. Three epitopes from NP of FluB have been identified for CD8⁺ T cells. Robbins *et al* [128], identified the HLA-A*0201-restricted NP₈₂₋₉₄ (MVVKLGEFYNQMM) epitope by its low similarity (leucine before a glycine followed by hydrophobic residues) to the FluA A2⁺-M1₅₆₋₆₈ epitope. Two HLA-B8 restricted epitopes from the NP protein of FluB (NP₃₀₋₃₈ RPIIRPAT and NP₂₆₃₋₂₇₁ ADRGLLRDI) were also identified later [129]. However, in a recent study, only the NP₃₀₋₃₈ peptide could induce INF γ production from FluB polyclonal CTLs [130]. It has also been proposed that CTL responses against FluB are preferentially directed against HLA-B8 epitopes [131]. It was recently shown that FluB-specific polyclonal CTLs cross-react with viruses from the opposing lineage of FluB [130]. Some influenza A PB1 epitopes are conserved even in FluB PB1 sequence [132], but whether T cells against these epitopes are cross-reactive and cross-protective against both FluA and FluB remains to be determined experimentally. Although primary infection with FluB cannot protect from secondary infection with FluA and vice versa, in the BALB/c mouse model [133,134], immunization strategies that probe T-cell responses against conserved and cross-reactive epitopes could potentially confer universal heterologous immunity against Flu viruses. Thus, determining

the cross-reactivity of T cells against FluA and FluB is a key step in understanding universal anti-influenza T cell immunity.

Flu B Vaccines

Immunization against FluB is provided in the traditional Trivalent Inactivated Vaccine (TIV). However, until recently the annual formulation of this vaccine only contained a virus from either the Victoria or the Yamagata lineage depending on the year (Table 1). The high levels of co-circulation of the two lineages in the last decade results in frequent mismatch of predicted and circulating strains and has necessitated the inclusion of viruses from both lineages in the formulation of a quadrivalent vaccine [27]. The quadrivalent vaccine is available as an inactivated intramuscular vaccine or as a LAIV in a nasal spray.

High yields of virus are required during the production of inactivated vaccines. For FluA, this is achieved by reassortment, resulting in the combination of the antigenic surface proteins in the PR8 (A/Puerto Rico/8/34) backbone, which grows to high titers during production in eggs. There is no established 'high-yield' strain for FluB and the yield of FluB during annual production varies greatly. Ways to generate high yields of FluB for vaccine purposes as well as studies into the attenuation mechanisms of influenza B LAIV have been reviewed [2]. Recently Le *et al* [135], generated reassortant viruses with B/Lee/40 as a donor strain and found that the NP gene from the B/Lee/40 strain was central for high-yield production. However, overall there is lack of a 'high-yield' FluB strain.

Antiviral Strategies against FluB

Two classes of licensed anti-influenza drugs exist, adamantanes (eg. amantadine and rimantadine) and NAIs, like oseltamivir and zanamivir. Adamantanes block the AM2 ion channel activity but not of BM2. This is attributed to the polar amino acid residues in the BM2 ion channel cavity, as they reduce accessibility of the drug to the ion channel pore [69]. Consequently, these drugs are ineffective against FluB.

NAIs can inhibit FluB replication, although with variable efficiency. The inhibitory effect of NAIs against both FluA and FluB can be attributed to the high conservation of the catalytic site across both FluA and FluB. The efficacy of NAIs and the resistance mutations of FluBs were extensively reviewed recently by Burnham et al [46]. Overall, oseltamivir shows lower efficacy against FluB than FluA, while zanamivir shows comparable efficacy.

Mutations conferring resistance to NAIs have been reported in clinical isolates, surveillance isolates as well as reverse genetics studies. These mutations can be found either in the catalytic site of the enzyme (eg. R152K) or in framework residues (eg D198N, I222T). Studies on the fitness cost of NAI resistance mutations have shown that, contrary to most resistance mutations, the E119A and H274Y mutations have little impact on viral replication and fitness [136,137]. Interestingly, it appears that different mutations have different frequencies and effects in the NA gene from the two FluB lineages [138]. Studies in NAI resistance by FluB highlight the possibility of uncompromised NAI-resistant viruses emerging in human populations, suggesting the need for increased surveillance and potentially novel antiviral strategies.

Host Range of FluB

Influenza B viruses are restricted to human hosts [2]. In 1999, a FluB outbreak in seals was reported in the Netherlands. Serological and virological data indicated that a strain of FluB circulating in humans was introduced to the seal population in 1995 [139] and further evidence of seal FluB infections were reported later [140,141]. Sequence analysis of the B/Seal/1999 virus revealed high identity to a human FluB strain, suggesting no adaptation of the virus in seals. A serological survey in swine farms across the US revealed antibodies against FluB in pig serum samples but virus was only detected by RT-PCR of the NS segment in a limited number of nasal swabs. Unfortunately, virus could not be isolated. The partial sequences derived from the RT-PCR were of high homology to human isolates, suggesting no adaptation. In an experimental setting, FluB could replicate in the respiratory tracts of pigs and infected animals exhibited flu-like symptoms, lung lesions and seroconversion. FluB transmitted via direct contact from infected to sentinel animals, although transmission was limited [142]. Antibodies against FluB have also been detected in guinea pigs [143], and FluB replicates and transmits efficiently during experimental infection of guinea pigs [144]. In addition, FluB replicates in ferret airway epithelial cells as well as infected animals and transmits between ferrets, which are considered the gold standard animal model for human influenza infections [145](unpublished observations Elderfield, Koutsakos & Barclay). Overall, while endemic circulation of FluB is restricted to humans, there are indications of FluB cross overs from humans to other species. However, there is currently

no evidence of sustained animal reservoirs of FluB. As discussed above, this restriction of FluB in humans may, at least to an extent, be attributed to the human host specific interactions of NS1[5,87,94] and M1 [58] proteins.

Conclusions & Future Perspectives

Over the last decade, it has become apparent that FluB constitutes a substantial part of annual flu epidemics. The particular pathogenicity of FluB in children, the co-circulation of the two lineages, which results in frequent mismatches of vaccine and circulating strains, and the emergence of NAI-resistance mutations with no apparent fitness cost further highlight the importance of FluB as a human pathogen.

Although it has generally been assumed that FluA and FluB viruses are similar, studies from the last decade emphasize multiple levels of uniqueness in FluB virology, immunology and evolution (Table 2). Some FluB proteins, like NP, have unique domains, others, like NB, are uniquely encoded by FluB viruses and still others can work in unique ways, like the cap-snatching mechanism of the FluB polymerases.

The numerous gaps of knowledge regarding FluB mean that multiple avenues of research can prove fruitful in the future. A better understanding of the evolution and phylodynamics of FluB viruses, such as the interactions between the two lineages and the interactions between FluA and FluB, can aid surveillance and vaccine strain determination and may inform epidemiology [39]. A clearer picture of the molecular virology of FluB could provide a better understanding of FluB pathogenicity, adaptation and design of novel antiviral strategies. A more comprehensive understanding of FluB

immunology would be an important step towards more effective vaccines and possibly antibody- and/or T cell-mediated universal influenza immunity. Overall, FluB is a research field waiting to explode.

Executive Summary

Clinical Presentation and Epidemiology

- FluB strains dominate every 2-4 years.
- Clinical presentations are similar to FluA, although usually more severe in children and adolescents.

Evolution and Phylodynamics

- Two lineages of FluB co-circulate globally, although usually one predominates in any given year..
- The two lineages exhibit contrasting phylodynamics.
- Re-assortment between the same lineage or two lineages contributes to the evolution of FluB.

Virology

- FluB has 8 gene segments encoding for 11 proteins.
- These generally have low sequence similarity to FluA and can exhibit important functional differences.
- NB is uniquely encoded by FluB and its function is unknown.
- NS1 inhibits the IFN $\alpha\beta$ response as well as ISG15 in a species-specific manner.
- The exact contributions of FluB proteins to pathogenicity and host-adaptation are unclear.

Immunology

- Mainly unexplored.
- FluB stimulates IFN $\alpha\beta$ and cytokine antiviral responses, although with kinetics different to FluA.
- Antibody responses are directed against HA and universally neutralizing antibodies against HA have been identified.
- T cells confer protection against primary and secondary infection in a mouse model.
- Only a handful CD8⁺ T cell epitopes have been identified and T cell cross-reactivity between two lineages has been described.
- Trivalent (two FluA and one FluB strain) and quadrivalent (two FluA and two FluB strains) vaccines are available and updated annually.

Antiviral Strategies

- Adamantanes are ineffective against FluB.
- Neuranimidase inhibitors work but not as efficiently as for FluA.
- Resistance mutations with no apparent fitness cost have been detected.

Host Range

- There is limited evidence of reverse zoonosis of FluB but no established animal reservoir.

References

1. Francis T, Jr. A New Type of Virus from Epidemic Influenza. *Science*, 92(2392), 405-408 (1940).
2. Jackson D, Elderfield RA, Barclay WS. Molecular studies of influenza B virus in the reverse genetics era. *The Journal of General Virology*, 92(Pt 1), 1-17 (2011).
- ** Thorough review of the use of reverse genetics to study Influenza B viruses.
3. Peter Palese MLS. Orthomyxoviridae: The Viruses and their Replication. In: *Fields Virology*. David M. Knipe, PMH (Ed. (LIPPINCOTT WILLIAMS & WILKINS, a WOLTERS KLUWER BUSINESS, USA, 2007) 1647-1689.
4. Rota PA, Wallis TR, Harmon MW, Rota JS, Kendal AP, Nerome K. Cocirculation of two distinct evolutionary lineages of influenza type B virus since 1983. *Virology*, 175(1), 59-68 (1990).
5. McCullers JA, Saito T, Iverson AR. Multiple genotypes of influenza B virus circulated between 1979 and 2003. *Journal of virology*, 78(23), 12817-12828 (2004).
6. Vijaykrishna D, Holmes EC, Joseph U *et al.* The contrasting phylodynamics of human influenza B viruses. *eLife*, 4, e05055 (2015).

- 686 **Detailed analysis of the phylogenetics and evolution of the two lineages.
- 687 7. Thompson WW, Shay DK, Weintraub E *et al.* Mortality associated with
688 influenza and respiratory syncytial virus in the United States. *Jama*,
689 289(2), 179-186 (2003).
- 690 8. Lin YP, Gregory V, Bennett M, Hay A. Recent changes among human
691 influenza viruses. *Virus research*, 103(1-2), 47-52 (2004).
- 692 9. W. Paul Glezen JKS, Carrie M. Kuehn, Kellie J. Ryan, and John
693 Oxford. The Burden of Influenza B: A Structured Literature Review.
694 *American Journal of Public Health*, 103(3), e43-e51 (2013).
- 695 * Interesting meta-analysis of Influenza B clinical data.
- 696 10. Gutierrez-Pizarra A, Perez-Romero P, Alvarez R *et al.* Unexpected
697 severity of cases of influenza B infection in patients that required
698 hospitalization during the first postpandemic wave. *The Journal of*
699 *infection*, 65(5), 423-430 (2012).
- 700 11. Su S, Chaves SS, Perez A *et al.* Comparing clinical characteristics
701 between hospitalized adults with laboratory-confirmed influenza a and
702 B virus infection. *Clinical infectious diseases : an official publication of*
703 *the Infectious Diseases Society of America*, 59(2), 252-255 (2014).
- 704 12. Thompson WW, Moore MR, Weintraub E *et al.* Estimating influenza-
705 associated deaths in the United States. *Am J Public Health*, 99 Suppl
706 2, S225-230 (2009).
- 707 13. Matias G, Taylor R, Haguenet F, Schuck-Paim C, Lustig R, Shinde V.
708 Estimates of mortality attributable to influenza and RSV in the United
709 States during 1997-2009 by influenza type or subtype, age, cause of
710 death, and risk status. *Influenza and other respiratory viruses*, 8(5),
711 507-515 (2014).
- 712 14. Kaczmarek M, Owen R, Barr IG. Annual report of the National
713 Influenza Surveillance Scheme, 2008. *Communicable diseases*
714 *intelligence quarterly report*, 34(1), 8-22 (2010).
- 715 15. Pubic Health England. Surveillance of influenza and other respiratory
716 viruses, including novel respiratory viruses, in the United Kingdom:
717 Winter 2012/13. (2013)

- 718 16. Tewawong N, Suwannakarn K, Prachayangprecha S *et al.* Molecular
719 epidemiology and phylogenetic analyses of influenza B virus in
720 Thailand during 2010 to 2014. *PloS one*, 10(1), e0116302 (2015).
- 721 17. Peter F. Wright GN, Yoshihiro Kawaoka. Orthomixoviruses. In: *Fields*
722 *Virology*. David M. Knipe, PMH (Ed. (LIPPINCOT WILLIAMS &
723 WILKINS, a WOLTERS KLUWER BUSINESS, USA, 2007)
- 724 18. Irving SA, Patel DC, Kieke BA *et al.* Comparison of clinical features
725 and outcomes of medically attended influenza A and influenza B in a
726 defined population over four seasons: 2004-2005 through 2007-2008.
727 *Influenza and other respiratory viruses*, 6(1), 37-43 (2012).
- 728 19. Moon JH, Na JY, Kim JH *et al.* Neurological and muscular
729 manifestations associated with influenza B infection in children.
730 *Pediatric neurology*, 49(2), 97-101 (2013).
- 731 20. Thabet FI, Khalil S, Naz F, Dyme IZ. Cerebellar mutism and reversible
732 cytotoxic edema in influenza B-associated encephalopathy. *Pediatric*
733 *neurology*, 49(6), 489-492 (2013).
- 734 21. Frank H, Wittekind C, Liebert UG *et al.* Lethal influenza B myocarditis
735 in a child and review of the literature for pediatric age groups. *Infection*,
736 38(3), 231-235 (2010).
- 737 22. Paddock CD, Liu L, Denison AM *et al.* Myocardial injury and bacterial
738 pneumonia contribute to the pathogenesis of fatal influenza B virus
739 infection. *The Journal of infectious diseases*, 205(6), 895-905 (2012).
- 740 23. Taremi M, Amoroso A, Nace HL, Gilliam BL. Influenza B-induced
741 refractory cardiogenic shock: a case report. *BMC infectious diseases*,
742 13, 452 (2013).
- 743 24. Aebi T, Weisser M, Bucher E, Hirsch HH, Marsch S, Siegemund M.
744 Co-infection of Influenza B and Streptococci causing severe
745 pneumonia and septic shock in healthy women. *BMC infectious*
746 *diseases*, 10, 308 (2010).
- 747 25. Scaber J, Saeed S, Ihekweazu C, Efstratiou A, McCarthy N, O'Moore
748 E. Group A streptococcal infections during the seasonal influenza
749 outbreak 2010/11 in South East England. *Euro surveillance : bulletin*
750 *Europeen sur les maladies transmissibles = European communicable*
751 *disease bulletin*, 16(5) (2011).

26. Erhard van der Vries DKMI, Benjamin J. Cowling, Jitao D. Zhang, Xiao Tong, Krzysztof Wojtowicz, Martin Schutten, and Charles A. Boucher. Outcomes and Susceptibility to Neuraminidase Inhibitors in Individuals Infected with Different Influenza B Lineages: the Influenza Resistance Information Study. *Journal of Infectious Diseases*, (2015).
27. Belshe RB. The need for quadrivalent vaccine against seasonal influenza. *Vaccine*, 28 Suppl 4, D45-53 (2010).
28. Hite LK, Glezen WP, Demmler GJ, Munoz FM. Medically attended pediatric influenza during the resurgence of the Victoria lineage of influenza B virus. *International journal of infectious diseases : IJID : official publication of the International Society for Infectious Diseases*, 11(1), 40-47 (2007).
29. Olson DR, Heffernan RT, Paladini M, Konty K, Weiss D, Mostashari F. Monitoring the impact of influenza by age: emergency department fever and respiratory complaint surveillance in New York City. *PLoS medicine*, 4(8), e247 (2007).
30. Socan M, Prosenc K, Ucakar V, Berginc N. A comparison of the demographic and clinical characteristics of laboratory-confirmed influenza B Yamagata and Victoria lineage infection. *Journal of clinical virology : the official publication of the Pan American Society for Clinical Virology*, 61(1), 156-160 (2014).
31. Tan Y, Guan W, Lam TT *et al.* Differing epidemiological dynamics of influenza B virus lineages in Guangzhou, southern China, 2009-2010. *Journal of virology*, 87(22), 12447-12456 (2013).
32. Wang YF, Chang CF, Chi CY, Wang HC, Wang JR, Su IJ. Characterization of glycan binding specificities of influenza B viruses with correlation with hemagglutinin genotypes and clinical features. *Journal of medical virology*, 84(4), 679-685 (2012).
33. Skowronski DM, Janjua NZ, Sabaiduc S *et al.* Influenza A/subtype and B/lineage effectiveness estimates for the 2011-2012 trivalent vaccine: cross-season and cross-lineage protection with unchanged vaccine. *The Journal of infectious diseases*, 210(1), 126-137 (2014).
34. Chen R, Holmes EC. The evolutionary dynamics of human influenza B virus. *Journal of molecular evolution*, 66(6), 655-663 (2008).

- 786 ** Thorough analysis of Influenza B evolution and epidemiology over and
787 extended time period.
- 788 35. Bedford T, Riley S, Barr IG *et al.* Global circulation patterns of seasonal
789 influenza viruses vary with antigenic drift. *Nature*, 523(7559), 217-220
790 (2015).
- 791 ** Detailed comparative analysis of Influenza A and B global spatio-temporal
792 patterns over 12 years.
- 793 36. Patterson Ross Z, Komadina N, Deng YM *et al.* Inter-Seasonal
794 Influenza is Characterized by Extended Virus Transmission and
795 Persistence. *PLoS pathogens*, 11(6), e1004991 (2015).
- 796 37. Nelson MI, Simonsen L, Viboud C *et al.* Stochastic processes are key
797 determinants of short-term evolution in influenza A virus. *PLoS*
798 *pathogens*, 2(12), e125 (2006).
- 799 38. Dudas G, Bedford T, Lycett S, Rambaut A. Reassortment between
800 influenza B lineages and the emergence of a coadapted PB1-PB2-HA
801 gene complex. *Molecular biology and evolution*, 32(1), 162-172 (2015).
- 802 ** Interesting bioinformatics analysis of Influenza B reassortment patterns over
803 28 years.
- 804 39. Laurie KL, Guarnaccia TA, Carolan LA *et al.* Interval Between
805 Infections and Viral Hierarchy Are Determinants of Viral Interference
806 Following Influenza Virus Infection in a Ferret Model. *The Journal of*
807 *infectious diseases*, (2015).
- 808 40. Wang Q, Tian X, Chen X, Ma J. Structural basis for receptor specificity
809 of influenza B virus hemagglutinin. *Proceedings of the National*
810 *Academy of Sciences of the United States of America*, 104(43), 16874-
811 16879 (2007).
- 812 41. Matrosovich MN, Gambaryan AS, Tuzikov AB *et al.* Probing of the
813 receptor-binding sites of the H1 and H3 influenza A and influenza B
814 virus hemagglutinins by synthetic and natural sialosides. *Virology*,
815 196(1), 111-121 (1993).
- 816 42. Wang Q, Cheng F, Lu M, Tian X, Ma J. Crystal structure of unliganded
817 influenza B virus hemagglutinin. *Journal of virology*, 82(6), 3011-3020
818 (2008).

- 819 43. Ni F, Mbawuike IN, Kondrashkina E, Wang Q. The roles of
820 hemagglutinin Phe-95 in receptor binding and pathogenicity of
821 influenza B virus. *Virology*, 450-451, 71-83 (2014).
- 822 44. Medina RA, Garcia-Sastre A. Influenza A viruses: new research
823 developments. *Nature reviews. Microbiology*, 9(8), 590-603 (2011).
- 824 45. Burmeister WP, Ruigrok RW, Cusack S. The 2.2 Å resolution crystal
825 structure of influenza B neuraminidase and its complex with sialic acid.
826 *The EMBO journal*, 11(1), 49-56 (1992).
- 827 46. Burnham AJ, Baranovich T, Govorkova EA. Neuraminidase inhibitors
828 for influenza B virus infection: efficacy and resistance. *Antiviral
829 research*, 100(2), 520-534 (2013).
- 830 ** Thorough review on the use and effectiveness of NAIs to treat Influenza B
831 infection.
- 832 47. Williams MA, Lamb RA. Effect of mutations and deletions in a
833 bicistronic mRNA on the synthesis of influenza B virus NB and NA
834 glycoproteins. *Journal of virology*, 63(1), 28-35 (1989).
- 835 48. Betakova T, Nermut MV, Hay AJ. The NB protein is an integral
836 component of the membrane of influenza B virus. *The Journal of
837 general virology*, 77 (Pt 11), 2689-2694 (1996).
- 838 49. Williams MA, Lamb RA. Polylactosaminoglycan modification of a small
839 integral membrane glycoprotein, influenza B virus NB. *Molecular and
840 cellular biology*, 8(3), 1186-1196 (1988).
- 841 50. Demers A, Ran Z, Deng Q *et al.* Palmitoylation is required for
842 intracellular trafficking of influenza B virus NB protein and efficient
843 influenza B virus growth in vitro. *The Journal of general virology*, 95(Pt
844 6), 1211-1220 (2014).
- 845 51. Hatta M, Kawaoka Y. The NB Protein of Influenza B Virus Is Not
846 Necessary for Virus Replication In Vitro. *Journal of virology*, 77(10),
847 6050-6054 (2003).
- 848 52. Shimbo K, Brassard DL, Lamb RA, Pinto LH. Viral and cellular small
849 integral membrane proteins can modify ion channels endogenous to
850 *Xenopus oocytes*. *Biophysical journal*, 69(5), 1819-1829 (1995).

- 851 53. Ewart GD, Sutherland T, Gage PW, Cox GB. The Vpu protein of
852 human immunodeficiency virus type 1 forms cation-selective ion
853 channels. *Journal of virology*, 70(10), 7108-7115 (1996).
- 854 54. Sunstrom NA, Premkumar LS, Premkumar A, Ewart G, Cox GB, Gage
855 PW. Ion channels formed by NB, an influenza B virus protein. *The*
856 *Journal of membrane biology*, 150(2), 127-132 (1996).
- 857 55. Fischer WB, Pitkeathly M, Wallace BA, Forrest LR, Smith GR, Sansom
858 MS. Transmembrane peptide NB of influenza B: a simulation, structure,
859 and conductance study. *Biochemistry*, 39(41), 12708-12716 (2000).
- 860 56. Lamb RA, Pinto LH. Do Vpu and Vpr of human immunodeficiency virus
861 type 1 and NB of influenza B virus have ion channel activities in the
862 viral life cycles? *Virology*, 229(1), 1-11 (1997).
- 863 57. Betakova T, Hay AJ. Evidence that the CM2 protein of influenza C
864 virus can modify the pH of the exocytic pathway of transfected cells.
865 *The Journal of general virology*, 88(Pt 8), 2291-2296 (2007).
- 866 58. McCullers JA, Hoffmann E, Huber VC, Nickerson AD. A single amino
867 acid change in the C-terminal domain of the matrix protein M1 of
868 influenza B virus confers mouse adaptation and virulence. *Virology*,
869 336(2), 318-326 (2005).
- 870 59. Cao S, Jiang J, Li J *et al.* Characterization of the nucleocytoplasmic
871 shuttle of the matrix protein of influenza B virus. *Journal of virology*,
872 88(13), 7464-7473 (2014).
- 873 60. Briedis DJ, Lamb RA, Choppin PW. Sequence of RNA segment 7 of
874 the influenza B virus genome: partial amino acid homology between
875 the membrane proteins (M1) of influenza A and B viruses and
876 conservation of a second open reading frame. *Virology*, 116(2), 581-
877 588 (1982).
- 878 61. Paterson RG, Takeda M, Ohigashi Y, Pinto LH, Lamb RA. Influenza B
879 virus BM2 protein is an oligomeric integral membrane protein
880 expressed at the cell surface. *Virology*, 306(1), 7-17 (2003).
- 881 62. Odagiri T, Hong J, Ohara Y. The BM2 protein of influenza B virus is
882 synthesized in the late phase of infection and incorporated into virions
883 as a subviral component. *The Journal of general virology*, 80 (Pt 10),
884 2573-2581 (1999).

- 885 63. Watanabe S, Imai M, Ohara Y, Odagiri T. Influenza B Virus BM2
886 Protein Is Transported through the trans-Golgi Network as an Integral
887 Membrane Protein. *Journal of virology*, 77(19), 10630-10637 (2003).
- 888 64. Wang J, Pielak RM, McClintock MA, Chou JJ. Solution structure and
889 functional analysis of the influenza B proton channel. *Nature structural*
890 *& molecular biology*, 16(12), 1267-1271 (2009).
- 891 * Solution of NMR structure of BM2 protein.
- 892 65. Imai M, Watanabe S, Ninomiya A, Obuchi M, Odagiri T. Influenza B
893 virus BM2 protein is a crucial component for incorporation of viral
894 ribonucleoprotein complex into virions during virus assembly. *Journal*
895 *of virology*, 78(20), 11007-11015 (2004).
- 896 66. Jackson D, Zurcher T, Barclay W. Reduced incorporation of the
897 influenza B virus BM2 protein in virus particles decreases infectivity.
898 *Virology*, 322(2), 276-285 (2004).
- 899 67. Hatta M, Goto H, Kawaoka Y. Influenza B virus requires BM2 protein
900 for replication. *Journal of virology*, 78(11), 5576-5583 (2004).
- 901 68. Mould JA, Paterson RG, Takeda M *et al.* Influenza B Virus BM2 Protein
902 Has Ion Channel Activity that Conducts Protons across Membranes.
903 *Developmental Cell*, 5(1), 175-184 (2003).
- 904 69. Ma C, Soto CS, Ohgashi Y *et al.* Identification of the pore-lining
905 residues of the BM2 ion channel protein of influenza B virus. *The*
906 *Journal of biological chemistry*, 283(23), 15921-15931 (2008).
- 907 70. Zhang H, Yu H, Wang J *et al.* The BM2 protein of influenza B virus
908 interacts with p53 and inhibits its transcriptional and apoptotic activities.
909 *Molecular and cellular biochemistry*, 403(1-2), 187-197 (2015).
- 910 71. Horvath CM, Williams MA, Lamb RA. Eukaryotic coupled translation of
911 tandem cistrons: identification of the influenza B virus BM2 polypeptide.
912 *The EMBO journal*, 9(8), 2639-2647 (1990).
- 913 72. Hatta M, Kohlmeier CK, Hatta Y, Ozawa M, Kawaoka Y. Region
914 required for protein expression from the stop-start pentanucleotide in
915 the M gene of influenza B virus. *Journal of virology*, 83(11), 5939-5942
916 (2009).
- 917 73. Powell ML, Naphine S, Jackson RJ, Brierley I, Brown TD.
918 Characterization of the termination-reinitiation strategy employed in the

- 919 expression of influenza B virus BM2 protein. *Rna*, 14(11), 2394-2406
920 (2008).
- 921 74. Powell ML, Leigh KE, Poyry TA, Jackson RJ, Brown TD, Brierley I.
922 Further characterisation of the translational termination-reinitiation
923 signal of the influenza B virus segment 7 RNA. *PloS one*, 6(2), e16822
924 (2011).
- 925 75. Stevens MP, Barclay WS. The N-terminal extension of the influenza B
926 virus nucleoprotein is not required for nuclear accumulation or the
927 expression and replication of a model RNA. *Journal of virology*, 72(6),
928 5307-5312 (1998).
- 929 76. Ng AK, Lam MK, Zhang H *et al.* Structural basis for RNA binding and
930 homo-oligomer formation by influenza B virus nucleoprotein. *Journal of*
931 *virology*, 86(12), 6758-6767 (2012).
- 932 77. Shen YF, Chen YH, Chu SY *et al.* E339...R416 salt bridge of
933 nucleoprotein as a feasible target for influenza virus inhibitors.
934 *Proceedings of the National Academy of Sciences of the United States*
935 *of America*, 108(40), 16515-16520 (2011).
- 936 78. Wanitchang A, Narkpuk J, Jongkaewwattana A. Nuclear import of
937 influenza B virus nucleoprotein: involvement of an N-terminal nuclear
938 localization signal and a cleavage-protection motif. *Virology*, 443(1),
939 59-68 (2013).
- 940 79. Sherry L, Smith M, Davidson S, Jackson D. The N terminus of the
941 influenza B virus nucleoprotein is essential for virus viability, nuclear
942 localization, and optimal transcription and replication of the viral
943 genome. *Journal of virology*, 88(21), 12326-12338 (2014).
- 944 80. Reich S, Guilligay D, Pflug A *et al.* Structural insight into cap-snatching
945 and RNA synthesis by influenza polymerase. *Nature*, 516(7531), 361-
946 366 (2014).
- 947 ****First crystal structure of Influenza heterotrimeric polymerase complex.**
- 948 81. Deng Q, Wang D, Xiang X *et al.* Nuclear localization of influenza B
949 polymerase proteins and their binary complexes. *Virus research*,
950 156(1-2), 49-53 (2011).
- 951 82. Wakai C, Iwama M, Mizumoto K, Nagata K. Recognition of cap
952 structure by influenza B virus RNA polymerase is less dependent on

- the methyl residue than recognition by influenza A virus polymerase. *Journal of virology*, 85(15), 7504-7512 (2011).
83. Liu Y, Yang Y, Fan J, He R, Luo M, Zheng X. The crystal structure of the PB2 cap-binding domain of influenza B virus reveals a novel cap recognition mechanism. *The Journal of biological chemistry*, 290(14), 9141-9149 (2015).
 84. Cauldwell AV, Long JS, Moncorge O, Barclay WS. Viral determinants of influenza A virus host range. *The Journal of general virology*, 95(Pt 6), 1193-1210 (2014).
 85. Dauber B, Heins G, Wolff T. The Influenza B Virus Nonstructural NS1 Protein Is Essential for Efficient Viral Growth and Antagonizes Beta Interferon Induction. *Journal of virology*, 78(4), 1865-1872 (2004).
 86. Hai R, Martinez-Sobrido L, Fraser KA, Ayllon J, Garcia-Sastre A, Palese P. Influenza B virus NS1-truncated mutants: live-attenuated vaccine approach. *Journal of virology*, 82(21), 10580-10590 (2008).
 87. Guan R, Ma LC, Leonard PG *et al.* Structural basis for the sequence-specific recognition of human ISG15 by the NS1 protein of influenza B virus. *Proceedings of the National Academy of Sciences of the United States of America*, 108(33), 13468-13473 (2011).
 88. Schneider J, Dauber B, Melen K, Julkunen I, Wolff T. Analysis of influenza B Virus NS1 protein trafficking reveals a novel interaction with nuclear speckle domains. *Journal of virology*, 83(2), 701-711 (2009).
 89. Noah DL, Twu KY, Krug RM. Cellular antiviral responses against influenza A virus are countered at the posttranscriptional level by the viral NS1A protein via its binding to a cellular protein required for the 3' end processing of cellular pre-mRNAs. *Virology*, 307(2), 386-395 (2003).
 90. Donelan NR, Dauber B, Wang X, Basler CF, Wolff T, Garcia-Sastre A. The N- and C-terminal domains of the NS1 protein of influenza B virus can independently inhibit IRF-3 and beta interferon promoter activation. *Journal of virology*, 78(21), 11574-11582 (2004).
 91. Dauber B, Schneider J, Wolff T. Double-stranded RNA binding of influenza B virus nonstructural NS1 protein inhibits protein kinase R but

- 986 is not essential to antagonize production of alpha/beta interferon.
987 *Journal of virology*, 80(23), 11667-11677 (2006).
- 988 92. Iwasaki A, Pillai PS. Innate immunity to influenza virus infection. *Nature*
989 *reviews. Immunology*, 14(5), 315-328 (2014).
- 990 93. Yuan W, Krug RM. Influenza B virus NS1 protein inhibits conjugation of
991 the interferon (IFN)-induced ubiquitin-like ISG15 protein. *The EMBO*
992 *journal*, 20(3), 362-371 (2001).
- 993 94. Sridharan H, Zhao C, Krug RM. Species specificity of the NS1 protein
994 of influenza B virus: NS1 binds only human and non-human primate
995 ubiquitin-like ISG15 proteins. *The Journal of biological chemistry*,
996 285(11), 7852-7856 (2010).
- 997 95. Paragas J, Talon J, O'Neill RE, Anderson DK, Garcia-Sastre A, Palese
998 P. Influenza B and C virus NEP (NS2) proteins possess nuclear export
999 activities. *Journal of virology*, 75(16), 7375-7383 (2001).
- 1000 96. Imai M, Watanabe S, Odagiri T. Influenza B virus NS2, a nuclear
1001 export protein, directly associates with the viral ribonucleoprotein
1002 complex. *Archives of virology*, 148(10), 1873-1884 (2003).
- 1003 97. Barclay WS, Palese P. Influenza B viruses with site-specific mutations
1004 introduced into the HA gene. *Journal of virology*, 69(2), 1275-1279
1005 (1995).
- 1006 98. Zhou B, Lin X, Wang W *et al.* Universal influenza B virus genomic
1007 amplification facilitates sequencing, diagnostics, and reverse genetics.
1008 *Journal of clinical microbiology*, 52(5), 1330-1337 (2014).
- 1009 99. Hoffmann E, Mahmood K, Yang CF, Webster RG, Greenberg HB,
1010 Kemble G. Rescue of influenza B virus from eight plasmids.
1011 *Proceedings of the National Academy of Sciences of the United States*
1012 *of America*, 99(17), 11411-11416 (2002).
- 1013 100. Jackson D, Cadman A, Zurcher T, Barclay WS. A Reverse Genetics
1014 Approach for Recovery of Recombinant Influenza B Viruses Entirely
1015 from cDNA. *Journal of virology*, 76(22), 11744-11747 (2002).
- 1016 101. Baker SF, Nogales A, Finch C *et al.* Influenza A and B virus intertypic
1017 reassortment through compatible viral packaging signals. *Journal of*
1018 *virology*, 88(18), 10778-10791 (2014).

- 1019 102. Loo YM, Fornek J, Crochet N *et al.* Distinct RIG-I and MDA5 signaling
1020 by RNA viruses in innate immunity. *Journal of virology*, 82(1), 335-345
1021 (2008).
- 1022 103. Dauber B, Martinez-Sobrido L, Schneider J *et al.* Influenza B virus
1023 ribonucleoprotein is a potent activator of the antiviral kinase PKR.
1024 *PLoS pathogens*, 5(6), e1000473 (2009).
- 1025 104. Kim MJ, Latham AG, Krug RM. Human influenza viruses activate an
1026 interferon-independent transcription of cellular antiviral genes: outcome
1027 with influenza A virus is unique. *Proceedings of the National Academy*
1028 *of Sciences of the United States of America*, 99(15), 10096-10101
1029 (2002).
- 1030 105. Osterlund P, Strengell M, Sarin LP *et al.* Incoming influenza A virus
1031 evades early host recognition, while influenza B virus induces
1032 interferon expression directly upon entry. *Journal of virology*, 86(20),
1033 11183-11193 (2012).
- 1034 *This study pinpoints interesting differences in innate recognition between FluA
1035 and FluB.
- 1036 106. Veckman V, Osterlund P, Fagerlund R, Melen K, Matikainen S,
1037 Julkunen I. TNF-alpha and IFN-alpha enhance influenza-A-virus-
1038 induced chemokine gene expression in human A549 lung epithelial
1039 cells. *Virology*, 345(1), 96-104 (2006).
- 1040 107. Ichinohe T, Pang IK, Iwasaki A. Influenza virus activates
1041 inflammasomes via its intracellular M2 ion channel. *Nature*
1042 *immunology*, 11(5), 404-410 (2010).
- 1043 108. Ponniah S, Doherty PC, Eichelberger M. Selective response of gamma
1044 delta T-cell hybridomas to orthomyxovirus-infected cells. *Journal of*
1045 *virology*, 70(1), 17-22 (1996).
- 1046 109. Nunes B, Pechirra P, Coelho A, Ribeiro C, Arraiolos A, Rebelo-de-
1047 Andrade H. Heterogeneous selective pressure acting on influenza B
1048 Victoria- and Yamagata-like hemagglutinins. *Journal of molecular*
1049 *evolution*, 67(4), 427-435 (2008).
- 1050 110. Wohlbold TJ, Nachbagauer R, Xu H *et al.* Vaccination with adjuvanted
1051 recombinant neuraminidase induces broad heterologous, but not

- heterosubtypic, cross-protection against influenza virus infection in mice. *mBio*, 6(2), e02556 (2015).
111. Levandowski RA, Gross PA, Weksler M, Staton E, Williams MS, Bonelli J. Cross-reactive antibodies induced by a monovalent influenza B virus vaccine. *Journal of clinical microbiology*, 29(7), 1530-1532 (1991).
 112. Janjua NZ, Skowronski DM, De Serres G *et al.* Estimates of influenza vaccine effectiveness for 2007-2008 from Canada's sentinel surveillance system: cross-protection against major and minor variants. *The Journal of infectious diseases*, 205(12), 1858-1868 (2012).
 113. Englund JA, Walter EB, Gbadebo A, Monto AS, Zhu Y, Neuzil KM. Immunization with trivalent inactivated influenza vaccine in partially immunized toddlers. *Pediatrics*, 118(3), e579-585 (2006).
 114. Walter EB, Neuzil KM, Zhu Y *et al.* Influenza vaccine immunogenicity in 6- to 23-month-old children: are identical antigens necessary for priming? *Pediatrics*, 118(3), e570-578 (2006).
 115. Ohmit SE, Thompson MG, Petrie JG *et al.* Influenza vaccine effectiveness in the 2011-2012 season: protection against each circulating virus and the effect of prior vaccination on estimates. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*, 58(3), 319-327 (2014).
 116. Skowronski DM, Hottes TS, De Serres G *et al.* Influenza B/Victoria antigen induces strong recall of B/Yamagata but lower B/Victoria response in children primed with two doses of B/Yamagata. *The Pediatric infectious disease journal*, 30(10), 833-839 (2011).
 117. Dreyfus C, Laursen NS, Kwaks T *et al.* Highly conserved protective epitopes on influenza B viruses. *Science*, 337(6100), 1343-1348 (2012).
- **First report of highly conserved and cross-reactive antibody epitopes of Influenza B, includes crystal structures.
118. Yasugi M, Kubota-Koketsu R, Yamashita A *et al.* Human monoclonal antibodies broadly neutralizing against influenza B virus. *PLoS pathogens*, 9(2), e1003150 (2013).
 119. Gravel C, Li C, Wang J *et al.* Qualitative and quantitative analyses of virtually all subtypes of influenza A and B viral neuraminidases using

- antibodies targeting the universally conserved sequences. *Vaccine*, 28(36), 5774-5784 (2010).
120. Doyle TM, Li C, Bucher DJ *et al.* A monoclonal antibody targeting a highly conserved epitope in influenza B neuraminidase provides protection against drug resistant strains. *Biochemical and biophysical research communications*, 441(1), 226-229 (2013).
121. McMichael AJ, Gotch FM, Noble GR, Beare PA. Cytotoxic T-cell immunity to influenza. *The New England journal of medicine*, 309(1), 13-17 (1983).
122. Gras S, Kedzierski L, Valkenburg SA *et al.* Cross-reactive CD8+ T-cell immunity between the pandemic H1N1-2009 and H1N1-1918 influenza A viruses. *Proceedings of the National Academy of Sciences of the United States of America*, 107(28), 12599-12604 (2010).
123. Sridhar S, Begom S, Bermingham A *et al.* Cellular immune correlates of protection against symptomatic pandemic influenza. *Nature medicine*, 19(10), 1305-1312 (2013).
124. Quinones-Parra S, Grant E, Loh L *et al.* Preexisting CD8+ T-cell immunity to the H7N9 influenza A virus varies across ethnicities. *Proceedings of the National Academy of Sciences of the United States of America*, 111(3), 1049-1054 (2014).
125. Wang Z, Wan Y, Qiu C *et al.* Recovery from severe H7N9 disease is associated with diverse response mechanisms dominated by CD8(+) T cells. *Nature communications*, 6, 6833 (2015).
126. Hillaire ML, van Trierum SE, Kreijtz JH *et al.* Cross-protective immunity against influenza pH1N1 2009 viruses induced by seasonal influenza A (H3N2) virus is mediated by virus-specific T-cells. *The Journal of general virology*, 92(Pt 10), 2339-2349 (2011).
127. Epstein SL, Lo CY, Mispion JA, Bennink JR. Mechanism of protective immunity against influenza virus infection in mice without antibodies. *Journal of immunology*, 160(1), 322-327 (1998).
- *First and, to our knowledge, only study on in vivo role of T cell during FluB infection.
128. P A Robbins LAL, P Rota, J Santos-Aguado, J Rothbard, A J McMichael, and J L Strominger. Comparison between two peptide

1120 epitopes presented to cytotoxic T lymphocytes by HLA-A2. Evidence
 1121 for discrete locations within HLA-A2. *J Immunology*, 143(4089), 103
 1122 (1989).

1123 129. Robbins PA, Rota PA, Shapiro SZ. A broad cytotoxic T lymphocyte
 1124 response to influenza type B virus presented by multiple HLA
 1125 molecules. *International immunology*, 9(6), 815-823 (1997).

1126 130. van de Sandt CE, Dou Y, Vogelzang-van Trierum SE *et al.* Influenza B
 1127 virus-specific CD8⁺ T lymphocytes strongly cross-react with viruses of
 1128 the opposing influenza B lineage. *The Journal of general virology*,
 1129 (2015).

1130 131. Boon AC, De Mutsert G, Fouchier RA, Sintnicolaas K, Osterhaus AD,
 1131 Rimmelzwaan GF. Preferential HLA usage in the influenza virus-
 1132 specific CTL response. *Journal of immunology*, 172(7), 4435-4443
 1133 (2004).

1134 132. Masanori Terajima JABB, Mary Dawn T Co and Francis A Ennis.
 1135 Cross-reactive human B cell and T cell epitopes between influenza A
 1136 and B viruses. *Virology Journal* 10(244) (2013).

1137 133. Epstein SL, Lo CY, Misplon JA *et al.* Mechanisms of heterosubtypic
 1138 immunity to lethal influenza A virus infection in fully immunocompetent,
 1139 T cell-depleted, beta2-microglobulin-deficient, and J chain-deficient
 1140 mice. *Journal of immunology*, 158(3), 1222-1230 (1997).

1141 134. Benton KA, Misplon JA, Lo CY, Brutkiewicz RR, Prasad SA, Epstein
 1142 SL. Heterosubtypic Immunity to Influenza A Virus in Mice Lacking IgA,
 1143 All Ig, NKT Cells, or T Cells. *The Journal of Immunology*, 166(12),
 1144 7437-7445 (2001).

1145 135. Le J, Orff EJ, Fulvini AA *et al.* Development of high yield reassortants
 1146 for influenza type B viruses and analysis of their gene compositions.
 1147 *Vaccine*, 33(7), 879-884 (2015).

1148 136. Burnham AJ, Armstrong J, Lowen AC, Webster RG, Govorkova EA.
 1149 Competitive fitness of influenza B viruses with neuraminidase inhibitor-
 1150 resistant substitutions in a coinfection model of the human airway
 1151 epithelium. *Journal of virology*, 89(8), 4575-4587 (2015).

1152 137. Burnham AJ, Baranovich T, Marathe BM, Armstrong J, Webster RG,
 1153 Govorkova EA. Fitness costs for Influenza B viruses carrying

neuraminidase inhibitor-resistant substitutions: underscoring the importance of E119A and H274Y. *Antimicrobial agents and chemotherapy*, 58(5), 2718-2730 (2014).

138. Farrukee R, Leang SK, Butler J *et al.* Influenza viruses with B/Yamagata- and B/Victoria-like neuraminidases are differentially affected by mutations that alter antiviral susceptibility. *The Journal of antimicrobial chemotherapy*, 70(7), 2004-2012 (2015).

139. A. D. M. E. Osterhaus GFR, B. E. E. Martina, T. M. Bestebroer, and R. A. M. Fouchier. Influenza B Virus in Seals. *Science*, 288(5468), 1051-1053 (2000).

**First record of Influenza B in a non-human host.

140. Andrea Blanc DR, Mario Clara, Federico Achaval, Alfredo Le Bas, and Juan Arbiza Serologic Evidence of Influenza A and B Viruses in South American Fur Seals (*Arctocephalus australis*). *Journal of Wildlife Diseases*, 45(2), 519-521 (2009).

141. Bodewes R, van de Bildt MW, van Elk CE *et al.* No serological evidence that harbour porpoises are additional hosts of influenza B viruses. *PloS one*, 9(2), e89058 (2014).

142. Ran Z, Shen H, Lang Y *et al.* Domestic pigs are susceptible to infection with influenza B viruses. *Journal of virology*, 89(9), 4818-4826 (2015).

143. Leyva-Grado VH, Mubareka S, Krammer F, Cardenas WB, Palese P. Influenza virus infection in guinea pigs raised as livestock, Ecuador. *Emerging infectious diseases*, 18(7), 1135-1138 (2012).

144. Pica N, Chou YY, Bouvier NM, Palese P. Transmission of influenza B viruses in the guinea pig. *Journal of virology*, 86(8), 4279-4287 (2012).

*Establishment of a FluB transmission system in guinea pigs.

145. Elderfield RA, Parker L, Stilwell P, Roberts KL, Schepelmann S, Barclay WS. Ferret airway epithelial cell cultures support efficient replication of influenza B virus but not Mumps virus. *The Journal of general virology*, (2015).

Acknowledgements

1187 MK is supported by a University of Melbourne International Fee Remission
1188 Scholarship (MIFRS) and a University of Melbourne International Research
1189 Scholarships (MIRS). KK is an NHMRC CDF Level2 Fellow. The authors
1190 declare no conflict of interest.

1191

1192

1193 Figure Legends

1194

1195 Figure 1. Overview of Influenza B viruses. A) Schematic of FluB virus. B)
1196 Genome organization of FluB. C) Non-coding sequences of FluB.

1197

1198 Table 1. Vaccines strains of FluB in the Northern and Southern Hemispheres.

1199 QV=Quadrivalent Vaccine, N/A= Not applicable.

1200

1201 Table 2. Comparison of FluA and FluB.

1202

A)

B)

C)

Figure 1 FluB Virus and Genome Organization

Northern Hemisphere

Flu Season	Strain	Lineage of TIV strain	To be included in QIV
1998-1999	B/Beijing/184/93	Yamagata	N/A
1999-2000	B/Beijing/184/93 or B/Shangdong/7/97	Yamagata	N/A
2000-2001	B/Beijing/184/93	Yamagata	N/A
2001-2002	B/Sichuan/379/99	Victoria	N/A
2002-2003	B/Hong Kong/330/2001	Victoria	N/A
2003-2004	B/Hong Kong/330/2001	Victoria	N/A
2004-2005	B/Shanghai/361/2002	Yamagata	N/A
2005-2006	B/Shanghai/361/2002	Yamagata	N/A
2006-2007	B/Malaysia/2506/2004	Victoria	N/A
2007-2008	B/Malaysia/2506/2004	Victoria	N/A
2008-2009	B/Florida/4/2006	Yamagata	N/A
2009-2010	B/Brisbane/60/2008	Victoria	N/A
2010-2011	B/Brisbane/60/2008	Victoria	N/A
2011-2012	B/Brisbane/60/2008	Victoria	N/A
2012-2013	B/Wisconsin/1/2010	Yamagata	B/Brisbane/60/2008
2013-2014	B/Massachusetts/2/2012	Yamagata	B/Brisbane/60/2008
2014-2015	B/Massachusetts/2/2012	Yamagata	B/Brisbane/60/2008
2015-2016	B/Phuket/3073/2013	Yamagata	B/Brisbane/60/2008

Southern Hemisphere

Flu Season	Strain	Lineage of TIV strain	To be included in QIV
1999	B/Beijing/184/93	Yamagata	N/A
2000	B/Beijing/184/93 or B/Shangdong/7/97	Yamagata	N/A
2001	B/Sichuan/379/99	Victoria	N/A
2002	B/Sichuan/379/99	Victoria	N/A
2003	B/Hong Kong/330/2001	Victoria	N/A
2004	B/Hong Kong/330/2001	Victoria	N/A
2005	B/Shanghai/361/2002	Yamagata	N/A
2006	B/Malaysia/2506/2004	Victoria	N/A
2007	B/Malaysia/2506/2004	Victoria	N/A
2008	B/Florida/4/2006	Yamagata	N/A
2009	B/Florida/4/2006	Yamagata	N/A
2010	B/Brisbane/60/2008	Victoria	N/A
2011	B/Brisbane/60/2008	Victoria	N/A
2012	B/Brisbane/60/2008	Victoria	N/A
2013	B/Wisconsin/1/2010	Yamagata	B/Brisbane/60/2008
2014	B/Massachusetts/2/2012	Yamagata	B/Brisbane/60/2008
2015	B/Phuket/3073/2013	Yamagata	B/Brisbane/60/2008

Table 1 FluB Vaccines Strains in Northern and Southern Hemispheres

FluA	FluB
<u>Phylogenetics</u>	
<ul style="list-style-type: none"> Subtyped by HA and NA (e.g. H1N1, H3N2) 	<ul style="list-style-type: none"> Two lineages according to HA: Victoria and Yamagata
<u>Epidemiology</u>	
<ul style="list-style-type: none"> More severe in children and elderly 	<ul style="list-style-type: none"> More severe in children and adolescents
<u>Clinical Presentation</u>	
<ul style="list-style-type: none"> Symptoms: soar throat, coughing, fever, nasal discharge Avian strains lead to severe disease and high mortality rates 	<ul style="list-style-type: none"> Symptoms: soar throat, coughing, fever, nasal discharge Can cause sever complications in children
<u>Virology</u>	
<ul style="list-style-type: none"> 8 segments, >13 known proteins Unique proteins: eg. PB1-F2, PB1-N40, PA-X 	<ul style="list-style-type: none"> 8 segments, 11 known proteins Unique proteins: NB
<u>Immunology</u>	
<ul style="list-style-type: none"> Generally, strong IFN$\alpha\beta$ and cytokine response HA is the major antibody target M1 and NP are the major CD8⁺ T cell antigens Two strains included in annual vaccine (one H3N2 and one H1N1) 	<ul style="list-style-type: none"> Strong IFN$\alpha\beta$ and cytokine response, details unclear HA is the major antibody target Major CD8⁺ T cell antigens are unknown One or two strains in annual vaccine depending on formulation
<u>Antivirals</u>	
<ul style="list-style-type: none"> Adamantanes used are effective NAIs effective, but resistance detected 	<ul style="list-style-type: none"> Adamantanes are ineffective NAIs effective, but resistance detected, in some cases without apparent fitness cost
<u>Host Range</u>	
<ul style="list-style-type: none"> Multiple animal reservoirs and possible animal hosts 	<ul style="list-style-type: none"> No established animal reservoirs

Table 2. Comparison of FluA and FluB