Extubation to facilitate mother-baby bonding in refractory acute respiratory distress syndrome

Graeme MacLaren^{1,2}

Kollengode R Ramanathan¹

Vitaly Sorokin¹

1: Cardiothoracic ICU and Department of Cardiac, Thoracic, and Vascular Surgery,
National University Health System, Singapore

2: Paediatric Intensive Care Unit, Department of Paediatrics, University of Melbourne,
The Royal Children's Hospital, Melbourne, Australia

Address for correspondence:

gmaclaren@iinet.net.au

Cardiothoracic ICU, National University Hospital, 5 Lower Kent Ridge Rd, Singapore
119074

Ph: (+65) 67725275

Fax: (+65) 67766475

Key words: extracorporeal life support, extracorporeal membrane oxygenation, awake,
influenza A(H1N1)

A 25 year old pregnant woman presented at 37 weeks gestation with worsening respiratory distress. A diagnosis of influenza A(H1N1) pneumonitis with acute respiratory distress syndrome (ARDS) was established. Following tracheal intubation, emergency Cesarean section, and delivery of a healthy baby boy, she developed progressive, severe respiratory failure with a $\text{PaO}_2/\text{FiO}_2$ ratio of 50. She was referred to our centre and cannulated onto venovenous extracorporeal membrane oxygenation (ECMO).

The ARDS persisted and the patient was unable to be weaned from ECMO after 7 days of extracorporeal support (Fig 1). However, influenza was no longer detectable on repeat polymerase chain reaction testing and she no longer posed an infectious risk to those in contact with her. In order to facilitate communication and to establish maternal-infant bonding, the patient was weaned from mechanical ventilation and extubated (Fig 2). Thereafter, she received daily physiotherapy and continuous positive airway pressure ventilation via face mask when asleep. The remainder of her intensive care stay was uncomplicated and she was successfully decannulated after 14 days of ECMO. She was discharged well from hospital 5 days later.

The authors declare that they have no conflict of interest.

The need for formal ethics application and review was waived by the institutional review board.

Acknowledgements:

We are grateful to the patient and her family for giving consent for publication.