The diagnostic value of a single measurement of superior vena cava flow in the first 24 hours of life in very preterm infants.

James R Holberton^{1,2}, Sandra M Drew¹, Rintaro Mori^{2,3}, Kai König^{1*}

1 Mercy Hospital for Women, Department of Paediatrics, Melbourne, Australia

2 London School of Hygiene and Tropical Medicine, London, UK

3 Collaboration for Research in Global Women's and Children's Health, Tokyo, Japan

* Corresponding author:

Kai König

Mercy Hospital for Women

Department of Paediatrics

163 Studley Road

Melbourne 3084

Victoria, Australia

Email: kaikonig@gmail.com

Phone: 0061-3-8458 4444

Fax: 0061-3-9459 5389

Conflict of interest: the authors declare no conflict of interest.

Funding: No specific funding.

Abstract

Low superior vena cava (SVC) flow has been associated with intraventricular haemorrhage (IVH) in very preterm infants. We studied the diagnostic value of a single measurement of SVC flow within the first 24 hours of life in very preterm infants and its association with occurrence or extension of IVH in a setting of limited availability of neonatal echocardiography. Preterm infants who were born at less than 30 weeks gestation, and who had an echocardiogram within 24 hours after birth were eligible.

Baseline, clinical, and ultrasound data were collected. 165 preterm infants were included. Low SVC flow (<41 ml/kg/min) occurred in 6 infants, and was associated with severe IVH and extension of IVH although this was not significant after adjusting for confounders. The only independently associated variable with low SVC flow was admission temperature (Odds Ratio 0.27, $p=0.001$). A review of SVC flow values shows values are higher now than initially reported.

Conclusion: This study does not show an association of low SVC flow and severe IVH or extension of IVH after adjusting for confounders as a single measurement of SVC flow did not add any diagnostic value in this cohort. Thus, the exact role of SVC flow measurements in the circulatory assessment of preterm infants remains to be elucidated. However, admission temperature may have an effect on systemic blood flow in very preterm infants.

Keywords:

Preterm infants, intraventricular haemorrhage, neonatal functional echocardiography, superior vena cava flow

Introduction

The newborn circulation represents a unique and complex situation due to its transitional character during the first days of life, including changing systemic and pulmonary vascular resistances and the existence of intra- and extracardiac shunts. Conventional parameters such as blood pressure, heart rate, or capillary refill time only give limited information, and do not correlate well with echocardiographically assessed systemic blood flow [10]. Therefore the use of functional echocardiography in the assessment of cardiovascular status in critically ill newborns has become more popular in recent years [3].

A decade ago, Kluckow and Evans published a new technique to assess systemic blood flow in very preterm infants. Using Doppler, they performed serial measurements of blood flow in the superior vena cava (SVC) as surrogate marker for systemic blood flow [4]. In their initial and subsequent cohorts using multiple measurements in the first 24 hours of life, their group demonstrated a strong association of low SVC flow and occurrence of intraventricular haemorrhage (IVH) in preterm babies born less than 30 weeks gestation [5].

Functional neonatal echocardiography requires sufficient training to be performed independently by neonatologists and is therefore still a limited resource in many neonatal intensive care units (NICU).

We aimed to investigate if a single measurement of SVC flow within the first 24 hours of life in preterm infants less than 30 weeks gestation is associated with occurrence or extension IVH in a setting of limited neonatal echocardiography availability.

Materials and Methods

This was a single-centre retrospective study performed in the NICU of the Mercy Hospital for Women, a tertiary perinatal centre in Melbourne, Australia, with approximately 180-200 very low birthweight infants admitted per year. The study was approved by both the institutional human research ethics committee and the ethics committee of the London School of Health and Tropical Medicine. Informed written consent was obtained from the parents of studied infants.

Patients

Infants born before 30 weeks gestation, and who were admitted to the NICU within 24 hours of age were eligible. Infants with major chromosomal, congenital or cardiac anomalies were excluded.

Echocardiography

Echocardiograms were requested at the treating neonatologist's discretion. SVC flow was determined by the described method of Kluckow and Evans [4]. Low SVC flow was defined as SVC flow less than 41 ml/kg/min [11].

All scans were performed by a single experienced echocardiographer (S. M. D.) who has over 25 years of experience in paediatric and neonatal echocardiography, and is an accredited supervisor in neonatal echocardiography for the 'Certificate in Clinician Performed Ultrasound' of the Australasian Society for Ultrasound in Medicine. The echocardiographer's regular working hours during the study period were half days on four days per week, with an on-call service after hours and on the other days. All echocardiogram recordings were reviewed and approved by a consultant paediatric cardiologist. The findings in the study population did not change following the cardiologist's review.

Cranial ultrasound

Cranial ultrasounds were performed at the treating neonatologist's request otherwise they were routinely performed on day 3, day 14 and day 42 of life by experienced sonographers who were not involved in the study. IVH was classified according to Papile's classification [15]. Infants diagnosed with an IVH usually have daily scans until the IVH appears to be stable. In the event of an IVH grade III or IV, once stable the scan interval would then change to every 2-3 days, and later weekly to assess for post haemorrhagic hydrocephalus. IVH was considered as "extended" if the grade of IVH was higher on any later scan than that obtained on the scan closest to the echocardiogram. All ultrasound scans were reported by consultant radiologists.

Clinical data

Collected data included: gestation; birthweight; sex; temperature on admission; first pH; first base excess; first lactate; the Clinical Risk Index for Babies II (CRIB II) score [18]; type of delivery; date and time and place of birth; time of echocardiogram; respiratory support and mean airway pressure at

time of echocardiogram; cardiovascular management both to the time of echocardiogram and in the first 24 hours of life; and survival to nursery discharge. Ventilator and cardiovascular information was recorded at the time of the echocardiogram and other information obtained from note review.

Statistical analysis

Initial data analysis consisted of comparing the group in our cohort who did have SVC flow measured with those who did not. T test for categorical data (with Fischer correction where appropriate), and Chi square for proportions was utilised. Analysis of data for the SVC cohort consisted of linear regression when assessing SVC flow as a continuous variable, whilst T test was used for binary variables. Co-variables for the models for severe IVH (grade III or IV) and extension of IVH were selected upon both clinical and statistical significance on univariate analysis in a logistic regression model and removing variables stepwise. Statistical significance was taken at the $P < 0.05$ level. A power calculation showed that a power at 0.8 with a significance level of 0.05 to detect a difference of low SVC flow from anticipated 38% (based on the original report of the Sydney group in 2000 [5]) to 19%, 165 babies were required.

Results

Over the 2 ½ year study period, 322 infants born less than 30 weeks gestation were admitted to NICU within 24 hours of birth. Four patients were excluded for chromosomal (2) or cardiac (1) or congenital (1) abnormalities, leaving 318 eligible infants. 165 (52%) infants had an echocardiography including SVC flow measurement within the first 24 hours of postnatal life and were included in the study.

Baseline characteristics demonstrate that babies who had echocardiographic assessment were smaller and sicker, as evidenced by lower gestation, lighter birthweight, and an increased request for echocardiography after hours. They also had a trend to higher CRIB II scores, IVH and death (table 1).

SVC flow

SVC flow was measured in 165 infants, 88 infants were male (53.3%); average echocardiography examination time was at 12.8 hours of life. The median SVC flow of all babies in the study within the

first 24 hours of life was 114 ml/kg/min (interquartile range 82, 150). Six infants had low SVC flow of less than 41 ml/kg/min. The median SVC flow of the remaining 159 babies was 116 ml/kg/min.

Low SVC flow was associated with CRIB II score, temperature at admission, earliest base excess, earliest lactate, volume of fluid bolus given to time of echocardiogram, and mean airway pressure at time of echocardiography. No association was found for gestation and birthweight (table 2). The only independently associated variable with low SVC flow was temperature on arrival into the unit, with an Odds Ratio (OR) (P values) of 0.27 (0.001) per degree centigrade. Given the low incidence of low SVC flows, we re-analysed the IVH outcomes looking at the group of infants with the lowest quartile of SVC flow. This analysis did not show an independent association between lowest quartile of SVC flow and extension of IVH, severe IVH or death (data not shown).

IVH

33 (20%) of 165 infants had an IVH. Of these, 11 (33.3%) were grade 1, none was grade 2, 9 (27.3%) were grade 3, and 13 (39.4%) were grade 4. 20 infants had extending IVH. Variables associated with severe IVH are shown in table 3.

A final model for either grade 3 or 4 IVH starting with the clinically and significantly associated variables was generated as described in the methods section. The final result was a model with only two independent variables: (I) gestation and (II) fluid bolus volume to time of echocardiogram remaining with adjusted ORs (P value) of 0.60 (0.001) per week of increasing gestation and 1.06 (0.002) per ml/kg of fluid respectively. Low SVC flow after adjusting for gestation and fluid bolus was no longer significant.

Similar as above, a final model for extended IVH starting with the clinically and significantly associated variables was generated. The final result was a model with three independent variables: (I) gestation, (II) fluid bolus to time of echocardiogram and (III) earliest base excess remaining with adjusted ORs (P value) of: 0.63 (0.003) per week of increasing gestation, 1.04 (0.04) per ml/kg of fluid and 0.86 (0.03) per decrease in mmol/L of base excess respectively. Low SVC flow after adjusting for gestation, fluid bolus and earliest base excess was not significant.

SVC flow and IVH

Comparing median SVC flows between groups with and without IVH did not show statistical significance (93 ml/kg/min for infants with grade 3 or 4 P/IVH vs. 117 ml/kg/min for infants without IVH; $p=0.06$), neither did the groups of babies with any IVH vs. no IVH (102 ml/kg/min vs. 117 ml/kg/min; $p=0.28$). Infants who had extension of IVH had a median SVC flow of 103.5 ml/kg/min, similar to the group of infants with any IVH.

Circulatory management

Given our higher SVC flows compared to initial reports, we looked at cardiovascular management in the first 24 hours of life of all study infants. In the first 24 hrs routine fluids delivered averaged 56 ml/kg/day. Fluid boluses were given to 124 infants (75%), of an average volume of 21.5 ml/kg. Inotropes were given to 128 infants, 74 receiving a single inotrope (usually dobutamine), 40 infants receiving two (usually dobutamine, then dopamine) and 14 infants receiving 3 (all dobutamine, dopamine and adrenaline).

109 infants were receiving inotropes at the time of the echocardiogram and 124 infants had received a fluid bolus prior to echocardiography of an average of 19.7 ml/kg.

Discussion

Since its introduction in 2000, SVC flow measurements in preterm infants have been studied by several research groups [1, 2, 4, 5, 7-14, 16, 17, 22, 23]. Our study confirms previous results that low SVC flow is on univariate analysis associated with grade 3 or 4 IVH and with extension of IVH, although it was not an independently significant variable after adjusting for confounders in this cohort.

This study describes the single largest cohort of SVC flow measurements in preterm infants to date. Although retrospective in its design it represents a 'real life' scenario given that the ability to do serial echocardiographic measurements at predetermined times is not commonly available in neonatal units, and is therefore limited to a research setting. To illustrate this, we calculated the number of echocardiograms to be performed, given the admission of 127 infants < 30 weeks gestation per year to our unit. To assess SVC flow at 5, 12, and 24 hours of age according to the original study, 381 scans

would be required. If the time of birth was similar to our 2.5 year cohort, then 298 of those scans would have been scheduled out of business hours, including 136 (one every 2.7 days) required between 10pm and 6am and one scan every second weekend, causing a tremendous extra workload, particularly for a single scanner. While serial assessment with functional neonatal echocardiography is desirable, it is only feasible in selected cases.

In this study, only one examiner performed echocardiograms. While this limited us in the total number of scans due to her availability, it strengthens the study results because inter-observer variability is excluded. In their original article, SVC flow measurements were performed by two operators and had differences of up to 57%, and 10% of paired readings were different by more than 33% [4]. Inter-observer variability has been discussed more recently by Groves and co-workers in their study. The 95% confidence limits of the inter-observer repeatability coefficients in their study were wide (35-136ml/kg/min), making interpretation of cut off values particularly difficult [1].

The median SVC flow of all babies in our study was 114 ml/kg/min, with an average time of echocardiography at 12.8 hours of life. This is similar to West and co-workers who reported an average SVC flow of 97 at 12 hours of age, and Groves and colleagues who reported a median flow of 101 ml/kg/min at 12 hours of age in their study [1, 23]. In their original study, Kluckow and Evans reported a median SVC flow of 56 ml/kg/min in their original cohort at 12 hours of life, and they defined a cut-off normal value of above 34 ml/kg/min at that time of echocardiography [4, 5].

Table 4 shows the averaged SVC flows for the first 24 hours of life, in all publications in which it has been reported in enough detail to be calculated [1, 2, 4, 5, 9, 12, 16, 17, 23]. Average SVC flows were calculated by obtaining the average of all readings before 24 hours of age. Two studies did not report SVC flow data in detail [7, 8] though a proportion of these infants have data reported in a subsequent study [9]. One study reported low SVC flow values only and was not included in the table [13].

Because our study had a low incidence of infants with low SVC flow as per the original definition, we re-analysed the IVH outcomes for those infants with the lowest quartile of SVC flow. There was no independent association between lowest quartile of SVC flow and severe IVH, extension of IVH, or death.

Although speculative, the increase in average SVC flow over the years since its introduction may be a result of improved neonatal circulatory management of very preterm babies over a decade's time. It is noteworthy that in a recent study from the Sydney group comparing milrinone versus placebo for prevention of low SVC flow, the number of infants with low SVC flow had decreased in both study arms: to 17% in the milrinone arm and 19% in the placebo arm. The authors also proposed changes in NICU management over time as the underlying reason for the decrease in low SVC numbers. Another reason for the lower than expected incidence in low SVC flow is the number of echocardiograms performed. The Sydney group's studies included three scans within the first 24 hours. Any infant with one measurement of low SVC flow, is classified as "low flow", so serial measurements would pick up more low SVC flows and therefore increase the incidence [5,11,17]. While the interpretation of prospectively and retrospectively obtained data needs careful consideration, the trend towards an increase in SVC flow over time is observed from the prospective studies alone. There is a need to revisit the definition of 'low SVC flow'.

There are some limitations to our study. The study population is a selected cohort of infants deemed unwell enough by the treating clinician to have an echocardiogram performed. Table 1 shows that infants who received echocardiograms were of lower gestation and lower birthweight. A selection bias is likely to have occurred, however Paradisis noted that low SVC flow is increasingly likely in lower gestation infants and indeed had a protocol to target this [17]. 58% of the non-studied infants were 28 or 29 weeks gestation, compared with 42% in the studied population. So while it is likely that more infants with lower flows were included in the study cohort this is by no means certain.

Generalisability of the findings is limited by the inherent variability of this technique, and differences in demographics and management of cardiovascular support between units. SVC flow measurement has a high inter-observer variation as mentioned above making comparisons of exact values problematic, however we looked at infants in the lowest SVC quartile as well and still found no evidence for association of severe IVH occurrence or IVH extension [1, 4].

A previously unreported finding is the association of low admission temperature and low SVC flow. Neonatal hypothermia has long been known to be associated with increased neonatal mortality [19].

Several randomised controlled trials showed that the use of polyethylene plastic bags prevents hypothermia at delivery in very preterm babies [6, 21]. Although no study has investigated longterm outcomes, current ILCOR neonatal resuscitation guidelines recommend the use of plastic bags or wraps in very low birthweight infants [20]. It is possible that increasing SVC flows more recently described are in part due to improved thermal care of premature infants, including but not limited to plastic wrapping. However we acknowledge that a low admission temperature could also reflect more extensive resuscitation of a sick infant, requiring more intervention at the cost of less sufficient temperature control.

The true value of SVC flow measurement remains to be elucidated. Several studies have shown that low SVC flow is associated with poorer outcome in preterm infants. It is acknowledged that the introduction of SVC flow measurement has substantially increased the awareness of the importance of cardiovascular assessment in the preterm neonate. This study however did not reveal any additional diagnostic value of a single measurement of SVC flow in very preterm infants in an echocardiographically limited setting. A randomised controlled trial investigating functional echocardiography guided cardiovascular management in the preterm neonate may be indicated.

References

- [1] Groves AM, Kuschel CA, Knight DB, Skinner JR (2008) Echocardiographic assessment of blood flow volume in the superior vena cava and descending aorta in the newborn infant. *Arch Dis Child Fetal Neonatal* Ed 93:F24-8
- [2] Hart J, Vemgal P, Cocks-Drew S, Harrison C, Andersen C (2006) The relation between inferior vena cava oxygen saturation, superior vena cava flow, fractional oxygen extraction and haemoglobin affinity in sick newborns: a pilot study. *Acta Paediatr* 95:50-5
- [3] Kluckow M, Seri I, Evans N (2007) Functional echocardiography: an emerging clinical tool for the neonatologist. *J Pediatr* 150:125-30
- [4] Kluckow M, Evans N (2000) Superior vena cava flow in newborn infants: a novel marker of systemic blood flow. *Arch Dis Child Fetal Neonatal* Ed 82:F182-7
- [5] Kluckow M, Evans N (2000) Low superior vena cava flow and intraventricular haemorrhage in preterm infants. *Arch Dis Child Fetal Neonatal* Ed 82:F188-94
- [6] Knobel RB, Wimmer JE, Jr., Holbert D (2005) Heat loss prevention for preterm infants in the delivery room. *J Perinatol* 25:304-8
- [7] Miletin J, Dempsey EM (2008) Low superior vena cava flow on day 1 and adverse outcome in the very low birthweight infant. *Arch Dis Child Fetal Neonatal* Ed 93:F368-71
- [8] Miletin J, Pichova K, Dempsey EM (2009) Bedside detection of low systemic flow in the very low birth weight infant on day 1 of life. *Eur J Pediatr* 168:809-13
- [9] Moran M, Miletin J, Pichova K, Dempsey EM (2009) Cerebral tissue oxygenation index and superior vena cava blood flow in the very low birthweight infant. *Acta Paediatr* 98:43-6

- [10] Osborn DA, Evans N, Kluckow M (2004) Clinical detection of low upper body blood flow in very premature infants using blood pressure, capillary refill time, and central-peripheral temperature difference. *Arch Dis Child Fetal Neonatal Ed* 89:F168-7
- [11] Osborn DA, Evans N, Kluckow M (2003) Hemodynamic and antecedent risk factors of early and late periventricular/intraventricular hemorrhage in premature infants. *Pediatrics* 112:33-9
- [12] Osborn DA, Evans N (2003) Randomized trial of high-frequency oscillatory ventilation versus conventional ventilation: effect on systemic blood flow in very preterm infants. *J Pediatr* 143:192-8
- [13] Osborn D, Evans N, Kluckow M (2002) Randomized trial of dobutamine versus dopamine in preterm infants with low systemic blood flow. *J Pediatr* 140:183-91
- [14] Osborn DA, Evans N, Kluckow M, Bowen JR, Rieger I (2007) Low superior vena cava flow and effect of inotropes on neurodevelopment to 3 years in preterm infants. *Pediatrics* 120:372-80
- [15] Papile LA, Burstein J, Burstein R, Koffler H (1978) Incidence and evolution of subependymal and intraventricular hemorrhage: a study of infants with birthweights less than 1,500 gm. *J Pediatr* 92:529-34
- [16] Paradisis M, Evans N, Kluckow M, Osborn D, McLachlan AJ (2006) Pilot study of milrinone for low systemic blood flow in very preterm infants. *J Pediatr* 148:306-13
- [17] Paradisis M, Evans N, Kluckow M, Osborn D (2009) Randomized trial of milrinone versus placebo for prevention of low systemic blood flow in very preterm infants. *J Pediatr* 154:189-95
- [18] Parry G, Tucker J, Tarnow-Mordi W (2003) CRIB II: an update of the clinical risk index for babies score. *Lancet* 361:1789-9
- [19] Silverman WA, Fertig JW, Berger AP (1958) The influence of the thermal environment upon the survival of newly born premature infants. *Pediatrics* 22:876-86
- [20] The International Liaison Committee on Resuscitation (ILCOR) consensus on science with treatment recommendations for pediatric and neonatal patients: neonatal resuscitation (2006). *Pediatrics* 117:e978-88
- [21] Vohra S, Roberts RS, Zhang B, Janes M, Schmidt B (2004) Heat Loss Prevention (HeLP) in the delivery room: A randomized controlled trial of polyethylene occlusive skin wrapping in very preterm infants. *J Pediatr* 145:750-3
- [22] de Waal K, Evans N, van der Lee J, van Kaam A (2009) Effect of lung recruitment on pulmonary, systemic, and ductal blood flow in preterm infants. *J Pediatr* 154:651-5
- [23] West CR, Groves AM, Williams CE, Harding JE, Skinner JR, Kuschel CA, Battin MR (2006) Early low cardiac output is associated with compromised electroencephalographic activity in very preterm infants. *Pediatr Res* 59:610-5

Table 1: Comparison of infants < 30 weeks gestation, who had SVC flow measured in the first 24 hours of life compared with those who did not.

	SVC flow measured	SVC flow not measured	P Value
Number	165	153	
Gestation (weeks)	26.6 (+/- 1.7)	27.3 (+/- 1.8)	0.003
Birth weight (grams)	940 (+/- 268)	1053 (+/- 281)	<0.001
1min Apgar	5 (+/- 1.8)	6 (+/- 2.1)	0.65
5 min Apgar	8 (+/- 1.5)	8 (+/- 1.9)	0.87
CRIB II	10 (+/- 3.3)	9.3 (+/- 3.5)	0.07
Any IVH	33 (20%)	19 (12%)	0.07
IVH (3 or 4)	22 (13%)	12 (8%)	0.08
Death	27 (16%)	14 (9%)	0.06
Scanned outside usual echocardiographers hours **	36 (22%)	3 (2%)	<0.001

SVC=Superior Vena Cava, CRIB II=Clinical Risk Index for Babies II, P/IVH=Periventricular/Intraventricular haemorrhage.

Values are actual numbers (proportions), or mean (standard deviation).

**After office hours outside 0800-1700, Mon, Wed, Thurs or Friday.

Table 2: Variable associations with low SVC flow.

	Normal SVC flow	Low SVC flow (<41 ml/kg/min)	P-Value
Number	159	6	
Gestation (weeks)	26.7 (+/- 1.8)	25.8 (+/- 1.5)	0.24
Birth weight (grams)	945 (+/- 267)	820 (+/- 273)	0.26
Inborn	137 (86%)	4 (67%)	0.62
CRIB II	10.0 (+/- 3.1)	13.5 (+/- 4.6)	0.009
Temperature at admission (degree Celsius)	36.0 (+/- 0.9)	34.6 (+/- 1.3)	<0.001
Earliest base excess (mmol/L)	-4.7 (+/- 3.6)	-10.9 (+/- 8.1)	<0.001
Earliest lactate (mmol/L)	4.0 (+/- 2.8)	9.0 (+/- 5.8)	<0.001
Volume of fluid to time of echo (ml/kg)	13.0 (+/- 11.4)	27.0 (+/- 17.5)	0.004
Mean airway pressure at time of echo (cmH ₂ O)	8.3 (+/- 2.3)	11.0 (+/- 2.5)	0.009

SVC=superior vena cava, CRIB II=Clinical Risk Index for Babies II

Table 3: Variables associated with grade 3 or 4 IVH.

	No IVH	Severe IVH (grade 3 or 4)	P Value
Number of infants	132	22	
Gestation (weeks)	27.0 (+/- 1.7)	25.2 (+/- 1.8)	<0.001
Birthweight (grams)	968 (+/- 261)	814 (+/- 298)	0.01
Male sex	65 (49%)	16 (73%)	0.05
CRIB II	9.5 (2.9)	13.1 (3.8)	<0.001
Inborn	117 (89%)	16 (73%)	0.09
Caesarean delivery	91 (69%)	10 (45%)	0.07
Temperature on admission (degree Celsius)	36.0 (+/- 0.8)	35.6 (+/- 1.3)	0.05
Earliest pH	7.29 (+/- 0.11)	7.23 (+/- 0.11)	0.03
Earliest base excess (mmol/L)	-4.3 (+/- 3.6)	-8.0 (+/- 5.0)	<0.001
Earliest lactate (mmol/L)	3.8 (+/- 2.7)	6.2 (+/- 4.2)	0.002
Mean airway pressure at TOE (cmH ₂ O)	8.2 (+/- 2.3)	9.4 (+/- 2.3)	0.04

Dobutamine dose at TOE (microgram/kg/min)	9.2 (+/- 8.6)	13.3 (+/- 7.5)	0.05
Fluid bolus in to TOE (ml/kg)	11.6 (+/- 10.4)	23.4 (+/- 15.4)	<0.001
Low SVC flow (<41ml/kg/min)	2 (1.5%)	4 (18%)	0.002

IVH= Intraventricular haemorrhage; CRIB II=Clinical Risk Index for Babies II; TOE=Time of echocardiogram; SVC=Superior vena cava

Table 4: Averaged values for SCV flow in the first 24 hours of life and population characteristics in published studies to date.

	Kluckow and Evans [4]	Kluckow and Evans [5]	Osborn and Evans [12]	Paradisis et al. [16]	Hart et al. [2]	West et al. [23]	Groves et al. [1]	Moran et al. [9]	Paradisis et al. [17]	This Study
Years studied	1995-96	1995-96	1999-2001	2002-03	N/R	2003-04	2002-04	2006-07	2003-06	2004-06
Number of infants	25	126	43	29	17	40	14	27	90	165
Neonatal Unit(s)	RPAH and RNSH, Sydney, Australia	RPAH and RNSH, Sydney, Australia	RPAH and RNSH, Sydney, Australia	RPAH and RNSH, Sydney, Australia	MHW, Melbourne, Australia	NWH, Auckland, New Zealand	NWH, Auckland, New Zealand	CWH, Dublin, Ireland	RPAH and RNSH, Sydney, Australia	MHW, Melbourne, Australia
Population	<30/40, <48 hours resp. support	<30/40, within 24 hours of life	23-28/40	<29/40	Infants requiring UAC and UVC	<31/40	<31/40, <48 hours resp. support	<1500g birth - weight	<30/40, <6 hours old	<30/40, within 24 hours of life
Gestation Median in weeks	29	27	26.3	25.6 (mean)	26.6	27	29	27.6 (mean)	26 (mean)	26.6

Cardio-vascular treatment	None before first scan	None before first scan	None before first scan-unless BP $\leq$ 20 mmHg	Milrinone at various doses, after a 15ml/kg bolus	As determined by treating clinician	Not reported	No cardio-Vascular support	NS 10 ml/kg, then adrenaline and/ or dobutamine based on various factors	NS 15 ml/kg, then milrinone or placebo, additional dopamine or NS if required	As determined by treating clinician
Inotrope with echo	N/R	23%	N/R	77%	N/R	N/R	N/R	13%	0 at first echo	66%
No. Echos in first 24 hours	3	3	3	4-5	2	3	3	1	3-4	1
Mean SVC flow in first 24 hours of life in ml/kg/min (SD)	73*	57*	81*	81*	79*	92*	101*	70 (40)	80	123 (58)
SVC flow at echo closest to 12 hours of life in ml/kg/min (SD)	75	56.5*	68*	75.4*	85 (66)	97	101	70 (40)	74	123 (58)

SVC = superior vena cava; RPA = Royal Prince Alfred Hospital; RNS = Royal North Shore Hospital; MHW = Mercy Hospital for Women; NWH = National Women's Hospital; CWH = Coombe Women's Hospital; UAC = umbilical artery catheter; UVC = umbilical venous catheter; BP = blood pressure; NS = normal saline; UO = urine output, SD = standard deviation

* = calculated value