Metallomics

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: L. Jin, D. M. Frazer, Y. Lu, S. J. Wilkins, S. Ayton, A. I. Bush and G. Anderson, *Metallomics*, 2019, DOI: 10.1039/C8MT00370J.

This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Significance to metallomics statement

View Article Online
DOI: 10.1039/C8MT00370J

Manganese is an essential metal, but how its trafficking is regulated within the body remains incompletely understood. Ferroportin is best known as an iron export protein, although it can also transport a number of other divalent metal ions, including potentially manganese. However, the literature is divided in this area, and in particular, there are few data on how ferroportin might contribute to manganese homeostasis under physiological conditions. By taking advantage of two knockout mouse models with altered ferroportin levels, we have been able to provide new insights into the contribution this transporter makes to manganese distribution in vivo.

Metallomics Accepted Manuscript

Metallomics

ARTICLE

Mice overexpressing hepcidin suggest ferroportin does not play a major role in Mn homeostasis

\Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Lian Jin^{a,b}, David M Frazer^a, Yan Lu^a, Sarah J Wilkins^a, Scott Ayton^c, Ashley Bush^c, Gregory J Anderson^{a,b,d}

Manganese is an essential metal that is required for a wide range of biological functions. Ferroportin (FPN), the only known cellular exporter of iron, has also been proposed to play a role in manganese export, but this relationship is incompletely understood. To investigate this in more detail *in vivo*, we examined the relative distributions of manganese and iron in Tmprss6 deficient mice, which are characterized by constitutively high expression of the iron regulatory hormone hepcidin and, consequently, very low FPN levels in their tissues. *Tmprss6*^{-/-} mice showed frank iron deficiency and reduced iron levels in most tissues, consistent with FPN playing an important role in the distribution of this metal, but manganese levels were largely unaffected. Associated studies using intestine-specific FPN knockout mice showed that loss of FPN significantly reduced the dietary absorption of iron, but had no effect on manganese intake. Taken together, our data suggest that FPN does not play a major role in Mn transport *in vivo*. They do not exclude a minor role for FPN in manganese homeostasis, nor the possibility that the transporter may be relevant at high Mn levels, but at physiological levels of this metal, other transport proteins appear to be more important.

Introduction

Metals such as iron (Fe) and manganese (Mn) are essential for a wide range of biological functions. Iron, for example, is required for oxygen delivery, energy production and DNA synthesis,¹ and manganese is required for the activity of many enzymes, including those involved in glucose and lipid metabolism and antioxidant defence.² Both Fe and Mn levels in the body must be strictly controlled as a deficiency or excess of either metal can be harmful. Iron deficiency can lead to anemia and its associated pathological consequences, including compromised neurological development when present in infancy.^{3, 4} In contrast, since Fe can catalyze the formation of toxic oxygen radicals, excess iron can damage vital organs including the liver and heart. Mn is also toxic when present in excess, and its neurotoxicity is particularly notable.² Significantly, high Mn levels have been associated with neurological disorders such as hepatic encephalopathy and parkinsonism.⁵⁻⁷ Because of this dual nature, being essential, yet being toxic in excess, sophisticated mechanisms have developed to maintain the cellular and body levels of Fe and Mn within the optimum physiological range. This is largely achieved by regulating the cellular intake and efflux of the

metals. For both Fe and Mn, the liver plays a central regulatory role.^{8, 9}

In broad terms, the regulation of Fe traffic in the body is better studied than that of Mn. Dietary Fe is taken up by intestinal enterocytes, then released into the circulation across the enterocyte basolateral membrane through ferroportin (FPN; SLC40A1), the only known mammalian iron export protein.¹⁰ In the circulation, iron binds to transferrin and is then distributed to cells throughout the body. In addition to being important for intestinal Fe absorption, FPN is required for Fe export from most body cells. This iron export process is regulated by the hepatic hormone hepcidin, which binds to FPN and leads to its internalization and degradation.¹¹ When body iron levels rise, hepcidin production is increased and FPN levels decline.¹² This in turn reduces iron absorption and iron accumulates in intestinal enterocytes. Iron also accumulates in other cells in the body. Conversely, when body iron levels are depleted, hepcidin expression is downregulated to allow for increased dietary Fe absorption and mobilization of body Fe stores, both of which require active FPN on the cell surface.^{13, 14} Pathological disturbances leading to either increased or decreased hepcidin expression lead to anemia and hemochromatosis respectively.¹⁵⁻¹⁷

FPN has been shown to transport several divalent metal ions other than Fe²⁺, including Co²⁺ and Zn²⁺, but not others such as Cu²⁺ or Cd²⁺.¹⁸ The literature on whether FPN is able to transport Mn is confusing and apparently contradictory. Some *in vitro* studies have suggested that FPN transports minimal amounts of Mn,^{18, 19} but others have suggested otherwise.^{20, 21} *In vivo* studies have generally suggested that FPN is able to

^a Iron Metabolism Laboratory, QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia.

^b Faculty of Medicine, University of Queensland, Brisbane, Australia

^c Melbourne Dementia Research Centre, Florey Institute of Neuroscience and Mental Health, University of Melbourne, Melbourne, Victoria, Australia

^d School of Chemistry and Molecular Biosciences, University of Queensland, Brisbane, Australia.

transport Mn, but its influence is usually modest. For example, it has been reported that the flatiron mouse (*ffe/+*), a model of genetic FPN deficiency, albeit with an unusual underlying mechanism, has a reduced capacity to absorb orally administered Mn and has reduced Mn levels in the blood, liver and bile.²² In addition, there is evidence that intestinal absorption of Mn is increased in *Hfe* knockout mice, a mouse model of HFE-related hemochromatosis²³ where FPN expression is increased secondary to diminished hepcidin expression.^{24, 25} While these studies provide some evidence to implicate FPN in Mn homeostasis, its relative contribution at physiological levels of Mn remains unclear.

FPN is only one of more than ten transmembrane metal transport proteins that have been implicated in Mn homeostasis.²⁶ These include proteins involved in Mn import, such as DMT1 (NRAMP2),^{27, 28} SLC39A8 (ZIP8)²⁹ and SLC39A14 (ZIP14),³⁰⁻³³ and those involved in Mn export, such as SLC30A10 (ZNT10),³⁴⁻³⁵ ATP2C1 (SPCA1)³⁶ and ATP13A2 (PARK9).³⁷⁻⁴⁰ Recent studies have suggested that the plasma membrane proteins SLC30A10 and SLC39A14 play particularly important roles in coordinately regulating body Mn levels. Patients with loss of function mutations in SLC30A10,^{41, 42} or mice lacking this protein,^{43, 44} show accumulation of Mn in the blood, liver and brain, and associated parkinsonian-like symptoms. The localization of SLC30A10 to the canalicular membrane of HepG2 cells⁴⁴ is consistent with it being involved in Mn excretion into the bile. Mice with a liver-specific knockout of *Slc30a10* show minimal alterations in body Mn,⁴⁵ implying that other Mn export pathways exist. The gut is a strong candidate for an alternative excretion route as SLC30A10 is also expressed on the apical membrane of mature enterocytes and appears to be responsible for effluxing Mn into the intestinal lumen.⁴⁵ SLC30A10 does not appear to be involved in regulating brain Mn levels under basal conditions, but it can protect against neurotoxicity when levels of the metal are increased.⁴⁵ Disruption of the metal importer SLC39A14 in both humans and mice has also been associated with body Mn accumulation in multiple organs, including the brain, with associated adverse neurological consequences.³⁰⁻³³ However, in this case, Mn levels in the liver are low. Radioactive ⁵⁴Mn distribution studies comparing *Slc39a14* null and wild-type mice indicate that this protein is important for hepatic and pancreatic Mn uptake, and for its gastrointestinal excretion.³¹ Taken together, these data suggest that SLC39A14 and SLC30A10 work together in hepatocytes to facilitate Mn excretion into the bile, and studies with *Slc39a14/Slc30a10* double knockout mice are consistent with this.⁴⁴ The loss of this major excretion pathway leads to the accumulation of Mn in the body and its associated adverse consequences. Interestingly, loss of SLC39A14 specifically in the liver in mice does not lead to substantive extrahepatic Mn loading,^{33, 44} implying that Mn can be excreted through other routes, most likely via the pancreas and intestinal epithelium.

In this study, we have taken advantage of transmembrane serine protease 6 (TMPRSS6) knockout mouse to further explore the links between FPN and Mn metabolism and to compare this to the homeostasis of iron. TMPRSS6 is a

protease which acts upstream of hepcidin to reduce its expression.⁴⁶ *Tmprss6*^{-/-} mice lack the protease and consequently hepcidin is expressed at a constitutively high level.⁴⁷ This in turn reduces FPN expression and leads to iron deficiency in the mice. These animals represent an ideal model in which to study the effects of increased hepcidin (and hence reduced FPN) on Mn metabolism. Using both these mice and mice carrying an intestine-specific deletion of FPN, we showed that FPN likely plays a minimal role in body Mn homeostasis. Metal transporters other than FPN appear to be more important in regulating Mn flux in the body under normal physiological conditions.

Experimental

Animal care and procedures

All experimental procedures were approved by the QIMR Berghofer Medical Research Institute Animal Ethics Committee. Homozygous *Tmprss6* knockout mice (*Tmprss6*^{-/-})⁴⁷ on a C57BL/6 background were used as a model of constitutively high hepcidin production. This strain has been validated previously in our hands to show that *Hamp1* mRNA levels and plasma hepcidin levels are constitutively high and that FPN protein is reduced in the proximal small intestine⁴⁸. Heterozygous littermates (*Tmprss6*^{+/-}) were used as controls as we observed no phenotypic differences between wild-type mice and heterozygotes.⁴⁸ In some studies, we also used mice lacking FPN specifically in the enterocytes which were created and characterized as previously described.⁴⁸ Briefly, they were produced by crossing FPN floxed mice with *vil-Cre-ERT2* mice (both on a C57BL/6 background), which express tamoxifen-inducible Cre recombinase under the control of the villin promoter. Cre recombinase expression was induced in these mice by injecting tamoxifen (75 µg/g body weight) subcutaneously at 8-10 days of age. Ferroportin floxed littermates lacking Cre recombinase also were injected with tamoxifen as the control group. Our previous work has shown that these animals have a 90% reduction in intestinal iron absorption and that this is associated with loss of FPN protein in the small intestine.⁴⁸ For consistency in our experiments, only male mice were used. Some sex differences in Fe and Mn homeostasis have been observed in mice,^{31, 43, 49} but, where present, these are almost always quantitative in nature and do not represent fundamental differences between the responses of males and females. Thus, we anticipate that the findings will be equally applicable to females. The animals were weaned 3 weeks after birth, then provided with unlimited access to a standard pellet diet (Norco Stockfeed, Lismore, NSW, Australia; Metal ion content: Fe 120mg/kg; Mn 80mg/kg; Zn 200mg/kg; Cu 16mg/kg) and tap water.

In initial studies, *Tmprss6*^{-/-} mice and *Tmprss6*^{+/-} controls were examined at 2, 4, 8 and 12 weeks of age. Animals were anesthetized (200mg/kg ketamine and 10mg/kg xylazine) before euthanasia. Blood was taken by cardiac puncture and mice were perfused with trace-metal grade saline (Sigma). Organs, including the liver, brain, heart, kidney and spleen

were collected, snap frozen in liquid nitrogen and stored at -80°C for subsequent investigation. Intestine-specific *Fpn* knockout mice were processed similarly at the age of 8-10 weeks. Perfusion with trace-metal grade saline was performed. In this case the liver was the only tissue collected.

In some experiments Mn was administered by oral gavage (100mg Mn/kg body weight) for 5 consecutive days. This represents approximately 5 times the daily intake of Mn each mouse would receive from a standard pellet diet. Both *Tmprss6*^{-/-} and *Tmprss6*^{+/-} mice were studied (n=6). Five days after the final gavage, mice were euthanised and blood and other tissues collected as described above.

Real time quantitative PCR

Trizol reagent (Thermo Fisher Scientific, Scoresby, Victoria, Australia) was used to extract total RNA according to the manufacturer's instructions. Complementary DNA was synthesized using M-MLV Reverse Transcriptase (Thermo Fisher Scientific) and an oligo (dT) primer according to the manufacturer's instructions. Real-time quantitative PCR was performed on a CFX384 Touch Real-Time PCR Detection System (Bio-Rad, Gladesville, NSW, Australia). Each sample was analyzed in triplicate, and gene expression was calculated from the Ct value using the standard curve method. Gene expression levels were expressed relative to the housekeeping gene, hypoxanthine guanine phosphoribosyl transferase (*Hprt*). Validation of primers and analyses were consistent with the MIQE guidelines.⁵⁰ The primer pairs used were: *Hamp1* forward: AGAGCTGCAGCCTTTGCAC; *Hamp1* reverse: AACTGGGAATTGTTACAGCATT; *Hprt* forward: GGACTGATTATGGACAGGA; *Hprt* reverse: GAGGGCCACAATGTGATG.

ICP-MS

Tissue for ICP-MS was dried at 110°C, weighed, and wet-digested²¹ prior to metal analysis. Briefly, 50 µL of HNO₃ (14.4M, 65% Suprapur, Merck) was added to the dried sample and digestion was allowed to proceed overnight at room temperature. The temperature was then raised to 90°C using a heating block and incubation continued 20 mins. After the addition of an equivalent volume of hydrogen peroxide (9.9 M, 30% Aristar, BDH), the samples were incubated for 30 mins at room temperature, followed by a further 15 mins at 70°C. The volume of each sample was then determined and 1% HNO₃ (0.22M) added to a total volume of 500 µL. Trace element levels were measured using an Agilent 7700 series ICP-MS instrument under routine multi-element operating conditions using a Helium Reaction Gas Cell. Metal concentrations were expressed as µg/g dry tissue for tissue samples or µM for serum samples.

Statistical analysis

Each experimental group contained 5-10 animals, and data are presented as the mean ± SEM. Statistical differences between

treatment groups were determined using t tests when comparing two groups or one-way analysis of variance (ANOVA) followed by Sidak's multiple comparisons test when three or more groups were involved. Differences were considered significant when $p < 0.05$. All analyses were carried out with GraphPad Prism 7 software (GraphPad Software, San Diego, CA, USA).

Results and discussion

Tmprss6 deficiency affects iron and manganese metabolism differently

In this study, we systemically compared iron and manganese homeostasis in *Tmprss6* deficient mice. We first confirmed that these animals had constitutively high hepcidin expression by examining *Hamp1* mRNA levels in mice aged 2, 4, 8 and 12 weeks (Fig. 1) and we have previously shown that *Hamp1* mRNA expression correlates with plasma hepcidin protein levels and intestinal FPN protein in these mice⁴⁸. Hepatic *Hamp1* expression increased in both *Tmprss6*^{-/-} and littermate controls as the mice aged, but at all ages hepcidin expression was significantly elevated in *Tmprss6* knockout mice compared with their heterozygous littermates. These results are consistent with published data on hepcidin expression in this mouse strain.⁴⁷

Fig. 1 *Hamp1* gene expression in *Tmprss6* knockout mice. *Hamp1* (hepcidin) mRNA levels were assessed in the liver of *Tmprss6* knockout mice (KO)(open squares) and heterozygous littermates (Het)(solid circles) at 2, 4, 8 and 12 weeks of age. Gene expression was analysed using qPCR and the data were normalized to the *Hamp1* mRNA level of 8 week-old heterozygous mice. Data are shown as means ± SEM, n=6 and differences between *Tmprss6* knockout and heterozygous mice were determined by t test separately at each age. (** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$)

Tmprss6 knockout mice also showed obvious anemia as indicated by reduced hemoglobin concentration, hematocrit and mean corpuscular volume (MCV) at all ages studied (Fig. 2A-C). Red blood cell (RBC) number was not influenced by the loss of TMPRSS6 (Fig. 2D). This anemia is consistent with

elevated hepcidin expression and with previous descriptions of *Tmprss6*^{-/-} mice.^{47,48}

Fig. 2 Hematology and iron status of *Tmprss6*^{-/-} mice (open squares) and heterozygous littermate controls (solid circles) at various postnatal ages. (A) Hgb: hemoglobin concentration; (B) Hct: hematocrit; (C) MCV: mean corpuscular volume; (D) RBC: red blood cell count; (E) serum iron; and (F) hepatic iron. Data are shown as means $\pm$ SEM, $n=6$ and differences between *Tmprss6* knockout and heterozygous mice were determined by t test separately at each age. (** $p < 0.01$; **** $p < 0.0001$).

Tmprss6^{-/-} mice also had lower serum and hepatic iron levels at all ages (Fig. 2E,F) which is to be expected in the presence of elevated hepcidin as the peptide binds to FPN on the surface of target cells, thereby reducing the donation of iron to the plasma. This might be expected to lead to iron accumulation within the tissues, but since intestinal enterocytes are a major site of hepcidin action, dietary iron absorption is reduced when hepcidin levels are high, and hence the iron content of most tissues is reduced (Figs 2F and 3A-C). An exception to this pattern is seen in 2 week-old mice when *Tmprss6* deficiency exerts little effect on tissue iron levels. This apparent anomaly can be explained by the finding that hepcidin is unable to bind to enterocyte FPN when the mice are suckling (up to 3 weeks of age) and thus iron continues to enter the body at the same rate as in animals with normal TMPRSS6 levels.⁴⁸ However, the extra-intestinal tissues remain responsive to high hepcidin and this explains why hepatic iron levels were slightly raised in two week-old *Tmprss6*^{-/-} mice (Fig. 2F). After weaning, mice have access to much larger amounts of iron in their diet, leading to a dramatic increase in liver iron levels between 2 and 4 weeks postnatally in wild-type animals.

We also examined the effects of increased hepcidin expression on Fe levels in several tissues other than the liver. A marked decline in Fe levels was observed in the brain and heart of *Tmprss6*^{-/-} mice (Fig. 3A,B), although the decrease was not as striking as it was in the liver (Fig. 3B). In the kidney, a significant drop in Fe level was only observed in 12 week-old *Tmprss6* knockouts (Fig. 3C), and no significant differences in splenic iron were observed between the two genotypes at any of the ages studied (Fig. 3D). The latter results may reflect slower iron turnover in the kidney and spleen.

Some studies have suggested that FPN contributes to Mn homeostasis and may be involved in cellular Mn export,^{18, 19} but others have suggested otherwise.^{20, 21} Since *Tmprss6*^{-/-} mice have greatly reduced tissue FPN levels as a result of hepcidin overexpression, we were able to use them as an *in vivo* model to investigate whether altered FPN expression affected Mn distribution. We thus measured Mn levels in various tissues of *Tmprss6*^{-/-} mice. In most cases, no significant differences were observed in Mn concentrations between *Tmprss6*^{-/-} mice and controls in the serum, brain, heart, kidney, spleen and liver (Fig. 4A-F). Small, but significant, differences were observed in the heart of 8 week old mice (Fig. 4C), the spleen of 12 week old mice (Fig. 4E) and the liver of 8 week old mice (Fig. 4F), but in each case Mn levels were increased. (In the same samples, Fe levels were decreased.) Whether these changes are sufficiently large to be physiologically relevant is difficult to determine.

Interestingly, Mn levels in the livers of 2 week old *Tmprss6*^{-/-} mice were significantly decreased (Fig. 4F). Mn concentrations have previously been reported to be similar between fetal and suckling mice,⁵¹ suggesting that the amount of Mn in the suckling pup depends largely on how much of the metal was transferred across the placenta before birth. Placental transport of Mn to the fetus could be impaired by a reduction in FPN caused by high fetal hepcidin in knockout fetuses, but this seems unlikely since Fe levels are far more sensitive to hepcidin action and they were not reduced. Further studies are required to examine the mechanisms operating in suckling mice. Since no decreases in Mn levels were seen in older mice, our data suggest that intestinal Mn absorption is not impaired in response to increased hepcidin and argue against a strong role for FPN in Mn transport in the gut. These data highlight a significant difference between Mn and Fe.

Fig. 3 Iron levels in various tissues of *Tmprss6*^{-/-} mice (open squares) and heterozygous littermate controls (solid circles) at various postnatal ages. (A) Brain, (B) heart, (C) kidney and (D) spleen. Data are shown as means $\pm$ SEM, n=6 and differences between *Tmprss6* knockout and heterozygous mice were determined by t test separately at each age. (*p < 0.05; **p < 0.01; ***p < 0.001; ****p < 0.0001)

Other metals such as copper (Fig. 5) and zinc (Fig. 6) showed similar behaviour to Mn, with no or minimal differences between *Tmprss6*^{-/-} and wild-type mice. A previous *in vitro* study¹⁶ showed that FPN was unable to transport significant amounts of Cu²⁺, but it was able to transport Zn²⁺, albeit far less efficiently than iron. These findings are largely in line with our data.

Taken together, our findings with *Tmprss6*^{-/-} mice suggest that FPN levels are not a major contributor to Mn distribution, as they are for Fe. These data do not exclude some role for FPN in the cellular export of Mn, but suggest that it is likely to be a minor role and that other (and potentially Mn-specific) Mn transport pathways are involved. Further investigations into the contributions of known or suspected Mn transporters are required to clarify the mechanisms underlying the physiological distribution of this metal.

Fig. 4 Mn levels in the serum (A), brain (B), heart (C), kidney (D), spleen (E) and liver (F) of *Tmprss6*^{-/-} mice (open squares) and heterozygous littermate controls (solid circles) at various postnatal ages. Data are shown as means $\pm$ SEM, n=6 and differences between *Tmprss6* knockout and heterozygous mice were determined by t test separately at each age. (*p < 0.01; ****p < 0.0001)

Fig. 5 Copper (Cu) levels in the serum (A), brain (B), heart (C), kidney (D), spleen (E) and liver (F) of *Tmprss6* deficient mice (open squares) and heterozygous control mice (solid circles). Data are shown as means $\pm$ SEM, n=6 and differences between *Tmprss6*

ARTICLE

Journal Name

knockout and heterozygous mice were determined by t test separately at each age. (** $p < 0.01$)

To provide some support for the data obtained with mice overexpressing hepcidin, we examined Fe and Mn levels in mice with specific deletion of FPN in the intestine.³² In intestinal *Fpn*^{-/-} mice, Fe levels in the liver were strongly decreased relative to those of control mice, whereas the hepatic Mn concentration was unchanged (Fig. 7). These findings are consistent with our observations in *Tmprss6* deficient animals.

Fig. 6 Zinc (Zn) levels in the serum (A), brain (B), heart (C), kidney (D), spleen (E) and liver (F) of *Tmprss6* deficient mice (open squares) and heterozygous control mice (solid circles). Data are shown as means $\pm$ SEM, $n=6$ and differences between *Tmprss6* knockout and heterozygous mice were determined by t test separately at each age. (* $p < 0.05$; ** $p < 0.01$)

Fig. 7 Mn (A) and Fe (B) levels in the liver of mice lacking FPN only in the intestinal tract. At 8-10 days of age, mice carrying a floxed *Fpn* allele and a tamoxifen-inducible Cre recombinase under the control of the villin promoter were injected with tamoxifen to inactivate the gene encoding FPN (open squares). Mice carrying a floxed *Fpn* allele but no Cre recombinase gene were also treated with tamoxifen and these animals acted as FPN positive controls (solid circles). Fe and Mn levels in the liver were measured when the mice were 8-10 weeks old. Data are shown as means $\pm$ SEM, $n=6$ and differences between groups were determined by t test. (**** $p < 0.0001$)

Effects of loss of *Tmprss6* in the presence of supraphysiological amounts of Mn

As loss of *Tmprss6* did not affect Mn distribution under normal physiological conditions where body Mn levels are relatively low, we investigated whether we could see any effects when wild-type and *Tmprss6*^{-/-} mice were challenged with a large dose of Mn. When Mn was administered by oral gavage at a dose of 100 mg/kg, both *Tmprss6*^{-/-} and heterozygous control mice accumulated more Mn in their liver, kidney and brain than saline treated mice (Fig. 8A-C). However, only small increases were observed in the serum, spleen and heart, and these were not statistically significant (Fig. 8D-F). Previous studies have shown that in Mn overload situations, the metal preferentially accumulates in the liver and brain, consistent with our findings.⁵² With one exception (see below) tissue Mn concentrations did not differ between *Tmprss6*^{-/-} and heterozygous control mice, consistent with our earlier observations that FPN does not appear to play a significant role in Mn distribution *in vivo*. The exception was the brain, where Mn levels in *Tmprss6* deficient mice were significantly lower than those of heterozygous control animals, but only under Mn overload conditions (Fig. 8B). The reason for this surprising finding (which is not consistent with the effects of elevated hepcidin expression on FPN) is not immediately obvious. However, Mn transport in the brain is complicated, with a range of transporters likely to be involved and the potential for crosstalk between these transporters.^{26, 53, 54}

Fig. 8 Mn concentrations in the liver (A), brain (B), kidney (C), serum (D), heart (E) and spleen (F) in the *Tmprss6*^{-/-} and heterozygous (control) mice following Mn administration by oral gavage. Data are shown as means $\pm$ SEM, n=6 and differences between groups were determined by ANOVA. (*p < 0.05; **p < 0.01; ***p < 0.001; ****p < 0.0001)

After being absorbed in the gastrointestinal tract, Mn enters the bloodstream and most is quickly transported to the liver which plays an important role in Mn storage, redistribution and elimination.^{2, 55} Mn that is not required for physiological functions within the body is sequestered by hepatocytes and excreted via the bile.^{56, 57} If FPN is involved in the export of Mn from hepatocytes, as it is for Fe, it might be expected that the amount of Mn reaching other body tissues would be compromised in *Tmprss6* knockout mice, but we did not find this to be the case. Interestingly, in the flatiron mouse, there was a significant reduction in the amount of Mn in the bile,²² suggesting that FPN could be involved in the biliary excretion of the metal. This will require further investigation.

There is evidence from a range of *in vitro* studies that FPN is capable of transporting Mn,^{20,21} although not all studies have supported this conclusion.^{18,19} Differences in experimental systems (e.g. mammalian cells vs *Xenopus* oocytes), Mn concentrations and details of the incubation conditions could account for these discrepancies. However, the *in vivo* situation is far less clear, and whether the *in vitro* studies can be used to reliably infer what happens in a normal physiological environment is questionable. The *in vitro* studies are almost entirely dependent on the overexpression of FPN protein, and

this, combined with high local concentrations of Mn in the incubation solutions, likely does not accurately reflect *in vivo* conditions. However, one cellular study where endogenous FPN was knocked down suggested that this led to a reduction in Mn-associated toxicity, but neither Mn levels nor transport *per se* were investigated²⁰. Our work with the *Tmprss6* and *Fpn* knockout mice is one of the very small number of studies to address this issue *in vivo*. Experiments with the flatiron mouse, which has dominant negative mutation in FPN that leads to altered body iron distribution,⁵⁸ provided some evidence that FPN could contribute to Mn export, although the differences observed were smaller than those generally found in *in vitro* studies.²² In a mouse model of HFE-related hemochromatosis, there were either no or only very small effects on Mn.²³ In this model, levels on FPN are reduced, but some of the transporter is likely to remain, so the weakness of the effect may not be considered surprising. In our own work, we found no clear evidence that overexpression of hepcidin, the normal physiological repressor of FPN, had a significant influence on Mn distribution, nor that loss of FPN itself in the gut. Taken together, these data suggest that if FPN does contribute to Mn transport in the body, it only does so in a very limited way.

If FPN does not play a major role as a Mn export protein, at least at physiological concentrations of the metal, then other transporters must be involved in Mn traffic around the body. As noted above, a range of other transporters have either been demonstrated to export Mn from cells/organelles or are likely to perform this function. Current attention is particularly focused on the coordinated roles of SLC30A10 and SLC39A14 in maintaining body Mn homeostasis, with SLC39A14 acting to transport Mn into hepatic, intestinal and pancreatic cells from which it is excreted via SLC30A10.⁴⁴ The roles of other Mn transporters, such as the lysosome protein ATP13A2 and the Golgi protein ATP2C1 in Mn efflux have yet to be fully explored.

Conclusions

In this study we have demonstrated that Mn and Fe behave quite differently in the presence of constitutively high hepcidin expression (and hence low tissue FPN levels). Fe export into the body from the intestinal epithelium is significantly reduced when hepcidin is high, and hence Fe levels in internal tissues are low. In contrast, *Tmprss6* deletion had little effect on Mn levels in any of the tissues studied, consistent with the conclusion that FPN does not make a major contribution to Mn distribution under these conditions. The use of intestine-specific FPN knockout mice confirmed that FPN had little effect on dietary Mn absorption. Several other Mn transport proteins have been identified and these are likely to be more important than FPN in the homeostasis of this metal, but further investigations are required to determine the relative contributions of these transporters and whether they have tissue-specific effects.

Conflicts of interest

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ARTICLE

There are no conflicts to declare.

Acknowledgements

This work was supported by Project Grant APP1051882 from the National Health and Medical Research Council of Australia (NHMRC). LJ was supported by a University of Queensland International Scholarship and GJA was supported by a Senior Research Fellowship from the NHMRC.

References

1. M. W. Hentze, M. U. Muckenthaler, B. Galy and C. Camaschella, Two to tango: regulation of mammalian iron metabolism, *Cell*, 2010, 142, 24-38.

2. C. L. Keen, J. L. Ensunsa and M. S. Clegg, Manganese metabolism in animals and humans including the toxicity of manganese, *Met Ions Biol Syst*, 2000, 37, 89-121.

3. B. Lozoff and M. K. Georgieff, Iron deficiency and brain development, *Semin Pediatr Neurol*, 2006, 13, 158-165.

4. T. Moos, T. Skjorringe and L. L. Thomsen, Iron deficiency and iron treatment in the fetal developing brain - a pilot study introducing an experimental rat model, *Reprod Health*, 2018, 15, 93.

5. D. Krieger, S. Krieger, O. Jansen, P. Gass, L. Theilmann and H. Lichtnecker, Manganese and chronic hepatic encephalopathy, *Lancet*, 1995, 346, 270-274.

6. J. L. Ordonez-Librado, V. Anaya-Martinez, A. L. Gutierrez-Valdez, L. Colin-Barenque, E. Montiel-Flores and M. R. Avila-Costa, Manganese inhalation as a Parkinson disease model, *Parkinsons Dis*, 2010, 2011, 612989.

7. A. C. Pfalzer and A. B. Bowman, Relationships between essential manganese biology and manganese toxicity in neurological disease, *Current Env Health Rep*, 2017, 4, 223-228.

8. G. Rishi and V. N. Subramaniam, The liver in regulation of iron homeostasis, *Am J Physiol Gastrointest Liver Physiol*, 2017, 313, G157-G165.

9. J. A. Roth, Homeostatic and toxic mechanisms regulating manganese uptake, retention, and elimination, *Biol Res*, 2006, 39, 45-57.

10. H. Drakesmith, E. Nemeth and T. Ganz, Ironing out ferroportin, *Cell Metab*, 2015, 22, 777-787.

11. E. Nemeth, M. S. Tuttle, J. Powelson, M. B. Vaughn, A. Donovan, D. M. Ward, T. Ganz and J. Kaplan, Heparin regulates cellular iron efflux by binding to ferroportin and inducing its internalization, *Science*, 2004, 306, 2090-2093.

12. A. Donovan, A. Brownlie, Y. Zhou, J. Shepard, S. J. Pratt, J. Moynihan, B. H. Paw, A. Drejer, B. Barut, A. Zapata, T. C. Law, C. Brugnara, S. E. Lux, G. S. Pinkus, J. L. Pinkus, P. D. Kingsley, J. Palis, M. D. Fleming, N. C. Andrews and L. I. Zon, Positional cloning of zebrafish ferroportin1 identifies a conserved vertebrate iron exporter, *Nature*, 2000, 403, 776-781.

13. L. Kautz, G. Jung, X. Du, V. Gabayan, J. Chapman, M. Nasoff, E. Nemeth and T. Ganz, Erythroferrone contributes to hepcidin suppression and iron overload in a mouse model of beta-thalassemia, *Blood*, 2015, 126, 2031-2037.

14. Y. Tang, Y. Li, H. Yu, C. Gao, L. Liu, S. Chen, M. Xing, L. Liu and P. Yao, Quercetin prevents ethanol-induced iron overload by regulating hepcidin through the BMP6/SMAD4 signaling pathway, *J Nutr Biochem*, 2014, 25, 675-682.

15. G. N. Ioannou and K. V. Kowdley, Heparin in HFE-associated hemochromatosis: another piece of the "iron" puzzle, *Gastroenterology*, 2004, 126, 615-616; discussion 616.

16. G. Mahajan, S. Sharma, J. Chandra and A. Nangia, Heparin and iron parameters in children with anemia of chronic disease and iron deficiency anemia, *Blood Res*, 2017, 52, 212-217.

17. E. Nemeth, A. Roetto, G. Garozzo, T. Ganz and C. Camaschella, Heparin is decreased in TFR2 hemochromatosis, *Blood*, 2005, 105, 1803-1806.

18. C. J. Mitchell, A. Shawki, T. Ganz, E. Nemeth and B. Mackenzie, Functional properties of human ferroportin, a cellular iron exporter reactive also with cobalt and zinc, *Am J Physiol Cell Physiol*, 2014, 306, C450-459.

19. K. J. Thompson, J. Hein, A. Baez, J. C. Sosa and M. Wessling-Resnick, Manganese transport and toxicity in polarized WIF-B hepatocytes, *Am J Physiol Gastrointest Liver Physiol*, 2018, 315, G351-G363.

20. E. K. Choi, T. T. Nguyen, S. Iwase and Y. A. Seo, Ferroportin disease mutations influence manganese accumulation and cytotoxicity, *FASEB J*, 2019, 33, 2228-2240.

21. Z. Yin, H. Jiang, E. S. Lee, M. Ni, K. M. Erikson, D. Milatovic, A. B. Bowman and M. Aschner, Ferroportin is a manganese-responsive protein that decreases manganese cytotoxicity and accumulation, *J Neurochem*, 2010, 112, 1190-1198.

22. Y. A. Seo and M. Wessling-Resnick, Ferroportin deficiency impairs manganese metabolism in flatiron mice, *FASEB J*, 2015, 29, 2726-2733.

23. J. Kim, P. D. Buckett and M. Wessling-Resnick, Absorption of manganese and iron in a mouse model of hemochromatosis, *PLoS One*, 2013, 8, e64944.

24. K. R. Bridle, D. M. Frazer, S. J. Wilkins, J. L. Dixon, D. M. Purdie, D. H. Crawford, V. N. Subramaniam, L. W. Powell, G. J. Anderson and G. A. Ramm, Disrupted hepcidin regulation in HFE-associated haemochromatosis and the liver as a regulator of body iron homeostasis, *Lancet*, 2003, 361, 669-673.

25. M. Muckenthaler, C. N. Roy, A. O. Custodio, B. Minana, J. deGraaf, L. K. Montross, N. C. Andrews and M. W. Hentze, Regulatory defects in liver and intestine implicate abnormal hepcidin and Cybrd1 expression in mouse hemochromatosis, *Nat Genet*, 2003, 34, 102-107.

26. T. V. Peres, M. R. Schettinger, P. Chen, F. Carvalho, D. S. Avila, A. B. Bowman and M. Aschner, Manganese-induced neurotoxicity: a review of its behavioral consequences and neuroprotective strategies, *BMC Pharmacol Toxicol*, 2016, 17, 57.

27. A. C. Illing, A. Shawki, C. L. Cunningham and B. Mackenzie, Substrate profile and metal ion selectivity of human divalent metal-ion transporter-1, *J Biol Chem*, 2012, 287, 30485-30496.

28. N. A. Wolff, M. D. Garrick, L. Zhao, L. M. Garrick, A. J. Ghio and F. Thévenod, A role for divalent metal transporter (DMT1) in mitochondrial uptake of iron and manganese, *Sci Rep*, 2018, 8, 211.

Metallomics Accepted Manuscript

29. W. Lin, D. R. Vann, P. T. Doulias, T. Wang, G. Landesberg, X. Li, E. Ricciotti, R. Scalia, M. He, N. J. Hand and D. J. Rader, Hepatic metal ion transporter ZIP8 regulates manganese homeostasis and manganese-dependent enzyme activity, *J Clin Invest*, 2017, 127, 2407-2417.
30. T. B. Aydemir, M. H. Kim, J. Kim, L. M. Colon-Perez, G. Banan, T. H. Mareci, M. Febo and R. J. Cousins, Metal transporter Zip14 (Slc39a14) deletion in mice increases manganese deposition and produces neurotoxic signatures and diminished motor activity, *J Neurosci*, 2017, 37, 5996-6006.
31. S. Jenkitkasemwong, A. Akinyode, E. Paulus, R. Weiskirchen, S. Hojyo, T. Fukada, G. Giraldo, J. Schrier, A. Garcia, C. Janus, B. Giasson and M. D. Knutson, SLC39A14 deficiency alters manganese homeostasis and excretion resulting in brain manganese accumulation and motor deficits in mice, *Proc Natl Acad Sci U S A*, 2018, 115:E1769-E1778.
32. K. Tuschl, E. Meyer, L. E. Valdivia, N. Zhao, C. Dadswell, A. Abdul-Sada, C. Y. Hung, M. A. Simpson, W. K. Chong, T. S. Jacques, R. L. Woltjer, S. Eaton, A. Gregory, L. Sanford, E. Kara, H. Houlden, S. M. Cuno, H. Prokisch, L. Valletta, V. Tiranti, R. Younis, E. R. Maher, J. Spencer, A. Straatman-Iwanowska, P. Gissen, L. A. Selim, G. Pintos-Morell, W. Coroleu-Lletget, S. S. Mohammad, S. Yoganathan, R. C. Dale, M. Thomas, J. Rihel, O. A. Bodamer, C. A. Enns, S. J. Hayflick, P. T. Clayton, P. B. Mills, M. A. Kurian and S. W. Wilson, Mutations in SLC39A14 disrupt manganese homeostasis and cause childhood-onset parkinsonism-dystonia, *Nat Commun*, 2016, 7, 11601.
33. Y. Xin, H. Gao, J. Wang, Y. Qiang, M. U. Imam, Y. Li, J. Wang, R. Zhang, H. Zhang, Y. Yu, H. Wang, H. Luo, C. Shi, Y. Xu, S. Hojyo, T. Fukada, J. Min and F. Wang, Manganese transporter Slc39a14 deficiency revealed its key role in maintaining manganese homeostasis in mice, *Cell Discov*, 2017, 18, 17025.
34. D. Leyva-Illades, P. Chen, C. E. Zogzas, S. Hutchens, J. M. Mercado, C. D. Swaim, R. A. Morrisett, A. B. Bowman, M. Aschner and S. Mukhopadhyay, SLC30A10 is a cell surface-localized manganese efflux transporter, and parkinsonism-causing mutations block its intracellular trafficking and efflux activity, *J Neurosci*, 2014, 34, 14079-14095.
35. C. E. Zogzas, M. Aschner and S. Mukhopadhyay, Structural elements in the transmembrane and cytoplasmic domains of the metal transporter SLC30A10 are required for its manganese efflux activity, *J Biol Chem*, 2014, 289, 15940-15957.
36. Z. Xia, J. Wei, Y. Li, J. Wang, W. Li, K. Wang, X. Hong, L. Zhao, C. Chen, J. Min and F. Wang, Zebrafish slc30a10 deficiency revealed a novel compensatory mechanism of Atp2c1 in maintaining manganese homeostasis, *PLoS Genet*, 2017, 13, e1006892.
37. S. M. Fleming, N. A. Santiago, E. J. Mullin, S. Pamphile, S. Karkare, A. Lemkuhl, O. R. Ekhaton, S. C. Linn, J. G. Holden, D. S. Aga, J. A. Roth, B. Liou, Y. Sun, G. E. Shull and P. J. Schultheis, The effect of manganese exposure in Atp13a2-deficient mice, *Neurotoxicology*, 2018, 64, 256-266.
38. J. P. Covy, E. A. Waxman and B. I. Giasson, Characterization of cellular protective effects of ATP13A2/PARK9 expression and alterations resulting from pathogenic mutants, *J Neurosci Res*, 2012, 90, 2306-2316.
39. A. D. Gitler, A. Chesi, M. L. Geddie, K. E. Strathearn, S. Hamamichi, K. J. Hill, K. A. Caldwell, G. A. Caldwell, A. A. Cooper, J. C. Rochet and S. Lindquist, Alpha-synuclein is part of a diverse and highly conserved interaction network that includes PARK9 and manganese toxicity, *Nat Genet*, 2009, 41, 308-315.
40. J. Tan, T. Zhang, L. Jiang, J. Chi, D. Hu, Q. Pan, D. Wang and Z. Zhang, Regulation of intracellular manganese homeostasis by Kufor-Rakeb syndrome-associated ATP13A2 protein, *J Biol Chem*, 2011, 286, 29654-29662.
41. M. Quadri, A. Federico, T. Zhao, G. J. Breedveld, C. Battisti, C. Delnooz, L. A. Severijnen, L. Di Toro Mammarella, A. Mignarri, L. Monti, A. Sanna, P. Lu, F. Punzo, G. Cossu, R. Willemsen, F. Rasi, B. A. Oostra, B. P. van de Warrenburg and V. Bonifati, Mutations in SLC30A10 cause parkinsonism and dystonia with hypermanganesemia, polycythemia, and chronic liver disease, *Am J Hum Genet*, 2012, 90, 467-477.
42. K. Tuschl, P. T. Clayton, S. M. Gospe Jr, S. Gulab, S. Ibrahim, P. Singhi, R. Aulakh, R. T. Ribeiro, O. G. Barsottini, M. S. Zaki, M. L. Del Rosario, S. Dyack, V. Price, A. Rideout, K. Gordon, R. A. Wevers, W. K. Chong and P. B. Mills, Syndrome of hepatic cirrhosis, dystonia, polycythemia, and hypermanganesemia caused by mutations in SLC30A10, a manganese transporter in man, *Am J Hum Genet*, 2012, 90, 457-466.
43. S. Hutchens, C. Liu, T. Jursa, W. Shawlot, B. K. Chaffee, W. Yin, A. C. Gore, M. Aschner, D. R. Smith and S. Mukhopadhyay, Deficiency in the manganese efflux transporter SLC30A10 induces severe hypothyroidism in mice, *J Biol Chem*, 2017, 292, 9760-9773.
44. C. Liu, S. Hutchens, T. Jursa, W. Shawlot, E. V. Polishchuk, R. S. Polishchuk, B. K. Dray, A. C. Gore, M. Aschner, D. R. Smith and S. Mukhopadhyay, Hypothyroidism induced by loss of the manganese efflux transporter SLC30A10 may be explained by reduced thyroxine production, *J Biol Chem*, 2017, 292, 16605-16615.
45. C. A. Taylor, S. Hutchens, C. Liu, T. Jursa, W. Shawlot, M. Aschner, D. R. Smith and S. Mukhopadhyay, SLC30A10 transporter in the digestive system regulates brain manganese under basal conditions while brain SLC30A10 protects against neurotoxicity, *J Biol Chem*, 2019, 294, 1860-1876.
46. C. Y. Wang, D. Meynard and H. Y. Lin, The role of TMPRSS6/matrilysin-2 in iron regulation and anemia, *Front Pharmacol*, 2014, 5, 114.
47. A. R. Folgueras, F. M. de Lara, A. M. Pendas, C. Garabaya, F. Rodriguez, A. Astudillo, T. Bernal, R. Cabanillas, C. Lopez-Otin and G. Velasco, Membrane-bound serine protease matrilysin-2 (Tmprss6) is an essential regulator of iron homeostasis, *Blood*, 2008, 112, 2539-2545.
48. D. M. Frazer, S. J. Wilkins, D. Darshan, C. S. G. Mirciov, L. A. Dunn and G. J. Anderson, Ferroportin is essential for iron absorption during suckling, but is hyporesponsive to the regulatory hormone hepcidin, *Cell Mol Gastroenterol Hepatol*, 2017, 3, 410-421.
49. S. McLachlan, K. E. Page, S. M. Lee, A. Loguinov, E. Valore, S. T. Hui, G. Jung, J. Zhou, A. J. Lusis, B. Fuqua, T. Ganz, E. Nemeth and C. D. Vulpe, Hamp1 mRNA and plasma hepcidin levels are influenced by sex and strain but do not predict tissue iron levels in inbred mice, *Am J Physiol Gastrointest Liver Physiol*, 2017, 313, G511-G523.

ARTICLE	Journal Name
50. S. A. Bustin, V. Benes, J. A. Garson, J. Hellemans, J. Huggett, M. Kubista, R. Mueller, T. Nolan, M. W. Pfaffl, G. L. Shipley, J. Vandesompele and C. T. Wittwer, The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments, <i>Clin Chem</i> , 2009, 55, 611-622.	View Article Online DOI: 10.1039/C8MT00370J
51. M. A. Okada, F. F. Neto, C. H. Noso, C. L. Voigt, S. X. Campos and C. Alberto de Oliveira Ribeiro, Brain effects of manganese exposure in mice pups during prenatal and breastfeeding periods, <i>Neurochem Int</i> , 2016, 97, 109-116.	
52. S. L. O'Neal and W. Zheng, Manganese toxicity upon overexposure: a decade in review, <i>Curr Environ Health Rep</i> , 2015, 2, 315-328.	
53. P. Chen, J. Bornhorst and M. Aschner, Manganese metabolism in humans, <i>Front Biosci</i> , 2018, 23, 1655-1679.	
54. K. Tuschl, P. B. Mills and P. T. Clayton, Manganese and the brain, <i>Int Rev Neurobiol</i> , 2013, 110, 277-312.	
55. A. J. Bertinchamps, S. T. Miller and G. C. Cotzias, Interdependence of routes excreting manganese, <i>Am J Physiol</i> , 1966, 211, 217-224.	
56. C. D. Davis, L. Zech and J. L. Greger, Manganese metabolism in rats: an improved methodology for assessing gut endogenous losses, <i>Proc Soc Exp Biol Med</i> , 1993, 202, 103-108.	
57. H. A. Schroeder, J. J. Balassa and I. H. Tipton, Essential trace metals in man: manganese. A study in homeostasis, <i>J Chronic Dis</i> , 1966, 19, 545-571.	
58. I. E. Zohn, I. De Domenico, A. Pollock, D. M. Ward, J. F. Goodman, X. Liang, A. J. Sanchez, L. Niswander and J. Kaplan, The flatiron mutation in mouse ferroportin acts as a dominant negative to cause ferroportin disease, <i>Blood</i> , 2007, 109, 4174-4180.	

Metallomics Accepted Manuscript

Knockout mice with constitutively low ferroportin show that ferroportin does not make a major contribution to manganese homeostasis *in vivo*