Chapter 17

Enrichment of the Plant Cytosolic Fraction

**Jeemeng Lao, Andreia M. Smith-Moritz, Jennifer C. Mortimer and
Joshua L. Heazlewood**

Abstract

The cytosol is at the core of cellular metabolism and contains many important metabolic pathways, including glycolysis, gluconeogenesis and the pentose phosphate pathway. Despite the importance of this matrix, few attempts have sought to specifically enrich this compartment from plants. Although a variety of biochemical pathways and signaling cascades pass through the cytosol, much of the focus has usually been targeted at the reactions that occur within membrane bound organelles of the plant cell. In this chapter, we outline a method for the enrichment of the cytosol from rice suspension cell cultures which includes sample preparation and enrichment as well as validation using immunoblotting and fluorescence tagged proteins.

Key words Cytosol, Cytoplasm, Rice, Cell culture.

1 Introduction

The cytosol or aqueous cytoplasm is the intracellular fluid found within the cell that contains a multitude of metabolic pathways, protein complexes, macromolecular structures and organelles [1]. The term ‘cytosol’ was initially used to define a solution derived from cellular fractionation [2]. However, more recently the term has been widely used to describe a distinct component of intact cells [3]. Thus, along with the term aqueous cytoplasm, the cytosol is now used to describe the contents of an intact cell including structures such as the cytoskeleton, but excluding organelles.

The plant cytosol is the site of a range of important metabolic activities and enables biochemical crosstalk to occur within the cell. However, there is a distinct lack of research focused on this compartment, with much of the literature concentrated on biochemical and signaling processes associated with the mitochondria, nucleus, peroxisome, plasma membrane and plastid [4]. To some extent, this may be due to

some complications in the enrichment of this compartment due to contamination from organelle rupture. However it is more likely that the cytosol represents an overlooked biochemical compartment, despite its clear importance in providing a partitioning mechanism for numerous biochemical processes within the cell [5].

The purification of subcellular compartments and organelles can require complex and involved separation and purification techniques. In the case of the plant endoplasmic reticulum (ER), mitochondrion, peroxisome and Golgi apparatus, the combination of density centrifugation and free-flow electrophoresis can dramatically increase the enrichment of an organelle fraction [6-9]. However, even with additional purification procedures, it may still be necessary to employ a subtraction method to remove contaminants to obtain a more accurate proteome [7, 10]. The utilization of a Potter-Elvehjem tissue homogeniser can be used to gently disrupt plant protoplasts and results in an increased yield of intact organelles [11]. The homogenization buffer contains sucrose to act as a cushion during cellular disruption preventing osmotic rupture and thus maintaining organelle integrity [12, 13]. For the enrichment of the cytosolic fraction, this is followed by centrifugation to remove unbroken cells, organelles and cell wall material [10].

The enrichment and isolation of plant compartments has been extensively detailed in a range of plant species for a select group of commonly investigated organelles and membranes [14-16]. More recently, with the utilization of mass spectrometry-based proteomic surveys, isolation techniques have been heavily focused on the reference plant *Arabidopsis* [4]. This is indeed the case when it comes to the enrichment of the plant cytosolic fraction, where only two examples are available, from *Arabidopsis* [10, 17] and soybean [18]. The method outlined here employs plant material from the important crop species of *Oryza sativa* (rice). The cytosolic enrichment from rice was performed similarly to approaches utilized with *Arabidopsis* material [17]. However, some optimization was still necessary to produce protoplasts, since different plant species and cell types have different cell wall compositions. Thus, the creation of a rice protoplast required the optimization of enzyme ratios in addition to varying enzyme cocktails. Following enrichment, the samples can be assessed for purity by immunoblotting, mass spectrometry and enzyme assays. Finally, proteins identified by mass spectrometry can be verified by employing transient subcellular localization techniques involving fluorescent proteins.

2 Materials

Prepare solutions with ultrapure water (18 M Ω cm at 25° C) and analytical grade reagents. Prepare all reagents at room temperature.

2.1 Plant Material

1. Plant Material: rice cell cultures (*see Note 1*).
2. Rice Growth Media: Murashige & Skoog without Nitrogen (without vitamins) 3.31 g/L, Sucrose 30 g/L, pH with KOH to 5.8, autoclave. After media is cooled, add vitamin mix to 1X.
3. 10X Vitamin Mix: 2,4-Dichlorophenoxyacetic acid 20 mg/L, Kinetin 2 mg/L, Gibberellic acid 1 mg/L, Murashige & Skoog Vitamin Mixture 10X/L, Coconut Water 200 mL, Glycine 0.75 g/L, L-Glutamine 8.77 g/L, L-Aspartic Acid 2.66 g/L, L-Arginine 2.28 g/L, filter sterilize (*see Note 2*).
4. Disposable 0.2 μ m solution filter system, such as the Nalgene™ Rapid-Flow™ Sterile Disposable Bottle Top Filters with SFCA Membrane. .
5. Erlenmeyer flasks, 250 mL.
6. Plant growth incubator or chamber (*see Note 3*).

2.2 Rice Cell Culture Protoplasts

1. Miracloth.
2. Enzyme Buffer: 0.4 M Mannitol, 3.6 mM MES-KOH, pH 5.7, 2.0% (w/v) cellulase “ONOUKA” RS, 0.5 % (w/v) pectolyase Y-23, 1.0 % (w/v) Driselase (*see Note 4*).
3. Variable speed benchtop orbital shaker.
4. Wash Buffer: 0.4 M mannitol, 3.6 mM MES-KOH, pH 5.7.
5. Homogenization Buffer: 0.4 M sucrose, 50 mM Tris-HCl, pH 7.5, 3 mM EDTA, 2 mM dithiothreitol (DTT). Add DTT just prior to homogenization (*see Note 5*).
6. Glass-Teflon Potter-Elvehjem tissue homogeniser (30-50 mL capacity), keep on ice (*see Note 6*).
7. Preparative centrifuge with rotors capable of processing 30 mL sample at 800 x g and 10,000 x g such as an Avanti J25 centrifuge (Beckman Coulter) with a JA-25.50 rotor (Beckman Coulter).
8. Ultracentrifuge with fixed angle rotor capable of processing 20 mL sample at 100,000 x g such as an Optima™ XE (Beckman Coulter) with a Type 70 Ti rotor (Beckman Coulter).

2.3 Protein Precipitation

1. Trichloroacetic acid (TCA) ~100% (w/v).
2. 100% (v/v) acetone.

3. Vacuum concentrator.
4. General Resuspension Buffer: 50 mM Tris-HCl, pH 8, 10 mM EDTA, 10 mM DTT.
5. Bradford protein assay kit, such as the Pierce Coomassie Plus (Bradford) Assay Kit (Thermo Fisher Scientific) (*see Note 7*).

2.4 Analysis of the Cytosolic Fraction by Immunoblotting

1. Electrophoresis chamber for protein separation, such as the Mini-PROTEAN® Tetra Cell (Bio-Rad).
2. Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) precast gels compatible with the electrophoresis chamber, such as 12% Mini-PROTEAN® TGX™ Gel (Bio-Rad).
3. Electrophoresis Buffer: 25 mM Tris-HCl (do not adjust pH), 192 mM glycine, 0.1% (w/v) SDS.
4. 4 x Sample Buffer: 200 mM Tris-HCl, pH 6.8, 400 mM DTT, 8% (w/v) SDS, 0.4% (w/v) bromophenol blue, 40% (v/v) glycerol.
5. Protein molecular weight markers.
6. Protein transfer apparatus, such as TE 70 PWR Semi-Dry Transfer Unit (GE Healthcare Life Sciences).
7. Transfer Buffer: Electrophoresis Buffer with 10% (v/v) methanol (*see Note 8*).
8. Whatman blotting paper.
9. Nitrocellulose blotting membrane.
10. TTBS Buffer: 20 mM Tris-HCl, pH 7.6 150 mM NaCl, 0.1% (v/v) Tween-20 (*see Note 9*).
11. Blocking solution, such as Blocker™ BLOTTO in TBS (Thermo Fisher Scientific).
12. Subcellular marker antibodies, such as UGPase (Agrisera AB).
13. Secondary antibody, such as Anti-Rabbit IgG Peroxidase (Sigma-Aldrich).
14. Detection reagent containing luminol and peroxide solution, such as Amersham ECL Prime Western Blotting Detection Reagent (GE Healthcare Lifesciences).
15. Chemiluminescence imager, such as Amersham Imager 600 (GE Healthcare Lifesciences).

2.5 Analysis of the Cytosolic Fraction by Mass Spectrometry

1. MS Resuspension Buffer: 50 mM Tris-HCl, pH 8, 10 mM EDTA, 5 M Urea, 10 mM DTT.
2. Dilution Solution: 50 mM Tris-HCl, pH 8 (*see Note 10*).
3. 50 mM iodoacetamide (IAA) (*see Note 11*).
4. High grade trypsin, such as Trypsin, from Porcine pancreas (Sigma-Aldrich).
5. Vacuum concentrator.
6. ACN1 solution: 80% (v/v) acetonitrile with 0.1% (v/v) trifluoroacetic acid.

7. ACN2 solution: 2% (v/v) acetonitrile with 0.1% (v/v) trifluoroacetic acid.
8. C₁₈ spin columns, such as Ultra-micro SpinColumns with C₁₈ (Harvard Apparatus).
9. Tandem mass spectrometer (MS/MS) with online liquid chromatography (LC) capabilities (nanoflow or capillary flow rates) capable of data dependent acquisitions.
10. Search engine for analyzing mass spectrometry data to identify proteins, such as. Mascot (Matrix Science).

2.6 Validation of cytosolic proteins by fluorescent tags

1. cDNA from plant material (*see Note 12*).
2. Thermocycler (PCR machine).
3. Gene specific primers containing Gateway® *attB1* and *attB2* recombination sequences.
4. PCR Master Mix containing Taq DNA polymerase, dNTPs, MgCl₂ and reaction buffers (*see Note 13*).
5. Gel extraction kit, such as QIAquick Gel Extraction Kit (Qiagen).
6. Plasmid purification kit, such as QIAprep Spin Miniprep Kit (Qiagen).
7. pDONR™/Zeo vector (Life Technologies).
8. Gateway® BP Clonase® II Enzyme Mix (Life Technologies).
9. Gateway® LR Clonase® II Enzyme mix (Life Technologies).
10. Competent *ccdB* Survival™ 2 *Escherichia coli* (*E. coli*).
11. Gateway compatible vector with fluorescent protein, such as pBullet series (*see Note 14*).
12. Fresh, medium sized yellow onions.
13. DNA LoBind 1.5 mL microfuge tubes (*see Note 15*).
14. Biolistic particle delivery system, such as PDS-1000/He™ Hepta System (Bio-Rad).
15. 1.0 µm gold microcarriers (Bio-Rad).
16. Macrocarriers (Bio-Rad).
17. 1,100 psi Rupture Disks (Bio-Rad).
18. 50% (v/v) sterile glycerol.
19. 70% (v/v) ethanol.
20. 100% (v/v) ethanol.
21. 2.5 M calcium chloride.
22. 0.1 M spermidine.
23. Vortex mixer.
24. Laser scanning confocal microscope (*see Note 16*).
25. Glass slides and coverslips (#1.5; 2x50 mm).

3 Methods

Perform all reactions at room temperature unless noted. A workflow is outlined in figure 1 highlighting the crucial steps involved in the enrichment of the cytosolic fraction. This method will detail the enrichment and validation of the cytosolic fraction from rice cell cultures but could be expanded to include any plant species or tissue type.

3.1 *Enrichment of the Rice Cytosolic Fraction*

3.1.1 *Rice Suspension Cell Culture*

1. Rice suspension cell cultures were grown in 100 mL of Rice Growth Media with 1X Vitamin Mix in a 250 mL Erlenmeyer flask shaking at 125 rpm in the dark at 30° C.

3.1.2 *Protoplast Preparation*

1. Collect cells and remove media by passing the solution through one piece of Miracloth. Squeeze out excess media by hand using a gentle pressure.
2. Use a 1:5 ratio of cells (g) to Enzyme Buffer (mL) and incubate in a large flask for 3 hours with gentle rotation (~35 to 45 rpm) in the dark and at room temperature (22° C) (*see Note 17*).
3. After 2 to 3 hours, ensure adequate digestion of cell walls by examining an aliquot by light microscopy (Figure. 2) (*see Note 18*).
4. After digestion of cell walls, centrifuge cells at 800 x g for 5 minutes at 4° C and discard supernatant (*see Note 19*).
5. Carefully resuspend cells with cooled Wash Buffer, use approximately a 1:5 ratio of cells (g) to Wash Buffer (mL) to ensure an adequate removal of digestion enzymes. Centrifuge washed cells at 800 x g for 5 minutes at 4° C and discard supernatant. Repeat the wash step.

3.1.3 *Homogenization*

1. Resuspend pellet in Homogenization Buffer (2 mL for every 1 g of cell) (*see Note 20*).
2. Disrupt cells by applying four strokes of a Potter-Elvehjem Tissue homogenizer maintaining a temperature of 4° C by keeping apparatus on ice (*see Note 21*).

3.1.4 *Cytosol Enrichment*

1. Centrifuge the disrupted cell lysate at 800 x g for 15 minutes at 4° C (*see Note 22*).
2. Transfer supernatant to a centrifuge tube and centrifuge at 10,000 x g for 15 minutes at 4° C (*see Note 23*).

3. Transfer the resultant supernatant to an ultracentrifuge tube and centrifuge at 100,000 x g for 1 hour at 4° C (*see Note 24*).
4. Remove the lipid layer on top and carefully collect supernatant (*see Note 25*).
5. Estimate protein concentration by conducting a Bradford assay [19] (*see Note 26*).
6. Transfer and store protein as 0.5 mg aliquots at -80° C (*see Note 27*).

3.1.5 *Removal of homogenization buffer from sample*

1. Add trichloroacetic acid to the sample to a final concentration of 20% (v/v) (*see Note 28*).
2. Incubate solution for 30 minutes on ice and centrifuge at 20,000 x g at 4° C.
3. Discard supernatant.
4. Wash pellet with cold acetone and then centrifuge at 20,000 x g at 4° C for 5 minutes and discard supernatant. Repeat wash.
5. Dry pellet by using a vacuum concentrator or by leaving in a fume hood for a few hours.
6. Resuspend the protein pellet in General Resuspension Buffer (*see Note 30*).
7. Estimate protein concentration by conducting a Bradford assay [19] (*see Note 31*).

3.2 **Assessment of Sample Purity by Immunoblotting**

1. An assessment of purity by immunoblotting can be performed on the various fractions produced during the cytosolic preparation (Figure 3) (*see Note 32*).
2. Use approximately 5 µg protein from each fraction, add Sample Buffer (1X) and water to a final volume of approximately 20 µL.
3. Incubated at 70° C for 10 minutes.
4. Briefly centrifuge to remove insoluble material and load samples into the wells of a previously assembled SDS-PAGE precast gel cassette, with a protein molecular weight marker loaded in the far left well (*see Note 33*).
5. Set the electrophoresis chamber at 125 V and run for around 90 min or until the dye front migrates to the bottom of the gel.
6. Remove the gel cassette from the assembly and pry it open to access the gel (*see Note 34*).
7. Prepare semi-dry transfer unit by rinsing the anode and cathode with ultrapure water.
8. Cut 6 pieces of blotting paper and one nitrocellulose membrane to the size of the gel and soak in transfer buffer for 2 minutes.
9. Place and align three sheets of blotting paper on the lower electrode (anode), followed by the membrane, gel, and 3 more blotting paper (*see Note 35*).
10. Set the semi-dry transfer unit to 38 mA and run for 1 hour for a 6 x 8 cm gel (*see Note 36*).

11. Remove the membrane and briefly rinse for 5 minutes with TTBS Buffer.
12. Incubate the membrane with Blocker™ BLOTTO for 1 hour with gentle shaking and then discard.
13. Wash with TTBS Buffer using gentle shaking for 5 minutes. Repeat wash three additional times.
14. Add primary antibody in TTBS Buffer with gentle shaking for 1 hour then discard (*see Note 37*).
15. Wash with TTBS Buffer using gentle shaking for 5 minutes. Repeat wash three additional times.
16. Add secondary antibody in TTBS Buffer with gentle shaking for 1 hour.
17. Wash with TTBS Buffer using gentle shaking for 5 minutes. Repeat wash three additional times.
18. Drain membrane of excess buffer and ensure the side of the membrane with the protein is facing up.
19. Mix 500 μ L of the luminol solution and 500 μ L of the peroxide solution together and add on top of the membrane (*see Note 38*).
20. Incubate at room temperature for 5 minutes.
21. Drain off excess reagent and visualize using a chemiluminescence imager (*see Note 39*).

3.3 Assessment of Sample Purity by Mass Spectrometry

1. Use approximately 20 μ g protein from the TCA resuspended pellet, minimize the volume required for resuspension (*see Note 40*).
2. Add MS Resuspension Solution ensuring the concentration of urea remains close to 5 M.
3. Add IAA to a final concentration of 10 mM and incubate for 30 minutes in the dark and room temperature (*see Note 41*).
4. Dilute samples to 1 M urea with Dilution Solution (*see Note 42*).
5. Add 10% (w/v) trypsin to the sample.
6. Incubate overnight at 37° C.
7. Perform a peptide clean up using C₁₈ spin columns (25 to 75 μ L capacity). After initial hydration of the column matrix with water (100 μ L) centrifuge (1000 x g, 2 minutes). Wash the C₁₈ spin column with 100 μ L ACN1 buffer and centrifuge (1000 x g, 2 minutes), then prime twice with 100 μ L ACN2 buffer, centrifuging (1000 x g, 2 minutes) after each step.
8. Add digested sample to the C₁₈ spin column and centrifuge (1000 x g, 2 minutes), wash twice with 100 μ L ACN2 buffer, centrifuging (1000 x g, 2 minutes) after each step. Elute with 100 μ L ACN1 buffer, centrifuging 1000 x g for 2 minutes. Repeat elution for total volume of 200 μ L. Concentrate using a vacuum concentrator until 1 to 5 μ L of the solution remains (*see Note 43*).
9. Resuspend or dilute peptides with ACN2 buffer to a concentration of about 0.25 μ g / μ L (*see Note 44*).
10. Samples (1 to 20 μ g) can now be analyzed by tandem mass spectrometry (*see Note 44*).

11. Analyze resultant tandem spectra against a plant protein database using a proteomic search engine (*see Note 45*).
12. Matched proteins derived from the cytosolic fraction can be profiled using Gene Ontologies (GO) [20] or by homology mapping the proteins to collections of previously assigned subcellular proteomes [21].

3.4 Assessment of Cytosolic Localization by Particle Bombardment

3.4.1 DNA Preparation

1. The localization of proteins identified by tandem mass spectrometry can be verified using fluorescence tagged protein co-localization (Figure 4).
2. Prepare cDNA from the relevant species using a commercially available kit (*see Note 46*).
3. Amplify gene of interest from cDNA using a PCR master mix with gene specific primers containing Gateway® attB1 and attB2 recombination sequences (*see Note 47*).
4. Add 1 x DNA loading dye to PCR reaction and run on a 1% (w/v) agarose gel.
5. Cut out the band corresponding to the gene of interest and perform gel cleanup using a gel extraction kit.
6. Perform a BP reaction of the PCR gene product with a pDONR™/Zeo vector using Gateway® BP Clonase® II Enzyme Mix (15-150 ng attB-PCR product, 150 ng pDONR™/Zeo, 2 µL Gateway® BP Clonase® II Enzyme Mix, and with water to a final reaction volume of 10 µL) and incubate at room temperature for 1 hour (*see Note 48*).
7. Transform the reaction product into competent *E. coli* and isolate the entry clone plasmid using plasmid isolation kit. Typical transformation reaction of chemically competent *E. coli* involves thawing on ice, addition of 1-100 ng of plasmid with gentle hand mixing, 30 minutes incubation on ice, 30 second heat shock at 42° C, 2 minutes on ice, addition of 500 µL Luria-Bertani (LB) broth, 1 hour recovery at 37° C, plating onto LB agar plates with appropriate selection, and overnight incubation at 37° C. Single colonies are isolated, grown in LB with appropriate selection, and incubated overnight at 37° C before being used for plasmid extraction.
8. Perform a LR reaction of the entry clone and pBullet-cyt-c using Gateway® LR Clonase® II Enzyme mix (50-150 ng entry clone, 150 ng pBullet-cyt-c, 2 µL LR Clonase® II Enzyme mix, and water to a final reaction volume of 10 µL) and incubate at room temperature for 1 hour (*see Note 49*).
9. Transform the reaction product into competent *E. coli* and isolate the plasmid using a plasmid isolation kit.

3.4.2 Particle Bombardment

1. Weigh out 15 mg of gold microcarrier into a microfuge tube (*see Note 50*).
2. Add 1 mL of 70% (v/v) ethanol.
3. Vortex for 3 minutes.
4. Let the microcarriers soak for 15 minutes.
5. Sediment the microcarriers with a brief centrifuge (5 seconds) (*see Note 51*).
6. Discard supernatant and wash microcarriers 3 times with sterile water.
7. Resuspend microcarriers in 500 μ L sterile 50% (v/v) glycerol. Aliquot microcarriers into microfuge tubes (25 μ L per tube) (*see Note 52*).
8. Add 2.5 μ L (approximately 500 ng) of the plasmid containing the fluorescently tagged clone to one of the microcarrier aliquots (*see Note 53*).
9. Add 25 μ L of 2.5 M CaCl_2 .
10. Add 10 μ L of 0.1 M spermidine (*see Note 54*).
11. Vortex for 10 minutes.
12. Centrifuge for 5 seconds.
13. Discard supernatant and wash microcarriers with 140 μ L of 100% (v/v) ethanol.
14. Resuspend with 20 μ L of 100% ethanol.
15. Wash macrocarrier and pressure disk (1100 PSI) with 100% (v/v) ethanol.
16. Load 20 μ L microcarrier onto macrocarrier (*see Note 55*).
17. Allow ethanol to evaporate (*see Note 56*).
18. Set up the biolistic particle delivery system according to manufacturer's recommendations if using PDS-1000/HeTM Hepta System (Bio-Rad), vacuum of 28 inch/Hg, target distance of 6 cm, and pressure disk of 1100 PSI. (*see Note 57*).
19. Remove onion layers from mid-section of the onion and place in a petri dish and put in stage of the biolistic particle delivery system.
20. After bombardment, add water to the petri dish to keep onion layer moist, cover with Parafilm M and keep in the dark. Leave overnight at room temperature (22° C) (*see Note 58*).

3.4.3 Confocal Laser Scanning Microscopy

1. Cut the onion layer to the size of a coverslip (22 x 50 mm) and remove the epidermal cells by peeling with a pair of tweezers (*see Note 59*).
2. Add water to the slide and place the epidermal peel in the water.
3. Add more water on top of the epidermal peel and place a coverslip on top (*see Note 60*).

4. Place the slide onto the confocal stage and initially focus using brightfield imaging and low magnification (*see Note 61*).
5. Initially use an appropriate filter, such as a DAPI filter to find transformed cells with a cyan fluorescent protein (CFP) signal. After locating a cell, a higher magnification can be used if required (*see Note 62*).
6. Configure the confocal for sequential acquisition mode, in this case for CFP and yellow fluorescent protein (YFP) (*see Note 63*).
7. Perform a live scan of the onion cell to detect one of the fluorescent molecules (such as, YFP) and fine tune focus (*see Note 64*).
8. Image the onion cells for the other fluorescent molecule (such as, CFP) to verify expression (*see Note 65*).
9. Take an image of the cell using both the fluorescent imaging configurations (*see Note 66*).
10. Images can be analyzed and processed with the microscope software or ImageJ [22].

4 Notes

1. One flask of 7-day grown suspension cell culture (100 mL) should provide 5 to 10 g of plant material. This amount is sufficient for most downstream applications. Different types of plant species and material can be employed, but the fractionation process may require extensive optimization. Cell suspension cultures can be created from callus generated by slicing sterile plant material (*e.g.* roots and leaves) and grown in sterile agar media plates with hormones (3.2 g/L Gamborg's B5 Medium including vitamins (Sigma-Aldrich), 20 g/L glucose, 0.5 g/L MES-KOH, pH 5.7, 8.5 g/L agar, 250 µg/L 2,4-Dichlorophenoxyacetic acid, and 100 µg/L kinetin).
2. Coconut water can be pre-filtered with Whatman paper before filter sterilization with 0.2 µm bottle top filter. This reduces clogging of the 0.2 µm filter.
3. The type of incubator or chamber will depend on the species or plant material being used. In the case of the rice suspension cell cultures, a shaking incubator with no light at 30° C is required.
4. It is important to dissolve all enzymes in the buffer prior to their addition to the cells. Enzyme ratios and different enzyme combinations are necessary for optimal protoplast generation. In this instance, the addition of Driselase was necessary to obtain suitable protoplasts from our rice suspension cultures. In comparison, the addition of Driselase was not required when preparing protoplast from Arabidopsis cell cultures [10].
5. DTT prevents the oxidation of proteins by reducing free sulfhydryl groups. It should be added to the buffer on the day it is used as it has a half-life of about 10 hours in a solution at pH 7.5 at 20° C.

6. The selection of an appropriate homogenizer is an important component of efficient protoplast rupturing. Generally, the negative pressure on the up stroke should result in an approximate 2 cm space or air bubble between the plunger and the homogenate. The key is to ensure enough mechanical stress to disrupt the protoplasts without disrupting subcellular organelles.
7. The selection of the appropriate protein assay reagents is important due to incompatibilities with some buffer. For example Coomassie is not compatible with urea buffers above 6 M. After sample suspension, it is possible to dilute the buffer components to achieve a more accurate estimation of protein concentration.
8. Methanol can enhance protein binding to the membrane.
9. Tween-20 helps prevent non-specific binding of antibodies.
10. The dilution of the urea concentration from 5 M in the MS Resuspension Buffer is necessary for optimal trypsin activity, which is retained in solutions up to 2 M urea.
11. The IAA is used to alkylate cysteine residues on proteins to prevent reformation of disulphide bonds which can reduce the effectiveness of protein digestion by trypsin.
12. There are many commercial kits available that have been optimized for the isolation of plant mRNA and subsequent synthesis of cDNA.
13. The utilization of a high fidelity polymerase reduces amplification errors and these enzymes are often more robust.
14. The pBullet collection of vectors has been optimized for the transient transformation of plant cells using particle bombardment approaches. They have been purpose built to enable the efficient localization of a candidate gene linked to YFP and the simultaneous co-localization of a subcellular marker protein linked to CFP [23]. It is also possible to employ other vectors containing fluorescent proteins to transiently localize a gene of interest using this method.
15. Gold microcarriers may adhere to the walls of standard microfuge tubes. This results in a loss of microcarriers when loading onto the macrocarriers. Microcarriers do not bind to DNA LoBind tubes.
16. The Laser Scanning Confocal Microscope (LSCM) must contain the capability to excite at the appropriate wavelengths. For example the pBullet vectors require 405 nm for cyan fluorescent protein (CFP) or 514 nm for yellow fluorescent protein (YFP).
17. Gentle rotation and a wide-base flask help reduce damage to the protoplasts which are quite delicate. For 5 g fresh weight (FW) of cells, a regular 250 mL Erlenmeyer culture flask is appropriate.

18. When cell walls are removed from plant cells by digestion, the cell will enlarge and the shape will become more spherical. The cells are also released from clumps and are found as individual cells. The edge of the cell (plasma membrane) becomes more difficult to discern by light microscopy.
19. The wash step removes residual enzymes from the protoplasts. The rice cell cultures are more compact after centrifugation than occurred with the Arabidopsis cell cultures [17]. This means it may be necessary to use a fine paintbrush to get the pellet loose after washing and centrifugation.
20. The sucrose in the buffer helps maintain subcellular organelle integrity during disruption. Generally concentrations of sucrose from 0.3 to 0.45 M are required.
21. For complete cellular disruption, the cells may need to be homogenized additional times. Successful homogenization can be verified by examining the resultant lysate by light microscopy to look for the presence of intact cells (**Figure 2**).
22. This step should result in a pellet enriched in unbroken cells, cell wall material, and nuclei.
23. This step should result in a pellet enriched in organelles such as mitochondria and plastid.
24. This step should result in a pellet enriched in the secretory pathway, ER, Golgi, and the plasma membrane.
25. The supernatant contains the enriched cytosolic fraction.
26. The final fraction comprising the cytosolic preparation (after 100,000 x g centrifugation) usually contains around 5 µg/µl of protein.
27. These aliquots contain a high concentration of sucrose which can effect downstream applications such as immunoblotting and enzyme assays. Although there are several techniques that could be employed to remove the sucrose, such as reverse phase chromatography, centrifugal filter concentrators or acetone precipitation, we have found these methods to be suboptimal with regard to the removal of sucrose and recovery of protein.
28. Samples can be stored at -80° C. Storing aliquots samples prevents multiple freeze-thaw cycles when the samples are required.
29. We have found that TCA precipitation is the most reliable method for the removal of components found in the Homogenization Buffer, including sucrose that can inhibit downstream applications and processing of the sample.
30. The TCA precipitated/acetone washed protein pellet can be extremely problematic to solubilize. In this instance, we have suggested a benign aqueous buffer; however the selected resuspension buffer is highly dependent on the downstream application. If undertaking analysis and separation by SDS-PAGE, it would be advisable to resuspend the pellet with 1 x Sample Buffer. In contrast, if the sample is to be analyzed by mass spectrometry, resuspend the pellet with MS Resuspension Buffer.

The denaturing properties of SDS or urea will greatly assist in the solubilization of the protein pellet.

31. If buffers containing urea or SDS (or other components) are employed to resuspend the protein pellet, ensure their compatibility with the protein estimation assay. For example, urea concentrations above 3 M are known to interfere with the Bradford assay. Other compatible methods of protein estimation can be employed to circumvent this issue.
32. Unfortunately there are limited sources of commercial antibodies against plant proteins. One of the main suppliers of antibodies raised against plant proteins is Agrisera AB (<http://www.agrisera.com/>)
33. The precast SDS-PAGE gel contains a comb and tape that need be removed prior to use. The wells should be rinsed with running buffer. The gel should be loaded onto the electrode assembly, if one gel is used a buffer dam is required. The assembly is loaded into the electrophoresis tank and 1X Running Buffer added, which should be enough to fill to the top of the gel wells. All wells should be loaded with Sample Buffer to ensure even migration of the proteins; this can be accomplished by loading blanks.
34. The flat end of a spatula is effective for this procedure. In addition, the stacking gel (containing the wells) should be removed prior to immunoblotting.
35. Protein transfer begins immediately once the gel touches the membrane, so care should be taken to align them correctly.
36. Conditions may vary depending on the unit being used and the size of the gel. Longer run time (an additional hour) may be needed to effectively transfer larger proteins or thicker gels.
37. Primary antibodies should be diluted based on manufacturer's recommendations. A dilution range from 1:1,000 to 1:10,000 depending on the affinity of the antibody. Generally, monoclonal antibodies require a higher dilution than that employed by polyclonal antibodies as the entire population recognizes the same epitope.
38. The mixed reagent should be protected from light if not immediately used as it is light sensitive.
39. After visualization, the membrane can be re-probed with different antibodies after using a stripping buffer, such as Restore Western Blot Stripping Buffer (Thermo Fisher Scientific).
40. If the protein pellet was solubilized in the General Resuspension Buffer, care should be taken to reduce the dilution of urea in the MS Resuspension Buffer. For example, if 5 μ L of protein is used, employ around 100 μ L of buffer to ensure urea remains close to 5 M.
41. Urea readily breaks down at elevated temperatures or when subjected to multiple freeze-thaw cycles. This breakdown product (isocyanic acid) is highly reactive and will result in the

uncontrolled modification of protein (carbamylation), specifically arginine, lysine and the amino terminus.

42. Urea concentrations above 2 M can inhibit trypsin activity.
43. The Spin Column removes contaminants (*e.g.* salts from the buffers) that interfere with the mass spectrometer.
44. If analyzing samples by nanoLC-MS/MS, generally around 1 µg of digested protein is employed. If using higher flow rates to introduce the sample to the MS, such as capillary liquid chromatography or standard flow, then the amount of digested protein needs to be increased to 20 to 40 µg [24].
45. Nearly all mass spectrometers include proprietary software for analysis of data and assignment of proteins (*e.g.* Sequest, Protein Pilot, Spectrum Mill, MassLynx). There is also commercial software not affiliated with an instrument manufacturer (*e.g.* Mascot, Scaffold) as well as a variety of open source software (*e.g.* OMSSA, X! Tandem). On the current generation of tandem mass spectrometers, between 1000 and 2000 unique protein identifications can be confidently assigned. However, this will be greatly dependent on the available genomic information for the species analyzed.
46. RNA is initially isolated and cDNA is produced by reverse transcribing the mRNA.
47. PCR reaction consists of PCR reaction buffer to 1X, 200 µM dNTPs, 0.2 µM forward primer, 0.2 µM reverse primer, 1-1000 ng of template DNA, and 1 unit of DNA polymerase. Typical parameters for thermocycler are 30 seconds for initial denaturation at 98° C, denaturation for 15 seconds at 98° C, annealing for 30 seconds at 45 to 68° C (depending on primer melting temperature), extension for 1 minute per kb of DNA being extended at 68° C, final extension for 5 minutes at 68° C. These steps may vary depending on manufacturer and polymerase being used.
48. Gateway reactions (BP and LR) can be scaled down to 2.5 µL final volume to reduce costs. Depending on the competence of the bacteria being used, the Gateway reactions may need to be incubated overnight to obtain more colonies.
49. For each subcellular marker, the pBullet collection is available with a C-terminal or N-terminal CFP to ensure the fluorescent marker does not interfere with a localization signal. The ‘-c’ designates this vector as containing a C-terminal CFP [23].
50. 15 mg of gold microcarriers is enough to perform 20 independent bombardments.
51. A longer centrifugation can result in the gold microcarriers clumping.
52. Aliquots can be stored at -20° C for future use.
53. Standard particle bombardment protocols recommend the DNA be at a concentration of 1 µg/µL, which typically requires the DNA to be concentrated after a plasmid preparation. The DNA

concentration step can be skipped if optimal transformation rates are not required. A DNA concentration of around 200 ng/ μ L (typical for plasmid purification from *E. coli*) is still effective for this method when employing the pBullet vectors.

54. Spermidine is hydroscopic (absorbs moisture) and is recommended to be prepared and used fresh. Repeated freeze-thaw cycles should generally be avoided. However, in our hands particle bombardment experiments will still work despite repeated freeze-thaw and storage at -20°C for over three months.
55. The gold microcarriers should be loaded closer to the center of the macrocarrier. Microcarriers near the end of the macrocarrier will not pass through the stopping screen and into the onion cells.
56. This can take between 30 minutes to 1 hour.
57. These settings are tested for onion cells and may need to be optimized for other material. We have found that epidermal peels can be kept on the onion slice for particle bombardment and peeled when ready for visualization by confocal.
58. An overnight incubation (16 hours) should be enough time for transient protein expression.
59. The cells may need to be fixed with 4% (v/v) paraformaldehyde in 50 mM phosphate buffered saline (PBS) for 5-10 minutes to prevent organelle movement.
60. Take care if using an inverted confocal microscope as the coverslip can become dislodged. Dry the coverslip if wet.
61. Do not let the objective touch the slide. It is good practice to focus by moving the objective away from the slide.
62. Generally, using the DAPI (4',6-diamidino-2-phenylindole) filter is fast and efficient way to scan cells for a fluorescent signal compared to employing the confocal imaging. The DAPI filter has a 350-400 nm excitation and an emission range of 420-510 nm which is compatible with both CFP and GFP/YFP. If other fluorophores are employed, such as mCherry, other more suitable filters would need to be used.
63. The current generation of confocal microscopes is capable of performing simultaneous scans (fast, crosstalk between signals) and sequential scans (slow, but less crosstalk). Simultaneous scanning allows the visualization of two signals without the need for fixation. The emission range can be narrowed to reduce crosstalk, although a reduction in signal intensity occurs. Importantly, sequential scanning can help verify the signals from both fluorophores are valid.
64. To prevent photobleaching of the fluorophores, start initial visualization with high camera gain and low laser power. Once focusing is done, reduce camera gain to about 600 and start increasing laser power and pixel dwell time to increase signal to noise.

65. For optical resolution, the pinhole must be optimized. The pinhole should be set to Airy unit 1, which is calculated for specific objectives and wavelengths. It is important to verify you are on Airy unit 1 when switching between objectives. Increasing the pinhole diameter will give more signal but at the expense of resolution.
66. Various parameters can be used to improve image quality, such as higher averaging, higher frame size, and lower scan speed.

Acknowledgements

This work was supported by the U. S. Department of Energy, Office of Science, Office of Biological and Environmental Research, through contract [DE-AC02-05CH11231] between Lawrence Berkeley National Laboratory and the U. S. Department of Energy. JLH is supported by an Australian Research Council (ARC) Future Fellowship [FT130101165].

References

1. Ito J, Parsons HT, and Heazlewood JL (2014) The Arabidopsis cytosolic proteome: the metabolic heart of the cell. *Front Plant Sci* 5:21
2. Clegg JS (1984) Properties and metabolism of the aqueous cytoplasm and its boundaries. *Am J Physiol* 246:R133-R51
3. Gene Ontology Consortium (2004) The Gene Ontology (GO) database and informatics resource. *Nucleic Acids Research* 32:D258-D61
4. Millar AH, and Taylor NL (2014) Subcellular proteomics-where cell biology meets protein chemistry. *Front Plant Sci* 5:55
5. Plaxton WC (1996) The organization and regulation of plant glycolysis. *Annu Rev Plant Physiol Plant Mol Biol* 47:185-214
6. Eubel H, Lee CP, Kuo J, et al. (2007) Free-flow electrophoresis for purification of plant mitochondria by surface charge. *Plant J* 52:583-94
7. Eubel H, Meyer EH, Taylor NL, et al. (2008) Novel proteins, putative membrane transporters, and an integrated metabolic network are revealed by quantitative proteomic analysis of Arabidopsis cell culture peroxisomes. *Plant Physiology* 148:1809-29
8. Parsons HT, Christiansen K, Knierim B, et al. (2012) Isolation and proteomic characterization of the Arabidopsis Golgi defines functional and novel targets involved in plant cell wall biosynthesis. *Plant Physiology* 159:12-26
9. Parsons HT, González Fernández-Niño SM, and Heazlewood JL (2014) in "Plant Proteomics: Methods and Protocols, Second Edition" (Jorrín Novo, J., V., Komatsu, S., Weckwerth, W., and Weinkoop, S., Eds.), Vol. 1072, pp. 527-39, Humana Press, New York
10. Ito J, Batth TS, Petzold CJ, et al. (2011) Analysis of the Arabidopsis cytosolic proteome highlights subcellular partitioning of central plant metabolism. *J Proteome Res* 10:1571-82
11. Meyer EH, and Millar AH (2008) in "2D PAGE: Sample Preparation and Fractionation" (Posch, A., Ed.), Vol. 425, pp. 163-69, Humana Press, New York
12. Dounce AL, Witter RF, Monty KJ, et al. (1955) A method for isolating intact mitochondria and nuclei from the same homogenate, and the influence of mitochondrial destruction on the properties of cell nuclei. *J Biophys Biochem Cy* 1:139-53
13. Schneider WC (1948) Intracellular distribution of enzymes III. The oxidation of octanoic acid by rat liver fractions. *J Biol Chem* 176:259-66
14. Day DA, Neuburger M, and Douce R (1985) Biochemical-characterization of chlorophyll-free mitochondria from pea leaves. *Aust J Plant Physiol* 12:219-28

15. Poincelo RP, and Day PR (1974) Improved method for isolation of spinach chloroplast envelope membranes. *Plant Physiology* 54:780-83
16. Widell S, and Larsson C (1981) Separation of presumptive plasma-membranes from mitochondria by partition in an aqueous polymer 2-phase system. *Physiol Plantarum* 51:368-74
17. Estavillo GM, Verhertbruggen Y, Scheller HV, et al. (2014) *in* "Plant Proteomics: Methods and Protocols" (Jorrín-Novo, J., V., Komatsu, S., Weckwerth, W., and Weinkoop, S., Eds.), Vol. 1072, pp. 453-67, Humana Press, New York
18. Oehrle NW, Sarma AD, Waters JK, et al. (2008) Proteomic analysis of soybean nodule cytosol. *Phytochemistry* 69:2426-38
19. Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical biochemistry* 72:248-54
20. Lao J, Sharma M, Sharma R, et al. (2015) Proteome profile of the endomembrane of developing coleoptiles from switchgrass (*Panicum virgatum*). *Proteomics* doi: 10.1002/pmic.201400487
21. Parsons HT, Weinberg CS, Macdonald LJ, et al. (2013) Golgi enrichment and proteomic analysis of developing *Pinus radiata* xylem by free-flow electrophoresis. *PLOS ONE* 8:e84669
22. Schneider CA, Rasband WS, and Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 9:671-5
23. Lao J, Oikawa A, Bromley JR, et al. (2014) The plant glycosyltransferase clone collection for functional genomics. *The Plant Journal* 79:517-29
24. González Fernández-Niño SM, Smith-Moritz AM, Chan LJG, et al. (2015) A standard flow liquid chromatography workflow for shotgun proteomics in bioenergy research. *Frontiers in Bioengineering and Biotechnology* 3:1-7

Figure Legends

Figure 1. Workflow outlining the enrichment of cytosol from rice suspension cell cultures.

Figure 2. Images of rice cells under light microscopy. (a) Seven day old rice cells highlighting cell clumping. (b) Intact rice protoplasts produced after enzymatic digestion of the cell wall. Cell clumping is minimized and cells margins difficult to visualize after protoplast production. (c) Cellular components released from rice protoplasts after homogenization with the Potter-Elvehjem tissue homogeniser. Scale bars: 10 μm .

Figure 3. Immunoblot of 5 μg of proteins with various antibody markers from fractions produced during the enrichment of the cytosolic fraction. The different fractions were total protein from rice protoplasts; 10,000 x g crude mixed organelle pellet; 100,000 x g membrane pellet; and the cytosolic fraction. Polyclonal antibodies were, UGPase (cytosol), Bip2 (endoplasmic reticulum), histone H3 (nucleus), VDAC-1 (mitochondria), and H^+ ATPase (plasma membrane). All antibodies were obtained from Agrisera AB. The enrichment procedure reduces organelle and membrane contamination.

Figure 4. Confirmation of cytosolic localization for candidate rice proteins using particle bombardment of onion epidermal cells. The pBullet-cyt-c/n vectors (Genbank: KJ081785 and KJ081787) containing ECFP with a cytosolic marker for co-localization were used for transformation. Os06g44270.1-EYFP contains the EYFP attached to the C-terminus, and EYFP-Os06g44270.1 contains the EYFP attached to the N-terminus. Scale bars: 20 μm .

Fig. 1

Fig. 2.

Fig. 3.

Fig. 4.