[Click here to view linked References](#)

Effect of elevated temperature on the ripening of full fat Cheddar cheese

Kevany Soodam^{1,2,5}, Lydia Ong^{1, 2,4}, Ian B. Powell^{3, 4}, Sandra E. Kentish^{1,4} and Sally L. Gras^{1, 2, 4*}

1. Department of Chemical and Biomolecular Engineering, The University of Melbourne, Parkville, Vic 3010, Australia.

2. The Bio21 Molecular Science and Biotechnology Institute, The University of Melbourne, Parkville, Vic 3010, Australia.

3. Dairy Innovation Australia Limited, 180 Princes Highway, Werribee, Victoria 3030, Australia.

4. ARC Dairy Innovation Hub, Department of Chemical and Biomolecular Engineering, The University of Melbourne, Parkville, Vic 3010, Australia.

5. Moorepark Food Research Centre, Teagasc Moorepark, Fermoy, Co. Cork, Ireland (current address).

*Corresponding author: Sally L. Gras. Address: The Bio21 Molecular Science and Biotechnology Institute, The University of Melbourne, Parkville, Victoria 3010, Australia. Tel.: +61 38344 6281; fax: +61 3 8344 4153. E-mail address: sgras@unimelb.edu.au

Short title:

Effect of temperature during Cheddar ripening

Abstract

Elevated temperatures can accelerate cheese ripening and decrease energy and storage requirements but the effect of these temperatures on the microstructure and concentration of biogenic amines is not known. Cheddar cheese was matured at four different ripening temperatures: 8°C, 15°C, 20°C or a combination of 8°C and 15°C. The ripening temperature was shown to alter the protein network of the cheese after ripening for only a few weeks. Ripening at higher temperatures accelerated the ripening process leading to increased proteolysis and the formation of free whey. Ripening the cheese at an elevated temperature of 15°C for only 33 days, however, may be beneficial as this timeframe provides the acceleration of ripening desired, without significant negative effects on texture or whey release. While the concentration of biogenic amines tryptamine and tyramine increased with ripening temperature these concentrations were generally low, confirming that biogenic amines do not represent a health concern under the conditions examined.

Keywords

Cheddar cheese; ripening temperature; cheese microstructure; cryo-SEM; Confocal laser scanning microscopy; biogenic amines

1. Introduction

Ripening is a slow and lengthy process for rennet coagulated cheeses, typically lasting from three weeks to two or more years (Fox 2002). During this period microbiological and biochemical changes occur, leading to changes in cheese texture and the development of flavour. Any decrease in the time taken for ripening can reduce the energy required for temperature control during maturation and decrease the need for storage space, leading to increased profitability. Accelerated ripening can be accomplished by an increase in the ripening temperature, the use of additional cultures (derived from starters or non-starter lactic acid bacteria (NSLAB) in live or attenuated form), or the use of enzymes (in many cases products of genetic modification, where legislatively approved) or of cultures with high enzyme activities.

Of these options, the employment of a higher maturation temperature is the easiest to implement. One of the first trials to accelerate ripening using temperature was in 1946, where mature good quality Cheddar cheese was obtained after 3-4 months of ripening at 16°C (El Soda 1993), rather than the typical temperature of 5-8 °C. Given the significant interest in the topic, numerous studies have investigated accelerated ripening; most show a direct link between accelerated flavour development and exposure to higher ripening temperatures (Aston et al. 1985; Folkertsma et al. 1996).

Despite the apparent benefits, higher temperatures have not been widely implemented due to the fear of growth of spoilage microorganisms and the development of off-flavours, as a result of the acceleration of biochemical changes (Hannon et al. 2005). To reduce the potential for growth of spoilage organisms due to an elevated ripening temperature, it has been suggested that Cheddar cheese should have a salt in moisture (S/M) of 4-5%, pH less than 5.2 and a NSLAB population of less than 1000/g towards the end of the cheese making

process, (Fedrick 1987) although the maximum NSLAB population will often depend on the species of NSLAB present.

Elevated temperature is known to increase the population of bacteria in Cheddar cheese when temperatures higher than 8°C are used, although microbial analysis showed that the effect of temperature on the NSLAB count is not necessarily noticeable at the end of 9 months (Cromie et al. 1987; Folkertsma et al. 1996). Aston et al. (1985) found that higher proteolytic activity could be observed at a ripening temperature of 17.5°C or 20°C and that off flavours, described as unclean or onion-like, developed when the Cheddar cheese was ripened for too long at higher temperatures (20°C, >16 weeks / 17.5°C, >32 weeks). Folkertsma et al. (1996) similarly noticed an increase in the rate of proteolysis and lipolysis in Cheddar cheese ripened at 16°C, leading to a softer cheese after a period of 9 months. A temperature of 12°C was consequently recommended for commercial purposes (Folkertsma et al. 1996). Law (2001) also reported a reduction of 60-75% in the maturation time when the Cheddar cheese was ripened at 12°C. A further effect of elevated ripening temperatures is that cheeses were found to be more brittle and less springy when the temperature was increased to 20°C (Fedrick and Dulley 1984).

Another potential consequence of ripening at elevated temperatures is that changes in the bacterial population and proteolysis may affect the concentration of amino acids and biogenic amines, such as putrescine, cadaverine and tyramine. Both amino acids and biogenic amines were observed to increase significantly with temperature and storage in Dutch type cheeses (Pachlová et al. 2012; Komprda et al. 2007), potentially introducing a health hazard for consumers. The final concentration of amines in Cheddar cheeses ripened at high temperatures at the end of the accelerated ripening period would therefore be valuable data to assess the relative merits of incubation at high temperatures.

Scanning electron microscopy (SEM) or confocal laser scanning microscopy (CLSM) are useful techniques that can be used to observe changes in cheese for which ripening was accelerated, by means such as addition of lipase in Feta (Karami et al. 2009). These techniques have not been widely applied, however, for the study of cheese ripened at elevated temperatures, although one study has used CLSM to illustrate the similar structure within Cheddar cheeses stored at 8°C and 25°C for a limited period of 70 hours (O'Reilly et al. 2003). Cheese structure can yield important information about the arrangement of protein and fat within the cheese and can provide insights into other fundamental parameters such as texture. It is therefore timely that microscopy be used to help evaluate the potential of elevated temperature as a means to accelerate ripening.

The aims of the current study were to investigate the impact of elevated temperature on the microstructure, texture, proteolysis and content of organic acids, amino acids and biogenic amines of ripened Cheddar cheese. Four treatments were applied: ripening at 8°C, 15°C, 20°C, or a two-stage ripening at 15°C for 33 days followed by 8°C for the subsequent ripening period. This study provides the first direct analysis of the effects of elevated maturation temperature on Cheddar cheese structure with the goal of relating temperature-dependent development of cheese functionality to structural considerations..

2. Materials and Methods

2.1 Cheese treatment

Three 20 kg blocks of commercial full fat Cheddar cheese, made with pasteurised milk and a defined-strain *Lactococcus lactis* bulk starter culture, were obtained from the same vat of cheese and the blocks were then cut and packaged into at least 84 ~500 g block samples. The cheese samples were then stored at 8°C (T8), 15°C followed by 8°C (T15-8), 15°C (T15) or 20°C (T20) for the ripening period (330 days, 21 blocks of cheeses for each treatment). For

one treatment (T15-8), the cheese was stored at 15°C for 33 days and then 8°C for the remainder of the trial. At selected time points during ripening, three 500 g block samples of each treatment were analysed using the methods described below, except for the amine analysis which used six 500 g block samples.

2.2 Compositional analysis of cheese

Cheeses were analysed in triplicate for fat, protein (%N x 6.38), salt and moisture using the gravimetric method AS2300.1.3 (Australian Standard 2008), the Kjeldahl method AS2300.1.2.1 (Australian Standard 1991), a potentiometric method (AOAC 2012) and the method described in AS2300.1.1 (Australian Standard 1988) respectively. The pH of the fresh and mature cheese was determined by blending 10 g of cheese with 10 g of water purified to a resistivity of 18.2 mΩ (Millipore MilliQ, Billerica, MA, USA) and the pH of the resultant slurry was then measured using an electrode pH meter (Orion 720A, Orion Pacific, Frankston, Australia).

2.3 Microscopic techniques

Cheese samples were analysed during ripening at days 1, 12, 33, 98, 194 by confocal laser scanning microscopy (CLSM, Leica TCS SP2; Leica Microsystems, Heidelberg, Germany), using a method described previously (Soodam et al. 2015). Stock solutions of Nile Red and Fast Green FCF were prepared at a concentration of 1 mg/mL in dimethyl sulfoxide and water respectively. These solutions were then diluted 10 fold just prior to staining. A total of 6 images and three three-dimensional pictures were collected for each cheese treatment at each time point during ripening. Reconstruction and analysis of the three-dimensional images was carried out using Imaris image processing software (Bitplane, South Windsor, CT, USA) as described previously (Ong et al. 2012) and yielded parameters as described by Soodam et al. (2015).

Cheese samples were also analysed with cryo scanning electron microscopy (Cryo-SEM, Quanta; Fei, Hillsborro, OR, USA) using a method described in a previous study (Ong et al. 2011). A total of 6 images were taken at a magnification of 2000 x, 4000 x or 8000 x from one representative cheese sample for each treatment at each time-point (days 1, 12, 33, 98, 194) during ripening.

2.4 Proteolysis determination

The proteolytic pattern of the cheese during ripening was determined using sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). The cheese stock solution was prepared using the method of Ong et al. (2006), except that 25 mg samples of cheeses were used. The stock solutions produced were then diluted ~4-fold with Trisaminomethane (Tris, 10 mmol/L) - Ethylenediaminetetraacetic acid (EDTA 1 mmol/L) pH 8.0 buffer. Standards of α -casein (CN), β -CN and κ -CN (Sigma-Aldrich, Castle Hill, Australia) were also prepared at a concentration of 0.25 mg/mL. An aliquot of 6.5 μ L was mixed with 2.5 μ L of Bolt LDS Sample Buffer (4x) and 1 μ L Bolt Reducing Agent (10x) (Invitrogen, Mt Waverley, Australia). The solution was heated at 70 °C for 10 min and then loaded into a 12% acrylamide Bis-Tris Plus gel (Invitrogen). A set of molecular standards, SeeBlue Plus2 pre-stained (Invitrogen), was also loaded in each gel. Bolt 2-(*N*-morpholino) ethanesulphonic acid SDS Running Buffer (1x) (Invitrogen) was used as buffer and the system was run for 75 min at 125 V. The protein bands were then stained and destained as described by Ong et al. (2006) except 7.5% v/v acetic acid was used and the de-staining solution did not contain methanol. The gel was finally scanned using a Fujifilm Intelligent Dark Box II with LAS-3000 Lite V2.2 software and LAS-3000 Image Multi Gauge (Fujifilm, Brookvale, Australia) to quantify the intensity of α - and β -CN bands during ripening. The data presented are representative of three gels.

2.5 Determination of amine content

Extraction of samples

Hydrochloric acid (HCl, 1 mL, 0.1 M) was added to 0.1 g of grated cheese. The samples were then cryo-milled (Precellys 24, Bertin Technologies, Île-de-France, France) at -10°C (program 2, 3 min total cycle; per cycle: 30s spinning, 45s waiting), followed by centrifugation for 10 min at 10000 rpm (~9000 g). The fat layer was removed and the supernatant was then collected (10 µL) for derivatization, avoiding the casein fraction at the bottom of the tube.

Derivatization of samples

The derivatization was performed using a method described by Boughton et al. (2011). Borate buffer (200 mM boric acid (Univar, Ajax Finechem, Seven Hills, Australia), 35.7 µM ¹³C, ¹⁵N L-Valine, 10 mM Tris(2-carboxyethyl)-phosphine (Sigma Aldrich), 1 mM ascorbic acid (Sigma Aldrich) was prepared and adjusted to pH 8.8 using 2 M sodium hydroxide. The supernatant (10 µL) was diluted 10 times using 0.1% formic acid and 10 µL of the diluted sample was then mixed with 70 µL borate buffer and 20 µL of 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate (AQC, Sigma-Aldrich, 10mM in 100% acetonitrile) derivatization reagent. The mixture was vortexed for 30 s and mixed at 55°C using a thermomixer (Eppendorf, Macquarie Park, Australia) for 15 min. The samples were then centrifuged and transferred into HPLC vials for LC-MS analysis.

Liquid chromatography – mass spectrometry (LC-MS) analysis

Quantitative analysis was performed using a 1290 LC-system coupled to a 6490 Electrospray Ionisation-Triple Quadrupole MS (LC-QqQ-MS, Agilent, Santa Clara, CA, USA) and a modified version of a method previously described in the literature (Boughton et al, 2011). Samples and amine standards (2 µL) were injected into the system. Ions were monitored in

the positive mode using a dynamic multiple reaction monitoring (DMRM) method optimized for each analyte. The source, collision energies and fragmentor voltages were optimized for each analyte by infusing a derivatized standard with LC eluent. The source conditions used were: sheath gas temperature 315°C, gas flow 10 L.min⁻¹, nebulizer pressure 45 psi (310 kPa) and capillary voltage 3800 V. Profiling experiments were performed by multiple reaction monitoring (MRM) in electrospray ionisation (ESI) positive mode by collecting transition of each selected precursor ion to 171.1⁺ aminoquinoline (Amq) fragment ion using collision energy of 20 V within the mass range of 100-1000 *m/z*.

The mobile phases used were (A) 0.1% formic acid (Sigma Aldrich) in water and (B) 0.1% formic acid in acetonitrile. Liquid chromatography separation was achieved by injecting samples into an Agilent Porshell 120 SB-C18 2.1 x 100 mm, 2.7 µm column with a flow rate of 600 µL/min. The temperature was kept at 65°C, resulting in operating pressures below 1000 bar with a 13 min run time. The gradient profile used in the method is shown in **Online Resource 1**.

The limit of quantification, defined as the concentration at which quantitative results can be reported with reasonable confidence (Armbruster et al. 1994), was 0.001 µM.

Recovery experiments

The extraction was performed six times for the same sample of cheese. Formic acid (30µL) was added to the supernatant (from the cryo-mill) of three of the samples (500 µL), acting as a negative control. The supernatant (500 µL each) of the other three samples was spiked with a standard solution containing a mixture of all the amines monitored during this experiment (30 µL, 150 µM). The estimated recovery for each amine was then calculated as shown below:

Recovery of amine

$$= \frac{\text{Concentration in spiked sample} - \text{Concentration in non-spiked sample}}{\text{concentration of additional amine}} \times 100$$

The recovery was 82-105% of the amines monitored during the experiment.

2.6 Texture and organic acid analysis

A texture analyser TA-XT Plus (Stable Micro Systems, Surrey, England) was used to obtain the texture profile of cheese samples using the method described by Soodam et al. (2015). Texture analysis was performed at least three times for each cheese treatment.

A method described in a previous study (Soodam et al. 2015) was used to quantify organic acids in mature cheese samples. Analysis was carried out using high performance liquid chromatography with a Shimadzu Prominence system equipped with a Bio-Rad Aminex HPX 87H cation exchange column connected to a cation H⁺ guard column (Bio-Rad Laboratories, Hercules, CA, USA); analysis of the acids was performed at 2 wavelengths (220 nm and 285 nm) to allow quantification of acids that co-elute.

2.7 Statistical Analysis

Experimental analysis was performed at days 1, 12, 33, 98 and 194, except for the amine analysis, which was performed at day 330.

For all analyses except for the amine analysis, all cheese treatments were analysed in triplicate (three individual ~500 g cheese blocks) and the results presented are the mean of the three data points (n=3), where all replicate chemical and physical analyses were averaged for each cheese block. For the amine analysis, all cheese treatments were analysed six times (six individual ~500 g cheese blocks) and the data are presented as a mean of the six data points (n=6). Interactions between ripening time and ripening temperature were also analysed using a general analysis of variance model in Genstat (VSN International, Hertfordshire, UK)

with the assumption that the individual ~500 g cheese blocks are considered independent at the start of the study. If the effect of time or temperature was significant, the data were further analysed using Tukey's paired comparison, with a significance level of $\alpha = 0.05$. In some cases, data at individual time points may also be analysed using one way analysis of variance (ANOVA) and Tukey's test, with a significance level of 0.05, with respect to the temperature treatment. Correlations were also analysed using Genstat (VSN International).

3. Results and Discussion

3.1 Compositional analysis of cheese

The mean compositions of the cheeses used for the maturation study, determined one day after pressing, are shown in **Table 1**. The composition of the cheese was within the parameters recommended for Cheddar ripening at elevated temperatures (Folkertsma et al. 1996), which includes a moisture less than 37%, a moisture in fat-free-substance (MFFS) less than 55% and a fat in dry matter (FDM) greater than 50%.

Some changes occurred in the cheese at the later stages of ripening due to the release of a small volume of whey from the T15-8 and T20 cheeses from day 98 onwards, with more whey being released from the T20 cheeses ($4.3 \pm 0.1\%$ w/w for the T20 cheeses compared to $0.3 \pm 0.2\%$ w/w for the T15-8 cheeses at day 194). The release in moisture resulted in an effective increase in the fat concentration at day 194 ($37.5 \pm 0.2\%$ w/w) compared to day 1 ($36.0 \pm 0.4\%$ w/w) ($P < 0.05$) for the T20 cheeses.

3.2 Microstructure of Cheddar cheese throughout ripening

Qualitatively, the temperature treatments resulted in changes to the protein network. These changes are most apparent in the 3D microstructure of the cheese samples at day 33 (**Fig. 1**). The protein strands of the T8 samples appear to be thinner, compared to the T20 cheese. This

observation is also supported by image analysis. At day 33, the number of vertices in the rendered image, which may indicate the branching in the network on a comparative basis, was significantly higher for the T8 samples compared to the T20 samples ($P<0.05$) (**Fig. 2a**). The greater number of vertices is consistent with the observation of more numerous thinner protein strands in samples incubated at a lower temperature. The T15 treatments, however, had a statistically similar number of vertices ($P>0.05$) to both the T8 and T20 cheeses, showing the change is gradual with increasing temperature.

The solubilisation of calcium phosphate has possibly led to these structural changes. Cheese is known to soften within the first month due to such solubilisation and a breakdown of the casein network (Chevanan and Muthukumarappan 2007; Hassan et al. 2004; O'Mahony et al. 2005). This solubilisation can lead to an increase in the hydration and swelling of the *para*-casein matrix of the cheese (O'Mahony et al. 2006).

Later in maturation, at day 194, the 3D images of the protein network appear similar to the images at day 33, suggesting that no further changes occur within the structure during further ripening (**Fig. 1**). However, the number of vertices is observed to decrease significantly during ripening ($P<0.05$), with the overall number being significantly lower at day 194 when compared to day 1, suggesting that the protein was continuing to solubilise during ripening. While the differences between the temperature treatments are still visible qualitatively, differences in the number of vertices are no longer statistically significant at day 194 ($P>0.05$, **Fig. 2b**). This was not expected and may reflect the higher variability of the number of vertices at later time points.

The distribution of the fat within the cheese did not change during ripening, irrespective of the temperature used. The number, sphericity and diameter of the fat globules were not significantly affected by the ripening temperature, ripening time or their interactions ($P>0.05$,

data not shown). This observation is consistent with previous observations that no further clumping of fat globules occurs during the ripening of Cheddar cheese due to the solid nature of the fat at the typical ripening temperature of 7°C (Guinee et al. 2000). At 20°C, however, the proportion of milk fat known to be liquid in milk is more than three times the percentage present at 8°C (Norris et al. 1973). Interestingly, in the context of cheese at 20°C in this study, no further clumping or changes to fat were observed, possibly due to the constraints of the protein network, although whey was released by the T15 and T20 cheeses from day 98 onwards.

Only minor changes could be observed between the different treatments at the different stages of ripening by cryo-SEM (images not shown). Gaps were observed at the protein and fat interface towards the end of ripening (day 194) when compared to images taken at day 1, most likely as an indirect result of proteolysis, which causes a weakening of the protein network, making the network more susceptible to changes during sample preparation. Low magnification cryo SEM images also show that more granules, which may be calcium salts, could be observed at higher temperatures (T15 and T20 treatments, **Fig. 3**). These structures occur despite the increased salt solubility at these temperatures and may occur as a result of whey migration. More free whey is produced when cheeses are stored at a high temperature, which could result in more movement of salts. Johnson (2004) suggested that the movement of free whey within the cheese or to the surface of the cheese helps to bring calcium lactate to nucleation sites, from which granules (usually described as ‘crystals’) can then develop. Higher ripening temperatures may also increase the rate of calcium lactate deposition due to lactate racemisation by cheese microflora (Chou et al. 2003). The presence of free whey and increased calcium lactate deposits may prove a deterrent to the industrial use of these high temperatures for long maturation periods.

3.3 Proteolysis in Cheddar cheese throughout ripening

Proteolysis was measured in the cheese using SDS-PAGE, as shown in **Fig. 4**. Protein bands were identified by comparison to the casein standards, as well as molecular weight standards. The residual percentages of α_{s1} -CN, α_{s2} -CN and β -CN were significantly affected by both ripening time and temperature ($P<0.05$, **Fig. 4b-d**). At day 194, the degradation of α_{s1} -CN, α_{s2} -CN and β -CN was significantly higher for the T20 cheese, compared to the control cheese, T8 ($P<0.05$). This result is in agreement with Folkertsma et al. (1996) who reported increased degradation of α_{s1} -CN and β -CN in 9 month old Cheddar cheese ripened at 16°C compared to at 8°C. While the cheeses were ripened for a shorter interval here (~ 6.4 months), this trend was already apparent for the T20 cheeses and it is expected that an additional ripening period of 3 months will further increase the differences between the other treatments.

The residual percentage of β -CN was also significantly higher than that of α_{s1} -CN or α_{s2} -CN at day 194 ($P<0.05$, **Fig. 4b-d**), when analysed independent of treatment, showing that the rate of hydrolysis of this protein was significantly lower. This was expected, as typically ~ 50% of β -CN is left in mature Cheddar (Banks 2011). While T8 cheeses still had $80 \pm 6\%$ of the β -CN left at day 194, T20 cheeses only had $45 \pm 9\%$ left, again demonstrating the increased proteolysis when a higher ripening temperature is used. Consistent with previous studies, the concentration of proteolytic products of α_s -CN and β -CN such as γ -CN (regions A, B) also increased with increasing ripening temperature (**Fig. 4a** Lanes 2-9).

3.4 Amine concentration in Cheddar cheese after ripening for 330 days

The concentration of the most common amino acids in the four cheese treatments at day 330 is shown in **Fig. 5a, b**. All concentrations shown in this figure are significantly affected by temperature ($P<0.05$) indicating an acceleration of ripening. The T20 cheeses have

significantly higher concentrations of all amino acids compared to the T8 cheeses, with the exception of glutamine, which shows a decline. The cause for the decrease of glutamine as a function of temperature is unknown, although some lactobacilli bacteria are known to deamidate glutamine to raise the pH of their immediate environment, which in turn allows the bacteria to grow in an environment where pH is the limiting factor (Vermeulen et al. 2007). A past study showed that ripening of control Cheddar cheese at 15°C resulted in a higher glutamine concentration than ripening at 6°C (Puchades et al. 1989). Proteolysis, however, is quite complex and the lower glutamine concentration may be due to increased breakdown as a result of secondary biochemical changes.

It is worthwhile to note that the T8 and the T15-8 cheeses have statistically similar concentrations of amino acids at day 330 ($P>0.05$), despite the 33 days spent ripening at an elevated temperature for the T15-8 cheeses. The impact of a higher concentration of such amino acids on the flavour of aged Cheddar cheese is, however, unknown. Higher concentrations of amino acids did not intensify the cheese flavour in a previous study (McGarry et al. 1994).

Biogenic amines in cheese can increase with age and temperature and can become toxic at high concentrations. Typical biogenic amines monitored during cheese ripening include histamine, putrescine, cadaverine, tyramine, tryptamine, phenethylamine, spermine and spermidine (Innocente et al. 2007). Histamine and spermine were not analysed during this experiment but the concentrations of the other biogenic amines are shown in **Fig. 5c**. Only tryptamine and tyramine showed clear trends with respect to ripening temperature, with the T20 cheeses having significantly higher concentrations than the T8 cheeses. The concentrations of the amines obtained here are also lower than the concentrations obtained from literature (Laleye et al. 1987). The biogenic amines in this study all have a concentration less than 25 mg/kg cheese while, in Cheddar cheese aged more than 12 months,

concentrations of 140 – 580 mg/kg, 140 – 330 mg/kg and 140 – 250 mg/kg of cheese were found for tyramine, putrescine and cadaverine respectively (Laleye et al. 1987). The past literature has stated, however, that aged cheese made in good hygienic conditions would usually have small concentrations of biogenic amines, with spermidine as the amine present at the highest concentration (Velišek 2014). This study confirms that biogenic amines are not a concern in the industrial samples examined even when matured at higher temperatures, although the situation could vary depending on the microbial composition of the starter culture used and of the non-starter cheese flora.

3.5 Texture profiles of Cheddar cheese throughout ripening

The textural properties of the cheeses including hardness, cohesiveness and chewiness were significantly affected by the ripening temperature, ripening time and the interaction between these two variables ($P<0.05$). This is consistent with the changes detected in the microstructure of the cheeses at the early stages of ripening.

The cheeses softened significantly from day 1 to day 98 (**Fig. 6a**), when analysed independently of temperature treatment. This effect is likely due to calcium phosphate solubilisation and proteolysis, as cleavage of the caseins reduces the ability of the caseins to contribute to the protein network (Lawrence et al. 1987; O'Mahony et al. 2005). From day 98 to day 194, however, the hardness of the cheeses increased significantly ($P<0.05$). This has also been reported in the literature (Chevanan and Muthukumarappan 2007). One possibility is that as the cheese ages, further proteolysis into water soluble peptides results in the exposure of new ionic groups which then compete for the available free water in the cheese. The water previously available for partial solubilisation of the protein network becomes associated with the new ionic groups, making the cheeses harder (Lawrence et al. 1987). By

the end of ripening at day 194, the T20 cheeses were significantly softer than the T8 and T15-8 cheeses ($P<0.05$), probably due to the acceleration of proteolysis in T20.

The chewiness and cohesiveness of the cheeses decreased significantly with time ($P<0.05$, **Fig. 6b-c**). The interaction between ripening time and ripening temperature was also significant ($P<0.05$ for both cohesiveness and chewiness). From **Fig. 6b-c**, the difference in the rates of change for both textural parameters is particularly noticeable during the first 33 days, with T8 cheeses appearing to have a slower decline in these two textural properties. This observation is again consistent with the microstructure observations. These two textural changes may also have an impact on consumer preference, as at day 194 the cohesiveness was significantly lower ($P<0.05$) for the T20 and T15 cheeses when compared to the T8 and T15-8 cheeses.

The textural results showed that the T15-8 cheese would provide the desired acceleration in ripening, without drastically affecting the quality of the cheeses, as shown by the statistically similar chewiness and cohesiveness between the T8 and T15-8 cheeses after ripening for 194 days.

3.6 Organic acid content of Cheddar cheese throughout ripening

Ten organic acids were monitored during ripening as the changes in proteolysis and microbial growth with elevated temperatures may also alter the metabolic pathways and the metabolic end products of bacteria within the cheese. Hippuric acid was not detected in any sample from any treatment at any time of ripening, in agreement with Lues et al. (1998) who could not detect any hippuric acid in 10-month old Cheddar cheese. All other acids were detected and most were significantly affected by ripening time, temperature and the interaction between these two variables ($P<0.05$), with the exception of pyruvic and uric acids, which were only significantly affected by the ripening time ($P<0.05$) and the interaction between

time and temperature ($P < 0.05$). It should be noted, however, that the apparent rise in citric and propionic acids observed with ripening time may potentially be affected by the co-elution of these acids with other compounds. From past literature, acetoin and propionic acid are known to co-elute at 210 nm (Kocaoglu-Vurma et al. 2008) and co-elution may also be occurring at the wavelength used in this study (220 nm).

Generally, it can be observed that the T15-8 treatment would be optimal as the organic acid content is generally quite similar to that of T8 cheese after ripening for 194 days (**Online Resource 2**), thereby reducing the risks of off-flavour development whilst still providing the accelerated changes to proteolysis and texture observed in the previous sections.

4. Conclusion

In this study, the maturation of cheeses subjected to different ripening temperatures was assessed. This is the first paper that monitors the structure of Cheddar cheese subjected to extended ripening at elevated temperatures. Ripening temperature was found to affect the rate of change of most parameters measured during the ripening period. Increasing the temperature of storage has the benefit of accelerating ripening. The 3D CLSM images show that short periods of storage at an elevated temperature of 15°C or 20°C affect the protein network significantly. Cheeses stored at 8°C appear to have thinner and more numerous protein strands than cheeses stored at 15°C and 20°C, results which were supported by number of vertices in the rendered protein surface obtained through image analysis. Longer periods of storage of 194 days or more at high temperatures of 15°C or 20°C may have an adverse effect on the texture and free whey release. Cheese that is ripened at an elevated temperature of 15°C for 33 days followed by longer ripening of up to 194 days at 8°C may have the benefit of accelerating the ripening without too many adverse effects. This study may prove useful to dairy manufacturers, as it shows that ripening treatments can be used to

minimise ripening time, whilst still providing cheese with the desired texture. It also illustrates that microscopy is a useful tool that complements existing techniques for comparing cheeses during the development of modified-temperature maturation, protocols or during routine cheese maturation where fine control of cheese development is desired.

Acknowledgements

The authors acknowledge the Australian Government for providing the Australian Postgraduate Award (APA) scholarship and Dairy Innovation Australia and its member companies for financial and practical support for this project (project grant 08209C). We also thank the Particulate Fluids Processing Centre (PFPC), the Bio21 Institute for access to equipment and more specifically, the Electron Microscopy Unit and the Biological Optical Microscopy Platform and Mr Roger Curtain for his help in operating the scanning electron microscope in cryo mode. We also thank Dr Thusitha Rupasinghe, a research and development scientist of Metabolomics Australia, School of BioSciences, the University of Melbourne (Australia) who provided assistance in the design and interpretation of the amines data and facilitated access to the equipment We acknowledge Ms Rachel Sore, for her advice concerning the statistical analysis. Sally Gras and Lydia Ong are supported by The ARC Dairy Innovation Hub (IH20100005).

Conflict of interest

The authors declare no conflict of interest

References

- AOAC (2012). Official methods of analysis of AOAC International, Method 971.27. Gaithersburg, USA: AOAC International.
- Armbruster, D. A., Tillman, M. D., & Hubbs, L. M. (1994). Limit of detection (LQD)/limit of quantitation (LOQ): comparison of the empirical and the statistical methods

exemplified with GC-MS assays of abused drugs. *Clinical Chemistry*, 40(7), 1233-1238.

Aston, J. W., Giles, J. E., Durward, I. G., & Dulley, J. R. (1985). Effect of elevated ripening temperatures on proteolysis and flavour development in Cheddar cheese. *Journal of Dairy Research*, 52(04), 565-572.

Australian Standard (1988). Methods of chemical and physical testing for the dairying industry - General methods and principles - Determination of total solids and moisture. *Australian Standard AS 2300.1.1*.

Australian Standard (1991). Methods of chemical and physical testing for the dairying industry - General methods and principles - Determination of nitrogen - Reference Kjeldahl method *Australian Standard AS 2300.1.2.1*.

Australian Standard (2008). Methods of chemical and physical testing for the dairying industry - General methods and principles - Determination of fat - Gravimetric method. *Australian Standard AS 2300.1.3*.

Banks, J. M. (2011). Cheese | Cheddar-type cheeses. In J. W. Fuquay, P. F. Fox, & P. L. H. McSweeney (Eds.), *Encyclopedia of Dairy Sciences (Second Edition)* (pp. 706-711). San Diego: Academic Press.

Boughton, B. A., Callahan, D. L., Silva, C., Bowne, J., Nahid, A., Rupasinghe, T., et al. (2011). Comprehensive profiling and quantitation of amine group containing metabolites. *Analytical Chemistry*, 83(19), 7523-7530.

Chevanan, N., & Muthukumarappan, K. (2007). Effect of calcium and phosphorus, residual lactose, and salt-to-moisture ratio on the melting characteristics and hardness of Cheddar cheese during ripening. *Journal of Food Science*, 72(4), E168-E176.

- Chou, Y. E., Edwards, C. G., Luedecke, L. O., Bates, M. P., & Clark, S. (2003). Nonstarter lactic acid bacteria and aging temperature affect calcium lactate crystallization in Cheddar cheese. *Journal of Dairy Science*, 86(8), 2516-2524.
- Cromie, S. J., Giles, J. E., & Dulley, J. R. (1987). Effect of elevated ripening temperatures on the microflora of Cheddar cheese. *Journal of Dairy Research*, 54(1), 69-76.
- El Soda, M. (1993). Accelerated maturation of cheese. *International Dairy Journal*, 3(4-6), 531-544.
- Fedrick, I. (1987). Technology and economics of the accelerated ripening of Cheddar cheese. *Australian Journal of Dairy Technology*, 42(1-2), 33-36.
- Fedrick, I. A., & Dulley, J. R. (1984). The effect of elevated storage temperatures on the rheology of Cheddar cheese. *New Zealand Journal of Dairy Science and Technology*, 19(2), 141-150.
- Folkertsma, B., Fox, P. F., & McSweeney, P. L. H. (1996). Accelerated ripening of Cheddar cheese at elevated temperatures. *International Dairy Journal*, 6(11-12), 1117-1134.
- Fox, P. F. (2002). Cheese | Overview. In H. Roginski, J. W. Fuquay, & P. F. Fox (Eds.), *Encyclopedia of Dairy Sciences* (pp. 252-261). Amsterdam: Academic Press.
- Guinee, T. P., Auty, M. A. E., & Fenelon, M. A. (2000). The effect of fat content on the rheology, microstructure and heat-induced functional characteristics of Cheddar cheese. *International Dairy Journal*, 10(4), 277-288.
- Hannon, J. A., Wilkinson, M. G., Delahunty, C. M., Wallace, J. M., Morrissey, P. A., & Beresford, T. P. (2005). Application of descriptive sensory analysis and key chemical indices to assess the impact of elevated ripening temperatures on the acceleration of Cheddar cheese ripening. *International Dairy Journal*, 15(3), 263-273.

- Hassan, A., Johnson, M. E., & Lucey, J. A. (2004). Changes in the proportions of soluble and insoluble calcium during the ripening of Cheddar cheese. *Journal of Dairy Science*, 87(4), 854-862.
- Innocente, N., Biasutti, M., Padovese, M., & Moret, S. (2007). Determination of biogenic amines in cheese using HPLC technique and direct derivatization of acid extract. *Food Chemistry*, 101(3), 1285-1289.
- Johnson, M. (2004). Revisiting calcium lactate crystals in cheese. *Dairy Pipeline*, 16(1), 1-11.
- Karami, M., Ehsani, M. R., Mousavi, S. M., Rezaei, K., & Safari, M. (2009). Microstructural properties of fat during the accelerated ripening of ultrafiltered-Feta cheese. *Food Chemistry*, 113(2), 424-434.
- Kocaoglu-Vurma, N. A., Harper, W. J., Drake, M. A., & Courtney, P. D. (2008). Microbiological, chemical, and sensory characteristics of Swiss cheese manufactured with adjunct *Lactobacillus* strains using a low cooking temperature. *Journal of Dairy Science*, 91(8), 2947-2959.
- Komprda, T., Smělá, D., Novická, K., Kalhotka, L., Šustová, K., & Pechová, P. (2007). Content and distribution of biogenic amines in Dutch-type hard cheese. *Food Chemistry*, 102(1), 129-137.
- Laleye, L. C., Simard, R. E., Gosselin, C., Lee, B. H., & Giroux, R. N. (1987). Assessment of Cheddar cheese quality by chromatographic analysis of free amino acids and biogenic amines. *Journal of Food Science*, 52(2), 303-307.
- Law, B. A. (2001). Controlled and accelerated cheese ripening: the research base for new technologies. *International Dairy Journal*, 11(4-7), 383-398.
- Lawrence, R. C., Creamer, L. K., & Gilles, J. (1987). Texture development during cheese ripening. *Journal of Dairy Science*, 70(8), 1748-1760.

- 524 Lues, J. F. R., Botha, W. C., & Smit, E. J. (1998). Ion-exchange HPLC analysis of a broad
525 spectrum of organic acids from matured Cheddar cheese and assessment of extraction
526 methods. *Food Research International*, 31(6–7), 441-447.
- 527 McGarry, A., Law, J., Coffey, A., Daly, C., Fox, P. F., & Fitzgerald, G. F. (1994). Effect of
528 genetically modifying the lactococcal proteolytic system on ripening and flavor
529 development in Cheddar cheese. *Applied and Environmental Microbiology*, 60(12),
530 4226-4233.
- 531 Norris, G. E., Gray, I. K., & Dolby, R. M. (1973). Seasonal variations in the composition and
532 thermal properties of New Zealand milk fat: II. Thermal properties of milk fat and
533 their relation to composition. *Journal of Dairy Research*, 40(03), 311-321.
- 534 O'Mahony, J. A., Lucey, J. A., & McSweeney, P. L. H. (2005). Chymosin-mediated
535 proteolysis, calcium solubilization, and texture development during the ripening of
536 Cheddar cheese. *Journal of Dairy Science*, 88(9), 3101-3114.
- 537 O'Mahony, J. A., McSweeney, P. L. H., & Lucey, J. A. (2006). A model system for studying
538 the effects of colloidal calcium phosphate concentration on the rheological properties
539 of Cheddar cheese. *Journal of Dairy Science*, 89(3), 892-904.
- 540 O'Reilly, C. E., Kelly, A. L., Oliveira, J. C., Murphy, P. M., Auty, M. A. E., & Beresford, T.
541 P. (2003). Effect of varying high-pressure treatment conditions on acceleration of
542 ripening of cheddar cheese. *Innovative Food Science & Emerging Technologies*, 4(3),
543 277-284.
- 544 Ong, L., Dagastine, R. R., Kentish, S. E., & Gras, S. L. (2011). Microstructure of milk gel
545 and cheese curd observed using cryo scanning electron microscopy and confocal
546 microscopy. *LWT - Food Science and Technology*, 44(5), 1291-1302.

- Ong, L., Dagastine, R. R., Kentish, S. E., & Gras, S. L. (2012). The effect of pH at renneting on the microstructure, composition and texture of Cheddar cheese. *Food Research International*, 48(1), 119-130.
- Ong, L., Henriksson, A., & Shah, N. P. (2006). Development of probiotic Cheddar cheese containing *Lactobacillus acidophilus*, *Lb. casei*, *Lb. paracasei* and *Bifidobacterium* spp. and the influence of these bacteria on proteolytic patterns and production of organic acid. *International Dairy Journal*, 16(5), 446-456.
- Pachlová, V., Buňka, F., Flasarová, R., Válková, P., & Buňková, L. (2012). The effect of elevated temperature on ripening of Dutch type cheese. *Food Chemistry*, 132(4), 1846-1854.
- Puchades, R., Lemieux, L., & Simard, R. E. (1989). Evolution of free amino acids during the ripening of Cheddar cheese containing added lactobacilli strains. *Journal of Food Science*, 54(4), 885-888.
- Soodam, K., Ong, L., Powell, I. B., Kentish, S. E., & Gras, S. L. (2015). Effect of calcium chloride addition and draining pH on the microstructure and texture of full fat Cheddar cheese during ripening. *Food Chemistry*, 181(0), 111-118.
- Velíšek, J. (2014). Antinutritional, toxic and other bioactive compounds. In *The Chemistry of Food* (pp. 741-846). Chichester, UK: John Wiley & Sons.
- Vermeulen, N., Gänzle, M. G., & Vogel, R. F. (2007). Glutamine deamidation by cereal-associated lactic acid bacteria. *Journal of Applied Microbiology*, 103(4), 1197-1205.

Tables and Figures

Fig. 1 Three dimensional (3D) confocal microscopy images of the microstructure of the protein network of the ripened cheese at day 33 and day 194. Protein is stained green (FCF). The images are 3D reconstructions built from 40 layers, with each layer separated from the next by ~0.25 µm and a total depth of 9.8 µm. The scale bars are 10 µm in length. Images in

colour can be viewed in the electronic version of this manuscript. In the black and white version of this manuscript, protein is lighter coloured (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article)

Fig. 2 Number of vertices in the rendered protein network determined from 3D confocal microscopy images of the ripened cheese at a) day 33 and b) day 194 for cheeses T8, T15-8, T15 and T20. The results are expressed as the mean $\pm$ the standard deviation of the mean (n=3). The number of vertices given is for a cheese volume of 140,000 μm^3 with 40 layers, where each layer is separated by $\sim 0.25 \mu\text{m}$, giving a total depth of $9.8 \mu\text{m}$

Fig. 3 Cryo-SEM microstructure of the ripened cheeses T8 (a), T15-8 (b), T15 (c) and T20 (d) at day 194. Images were captured using a solid state detector at a magnification of 2000 x. The scale bars are $50 \mu\text{m}$ in length. Arrows in white indicate possible crystalline salts

Fig. 4 a) Image of a representative SDS-PAGE gel (4 - 12 %) of day 1 cheese (Lane 1) and ripened cheeses T8 (Lanes 2, 6), T15-8 (Lanes 3, 7), T15 (Lanes 4, 8) and T20 (Lanes 5, 9) at day 33 (Lanes 2-5) and day 194 (Lanes 6-9), and changes in the proportion of b) α_{s1} -CN, c) α_{s2} -CN and d) β -CN, expressed as a percentage of the concentration present at day 1 in T8 (■), T15-8 (●), T15 (▲) and T20 (◆) cheeses at days 1, 33 and 194. Lane 1 was used to indicate the cheeses at day 1, the starting point for all treatments. STD is the standard molecular weight marker and α -CN, β -CN and κ -CN are pure standards of these compounds. The results are expressed as the mean $\pm$ the standard deviation of the mean (n=3)

Fig. 5 Concentration of a, b) amino acids and c) biogenic amines, expressed as mg/kg of cheese, in T8 (□), T15-8 (□), T15 (▣) and T20 (▣) cheeses at day 330. The results are expressed as the mean $\pm$ the standard deviation of the mean (n=6, except for T15-8: n=5)

Fig. 6 Changes in the a) hardness, b) chewiness and c) cohesiveness of T8 (■), T15-8 (●), T15 (▲) and T20 (◆) cheeses at days 1, 12, 33, 98 and 194. The results are expressed as the mean $\pm$ the standard deviation of the mean (n=3)

Table 1 Composition of fresh cheese used in this study. The results are expressed as the mean $\pm$ the standard deviation of the mean (n=3)

Online Resource 1 Gradient used during LC-MS run

Online Resource 2 Changes in the concentration of a) lactic acid, b) acetic acid, c) citric acid, d) formic acid, e) propionic acid, f) butyric acid, g) pyruvic acid, h) orotic acid and i) uric acid in T8 (■), T15-8 (●), T15 (▲) and T20 (◆) cheeses measured at days 1, 12, 33, 98 and 194. The results are expressed as the mean $\pm$ the standard deviation of the mean (n=3)

Fig. 1 (coloured for web)

T8

T20

Day 33

T8

T20

Day 194

Fig. 1 (grey for printing)

Fig. 2

Day 194

Day 194

Fig. 3

a

MW

b

c

d

Fig.4

a

b

c

Fig. 5

Fig. 6

Food and Bioprocess Technology

Effect of elevated temperature on the ripening of full fat Cheddar cheese

Kevany Soodam^{1, 2, 5}, Lydia Ong^{1, 2, 4}, Ian B. Powell^{3, 4}, Sandra E. Kentish^{1, 4} and Sally L. Gras^{1, 2, 4*}

1. Department of Chemical and Biomolecular Engineering, The University of Melbourne, Parkville, Vic 3010, Australia.
2. The Bio21 Molecular Science and Biotechnology Institute, The University of Melbourne, Parkville, Vic 3010, Australia.
3. Dairy Innovation Australia Limited, 180 Princes Highway, Werribee, Victoria 3030, Australia.
4. ARC Dairy Innovation Hub, Department of Chemical and Biomolecular Engineering, The University of Melbourne, Parkville, Vic 3010, Australia.
5. Moorepark Food Research Centre, Teagasc Moorepark, Fermoy, Co. Cork, Ireland (current address).

*Corresponding author: Sally L. Gras. Address: The Bio21 Molecular Science and Biotechnology Institute, The University of Melbourne, Parkville, Victoria 3010, Australia. Tel.: +61 38344 6281; fax: +61 3 8344 4153. E-mail address: sgras@unimelb.edu.au

Table 1: Composition of fresh cheese used in this study. The results are expressed as the mean $\pm$ the standard deviation of the mean (n=3).

Component/Measure	% w/w
Fat	36.0 $\pm$ 0.4
Protein	24.8 $\pm$ 0.2
Salt	1.7 $\pm$ 0.1
Moisture	34.7 $\pm$ 0.4
Moisture in fat-free-substance (MFFS)	54.2 $\pm$ 0.3
Fat in dry matter (FDM)	55.1 $\pm$ 0.3

Food and Bioprocess Technology

Effect of elevated temperature on the ripening of full fat Cheddar cheese

Kevany Soodam^{1,2,5}, Lydia Ong^{1, 2,4}, Ian B. Powell^{3, 4}, Sandra E. Kentish^{1,4} and Sally L. Gras^{1, 2, 4*}

1. Department of Chemical and Biomolecular Engineering, The University of Melbourne, Parkville, Vic 3010, Australia.
2. The Bio21 Molecular Science and Biotechnology Institute, The University of Melbourne, Parkville, Vic 3010, Australia.
3. Dairy Innovation Australia Limited, 180 Princes Highway, Werribee, Victoria 3030, Australia.
4. ARC Dairy Innovation Hub, Department of Chemical and Biomolecular Engineering, The University of Melbourne, Parkville, Vic 3010, Australia.
5. Moorepark Food Research Centre, Teagasc Moorepark, Fermoy, Co. Cork, Ireland (current address).

*Corresponding author: Sally L. Gras. Address: The Bio21 Molecular Science and Biotechnology Institute, The University of Melbourne, Parkville, Victoria 3010, Australia. Tel.: +61 38344 6281; fax: +61 3 8344 4153. E-mail address: sgras@unimelb.edu.au

Online Resource 1: Gradient used during LC-MS run

Time	Solvent A (%)
0	99
2	99
6	90
8.2	80
9.2	60
10	0
11.8	0
13	99