Title: A psychological intervention (ConquerFear) for treating fear of cancer recurrence: Views of study therapists regarding sustainability

Authors: Phyllis Butow^{1,2,3}, Douglas Williams², Belinda Thewes², Stephanie Tesson^{1,3}, Louise Sharpe³, Allan ‘Ben’ Smith^{1,4}, Joanna E Fardell^{1, 5}, Jane Turner⁶, Jemma Gilchrist⁷, Afaf Girgis⁴, Jane Beith⁸ and the Conquer-Fear Authorship Group.

Affiliations:

1. Psycho-Oncology Co-operative Research Group (PoCoG), University of Sydney
2. Centre for Medical Psychology and Evidence-based Decision-making (CeMPED), University of Sydney
3. School of Psychology, University of Sydney
4. Centre for Oncology Education and Research Translation (CONCERT), Ingham Institute for Applied Medical Research, South Western Sydney Clinical School, University of NSW
5. Kids Cancer Centre, Sydney Children’s Hospital and School of Women’s and Children’s Health, Faculty of Medicine, University of NSW.
6. Mental Health Centre, Faculty of Medicine, University of Queensland
7. Crown Princess Mary Cancer Centre, Breast Cancer Institute, Westmead Hospital, Sydney
8. Dept. of Medical Oncology, Lifehouse, Missendon Rd Camperdown, NSW.

Corresponding author:

Phyllis Butow, Professor, School of Psychology,

Lifehouse Level 6 - North (C39Z), The University of Sydney

Email: phyllis.butow@sydney.edu.au

Phone: +61 2 93512859; Fax: 61 2 90365292

ABSTRACT

Purpose: Many evidence-based interventions are not implemented into routine care, thus failing to impact patient outcomes. A critical area of research is to investigate factors influencing intervention sustainability. ConquerFear is an effective intervention for fear of cancer recurrence (FCR) which resulted in greater reductions in FCR immediately and 6 months post treatment than a relaxation control intervention. This qualitative study aimed to determine the barriers and facilitators to sustainability of ConquerFear from the perspectives of study therapists.

Methods: Twenty-six therapists who had delivered ConquerFear in the randomised controlled trial were invited to participate in a semi-structured phone interview. Theoretical saturation was reached after 14 interviews. Thematic analysis was used to identify themes which were then mapped to the PAHRiS implementation framework.

Results: Participants comprised 13 males and 1 female, with on average 14 years of experience working in psycho-oncology. Nine over-arching themes falling into three domains were identified: Evidence (intervention credibility, experienced efficacy, perceived need for intervention); Context (attitude to and capacity for survivorship/ FCR care, therapist orientation and flexibility, limited referral) and Facilitation (intervention/service fit, intervention/patient fit, and training, support and provided resources).

Conclusions: Results suggest that ConquerFear is a sustainable intervention in routine clinical practice. Facilitators included a sound evidence-base, a context where survivorship care is incorporated into service goals, and a good fit between the intervention and both therapist orientation and patient need. Flexibility of delivery and ability to adapt to patient need would assist sustainability.

Implications for cancer survivors: ConquerFear has the potential to sustainably reduce the impact of fear of cancer recurrence. It is important to establish the importance of survivorship care within cancer systems. While this study focused on ConquerFear, we believe these results have implications for other psycho-oncology intervention implementation efforts.

Keywords: Fear of cancer recurrence; intervention; sustainability

INTRODUCTION

Fear of cancer recurrence (FCR) has been defined as the “fear, worry or concern relating to the possibility that cancer will come back or progress.” [1 p3265]. FCR is highly prevalent; up to 70% of cancer survivors report moderate to high FCR [2-4]. FCR is within the top three unmet needs reported by cancer survivors FCR [2]. Consequently, it has been a research priority to develop and evaluate effective interventions to support survivors to better manage FCR.

Our group recently evaluated a novel, theoretically grounded intervention to reduce FCR, ConquerFear [5]. Based on a framework guided by a review of relevant theoretical literature [6], ConquerFear took a primarily meta-cognitive approach [7], based on the Self-Regulatory Executive Function (S-REF) model of anxiety [8]. It did not challenge the content of cognitions (perceived risk of recurrence) but rather survivors’ beliefs about worry (over-valuing or fearing worry). Strategies to increase attentional control and enable disengagement from worrying thoughts were also taught. In addition, based on self-regulation theory [9], ConquerFear provided information and established behavioural contracts regarding surveillance where necessary; and based on relational frame theory [10], included values clarification and committed action.

In a randomised controlled trial (RCT) of ConquerFear [5], participants (n=222) were randomly assigned to five face-to-face sessions of either ConquerFear or an attention control (Taking-it-Easy relaxation therapy). ConquerFear participants had clinically and statistically greater improvements than control participants from baseline to immediately post-intervention on Fear of Cancer Inventory (FCRI) scores ($P < .001$) and most secondary outcomes; differences on FCRI scores were maintained at 3- and 6- months follow-up.

Despite clear evidence of efficacy however, interventions are often not implemented into routine care, are not sustainable [11] and thus do not lead to the real-world change they are

intended to create [12]. A number of frameworks have been developed to guide the design of strategies to optimise sustainability [13]. One such framework is the Promoting Action Research in Health Services determinant framework (PARiHS) [14], which posits that successful implementation occurs when the evidence is robust, the context supportive, and the intervention appropriately facilitated. Frameworks such as PARiHS emphasise the importance of early assessment of contextual factors likely to influence sustainability, through real world testing [14].

The ConquerFear RCT was specifically designed to optimise sustainability and enable a smooth transition to routine care after the trial. Twenty-six psychologists and psychiatrists working in 17 oncology settings around Australia with at least 5 years of experience in clinical practice, at least 2 in Oncology, and training in cognitive-behavioural therapy, were purposively recruited to deliver the intervention. After intensive training, with ongoing supervision, trial therapists treated on average 15 survivors (1-25) each within their normal work environment. Thus, at the end of the trial, we hoped they would have the skills, confidence and opportunity to continue using ConquerFear in routine practice.

An important real-world test of ConquerFear would be the extent to which therapists did incorporate it into routine care. Thus, the aim of the current study was to establish therapists' ongoing use of ConquerFear, as well as to elicit their views regarding the barriers and facilitators to making ConquerFear sustainable in routine clinical practice.

METHODS

Participants

Participants were 14 of the 26 ConquerFear therapists, who were available for interview within the study timeframe. There were no differences between participants and non-participants on age, gender and years of experience, and saturation was achieved within the included interviews.

Design and Procedure

This qualitative study comprised audiotaped and transcribed semi-structured phone interviews, conducted by an experienced qualitative researcher. Interview questions (see Appendix A) were guided by the PARiHS framework, and elicited aspects of evidence, context and facilitation that influenced likelihood of sustainability, attitudes to specific components of ConquerFear, and to different potential formats for ConquerFear delivery (e.g. group, online). The study was approved by the University of Sydney Human Research Ethics Committee.

Analyses:

The interviews were subjected to framework analysis. Key themes and relationships were identified inductively from the data [15]. DW and PB read the first four transcripts and collaboratively named and grouped emergent codes into themes and subthemes to form the preliminary analytical coding framework. DW applied the preliminary framework to all transcripts, modifying it where needed following discussion with PB. The final themes alongside exemplar quotations were discussed with the research team, including two ConquerFear therapists, as a form of member checking, and entered into an Excel spreadsheet to create a framework matrix, to enable understanding of patterns in the data. Finally, all themes were grouped in relation to the PARiHS framework.

Results

Participant Characteristics: Participants comprised 13 psychologists and 1 psychiatrist working in public hospitals. One was male and 13 female, with on average 14 years of experience in Psycho-Oncology.

All therapists intended to continue using ConquerFear in routine practice, some according to the manual (n=4) but most adapting it by either omitting components they found less useful, or more flexibly delivering components depending on each patient's needs (n=10). We identified nine key themes that impacted sustainability: intervention credibility (evidence/theory/face validity); experienced efficacy; perceived need for intervention; attitude to and capacity for survivorship/FCR care; therapist orientation and flexibility; expert referral; intervention/service fit; intervention/patient fit; and training and support and provided resources. These are presented below according to each domain of the PARiHS framework.

Table 1 about here

1. Evidence

The evidence and theory-base behind ConquerFear was well-recognized and accepted by almost all therapists, and this acted as a motivator and facilitator for sustainability. Acceptance of evidence of *need* for ConquerFear in their particular service (e.g. clinical FCR a common patient presentation; FCR requiring a unique approach) was however, less consistent.

1.1 Intervention credibility

Most therapists believed that ConquerFear came from well respected, rigorous researchers whose credentials could be depended upon. Most therapists discussed the perceived strength of both the evidence-base (provided by a well conducted RCT) and the theoretical foundation (provided by the clear theoretical framework underpinning ConquerFear). They felt the evidence-base both increased their own confidence and made it easy to “sell” the intervention to their oncologist and nurse colleagues.

“To be able to say that this is something that’s been robustly, um, designed and studied and we’ve got good results. That, you know, that stands you in good stead.”

“So I really love the whole model of having really expert...researchers who can say, you know, this is how we want to do things.”

1.2 Experienced efficacy

Most therapists also cited their own experience of delivering ConquerFear which had provided personal evidence of its efficacy as a strong motivator to continue using ConquerFear in routine practice. Therapists cited the powerful impact of seeing their patients benefit from the intervention.

“If I was seeing cancer, I would a hundred per cent be using it, because it really works, and I don’t need it to work on a statistical level, I saw in person that it worked.”

“I’m a huge advocate of that program. I, I saw it with my own eyes how it changed people in five sessions, it was just unbelievable.”

1.3 Perceived need for intervention

Most therapists felt there was a strong need for ConquerFear because of the prevalence and impact of FCR in their patients.

“Those sort of concepts and ideas would probably come up in possibly a quarter of the people that I see as outpatients.”

However, some therapists reported a low level of clinical FCR in their patients, not requiring an intensive intervention. One therapist was unconvinced that FCR is a distinct clinical diagnosis that needs a unique approach, rather seeing it as a manifestation of anxiety for which traditional therapeutic approaches would be appropriate.

“And there aren’t many people that come in with clinical levels of fear of recurrence.”

“I wondered sometimes whether it was ... a manifestation of a sort of anxiety anyway... obviously some people are more anxious or have a higher level of arousal in any situation, so whether you can break it down and say a fear of this, this, this and this, but, yeah, I’m a bit ambivalent about it really, I think.”

2. Context

2.1 Attitude to and capacity for working with survivorship/FCR issues

Many therapists noted that survivorship (and FCR, as a component of survivorship) was not a priority in their services, making it more difficult to justify spending time on treating FCR. Lack of resources and time contributed to the pressure to see patients perceived to have greater needs whilst receiving active treatment.

“... the focus is likely on new patients coming ... and getting them to treatment rather than the follow up patients who are maybe perceived to be ... not in so much need of urgent help.”

“... if I had to do it again I would have had to work a lot harder to justify why I was working with people who are basically cured medically.”

One therapist noted that fear of cancer recurrence was not a recognised diagnosis fitting neatly into currently health service funding models, making it more difficult to train others, such as private psychologists, to share the increasing patient load.

“But the private psychologists, it doesn’t necessarily meet the criteria for a mental health plan [Australian funding model].”

However, some therapists noted that their service recognises the need for a greater focus on survivorship, and perceived ConquerFear as an opportunity to meet that expectation.

2.2 Therapist orientation and flexibility

The match between therapists’ therapeutic orientation and that of ConquerFear, their familiarity with components of ConquerFear before participating in the trial, and their attitude to following a manualized therapy, all heavily influenced therapists’ intention to continue use of ConquerFear in full versus in part. Some therapists felt they lost creativity following a manual, which they felt reduced the nuance and personalisation of therapy.

“The model was very consistent with my own understanding of cognitive behavioural therapy models....”

“I felt like I lost the flexibility to be creative in therapy in response to how the patient was presenting or their nuances...”

Many therapists noted they would use aspects of ConquerFear with which they were familiar and comfortable, and which they believed would be most efficacious for their clients but would not use other aspects.

“I use parts of it. Probably the conceptualisation, so I present that. I mean I use the values cards but I used that before anyway.”

As patients commonly brought multiple issues, therapists prioritised different issues and some found it difficult to confine therapy to five sessions even when previously negotiated.

“I probably more often use it in a little bit more of a piecemeal approach, which is probably not ideal, but it’s the reality of clinical practice. ... the only moment we’ve got guaranteed with the patient is the moment we’ve got them in front of us, um, and you don’t want to rush it but you also want to work with what’s current in that particular session.”

Other therapists noted the importance of strictly following a proven manualised therapy for best, evidence-based results.

“For the next five sessions I literally have the manual on my lap.”

2.3 Limited referral

Some therapists noted they were reliant on referral from oncology colleagues to identify patients with high FCR, and that this was not always forthcoming. Some colleagues did not easily elicit or recognise FCR, and if screening was not in place to formally assess psychological wellbeing,

it was sometimes missed. If few patients were seen, it made it harder for therapists to maintain their skills in ConquerFear, making an additional barrier to continued use.

“We rely very heavily on the nurses and the doctors and other staff in the hospital to pick up on patient’s concerns, and so it’s really dependent on their – their own insight and their capacity to make a referral to psychology.”

3. Facilitation

3.1 Intervention/service fit

Therapists noted that implementing ConquerFear in their service was easier if it fitted with the usual patient flow, matched the number and length of therapy sessions usually conducted, and if their service had a sufficient number of psycho-oncologists with appropriate training. For example, if patients primarily came from rural/regional regions to which they returned after active treatment, a 5-session face-to-face survivorship therapy was often not feasible.

“One site I work in we have an hour assessment and half an hour reviews of patients. So it’s not a very long time to get through strategies, um, so we would really have to work out, um, ways of, um, breaking down the therapy.”

“We don’t have a lot of, um, psychologists that work in cancer care services. Across the state, I think there’s about 17 of us.”

Therapists were generally positive about alternative strategies to address these barriers, including delivering ConquerFear in group or online format

“... a good way of increasing the reach that, um, the minute it’s online it has the potential to reach a global community rather than just, um, one person at a time. So, um, yeah, that’s one obvious one.”

However some expressed concerns about moving away from an individually-delivered model, suggesting that aspects such as previous trauma and existential issues could not be fully and safely explored in a group or unsupported online environment.

“I don’t think that it would be, um, ethically as appropriate to talk about some of the vulnerability factors” [in either group or internet formats].

“I think you’d need to have a therapist, sort of still guiding or assisting or as a contact point.”

3.2 Intervention/patient fit

Many therapists noted that ConquerFear was more appropriate for some patients than others. Specifically, some felt that patients who were psychologically minded, open to experiencing emotion and less avoidant, and ready to commit to an extended therapy, would benefit more.

“I really have to admit, I think quite verbally intelligent, psychologically minded people would work well with this model....”

“So, I think that it is actually quite a confronting therapy and I think for people who ... are more likely to cope with their fear of cancer coming back through suppression and avoidance, rather than being hypervigilant uh, they were actually quite tricky to tap into because the therapy was making them face some of the stuff that they, yeah, weren’t really wanting to.”

However, some therapists believed that ConquerFear was appropriate for any patient, and indeed, expressed their surprise that patients who they had initially thought would not understand or engage with the therapy very much enjoyed and benefited from it.

“I think it doesn’t matter what, where, who you are, you would still benefit from this program.”

“The third lady again was quite elderly and ... I thought oh yeah, I’m really struggling to get this across to her, but she seemed to really like it and she actually did more of the ATT [attention training technique] training than I thought she would.”

3.3 Training, support and resources

Most therapists discussed the importance of adequate training, mentorship and ongoing support to build their confidence in delivering the intervention and ensure a high standard of delivery.

ConquerFear resources, including a printed manual, summaries of each session, printed patient handouts and a compact disc with attention training exercises, were all cited as immensely helpful in building and maintaining skills.

“It’s a very self-explanatory manual, it’s a very, you know, good - it’s pretty easy to follow.”

“People cannot be given the manual, to run with, they have to do the training”

“...ongoing peer supervision, so if we were having any problems implementing it or we weren’t sure about something we could contact somebody who knew the program....”

Discussion

The aim of this study was to investigate experienced therapists’ perceptions of factors influencing the sustainability of ConquerFear. We used the PARHiS model [14] to guide our

questioning and analysis, and indeed found that the identified themes fitted well within the three PAHRiS domains: evidence, contextual, and facilitation factors.

The PARiHS framework posits that intervention sustainability is most likely when staff perceive the evidence for the intervention to be robust, both in terms of evidence based on well-conducted studies with stringent designs, and personal experience of the need for and efficacy of interventions when delivered in real life, derived from observation or patient or staff feedback [14]. Both forms of evidence were evident in therapists' discourse during study interviews. The high quality of evidence provided by a rigorously designed randomized controlled trial conducted by a credible research team was cited as enabling therapists to both confidently deliver ConquerFear and convince oncology colleagues to refer eligible patients.

Therapists also spoke vividly and powerfully about their personal experience of the intervention's efficacy with individual patients. They had observed high patient engagement with the therapy, and its ability to transform patient outcomes, which was a strong motivator to continue using ConquerFear. Therapists' personal involvement in the ConquerFear trial also ensured early awareness of, and confidence in, trial results, which has been found in other studies to encourage early uptake of proven interventions [16].

There was more diversity in therapists' beliefs about the need for ConquerFear. While some reported that FCR was a common and significant problem in their practice, others reported that severe FCR was rarely seen, meaning that ConquerFear was not often required (although they used elements for sub-clinical FCR). The definition of clinical FCR is still controversial, although a recent Delphi study [1] involving experts in FCR identified the following characteristics as indicating clinical levels of FCR: a preoccupation with the cancer returning or progressing; unhelpful coping behaviours; impairment in daily function; clinically significant

distress; and limited capacity for making future plans. A score ≥ 13 on the Fear of Cancer Recurrence Inventory (FCRI) has been suggested as a clinical cut-off, although recent research suggests ≥ 22 may be more appropriate [17]. This confusion in the literature is undoubtedly contributing to confusion in the clinic regarding who to refer to ConquerFear. Our recent analysis of mediators and moderators of ConquerFear outcomes [18] showed that those with lower levels of FCR appeared to benefit equally from simple relaxation strategies, providing more support for a cut-off of 22. More work is required to build consensus around the assessment processes to guide clinicians in this decision.

The PARiHS framework posits that some contexts are more conducive to successful implementation of evidence into practice than others. For example, organizations that have strong leaders who act as champions for change, that are flexible and responsive to changing circumstances and have a process for monitoring progress through change [14] are more likely to implement new interventions successfully. In this study, contexts where the focus was on patients in active treatment and which did not provide resources, space or time for the care of survivors, were not conducive to routine implementation of ConquerFear. Survivorship has increasingly been recognized as a priority in cancer care, as the number of survivors living with long-term side-effects of complex and prolonged treatments has increased. Indeed, it has been noted that need for help with FCR in patients receiving novel treatments such as immunotherapy for which no long-term data exist to ameliorate uncertainty, may be even greater than in other cohorts [19]. While guidelines and mission statements are increasingly emphasizing clinical responsibility for survivorship, resource-poor health systems and clinical services have been slower to respond.

Similarly, services in which psychosocial support was less embedded, routine screening for distress was not routine, and where Oncology staff lacked the confidence and skills to recognize and refer patients with high FCR, struggled to implement ConquerFear. Therapists who were rarely called upon to deliver ConquerFear perceived rapid loss of the therapeutic skills they had learned, hindering delivery of the intervention. Furthermore, a low referral base also contributed to a perceived low need for the intervention.

Flexibility and openness to change was also an issue at the therapist level. Those with a similar theoretical orientation to ConquerFear, who were familiar and comfortable with the components of ConquerFear, were more likely to embrace these methods, and welcome the opportunity to improve their skills. Some therapists noted that additional training and support would overcome initial resistance and build confidence.

The PaHRIS framework also recognises the importance of facilitators to implementation, and indeed, the fit between requirements of ConquerFear, resources available and the patient base of the site, influenced uptake of ConquerFear in routine care. Face-to-face delivery and long sessions could not easily be accommodated at some sites due to insufficient psychosocial oncology staff and defined scheduling of appointments, requiring adaptation of the intervention. Clearly, intervention sustainability is enhanced if it is responsive to clinical needs and flexible in its delivery, for example, by dividing up sessions into components that can be delivered in a flexible number of sessions. Similarly, therapists noted that ConquerFear might be appropriate for some, but not all patients. Patients who used an avoidant coping style were thought to find the therapy more confronting, although this has not been born out in our mediator/moderator analysis of ConquerFear, where avoidance was not found to be associated with therapy uptake or outcome [18].

Other formats (group or internet-based, or delivery by community-based psychologists) were suggested as a way to overcome resource gaps. E-health has become increasingly popular over recent years [20], with evidence they can be equally effective as face-to-face formats. However, it remains to be seen whether elements of psychosocial interventions such as review of trauma and homework where sustained home practice is required, can be delivered effectively online or in group format where privacy and confidentiality are harder to ensure and a therapeutic bond is lacking. Trials of both an internet and group versions of ConquerFear are currently underway as blended interventions with initial face-to-face individual therapeutic contact, as per therapists' recommendation.

These findings have some important theoretical and clinical implications. Firstly, the impact of therapists' personal experience of ConquerFear's efficacy underscores the utility of designing pragmatic phase 3 randomised controlled trials with sustainability in mind. We ensured widespread exposure of key Psycho-oncology staff to ConquerFear by recruiting 26 therapists around Australia to deliver the intervention, potentially sacrificing tight control over consistent therapy delivery (although our fidelity instructions, cues and checks did result in high treatment consistency within the trial). Once exposed to ConquerFear, and its impact on patients, most therapists have become key champions of its ongoing implementation around Australia.

We need to acknowledge some limitations of the present study. First, while the perspectives of therapists from the ConquerFear trial are clearly informative, it would also be useful to explore implementation perspectives of therapists less familiar with the intervention, and of other key staff (such as medical oncologists and administrators).

These limitations notwithstanding, this study has a number of strengths. We employed rigorous methods in gathering and analysing data and reached saturation in our sample. We used

a well-recognised implementation framework (PAHRiS) to guide our analysis. The present results confirm that ConquerFear is sustainable, provided it can accommodate local resources and constraints. Our results suggest that a carefully designed sustainability strategy incorporating adequate training and support that elicits and addresses therapists' concerns, tailoring of intervention content and format to local needs, and communication of both objective and subjective evidence of the efficacy of interventions, is required to ensure their uptake into routine care. The PAHRiS framework proved to be a useful lens with which to understand sustainability issues and might be usefully employed to design such a strategy.

Compliance with Ethical Standards:

This study was unfunded.

Phyllis Butow declares that she has no conflict of interest. Douglas Williams declares that he has no conflict of interest. Belinda Thewes declares that she has no conflict of interest. Stephanie Tesson declares that she has no conflict of interest. Louise Sharpe declares that she has no conflict of interest. Allan 'Ben' Smith declares that he has no conflict of interest. Joanna E Fardell declares that she has no conflict of interest. Jane Turner declares that she has no conflict of interest. Jemma Gilchrist declares that she has no conflict of interest. Afaf Girgis declares that she has no conflict of interest. Jane Beith declares that she has no conflict of interest.

Ethical approval:

This article does not contain any studies with animals performed by any of the authors.

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent was obtained from all individual participants included in the study.

REFERENCES

1. Lebel S, Ozakinci G, Humphris G, Mutsaers B, Thewes B, Prins J, et al. From normal response to clinical problem: definition and clinical features of fear of cancer recurrence. *Support Care Cancer*. 2016;24(8):3265-8.
2. Simard S, Thewes B, Humphris G, et al: Fear of cancer recurrence in adult cancer survivors: a systematic review of quantitative studies. *J Canc Surviv* 7(3): 300-322, 2013.
3. Thewes B, Butow P, Bell M, et al: Fear of cancer recurrence in young women with a history of early-stage breast cancer: a cross-sectional study of prevalence and association with health behaviours. *Support Care Cancer* 20(11): 2651-2659, 2012.
4. Koch L, Bertram H, Eberle A, et al: Fear of recurrence in long-term breast cancer survivors-still an issue. Results on prevalence, determinants, and the association with quality of life and depression from the cancer survivorship-a multi-regional population-based study. *Psychooncology* 23(5):547-54, 2014.
5. Butow P, Sharpe L, Thewes B, Turner J, Gilchrist J, Girgis A, et al. A randomized controlled trial (RCT) of a psychological intervention (Conquer Fear) to reduce clinical levels of fear of cancer recurrence in breast, colorectal and melanoma cancer survivors. *J Clin Oncol*, 2017; 35(36):4066-4077.
6. Fardell, J. E., Thewes, B., Turner, J., Gilchrist, J., Sharpe, L., Smith, A., . . . Butow, P. (2016). Fear of cancer recurrence: a theoretical review and novel cognitive processing formulation. *J Cancer Surviv*, 10: 663.
7. Wells, A. (1999). A metacognitive model and therapy for generalized anxiety disorder. *Clinical Psychology & Psychotherapy*, 6(2), 86-95.
8. Wells, A., & Matthews, G. (1996). Modelling cognition in emotional disorder: The S-REF model. *Behaviour research and therapy*, 34(11-12), 881-888.
9. Nerenz, D. R., & Leventhal, H. (1983). Self-regulation theory in chronic illness. *Coping with chronic disease: Research and applications*, 13-38.

10. Hayes, S. C., Luoma, J. B., Bond, F. W., Masuda, A., & Lillis, J. (2006). Acceptance and commitment therapy: Model, processes and outcomes. *Behaviour research and therapy*, 44(1), 1-25.
11. Scheirer MA, Dearing JW. An agenda for research on the sustainability of public health programs. *Am J Public Health*. 2011;101(11):2059.
12. Grol R. Successes and failures in the implementation of evidence-based guidelines for clinical practice. *Med Care*. 2001;39:S46-54.
13. Nilsen P. Making sense of implementation theories, models and frameworks. *Implementation Science* 2015;10.
14. Kitson A, Harvey G, McCormack B. Enabling the implementation of evidence-based practice: a conceptual framework. *Quality in Health Care*. 1998;7:149-58.
15. Gale NK, Heath G, Cameron E, Rashid S, Redwood S. Using the framework method for the analysis of qualitative data in multi-disciplinary health research. *BMC Med Res Methodol* 2013;13:117.
16. Majumdar SR, Roe MT, Peterson ED, et al. Better outcomes for patients treated at hospitals that participate in clinical trials, *Arch Intern Med*, 2008;168, 657-662.
17. Fardell JE, Jones G, Smith AB, Lebel S, Thewes B, Costa D et al. (2017). Exploring the screening capacity of the Fear of Cancer Recurrence Inventory - Short Form for clinical levels of fear of cancer recurrence. *Psychooncology*. doi:10.1002/pon.4516
18. Sharpe L, Turner J, Fardell J, Thewes B, Smith B, Gilchrist J, et al. ConquerFear for treating fear of cancer recurrence: Mediators and Moderators of treatment efficacy. Under review.
19. Thewes B, Husson O, Poort J, Custers JAE, Butow PN, McLachlan S-A, Prins JB. Fear of cancer recurrence in an era of personalized medicine. *JCO* 2017; 35 (29): 3275-3278.
20. McAlpine H, Joubert L, Martin-Sanchez F, Merolli M, Drummond KJ. A systematic review of types and efficacy of online interventions for cancer patients. *Patient Educ Couns* 2015; 98(3): 283-95.

Table 1. Qualitative themes as related to the PARiHS framework

Evidence	Context	Facilitation
Intervention credibility Experienced efficacy Perceived need	Attitude to and capacity for survivorship/FCR care Therapist orientation and flexibility Expert referral	Intervention fit Patient/intervention fit Training, support and resources