Manuscript Details

Manuscript number	PSY_2020_497_R1
Title	Muscarinic M1 and M4 receptors: Sites for hypothesis driven drug development for schizophrenia
Article type	Review Article

Abstract

The finding that the drug KarXT, a formulation of xanomeline and trospium which targets muscarinic receptors, has given a positive result in reducing the positive and negative symptoms of schizophrenia in a phase II trial suggests targeting muscarinic receptors is a new approach to treating the disorder. This review will detail the synergistic interplay between studies to understand the role of muscarinic receptors in the aetiology of schizophrenia and drug development and how this has supported the hypothesis that activating the muscarinic M1 and M4 receptors is critical to the efficacy of KarXT, in schizophrenia. The discovery of an intermediate phenotype within schizophrenia which is characterised by the presence of a marked loss of cortical muscarinic M1 receptors will be reviewed. Highlighted will be progress in understanding the biochemistry of that intermediate phenotype and evidence to suggest that those with the intermediate phenotype may resist treatment with agonist to the orthosteric site on the muscarinic M1 and M4 receptor. Finally, the possibility of using drugs targeting the allosteric binding sites on muscarinic receptors to treat schizophrenia will be discussed. This timely review will therefore highlight how research can influence hypothesis driven drug discovery that should produce new treatments for schizophrenia.

Keywords	schizophrenia, muscarinic receptor, cortex, drug development
Corresponding Author	Brian Dean
Order of Authors	Brian Dean, Elizabeth Scarr
Suggested reviewers	Claudia Vingerhoets, Thomas Raedler, Anantha Shekhar, Jeffrey Conn, David Volk

Submission Files Included in this PDF

File Name [File Type]

Covering Letter.docx [Cover Letter]

Rebuttal.docx [Response to Reviewers]

Highlights.docx [Highlights]

Abstract.docx [Abstract]

Title Page.docx [Title Page (with Author Details)]

CHRM Review_Post Review_Tracked Changes.docx [Manuscript File]

Figure 1.tif [Figure]

Figure 2.jpg [Figure]

Figure 3.jpg [Figure]

Conflicts of Interest.docx [Conflict of Interest]

Author Statement.docx [Author Statement]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

30 Royal Parade,
Parkville,
Victoria 3052,
Australia.

Telephone: 613-8344-3786
Facsimile: 613-9348-1707
Email: anddali@unimelb.edu.au

5 April 2020

Dr Lynn DeLisi
Editor-in-Chief
Psychiatry Research

Dear Dr DeLisi

Herewith find a copy of our revised manuscript entitled “Muscarinic M1 and M4 receptors: Sites for hypothesis driven drug development for schizophrenia” for consideration for publication in Psychiatry Research.

We look forward to hearing from you on this matter.

Yours sincerely,

On Behalf of the Authors

5 April 2020

Dr Lynn DeLisi
Editor-in-Chief
Psychiatry Research

Dear Dr DeLisi

We thank the Reviewers for their comments on our manuscript entitled “Muscarinic M1 and M4 receptors: Sites for hypothesis driven drug development for schizophrenia” (Ref: PSY_2020_497). We have used these comments extensively to modify the original manuscript we submitted as follows:

Reviewer 1:

1. the absence of GWAS data implicating muscarinic receptors. The authors carefully describe postmortem and clinical findings that implicate M1/M4 in schizophrenia, but without supporting GWAS data. This review would be an excellent venue to discuss the mechanisms by which the human brain in schizophrenia can manifest profound changes in cortical circuitry in the absence of GWAS findings.

Response: A paragraph has been added to the conclusions of our revised manuscript that addresses this important issue.

Reviewer 2

1. One of my concerns is that the results of this phase II study have not been published yet (other than in a press-release to investors). This fact needs to be clarified in the manuscript. While I share the authors’ enthusiasm about the positive results of a phase II study, these results need to be confirmed in larger phase III studies. As bitopertin and encenicline have shown, the results of a positive phase II study don’t necessarily translate in a positive phase III study. I would ask the authors to clarify this point and not to raise too many expectations.

Response: Sad but true comments, even though the main problem seems to be a growing placebo effect in phase III trials! We have added this cautionary message to our Concluding Remarks where we feel it will have greatest impact.

2. Unfortunately the submission has different abstracts. Please make sure that you use the same version of the abstract.

Response: The authors apologise for this error which is corrected in our re-submission.

3. The authors need to provide more information on KarXT in the abstract. Many readers may not know that this abbreviation stands for a commercial formulation of xanomeline and trospium.

Response: The authors acknowledge this important point and now define the formulation of KarXT in the abstract.

4. The authors also talk about ‘a large phase II study’. With a recruitment goal of a total of 180 subjects, this raises wrong expectations.
Response: The authors agree with this comment and therefore have deleted the word “large”.
5. Referring to the Shekhar study as a ‘drug trial’ is pushing things (the number of subjects enrolled was very small).
Response: This is a fair comment which is why the authors refer to the “clinical trial” of xanomeline in schizophrenia, we believe this is a reasonable name for this early trial of the drug in treating schizophrenia. We could find nowhere in the manuscript we referred to this study as a “drug trial”.
6. Please replace PANNS with PANSS!
Response: This modification has been made.
7. Page 3 / 4: The whole paragraph ‘Whilst early attempts’ is misleading from my perspective. The authors state that ‘activating, rather than blocking, muscarinic receptors could be beneficial in the treatment of schizophrenia’. The therapeutic goal of administering muscarinic antagonists is to ameliorate antipsychotic-induced motor side-effects.
Response: In response to the Reviewer finding this paragraph misleading we have edited the first sentence to clearly indicate the early attempts to treat patients with schizophrenia involved the use of atropine toxicity, not the use of muscarinic antagonists to control the extrapyramidal side-effects of typical antipsychotic drugs as covered in the second sentence of the revised manuscript.
8. The discussion of allosteric versus orthosteric M1/M4 agonists should mention that no allosteric agonist has yet proven effective in the treatment of schizophrenia.
Response: This is a fair comment and a sentence has been added to the relevant paragraph acknowledging this point.
9. The first sentence ‘The release of promising data showing the efficacy of KarXT’ is unacceptable. These are preliminary data (keep in mind that this manuscript is being submitted to scientific journal not a commercial publication).
Response: Acknowledged, the word “preliminary” now replaces “promising”.
10. I don’t fully understand how the reference to van Os’ paper improves this manuscript
Response: This point is made to warn that attempts to remove the diagnoses of schizophrenia, as has occurred in Japan, may be premature and could disadvantage ongoing studies as whatever the diagnosis “defines” does have a biological underpinning.
11. The Figures should clarify that they refer data from previous presentations.
Response: It has been made clear that Figure 1 and Figure 2A and 2B have been created using information relate to previous publications.

Reviewer 3:

1. Page 5; Please refer to the [123I]IQNB ligand as *single photon emission computed tomography ligand* (instead of “single photon computed emission tomography radioligand”).
Response: This has been corrected.
2. -Page 5: The authors write: “In addition, using the more selective SPECT ligand, 123I-IDEX, it was shown that lower levels of muscarinic M1 receptors were associated with poorer performance in verbal learning and memory and more severe negative symptoms in medication free patients with a psychotic disorder (Bakker et al., 2018)”. Can the authors please clarify that in this study no control group was included and that therefore, it is unknown whether M1 receptor density was decreased in these patients.
Response: This important point has been included in our revised manuscript.
3. -Page 7: “higher levels of miR-107 which targets muscarinic M1 receptor expression (Scarr et al., 2013a)”. Please insert the word and prio to higher....

Response: This has been corrected in our revised manuscript.

4. -Page 7: the authors describe a few molecular cytoarchitectural characteristics of the MRDS group. Are these changes they describe compared to non-MRDS schizophrenia patients or controls or both? If changes are compared to non-MRDS schizophrenia patients, is it known how non-MRDS schizophrenia patients relate to controls?

Response: In our revised manuscript we now make it clear the measures in question do not differ between non-MRDS and controls and include a comment on the only measure we have found that differentiates non-MRDS from controls and MRDS.

5. -Page 12: beginning of the paragraph. "Given the notion of using allosteric modulators to treat schizophrenia it is of interest that the ability of the muscarinic M1 receptor allosteric modulator to alter the affinity of that receptor can be measured using human cortex (Dean et al., 2016), a measurement that should be an indicator of the likely responsiveness of that receptor to allosteric modulation (Ma et al., 2009; Shirey et al., 2009)." This sentence is very long and difficult to read. Can the authors rephrase?

Response: In our revised manuscript, this sentence has been rephrased and made into two sentences in such a way we hope makes its meaning clearer.

In addition to the above, we have revised the highlights for this manuscript to reflect changes made in response to the Reviewers comments.

As we have been able to improve our manuscript by incorporating nearly all of the changes suggested by the Reviewers, we now submitted our revised manuscript for consideration for publication in Psychiatry Research.

We look forward to hearing from you on this matter.

Yours sincerely,

Brian Dean

On behalf of the authors.

Highlights

- Commentary on Two drug trials now reports that treating patients with schizophrenia with muscarinic M1 / M4 receptor agonists reduces the severity of positive and negative symptoms. On study also showed improvements in cognitive deficits.
- A review of Postmortem-postmortem and neuroimaging studies showing lower levels of muscarinic M1 and M4 receptors in the CNS of patients with schizophrenia, suggesting treating with muscarinic receptor agonists should is-restoreing cholinergic homeostasis in the CNS of those with the disorder.
- An update on the biochemical abnormalities in the cortex from ~~discovery of~~ a cholinergic intermediate phenotype within ~~schizophrenia and progress in understanding its underlying~~ aetiology.
- Discussion as to whether those in the cholinergic intermediate phenotype may be resistant to treatment with drugs that target the orthosteric, but not allosteric, site on the muscarinic m1 and M4 receptors.

The finding that the drug KarXT, a formulation of xanomeline and trospium which targets muscarinic receptors, has given a positive result in reducing the positive and negative symptoms of schizophrenia in a phase II trial suggests targeting muscarinic receptors is a new approach to treating the disorder. This review will detail the synergistic interplay between studies to understand the role of muscarinic receptors in the aetiology of schizophrenia and drug development and how this has supported the hypothesis that activating the muscarinic M1 and M4 receptors is critical to the efficacy of KarXT, in schizophrenia. The discovery of an intermediate phenotype within schizophrenia which is characterised by the presence of a marked loss of cortical muscarinic M1 receptors will be reviewed. Highlighted will be progress in understanding the biochemistry of that intermediate phenotype and evidence to suggest that those with the intermediate phenotype may resist treatment with agonist to the orthosteric site on the muscarinic M1 and M4 receptor. Finally, the possibility of using drugs targeting the allosteric binding sites on muscarinic receptors to treat schizophrenia will be discussed. This timely review will therefore highlight how research can influence hypothesis driven drug discovery that should produce new treatments for schizophrenia.

Muscarinic M1 and M4 receptors:

Sites for hypothesis driven drug development for schizophrenia.

Brian Dean^{a*} and Elizabeth Scarr^b

^a The Molecular Psychiatry Laboratory, The Florey Institute for Neuroscience and Mental Health,
Parkville, Victoria 3052, Australia,

^b Melbourne Veterinary School, Faculty of Veterinary and Agricultural Sciences, University of
Melbourne, Parkville, Victoria 3010, Australia.

* Corresponding Author: Professor Brian Dean, Head, The Molecular
Psychiatry Laboratory, The Florey Institute for
Neuroscience and Mental Health, 30 Royal Parade,
Parkville, Victoria 3052, Australia.

Email: brian.dean@florey.edu.au; Telephone: 613-8344-3786

Muscarinic M1 and M4 receptors:
Hypothesis driven drug development for schizophrenia.

Brian Dean^{a*} and Elizabeth Scarr^b

^a The Molecular Psychiatry Laboratory, The Florey Institute for Neuroscience and Mental Health,
Parkville, Victoria 3052, Australia,

^b Melbourne Veterinary School, Faculty of Veterinary and Agricultural Sciences, University of
Melbourne, Parkville, Victoria 3010, Australia.

* Corresponding Author: Professor Brian Dean, Head, The Molecular
Psychiatry Laboratory, The Florey Institute for
Neuroscience and Mental Health, 30 Royal Parade,
Parkville, Victoria 3052, Australia.

Email: brian.dean@florey.edu.au; Telephone: 613-8344-3786

Abstract

The finding that the drug KarXT, a formulation of xanomeline and trospium which targets muscarinic receptors, has given a positive result in reducing the positive and negative symptoms of schizophrenia in a phase II trial suggests targeting muscarinic receptors is a new approach to treating the disorder. This review will detail the synergistic interplay between studies to understand the role of muscarinic receptors in the aetiology of schizophrenia and drug development and how this has supported the hypothesis that activating the muscarinic M1 and M4 receptors is critical to the efficacy of KarXT, in schizophrenia. The discovery of an intermediate phenotype within schizophrenia which is characterised by the presence of a marked loss of cortical muscarinic M1 receptors will be reviewed. Highlighted will be progress in understanding the biochemistry of that intermediate phenotype and evidence to suggest that those with the intermediate phenotype may resist treatment with agonist to the orthosteric site on the muscarinic M1 and M4 receptor. Finally, the possibility of using drugs targeting the allosteric binding sites on muscarinic receptors to treat schizophrenia will be discussed. This timely review will therefore highlight how research can influence hypothesis driven drug discovery that should produce new treatments for schizophrenia.

Drugs that affect the cholinergic system have long been suggested to reduce the severity of symptoms in patients with schizophrenia (Cohen et al., 1944). The human cholinergic system is made up of basal-forebrain and medial septal nucleus projections, the pedunculo-pontine–lateral dorsal tegmental neurons, striatal interneurons, the cranial-nerve nuclei, the vestibular nuclei and the spinal-cord preganglionic and motoneurons (Perry et al., 1999; Scarr et al., 2013b). The components of the cholinergic system can influence many CNS regions, in particular neurons from the basal-forebrain and the medial septal nucleus extend into the cortex and hippocampus. The activity of the cholinergic system is regulated by acetylcholine acting through two families of receptors, the ion-gated nicotinic receptors and the G-protein coupled muscarinic receptors (Scarr et al., 2013b). There are 5 muscarinic receptors (M1 to 5) in the human CNS, the variation in their distribution and signalling are critical in mediating the overall responsiveness of the cholinergic system to activation by acetylcholine (Brann et al., 1993). As a result of recent developments, this review will focus on the roles of muscarinic receptors in the pathophysiology and treatment of schizophrenia, however, there is also evidence to implicate nicotinic receptors in the pathophysiology and treatment of the disorder; this has been reviewed elsewhere (Girgis et al., 2019).

Targeting muscarinic receptors to treat schizophrenia

Whilst early attempts to ameliorate the severity of the symptoms of schizophrenia using atropine toxicity, which involved ~~by~~ blockading muscarinic receptors with the that antagonist, ~~atropine~~, showed some initial promise (Forrer, 1951) this treatment had a high risk/benefit ratio and therefore was not viewed as clinically useful. However, there ~~was~~ has been the ~~a~~ widespread use of muscarinic receptor antagonists, such as benztropine, to lessen the movement disorders experienced

by people with schizophrenia treated with the first-generation antipsychotic drugs (Ogino et al., 2014). Whilst the ongoing use of muscarinic receptor antagonists to control the movement disorders associated with antipsychotic drug treatment suggested a clear clinical benefit such treatment was also associated with worsening of cognitive impairments and tardive dyskinesia as well as the onset of delirium (Minzenberg et al., 2004). These treatment outcomes contrasted to the finding that chewing betel nut lessened the severity of symptoms of schizophrenia (Sullivan et al., 2000), an effect thought to be due to betel nut containing arecoline, a non-selective muscarinic receptor agonist (Sullivan et al., 2007). Together, these data suggested activating, rather than blocking, muscarinic receptors could be beneficial in the treatment of schizophrenia.

Studies on muscarinic receptors in CNS in schizophrenia

Given studies suggesting activating muscarinic receptors would be beneficial in the treatment of schizophrenia it was significant that a number of studies reported lower levels of [³H]pirenzepine binding in the cortex (Crook et al., 2001; Zavitsanou et al., 2004), hippocampus (Crook et al., 2000) and striatum (Dean et al., 1996), but not the thalamus (Dean et al., 2004), from patients with schizophrenia because [³H]pirenzepine selectively binds to the muscarinic 1 and 4 receptors (Scarr and Dean, 2008). These binding data suggested that, compared to controls, there were lower levels of muscarinic M1 receptor, muscarinic M4 receptor or both of these receptors in the cortex, hippocampus and striatum from patients with schizophrenia. Moreover, lower levels of cortical [³H]pirenzepine binding had some specificity for schizophrenia because lower levels of binding was not found in the cortex of subjects with major depressive disorders, bipolar disorders (Gibbons et al., 2016; Zavitsanou et al., 2004), Alzheimer's disease (Scarr et al., 2017) or Parkinson's disease (McOmish et al., 2017).

One concern about post-mortem CNS findings was that they were an artefact due to autolysis or other factors occurring after death. It is therefore significant that the use of a pan-muscarinic receptor ~~single photon emission computed tomography ligand~~~~single photon computed emission tomography radioligand~~, [I-123]iodoquinuclidinyl benzilate ([¹²³I]IQNB), confirmed there were lower levels of CHRMs in many CNS regions in antipsychotic-drug free patients with schizophrenia (Raedler et al., 2003). In addition, using the more selective SPECT ligand, ¹²³I-IDEX, it was shown that lower levels of muscarinic M1 receptors were associated with poorer performance in verbal learning and memory and more severe negative symptoms in medication free patients with a psychotic disorder (Bakker et al., 2018). Importantly, this study did not have data from normal controls and therefore it was not possible to determine if the patients with schizophrenia had lower levels of CNS muscarinic M1 receptors compared to those who did not have the disorder. —These neuroimaging data added to the hypothesis, put forward after the emergence of post-mortem data showing lower levels of muscarinic M1 and / or M4 receptors in the CNS from subjects with schizophrenia, that activating the muscarinic M1 and / or M4 receptors with drugs would be of therapeutic benefit in patients with the disorder (Dean et al., 2003; Felder et al., 2001).

The use of [³H]pirenzepine binding and SPECT neuroimaging suggested there were lower levels of muscarinic M1 and / or M4 receptors in patients with schizophrenia. However, the use of receptor specific antibodies and the measurement of levels of mRNA showed that there were lower levels of muscarinic M1 receptor (Dean et al., 2002; Mancama et al., 2003), but not muscarinic 2, 3 (Scarr et al., 2006) or 4 (Dean et al., 2002) receptors, in the dorsolateral prefrontal cortex from patients with schizophrenia. By contrast, data from the measurement of mRNA suggested the level of muscarinic M1 receptor was not altered in the striatum (Dean et al., 2000) or hippocampus (Scarr et al., 2007) from patients with schizophrenia, rather, levels of muscarinic M4 receptor were lower in the

hippocampus from those with the disorder (Scarr et al., 2007). These data argued that stimulating the muscarinic M1 and / or 4 receptors could give therapeutic benefits by ensuring adequate signaling through the muscarinic M1 receptor in the cortex and muscarinic M4 receptors in the hippocampus and sub-cortical regions in patients with schizophrenia.

The use of a muscarinic M1 / M4 receptor agonist in the treatment of schizophrenia

Initially conceptualised to treat the symptoms of Alzheimer's disease, the drug xanomeline was designed to be a high affinity agonist at the muscarinic M1 receptor (Shannon et al., 1994). However, it was subsequently shown that xanomeline was an agonist at the muscarinic M1 receptor and muscarinic M4 receptor and to have antipsychotic like behavioural properties in rodents (Shannon et al., 2000) and primates (Andersen et al., 2003). These data, added to a growing understanding of changes in levels of muscarinic receptors in the CNS of patients with schizophrenia, led to the proposal that xanomeline would prove useful in treating the symptoms of that disorder (Bymaster et al., 2002). This hypothesis was supported by a clinical trial in patients with schizophrenia (10 xanomeline vs 10 placebo) which showed that after 4 weeks of treatment xanomeline improved the total Brief Psychiatric Rating Scale and Positive and Negative Syndrome scores and robustly improved measures of verbal learning and short term memory function (Shekhar et al., 2008). In this study adverse events, particularly involving the gastrointestinal system, were reported and hence xanomeline did not appear an attractive drug to take through to the clinic (Bender et al., 2017).

Whilst the side-effects of xanomeline made it an unattractive drug to take to the clinic as a monotherapy the results obtained in the trial of the drug in subjects with schizophrenia amplified ongoing interest in the use of muscarinic M1 / M4 receptor agonists to treat schizophrenia (Bender

et al., 2017). This led to the development of a coformulation of xanomeline and trospium which has been named KarXT (Miller et al., 2016). Trospium was included because it is a muscarinic receptor antagonist that has minimal, if any, penetration of the blood brain barrier (Staskin et al., 2010) which means its actions are limited to the periphery. Therefore, the hypothesis underpinning the formulation of xanomeline with trospium was that trospium would block the unwanted peripheral side-effects of xanomeline. In a recent announcement of the outcome of a Phase 2 clinical trial it was stated that KarXT gave a clinically meaningful and statistically significant reduction in total Positive and Negative Syndrome Scales compared to placebo that included significant reductions in both PANSS-Positive and PANSS-negative scores (<https://karunatx.com/>). Importantly, the number of discontinuations of treatment due to adverse effects in the KarXT and placebo study arms were equal. These data add strong support to the notion that activating CNS muscarinic M1 and M4 receptors will have significant therapeutic benefits for patients with schizophrenia.

A muscarinic M1 receptor-intermediate phenotype in schizophrenia

It has been suggested that studying intermediate phenotypes in disorders such as schizophrenia will improve understanding of the pathophysiology of the disorder (Tamminga et al., 2017). Establishing such intermediate phenotypes is dependent on identifying bio- or clinical markers that can be used to differentiate groups within syndromes such as schizophrenia. Hence, it was significant that within a large study on the levels of CHRMs in the cortex of patients with schizophrenia it was shown that 25% of those within the syndrome could be separated into a sub-group who, on average, had a 75% lower level of cortical muscarinic M1 receptors (Figure 1)(Scarr et al., 2009). Further study of the molecular cytoarchitecture of this sub-group, termed a muscarinic receptor deficit sub-group with schizophrenia (MRDS), has shown they have widespread decreases in cortical muscarinic M1 receptor (Gibbons et al., 2013; Seo et al., 2014), altered patterns of the

muscarinic M1 receptor gene promoter methylation (Scarr et al., 2013a), lower levels of muscarinic M1 receptor mRNA (Scarr et al., 2013a) and, higher levels of miR-107 which targets muscarinic M1 receptor expression (Scarr et al., 2013a) when compared to controls. Notably, non-MRDS do not differ in these measures from controls but, unlike MRDS, have changes in levels of pyruvate kinase in their striatum (Dean et al., 2016b) which could be inhibiting their ability to utilise glucose.

The alteration in gene promoter methylation and higher levels of miR-107 suggests part of the pathophysiology of MRDS involves mechanisms driving a decrease in muscarinic M1 receptor expression in patients with MRDS. Further evidence that patients with MRDS display some differences in pathophysiology from other subjects with schizophrenia comes from a transcriptomics study showing there are changes in gene expression that occur in the cortex of patients with MRDS but not Non-MRDS (Scarr et al., 2018c). Some of those changes in gene expression include genes that were able to be grouped into a gene expression interactome. A gene expression interactome includes networks of genes with proven molecular interactions that likely constitute pathways affected by the pathophysiology of a disorder (Caldera et al., 2017). Significantly, the pathway of genes identifies in the cortex of subjects with MRDS is also predicted to impact on muscarinic M1 receptor gene expression (Scarr et al., 2018c).

Moving to sub-cortical pathophysiology, patients with MRDS do not have lower levels of striatal pyruvate kinase found in subjects with schizophrenia that do not have an MRDS intermediate phenotype (non-MRDS) (Dean et al., 2015). Pyruvate kinase is a rate limiting enzyme in the Embden-Meyerhof pathway critical to glucose utilisation (Luo and Semenza, 2011) and hence the lower levels of that enzyme in the striatum of patients with non-MRDS could be contributing to the decreased ability to utilise glucose that has been reported using PET neuroimaging studies of schizophrenia (Buchsbaum and Hazlett, 1998; Horga et al., 2011).

Recently, it has been shown that muscarinic M1 receptors are highly expressed by pyramidal cells (Figure 2A) in laminae III and V in the human dorsolateral prefrontal cortex (Figure 2B). Notably, compared to controls, there are markedly lower numbers of muscarinic M1 receptor positive pyramidal neurons in both these cortical laminae in patients with MRDS and there a significantly lower levels of muscarinic M1 receptor positive neurons in both cortical laminae in those with non-MRDS (Scarr et al., 2018a). Pyramidal cells are glutamatergic neurons that are present in all laminae of the human cortex, except laminae I (Defelipe, 2011). Notably, activation of the muscarinic M1 receptor causes an increase in levels of extracellular glutamate (Sanz et al., 1997), presumably by causing an increased efflux of the neurotransmitter from the pyramidal neurons. Therefore, it is likely that reductions in muscarinic M1 receptor positive pyramidal cells would impact on cortical glutamate homeostasis. Whilst there were lower levels of muscarinic M1 receptor positive neurons in MRDS and non-MRDS, there was no decrease in total neuronal number in laminae III or V in either of those groups. This shows there must be higher levels of muscarinic M1 receptor negative neurons in both MRDS and non-MRDS. Hence, at this point, it would not be possible to predict whether such a complex breakdown in control of glutamate homeostasis would be associated with a hypo-glutamatergic state or a hyper-glutamatergic state. Being able to make such a delineation would be important as a hyper-glutamatergic state is predicted to be part of the pathophysiology of schizophrenia (Moghaddam and Javitt, 2012).

The changes in muscarinic M1 receptor positive neurons in laminae III from patients with MRDS and non-MRDS has implications for other hypotheses focussed on the pathophysiology of

schizophrenia. For example, it has been suggested that a breakdown in neuronal circuitry in laminae III of the dorsolateral frontal cortex, affecting glutamatergic and GABAergic neurotransmission, is an important component of the pathophysiology of schizophrenia (Lewis et al., 2012). The hypothesis has focussed on a breakdown in the control of the interaction of glutamatergic pyramidal cells, GABAergic and cholecystinin basket cells as well as chandelier cells in cortical laminae III in patients with schizophrenia. Therefore, the presence of muscarinic M1 receptors on pyramidal cells in laminae III of the human cortex (Figure 2C) and the loss of neurons positive for that receptor in MRDS and non-MRDS suggests that dysregulation of the cholinergic control of cortical homeostasis may also have to be factored into that model.

Abnormalities in cortical sub-cortical communications have been implicated in the pathophysiology of schizophrenia (Sheffield and Barch, 2016). It is therefore significant that pyramidal cells in human cortical laminar V are critical in cortical sub-cortical communications (Fuster, 2015). The loss of muscarinic M1 receptor positive pyramidal cells in laminae V from subjects with MRDS and non-MRDS suggests that altered cholinergic regulation of cortical sub-cortical communication could be a significant component of the pathophysiology in MRDS (Figure 2D). Finally, lower levels of primary and secondary basilar dendrites, consistent with a decrease in synaptic surface area, has been reported in either cortical laminae III only (Glantz and Lewis, 2000) or in cortical laminae III and V in patients with schizophrenia (Broadbelt et al., 2002). These data argue that the loss of muscarinic M1 receptor positive neurons is part of a complex disruption in cortical pyramidal cell function in schizophrenia.

Muscarinic M1 receptor-intermediate phenotype: Implications for cholinergic treatments for schizophrenia

The data on MRDS would seem to support the notion that studying intermediate phenotypes within the syndrome of schizophrenia will be advantageous in gaining a better understanding of the pathophysiology of the disorder (Tamminga et al., 2017). Importantly, it has also been shown that the potency of oxotremorine-M to stimulate Gq/11-[35S]-GTPγS binding using tissue from human dorsolateral prefrontal cortex was significantly decreased in the MRDS group compared to controls and non-MRDS (Salah-Uddin et al., 2009). Although oxotremorine-M is a pan-muscarinic receptor agonist (Ringdahl and Jenden, 1983), the assay conditions used in the study of schizophrenia meant only muscarinic M1 receptor G-protein recruitment would be detected (Salah-Uddin et al., 2009). Therefore, the data on oxotremorine-M mediated G-protein recruitment suggests that patients with MRDS may not respond optimally to treatment with muscarinic M1 receptor agonists like xanomeline. There was a trend toward decreased AC-42 stimulated Gq/11-[35S]-GTPγS binding using the tissue from patients with MRDS, which did not reach significance. AC-42 is a muscarinic M1 receptor allosteric agonist (Conn et al., 2009) and therefore data from the study of that drug using tissue from patients with MRDS could indicate that targeting allosteric sites on the muscarinic M1 receptor may be a more effective treatment for patients with that intermediate phenotype.

Allosteric sites on muscarinic receptors

Xanomeline is a drug which preferentially binds to the orthosteric binding site on the muscarinic M1 and M4 receptors (Bender et al., 2017). The orthosteric site is the site on the receptor activated by the endogenous ligand (Figure 3)(Gregory et al., 2010) which, in the case of the muscarinic receptors, is acetylcholine. Orthosteric sites on muscarinic receptors have been preserved through evolution and therefore have similar structures making it difficult to develop drugs specific to individual muscarinic receptors. However, allosteric binding sites, which can modulate the activity

of a muscarinic receptor but are distal to the orthosteric site and have not been preserved through evolution, have been identified for each muscarinic receptor. This means that drugs that have been synthesised to bind to the allosteric site on muscarinic receptors have very high, if not total, specificity for a single muscarinic receptor (Felder et al., 2018). Given the outcomes of treating patients with schizophrenia with muscarinic M1 / 4 receptor agonists it is now suggested that drugs that specifically modulate the allosteric site on those two receptors to alter receptor activity may provide a better therapeutic / side effect window compared to drugs acting at the orthosteric site on those receptors (Yohn and Conn, 2018). Importantly, such conclusions remain hypothetical until allosteric modulators are shown to give therapeutic benefits when treating schizophrenia.

One innovative use of human postmortem CNS is to measure ~~Given the notion of using allosteric modulators to treat schizophrenia it is of interest that the ability of the~~ ability of muscarinic M1 receptor allosteric modulators to alter the affinity of that receptor for acetylcholine using human cortex can be measured using human cortex (Dean et al., 2016a). This, a measurement is important as it ~~that~~ should be an indicator of the likely responsiveness of that receptor to allosteric modulation (Ma et al., 2009; Shirey et al., 2009). Hence, it is significant that it has been shown, using human dorsolateral prefrontal cortex, striatum and hippocampus, that patients with MRDS are less responsive to muscarinic M1 receptor allosteric activation than are non-MRDS and controls (Dean et al., 2016a; Hopper et al., 2019). It has been shown that drugs that target the muscarinic M1 receptor can differ *in vitro* and *in vivo* (Bakker et al., 2015). This raises the possibility that, unlike for findings on drug response *in vitro* (Dean et al., 2016a; Hopper et al., 2019; Salah-Uddin et al., 2009), patients with MRDS may show a good therapeutic response to drugs targeting the allosteric or orthosteric sites on muscarinic M1 and M4 receptors. However, if some patients are non-responsive, it could be because they have the MRDS intermediate phenotype. Hence, measuring levels of cortical muscarinic M1 receptors using newly available SPECT (Bakker et al.,

2015) or PET (Rowe et al., 2019) radioligands would be worthwhile to determine if there was an association between lower levels of cortical muscarinic M1 receptors and treatment response in patients with schizophrenia.

Outcomes from activating muscarinic M1 and / or M4 receptor

Initial findings from the trial of xanomeline on the treatment of schizophrenia suggested the drug was effective in lowering the severity of positive, negative and cognitive symptoms (Shekhar et al., 2008). It is thought that the underlying cause of the positive symptoms of schizophrenia is a sub-cortical hyperdopaminergic state (Abi-Dargham et al., 2000). It is therefore significant that muscarinic M4 receptor knock out mice were shown to have high levels of dopamine and dopamine efflux in the nucleus accumbens and enhanced sensitivity to dopaminergic psychostimulants (Tzavara et al., 2004). This suggests sub-cortical muscarinic M4 receptors are critical in maintaining dopaminergic homeostasis and that activating the receptor would counteract the hyperdopaminergic state in schizophrenia. Consistent with this hypothesis, pre-clinical studies with muscarinic M4 receptor positive allosteric modulators, which activate the muscarinic M4 receptor, consistently reversed dopaminergic hyperactive behaviour in rodents (Carruthers et al., 2015). Thus, it would seem reasonable to conclude that the targeting of the muscarinic M4 receptor is critical if a drug is to have antipsychotic properties. By contrast, it is less clear that targeting the muscarinic M4 receptor would reduce the negative symptoms and cognitive deficits associated with schizophrenia.

Acetylcholine has long been recognised as a key modulator of cognitive ability in humans (Perry et al., 1999). The use of a muscarinic M1 receptor allosteric agonist in a human study showed that the receptor played a critical role in modulating deficits in episodic memory in the nicotine abstinence

model of cognitive dysfunction (Nathan et al., 2013). In addition, studies have shown positive allosteric modulation can maintain cognitive function in the rat (Kucinski et al., 2020) and monkey (Lange et al., 2015). Therefore, it is reasonable to argue that the muscarinic M1 receptor is critical in maintaining cognitive functioning possibly by mechanisms that includes the actions of catechol-O-methyltransferase (Dean and Scarr, 2016; Parkin et al., 2018). Finally, the human cortex is critical in maintaining cognitive ability (Keefe and Nowakowski, 2020) and has high levels of muscarinic M1 receptor (Flynn et al., 1995) which again favours the argument that cognitive benefits will flow from targeting the muscarinic M1 receptor in the cortex.

Seemingly supporting a central role for the muscarinic M1 receptor in the cognitive deficits of schizophrenia are two studies that have reported that homozygotes at the muscarinic M1 receptor 267C/A with schizophrenia made a higher number of preservative errors compared to those who had the disorder and were heterozygotes (Liao et al., 2003; Scarr et al., 2012). Importantly, these studies also showed that genotype at 267C/A did not alter the risk for schizophrenia (Liao et al., 2003; Scarr et al., 2012) and is not associated with the levels of muscarinic M1 receptor in the human cortex (Scarr et al., 2012). By contrast, homozygotes at 267C/A muscarinic M1 receptor SNP with schizophrenia have been shown to have reduced grey matter volume in the right precentral gyrus compared to heterozygotes with the disorder (Cropley et al., 2015). The precentral gyrus is very important in motor hand control (Yousry et al., 1997) and therefore it seems likely the association between muscarinic M1 receptor 267C/A genotype and preservation was reflective of motor, rather than cognitive, abnormalities.

The biochemical underpinnings of negative symptoms have not been well understood but recent neuroimaging data have led to the suggestion that the underlying biochemistry of negative symptoms involves a breakdown in cortical dopaminergic and glutamatergic homeostasis (Howes et

al., 2015). It has also been noted that many data would support the hypothesis that a hypercholinergic state could contribute to the onset of the negative symptoms associated with schizophrenia (Tandon and Greden, 1989). This latter hypothesis makes it difficult to postulate why activating muscarinic receptors would have a benefit in the treatment of negative symptoms. However, it has been suggested that cognitive deficits and negative symptoms show correlated severity and shared features in patients with schizophrenia (Harvey et al., 2006). It could therefore be that lessening the severity of cognitive deficits by muscarinic M1 receptor activation will have a flow on effect to lessen negative symptoms in the same patient.

Concluding Remarks

The release of ~~preliminary promising~~ data showing the efficacy of KarXT as a treatment that may lessen the positive and negative symptoms of schizophrenia is one step closer to showing that activating muscarinic M1 and M4 receptors is a way of reducing the symptom burden for patients with schizophrenia (Scarr et al., 2013b). If such treatments reach the clinic, it will be the first time that the mechanisms by which drugs used to treat schizophrenia can be so fully described. However, the recent failures to show clinical efficacy of the glycine uptake inhibitor, bitopertin (Bugarski-Kirola et al., 2017), and the positive allosteric modulator of the α nicotinic receptor, AVL-3288 (Kantrowitz et al., 2020), in phase III trials after successful phase II trials shows there are still considerable hurdles to be cleared to transition muscarinic receptor activating drugs into widespread use for the treatment schizophrenia.

Notably, gene wide association studies (GWAS) on schizophrenia have failed to show a change in gene sequence at any single nucleotide polymorphism in the muscarinic M1 receptor is associated with a changed risk for schizophrenia (Schizophrenia Working Group of the Psychiatric Genomics

et al., 2014), a replication on what was reported using DNA from postmortem CNS (Scarr et al., 2009). It is proposed that in the post-GWAS error the only studies that will be able to contribute to understanding the complex aetiology of schizophrenia, and that will led to new treatments for the disorder, will be those that translate the GWAS findings into an understandable biology (Golov et al., 2020). However, the advances made in understanding the role of the muscarinic M1 receptor in the aetiology of schizophrenia and as a potential drug target would suggest a fixation on only GWAS related biology is over restrictive. As an example, recent findings on the loss of muscarinic M1 receptor positive pyramidal cells in cortical laminae III and V from patients with schizophrenia (Scarr et al., 2018a) strongly argues that the ongoing study of changes in cortical circuitry in schizophrenia (Hoftman et al., 2017) is one approach that will continue to advance our understanding of the molecular basis of that complex disorder.

It has been suggested that the diagnoses of schizophrenia defines a disorder where the efforts to develop better treatments and hence outcomes is hopeless and therefore should be discarded (Os, 2016). By contrast, studies on cortical gene expression suggest that, compared to mood disorders (Scarr et al., 2019), schizophrenia has a unique pathophysiology (Scarr et al., 2018b). In addition, data suggest that changes in cortical molecular cytoarchitecture can be used to identify distinct intermediate phenotypes within the syndrome of schizophrenia (Fillman et al., 2013; Scarr et al., 2009). This review describes the synergistic interplay between understanding the role of muscarinic receptors in the pathophysiology of schizophrenia and the development of drugs targeting those receptors to treat the illness. Such interactions would not have been possible in the absence of the diagnoses of schizophrenia. Thus, given advances been made in understanding the pathophysiology of MRDS, it may be better to continue to seek out intermediate phenotypes as an approach to better understanding the pathophysiology of what has always been recognised as a syndrome of schizophrenias (Bleuler, 1950).

|

Declaration of Interest: Nothing to declare

Figure Legends

Figure 1: The probability distribution of cortical muscarinic M1 receptor levels, measured using [³H]pirenzepine binding, in 80 patients with schizophrenia (blue line) and 74 controls (red line). The kernel density estimation of these binding data showed the patients with schizophrenia consisted of two groups and the controls one group. One group within schizophrenia consisted of 26 patients with a marked deficit in muscarinic M1 receptor (muscarinic receptor deficit schizophrenia: MRDS). By contrast, in the second group, levels of cortical muscarinic M1 receptors did not differ from controls (NON-MRDS).

Modified from Scarr et al. 2009.

Figure 2A: An example of the visualisation of muscarinic M1 receptors positive neurons, some highlighted by arrows, that are abundant in laminae III and V of the human cortex.

B: A cross section through the gray matter of the human dorsolateral prefrontal cortex showing the cortical lamina.

C: A schematic showing the circuitry proposed to be disrupted in laminae III of the cortex from subjects with schizophrenia. This model was based on, among other changes, changes in the levels of GABA receptors (GABA_A α1R and GABA_A α2R) on pyramidal cells in laminae III from patients with schizophrenia. Data showing a loss of muscarinic M1 positive pyramidal cells in patients with schizophrenia suggest cholinergic control of neuronal activity in laminae III could have a role in this model.

D: A schematic showing cholinergic innervation onto pyramidal cells in laminae V of the human cortex which are important in cortical sub-cortical communication. The loss of muscarinic M1 receptor positive pyramidal cells in cortical laminae V from subjects with schizophrenia suggests a breakdown in cholinergic homeostasis in schizophrenia

could be affecting cortical sub-cortical communication. [Figure 2A and 2B based on data published in Scarr et al. 2018a.](#)

Figure 3: A schematic showing the evolutionary conserved, and therefore structurally similar, orthosteric binding site on the muscarinic M1 and M4 receptors. The orthosteric site is where acetylcholine binds to the muscarinic M1 and M4 receptor which in turn increases signalling through Gq for the muscarinic M1 receptor and Gi for the muscarinic M4 receptor. Acetylcholine can bring about different outcomes by activating muscarinic receptors because of their different CNS localisation and signalling mechanisms. The allosteric sites on muscarinic receptors are distal to the orthosteric site, have not been preserved across evolution and are structurally different. Hence, it has been possible to design drugs that target specific allosteric targets thus such drugs show extremely high, if not total, specificity for a single muscarinic receptor.

References

- Abi-Dargham, A., Rodenhiser, J., Printz, D., Zea-Ponce, Y., Gil, R., Kegeles, L.S., Weiss, R., Cooper, T.B., Mann, J.J., Van Heertum, R.L., Gorman, J.M., Laruelle, M., 2000. Increased baseline occupancy of D2 receptors by dopamine in schizophrenia. *Proceedings of the National Academy of Sciences of the United States of America* 97 (14), 8104-8109.
- Andersen, M.B., Fink-Jensen, A., Peacock, L., Gerlach, J., Bymaster, F., Lundbaek, J.A., Werge, T., 2003. The muscarinic M1/M4 receptor agonist xanomeline exhibits antipsychotic-like activity in *Cebus apella* monkeys. *Neuropsychopharmacology* 28 (6), 1168-1175.
- Bakker, G., Vingerhoets, C., Boucherie, D., Caan, M., Bloemen, O., Eersels, J., Booij, J., van Amelsvoort, T., 2018. Relationship between muscarinic M1 receptor binding and cognition in medication-free subjects with psychosis. *Neuroimage Clin* 18, 713-719.
- Bakker, G., Vingerhoets, W.A., van Wieringen, J.P., de Bruin, K., Eersels, J., de Jong, J., Chahid, Y., Rutten, B.P., DuBois, S., Watson, M., Mogg, A.J., Xiao, H., Crabtree, M., Collier, D.A., Felder, C.C., Barth, V.N., Broad, L.M., Bloemen, O.J., van Amelsvoort, T.A., Booij, J., 2015. ¹²³I-iododexetimide preferentially binds to the muscarinic receptor subtype M1 in vivo. *Journal of Nuclear Medicine* 56 (2), 317-322.
- Bender, A.M., Jones, C.K., Lindsley, C.W., 2017. Classics in Chemical Neuroscience: Xanomeline. *ACS Chemical Neuroscience* 8 (3), 435-443.
- Bleuler, E., 1950. *Demetia Praecox or the groups of schizophrenias*. International Press, New York.

- Brann, M.R., Ellis, J., Jørgensen, H., Hill-Eubanks, D., Jones, S.V.P., 1993. Chapter 12: Muscarinic acetylcholine receptor subtypes: localization and structure/function, in: Cuello, A.C. (Ed.), *Progress in Brain Research*. Elsevier, pp. 121-127.
- Broadbelt, K., Byne, W., Jones, L.B., 2002. Evidence for a decrease in basilar dendrites of pyramidal cells in schizophrenic medial prefrontal cortex. *Schizophrenia Research* 58 (1), 75-81.
- Buchsbaum, M.S., Hazlett, E.A., 1998. Positron emission tomography studies of abnormal glucose metabolism in schizophrenia. *Schizophrenia Bulletin* 24 (3), 343-364.
- Bugarski-Kirola, D., Blaettler, T., Arango, C., Fleischhacker, W.W., Garibaldi, G., Wang, A., Dixon, M., Bressan, R.A., Nasrallah, H., Lawrie, S., Napieralski, J., Ochi-Lohmann, T., Reid, C., Marder, S.R., 2017. Bitopertin in Negative Symptoms of Schizophrenia-Results From the Phase III FlashLyte and DayLyte Studies. *Biol Psychiatry* 82 (1), 8-16.
- Bymaster, F.P., Felder, C., Ahmed, S., McKinzie, D., 2002. Muscarinic receptors as a target for drugs treating schizophrenia. *Current Drug Targets. CNS Neurological Disorders* 1 (2), 163-181.
- Caldera, M., Buphamalai, P., Müller, F., Menche, J., 2017. Interactome-based approaches to human disease. *Current Opinion in Systems Biology* 3, 88-94.
- Carruthers, S.P., Gurvich, C.T., Rossell, S.L., 2015. The muscarinic system, cognition and schizophrenia. *Neuroscience and Biobehavioral Reviews* 55, 393-402.

- Cohen, L.H., Thale, T., Tissenbaum, M.J., 1944. Acetylcholine treatment of schizophrenia. *Psychiatry Research* 51, 171-175.
- Conn, P.J., Jones, C.K., Lindsley, C.W., 2009. Subtype-selective allosteric modulators of muscarinic receptors for the treatment of CNS disorders. *TIPS* 30 (3), 148-155.
- Crook, J.M., Tomaskovic-Crook, E., Copolov, D.L., Dean, B., 2000. Decreased muscarinic receptor binding in subjects with schizophrenia: a study of the human hippocampal formation. *Biological Psychiatry* 48 (5), 381-388.
- Crook, J.M., Tomaskovic-Crook, E., Copolov, D.L., Dean, B., 2001. Low muscarinic receptor binding in prefrontal cortex from subjects with schizophrenia: a study of Brodmann's areas 8, 9, 10, and 46 and the effects of neuroleptic drug treatment. *Psychiatry Research* 158 (6), 918-925.
- Cropley, V.L., Scarr, E., Fornito, A., Bousman, C.A., Scott, R., Cairns, M., Tooney, P., The Australian Schizophrenia Research Bank, Pantelis, C., Dean, B., 2015. The effect of a muscarinic receptor 1 gene variant on grey matter volume in schizophrenia. *Psychiatry Research: Neuroimaging* In Press.
- Dean, B., Bymaster, F.P., Scarr, E., 2003. Muscarinic receptors in schizophrenia. *Current Molecular Medicine* 3 (5), 419-426.
- Dean, B., Crook, J.M., Opekin, K., Hill, C., Keks, N., Copolov, D.L., 1996. The density of muscarinic M1 receptors is decreased in the caudate-putamen of subjects with schizophrenia. *Molecular Psychiatry* 1 (1), 54-58.

- Dean, B., Crook, J.M., Pavey, G., Opeskin, K., Copolov, D.L., 2000. Muscarinic1 and 2 receptor mRNA in the human caudate-putamen: no change in m1 mRNA in schizophrenia. *Molecular Psychiatry* 5 (2), 203-207.
- Dean, B., Gray, L., Keriakous, D., Scarr, E., 2004. A comparison of M1 and M4 muscarinic receptors in the thalamus from control subjects and subjects with schizophrenia. *Thalamus and Related Systems* 2, 287-295.
- Dean, B., Hopper, S., Conn, P.J., Scarr, E., 2016a. Changes in BQCA allosteric modulation of [3H]NMS binding to human cortex within schizophrenia and by divalent cations. *Neuropsychopharmacology* 41, 1620-1628.
- Dean, B., McLeod, M., Keriakous, D., McKenzie, J., Scarr, E., 2002. Decreased muscarinic(1) receptors in the dorsolateral prefrontal cortex of subjects with schizophrenia. *Molecular Psychiatry* 7 (10), 1083-1091.
- Dean, B., Scarr, E., 2016. COMT genotype is associated with differential expression of muscarinic M1 receptors in human cortex. *Psychiatry Research* 171 (6), 784-789.
- Dean, B., Thomas, N., Lai, C.Y., Chen, W.J., Scarr, E., 2015. Changes in cholinergic and glutamatergic markers in the striatum from a sub-set of subjects with schizophrenia. *Schizophrenia Research* 169 (1-3), 83-88.
- Dean, B., Thomas, N., Scarr, E., Udawela, M., 2016b. Evidence for impaired glucose metabolism in the striatum, obtained postmortem, from some subjects with schizophrenia. *Translational Psychiatry* 6 (11), e949.

- Defelipe, J., 2011. The evolution of the brain, the human nature of cortical circuits, and intellectual creativity. *Frontiers in Neuroanatomy* 5, 29-29.
- Felder, C.C., Goldsmith, P.J., Jackson, K., Sanger, H.E., Evans, D.A., Mogg, A.J., Broad, L.M., 2018. Current status of muscarinic M1 and M4 receptors as drug targets for neurodegenerative diseases. *Neuropharmacology* 136, 449-458.
- Felder, C.C., Porter, A.C., Skillman, T.L., Zhang, L., Bymaster, F.P., Nathanson, N.M., Hamilton, S.E., Gomeza, J., Wess, J., McKinzie, D.L., 2001. Elucidating the role of muscarinic receptors in psychosis. *Life Sciences* 68 (22-23), 2605-2613.
- Fillman, S.G., Cloonan, N., Catts, S.V., Miller, L.C., Wong, J., McCrossin, T., Cairns, M., Weickert, C.S., 2013. Increased inflammatory markers identified in the dorsolateral prefrontal cortex of individuals with schizophrenia. *Molecular Psychiatry* 18 (2), 206-214.
- Flynn, D.D., Ferrari-Dileo, G., Mash, D.C., Levey, A.I., 1995. Differential regulation of molecular subtypes of muscarinic receptors in Alzheimer's disease. *Journal of Neurochemistry* 64 (4), 1888-1891.
- Forrer, G.R., 1951. Atropine toxicity in the treatment of mental disease. *Psychiatry Research* 108 (2), 107-112.
- Fuster, J.M., 2015. *The Prefrontal Cortex*, 5 ed. Academic Press, New York.

- Gibbons, A., Scarr, E., Boer, S., Money, T., Jeon, W.J., Felder, C., Dean, B., 2013. Widespread decreases in cortical muscarinic receptors in a subset of people with schizophrenia. *International Journal of Neuropsychopharmacology* 16, 37-46.
- Gibbons, A.S., Jeon, W.J., Scarr, E., Dean, B., 2016. Changes in Muscarinic M2 Receptor Levels in the Cortex of Subjects with Bipolar Disorder and Major Depressive Disorder and in Rats after Treatment with Mood Stabilisers and Antidepressants. *International Journal of Neuropsychopharmacology* 19.
- Girgis, R.R., Zoghbi, A.W., Javitt, D.C., Lieberman, J.A., 2019. The past and future of novel, non-dopamine-2 receptor therapeutics for schizophrenia: A critical and comprehensive review. *Journal of Psychiatric Research* 108, 57-83.
- Glantz, L.A., Lewis, D.A., 2000. Decreased dendritic spine density on prefrontal cortical pyramidal neurons in schizophrenia. *Psychiatry Research* 57 (1), 65-73.
- Golov, A.K., Kondratyev, N.V., Kostyuk, G.P., Golimbet, V.E., 2020. Novel Approaches for Identifying the Molecular Background of Schizophrenia. *Cells* 9 (1), 246.
- Gregory, K.J., Sexton, P.M., Christopoulos, A., 2010. Overview of Receptor Allostereism. *Current Protocols in Pharmacology* 51 (1), 1.21.21-21.21.34.
- Harvey, P.D., Koren, D., Reichenberg, A., Bowie, C.R., 2006. Negative symptoms and cognitive deficits: what is the nature of their relationship? *Schizophrenia Bulletin* 32 (2), 250-258.

- Hoftman, G.D., Datta, D., Lewis, D.A., 2017. Layer 3 Excitatory and Inhibitory Circuitry in the Prefrontal Cortex: Developmental Trajectories and Alterations in Schizophrenia. *Biological Psychiatry* 81 (10), 862-873.
- Hopper, S., Pavey, G.M., Gogos, A., Dean, B., 2019. Widespread Changes in Positive Allosteric Modulation of the Muscarinic M1 Receptor in Some Participants With Schizophrenia. *International Journal of Neuropsychopharmacology* 22 (10), 640-650.
- Horga, G., Parellada, E., Lomena, F., Fernandez-Egea, E., Mane, A., Font, M., Falcon, C., Konova, A.B., Pavia, J., Ros, D., Bernardo, M., 2011. Differential brain glucose metabolic patterns in antipsychotic-naïve first-episode schizophrenia with and without auditory verbal hallucinations. *Journal of Psychiatry and Neuroscience* 36 (5), 312-321.
- Howes, O., McCutcheon, R., Stone, J., 2015. Glutamate and dopamine in schizophrenia: an update for the 21st century. *Journal of Psychopharmacology (Oxford, England)* 29 (2), 97-115.
- Kantrowitz, J.T., Javitt, D.C., Freedman, R., Sehatpour, P., Kegeles, L.S., Carlson, M., Sobeih, T., Wall, M.M., Choo, T.-H., Vail, B., Grinband, J., Lieberman, J.A., 2020. Double blind, two dose, randomized, placebo-controlled, cross-over clinical trial of the positive allosteric modulator at the $\alpha 7$ nicotinic cholinergic receptor AVL-3288 in schizophrenia patients. *Neuropsychopharmacology*.
- Keefe, M.G., Nowakowski, T.J., 2020. Evolutionary Expansion of Human Cerebellar Germinal Zones. *Trends in Neurosciences* 43 (2), 75-77.

- Kucinski, A., Phillips, K.B., Koshy Cherian, A., Sarter, M., 2020. Rescuing the attentional performance of rats with cholinergic losses by the M1 positive allosteric modulator TAK-071. *Psychopharmacology* 237 (1), 137-153.
- Lange, H.S., Cannon, C.E., Drott, J.T., Kuduk, S.D., Uslaner, J.M., 2015. The M1 Muscarinic Positive Allosteric Modulator PQCA Improves Performance on Translatable Tests of Memory and Attention in Rhesus Monkeys. *Journal of Pharmacology and Experimental Therapeutics* 355 (3), 442-450.
- Lewis, D.A., Curley, A.A., Glausier, J.R., Volk, D.W., 2012. Cortical parvalbumin interneurons and cognitive dysfunction in schizophrenia. *Trends in Neurosciences* 35 (1), 57-67.
- Liao, D.L., Hong, C.J., Chen, H.M., Chen, Y.E., Lee, S.M., Chang, C.Y., Chen, H., Tsai, S.J., 2003. Association of muscarinic m1 receptor genetic polymorphisms with psychiatric symptoms and cognitive function in schizophrenic patients. *Neuropsychobiology* 48 (2), 72-76.
- Luo, W., Semenza, G.L., 2011. Pyruvate kinase M2 regulates glucose metabolism by functioning as a coactivator for hypoxia-inducible factor 1 in cancer cells. *Oncotarget* 2 (7), 551-556.
- Ma, L., Seager, M.A., Wittmann, M., Jacobson, M., Bickel, D., Burno, M., Jones, K., Graufelds, V.K., Xu, G., Pearson, M., McCampbell, A., Gaspar, R., Shughrue, P., Danziger, A., Regan, C., Flick, R., Pascarella, D., Garson, S., Doran, S., Kreatsoulas, C., Veng, L., Lindsley, C.W., Shipe, W., Kuduk, S., Sur, C., Kinney, G., Seabrook, G.R., Ray, W.J., 2009. Selective activation of the M1 muscarinic acetylcholine receptor achieved by allosteric potentiation. *Proceedings of the National Academy of Sciences of the United States of America* 106 (37), 15950-15955.

- Mancama, D., Arranz, M.J., Landau, S., Kerwin, R., 2003. Reduced expression of the muscarinic 1 receptor cortical subtype in schizophrenia. *Psychiatry Research* 119 (1), 2-6.
- McOmish, C., Pavey, G., McLean, C., Horne, M., Dean, B., Scarr, E., 2017. Muscarinic receptor binding changes in postmortem Parkinson's disease. *Journal of Neural Transmission* 124 (2), 227-236.
- Miller, A., Kavoussi, R., Breier, A., 2016. Xanomeline Plus Trospium: A Novel Strategy to Enhance Pro-Muscarinic Efficacy and Mitigate Peripheral Side Effects. *Neuropsychopharmacology* 41, S230.
- Minzenberg, M.J., Poole, J.H., Benton, C., Vinogradov, S., 2004. Association of anticholinergic load with impairment of complex attention and memory in schizophrenia. *Psychiatry Research* 161 (1), 116-124.
- Moghaddam, B., Javitt, D., 2012. From revolution to evolution: the glutamate hypothesis of schizophrenia and its implication for treatment. *Neuropsychopharmacology* 37 (1), 4-15.
- Nathan, P.J., Watson, J., Lund, J., Davies, C.H., Peters, G., Dodds, C.M., Swirski, B., Lawrence, P., Bentley, G.D., O'Neill, B.V., Robertson, J., Watson, S., Jones, G.A., Maruff, P., Croft, R.J., Laruelle, M., Bullmore, E.T., 2013. The potent M(1) receptor allosteric agonist GSK1034702 improves episodic memory in humans in the nicotine abstinence model of cognitive dysfunction. *International Journal of Neuropsychopharmacology* 16 (4), 721-731.

- Ogino, S., Miyamoto, S., Miyake, N., Yamaguchi, N., 2014. Benefits and limits of anticholinergic use in schizophrenia: focusing on its effect on cognitive function. *Psychiatry and Clinical Neurosciences* 68 (1), 37-49.
- Os, J.v., 2016. "Schizophrenia" does not exist. *BMJ* 352, i375.
- Parkin, G.M., Udawela, M., Gibbons, A., Scarr, E., Dean, B., 2018. Catechol-O-methyltransferase (COMT) genotypes are associated with varying soluble, but not membrane-bound COMT protein in the human prefrontal cortex. *Journal of Human Genetics* 63, 1251-1258.
- Perry, E., Walker, M., Grace, J., Perry, R., 1999. Acetylcholine in mind: a neurotransmitter correlate of consciousness? *Trends in Neuroscience* 22 (6), 273-280.
- Raedler, T.J., Knable, M.B., Jones, D.W., Urbina, R.A., Gorey, J.G., Lee, K.S., Egan, M.F., Coppola, R., Weinberger, D.R., 2003. In vivo determination of muscarinic acetylcholine receptor availability in schizophrenia. *Psychiatry Research* 160 (1), 118-127.
- Ringdahl, B., Jenden, D.J., 1983. Pharmacological properties of oxotremorine and its analogs. *Life Sciences* 32 (21), 2401-2413.
- Rowe, C., Dean, B., Ackermann, U., Goh, R., Guzman, R., Kanaan, R., Chong, L., Dore, V., Bozinovski, S., Masters, C., Villemagne, V., 2019. In vivo imaging of brain muscarinic receptors with 18F-Fluorobenzyl Dexetimide: First in human studies. *Journal of Nuclear Medicine* 60 (supplement 1), 1470.

- Salah-Uddin, H., Scarr, E., Pavey, G., Harris, K., Hagan, J.J., Dean, B., Challiss, R.A., Watson, J.M., 2009. Altered M(1) muscarinic acetylcholine receptor (CHRM1)-G α (q/11) coupling in a schizophrenia endophenotype. *Neuropsychopharmacology* 34 (9), 2156-2166.
- Sanz, B., Exposito, I., Mora, F., 1997. M1 acetylcholine receptor stimulation increases the extracellular concentrations of glutamate and GABA in the medial prefrontal cortex of the rat. *Neurochemical Research* 22 (3), 281-286.
- Scarr, E., Cowie, T.F., Kanellakis, S., Sundram, S., Pantelis, C., Dean, B., 2009. Decreased cortical muscarinic receptors define a subgroup of subjects with schizophrenia. *Molecular Psychiatry* 14 (11), 1017-1023.
- Scarr, E., Craig, J.M., Cairns, M.J., Seo, M.S., Galati, J.C., Beveridge, N.J., Gibbons, A., Juzva, S., Weinrich, B., Parkinson-Bates, M., Carroll, A.P., Saffery, R., Dean, B., 2013a. Decreased cortical muscarinic M1 receptors in schizophrenia are associated with changes in gene promoter methylation, mRNA and gene targeting microRNA. *Translational Psychiatry* 3, e230.
- Scarr, E., Dean, B., 2008. Muscarinic receptors: do they have a role in the pathology and treatment of schizophrenia? *Journal of Neurochemistry* 107 (5), 1188-1195.
- Scarr, E., Gibbons, A.S., Neo, J., Udawela, M., Dean, B., 2013b. Cholinergic connectivity: it's implications for psychiatric disorders. *Front Cell Neurosci.* 7, 55.
- Scarr, E., Hopper, S., Vos, V., Seo, M.S., Everall, I.P., Aumann, T.D., Chunam, G., Dean, B., 2018a. Low levels of muscarinic M1 receptor positive neurons in cortical layers III and V in

Brodmann's areas 9 and 17 from individuals with schizophrenia. *Journal of Psychiatry and Neuroscience* 43, 338-346.

Scarr, E., Keriakous, D., Crossland, N., Dean, B., 2006. No change in cortical muscarinic M2, M3 receptors or [35S]GTPgammaS binding in schizophrenia. *Life Sciences* 78 (11), 1231-1237.

Scarr, E., McLean, C., Dean, B., 2017. Higher levels of different muscarinic receptors in the cortex and hippocampus from subjects with Alzheimer's disease. *Journal of Neural Transmission* 124 (3), 273-284.

Scarr, E., Sundram, S., Deljo, A., Cowie, T.F., Gibbons, A.S., Juzva, S., Mackinnon, A., Wood, S.J., Testa, R., Pantelis, C., Dean, B., 2012. Muscarinic M1 receptor sequence: preliminary studies on its effects on cognition and expression. *Schizophrenia Research* 138 (1), 94-98.

Scarr, E., Sundram, S., Keriakous, D., Dean, B., 2007. Altered hippocampal muscarinic m4, but not m1, receptor expression from subjects with schizophrenia. *Biological Psychiatry* 61 (10), 1161-1170.

Scarr, E., Udawela, M., Dean, B., 2018b. Changed frontal pole gene expression suggest altered interplay between neurotransmitter, developmental, and inflammatory pathways in schizophrenia. *NPJ Schizophrenia* 4 (1), 4.

Scarr, E., Udawela, M., Dean, B., 2019. Changed cortical risk gene expression in major depression and shared changes in cortical gene expression between major depression and bipolar disorders. *Australian and New Zealand Journal of Psychiatry* 53, 1189-1198.

Scarr, E., Udawela, M., Thomas, E.A., Dean, B., 2018c. Changed gene expression in subjects with schizophrenia and low cortical muscarinic M1 receptors predicts disrupted upstream pathways interacting with that receptor. *Molecular Psychiatry* 23, 295-303.

Schizophrenia Working Group of the Psychiatric Genomics, C., Ripke, S., Neale, B.M., Corvin, A., Walters, J.T.R., Farh, K.-H., Holmans, P.A., Lee, P., Bulik-Sullivan, B., Collier, D.A., Huang, H., Pers, T.H., Agartz, I., Agerbo, E., Albus, M., Alexander, M., Amin, F., Bacanu, S.A., Begemann, M., Belliveau, R.A., Bene, J., Bergen, S.E., Bevilacqua, E., Bigdeli, T.B., Black, D.W., Bruggeman, R., Buccola, N.G., Buckner, R.L., Byerley, W., Cahn, W., Cai, G., Campion, D., Cantor, R.M., Carr, V.J., Carrera, N., Catts, S.V., Chambert, K.D., Chan, R.C.K., Chan, R.Y.L., Chen, E.Y.H., Cheng, W., Cheung, E.F.C., Chong, S.A., Cloninger, C.R., Cohen, D., Cohen, N., Cormican, P., Craddock, N., Crowley, J.J., Curtis, D., Davidson, M., Davis, K.L., Degenhardt, F., Del Favero, J., Demontis, D., Dikeos, D., Dinan, T., Djurovic, S., Donohoe, G., Drapeau, E., Duan, J., Dudbridge, F., Durmishi, N., Eichhammer, P., Eriksson, J., Escott-Price, V., Essioux, L., Fanous, A.H., Farrell, M.S., Frank, J., Franke, L., Freedman, R., Freimer, N.B., Friedl, M., Friedman, J.I., Fromer, M., Genovese, G., Georgieva, L., Giegling, I., Giusti-Rodríguez, P., Godard, S., Goldstein, J.I., Golimbet, V., Gopal, S., Gratten, J., de Haan, L., Hammer, C., Hamshere, M.L., Hansen, M., Hansen, T., Haroutunian, V., Hartmann, A.M., Henskens, F.A., Herms, S., Hirschhorn, J.N., Hoffmann, P., Hofman, A., Hollegaard, M.V., Hougaard, D.M., Ikeda, M., Joa, I., Julià, A., Kahn, R.S., Kalaydjieva, L., Karachanak-Yankova, S., Karjalainen, J., Kavanagh, D., Keller, M.C., Kennedy, J.L., Khrunin, A., Kim, Y., Klovins, J., Knowles, J.A., Konte, B., Kucinkas, V., Kucinskiene, Z.A., Kuzelova-Ptackova, H., Kähler, A.K., Laurent, C., Lee, J., Lee, S.H., Legge, S.E., Lerer, B., Li, M., Li, T., Liang, K.-Y., Lieberman, J., Limborska, S., Loughland, C.M., Lubinski, J., Lönngqvist, J., Macek, M., Magnusson,

P.K.E., Maher, B.S., Maier, W., Mallet, J., Marsal, S., Mattheisen, M., Mattingdal, M.,
 McCarley, R.W., McDonald, C., McIntosh, A.M., Meier, S., Meijer, C.J., Meleghe, B.,
 Melle, I., Meshulam-Gately, R.I., Metspalu, A., Michie, P.T., Milani, L., Milanova, V.,
 Mokrab, Y., Morris, D.W., Mors, O., Murphy, K.C., Murray, R.M., Myin-Germeys, I.,
 Müller-Myhsok, B., Nelis, M., Nenadic, I., Nertney, D.A., Nestadt, G., Nicodemus, K.K.,
 Nikitina-Zake, L., Nisenbaum, L., Nordin, A., O'Callaghan, E., O'Dushlaine, C., O'Neill,
 F.A., Oh, S.-Y., Olincy, A., Olsen, L., Van Os, J., Psychosis Endophenotypes International,
 C., Pantelis, C., Papadimitriou, G.N., Papiol, S., Parkhomenko, E., Pato, M.T., Paunio, T.,
 Pejovic-Milovancevic, M., Perkins, D.O., Pietiläinen, O., Pimm, J., Pocklington, A.J.,
 Powell, J., Price, A., Pulver, A.E., Purcell, S.M., Quesada, D., Rasmussen, H.B.,
 Reichenberg, A., Reimers, M.A., Richards, A.L., Roffman, J.L., Roussos, P., Ruderfer,
 D.M., Salomaa, V., Sanders, A.R., Schall, U., Schubert, C.R., Schulze, T.G., Schwab, S.G.,
 Scolnick, E.M., Scott, R.J., Seidman, L.J., Shi, J., Sigurdsson, E., Silagadze, T., Silverman,
 J.M., Sim, K., Slominsky, P., Smoller, J.W., So, H.-C., Spencer, C.C.A., Stahl, E.A.,
 Stefansson, H., Steinberg, S., Stogmann, E., Straub, R.E., Strengman, E., Strohmaier, J.,
 Stroup, T.S., Subramaniam, M., Suvisaari, J., Svrakic, D.M., Szatkiewicz, J.P., Söderman,
 E., Thirumalai, S., Toncheva, D., Tosato, S., Veijola, J., Waddington, J., Walsh, D., Wang,
 D., Wang, Q., Webb, B.T., Weiser, M., Wildenauer, D.B., Williams, N.M., Williams, S.,
 Witt, S.H., Wolen, A.R., Wong, E.H.M., Wormley, B.K., Xi, H.S., Zai, C.C., Zheng, X.,
 Zimprich, F., Wray, N.R., Stefansson, K., Visscher, P.M., Wellcome Trust Case-Control, C.,
 Adolfsson, R., Andreassen, O.A., Blackwood, D.H.R., Bramon, E., Buxbaum, J.D.,
 Børglum, A.D., Cichon, S., Darvasi, A., Domenici, E., Ehrenreich, H., Esko, T., Gejman,
 P.V., Gill, M., Gurling, H., Hultman, C.M., Iwata, N., Jablensky, A.V., Jönsson, E.G.,
 Kendler, K.S., Kirov, G., Knight, J., Lencz, T., Levinson, D.F., Li, Q.S., Liu, J., Malhotra,
 A.K., McCarroll, S.A., McQuillin, A., Moran, J.L., Mortensen, P.B., Mowry, B.J., Nöthen,

M.M., Ophoff, R.A., Owen, M.J., Palotie, A., Pato, C.N., Petryshen, T.L., Posthuma, D., Rietschel, M., Riley, B.P., Rujescu, D., Sham, P.C., Sklar, P., St Clair, D., Weinberger, D.R., Wendland, J.R., Werge, T., Daly, M.J., Sullivan, P.F., O'Donovan, M.C., 2014. Biological Insights From 108 Schizophrenia-Associated Genetic Loci. *Nature* 511 (7510), 421-427.

Seo, M.S., Scarr, E., Dean, B., 2014. An investigation of the factors that regulate muscarinic receptor expression in schizophrenia. *Schizophrenia Research* 158 (1-3), 247-254.

Shannon, H.E., Bymaster, F.P., Calligaro, D.O., Greenwood, B., Mitch, C.H., Sawyer, B.D., Ward, J.S., Wong, D.T., Olesen, P.H., Sheardown, M.J., 1994. Xanomeline: a novel muscarinic receptor agonist with functional selectivity for M1 receptors. *Journal of Pharmacology and Experimental Therapeutics* 269 (1), 271-281.

Shannon, H.E., Rasmussen, K., Bymaster, F.P., Hart, J.C., Peters, S.C., Swedberg, M.D., Jeppesen, L., Sheardown, M.J., Sauerberg, P., Fink-Jensen, A., 2000. Xanomeline, an M(1)/M(4) preferring muscarinic cholinergic receptor agonist, produces antipsychotic-like activity in rats and mice. *Schizophrenia Research* 42 (3), 249-259.

Sheffield, J.M., Barch, D.M., 2016. Cognition and resting-state functional connectivity in schizophrenia. *Neuroscience and Biobehavioral Reviews* 61, 108-120.

Shekhar, A., Potter, W.Z., Lightfoot, J., Lienemann, J., Dube, S., Mallinckrodt, C., Bymaster, F.P., McKinzie, D.L., Felder, C.C., 2008. Selective muscarinic receptor agonist xanomeline as a novel treatment approach for schizophrenia. *American Journal of Psychiatry* 165 (8), 1033-1039.

- Shirey, J.K., Brady, A.E., Jones, P.J., Davis, A.A., Bridges, T.M., Kennedy, J.P., Jadhav, S.B., Menon, U.N., Xiang, Z., Watson, M.L., Christian, E.P., Doherty, J.J., Quirk, M.C., Snyder, D.H., Lah, J.J., Levey, A.I., Nicolle, M.M., Lindsley, C.W., Conn, P.J., 2009. A selective allosteric potentiator of the M1 muscarinic acetylcholine receptor increases activity of medial prefrontal cortical neurons and restores impairments in reversal learning. *Journal of Neuroscience* 29 (45), 14271-14286.
- Staskin, D., Kay, G., Tannenbaum, C., Goldman, H.B., Bhashi, K., Ling, J., Oefelein, M.G., 2010. Trospium chloride has no effect on memory testing and is assay undetectable in the central nervous system of older patients with overactive bladder. *International Journal of Clinical Practice* 64 (9), 1294-1300.
- Sullivan, R.J., Allen, J.S., Otto, C., Tiobech, J., Nero, K., 2000. Effects of chewing betel nut (*Areca catechu*) on the symptoms of people with schizophrenia in Palau, Micronesia. *Br.J.Psychiatry* 177, 174-178.
- Sullivan, R.J., Andres, S., Otto, C., Miles, W., Kydd, R., 2007. The effects of an indigenous muscarinic drug, Betel nut (*Areca catechu*), on the symptoms of schizophrenia: a longitudinal study in Palau, Micronesia. *Psychiatry Research* 164 (4), 670-673.
- Tamminga, C.A., Pearlson, G.D., Stan, A.D., Gibbons, R.D., Padmanabhan, J., Keshavan, M., Clementz, B.A., 2017. Strategies for Advancing Disease Definition Using Biomarkers and Genetics: The Bipolar and Schizophrenia Network for Intermediate Phenotypes. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging* 2 (1), 20-27.

- Tandon, R., Greden, J.F., 1989. Cholinergic hyperactivity and negative schizophrenic symptoms. A model of cholinergic/dopaminergic interactions in schizophrenia. *Psychiatry Research* 46 (8), 745-753.
- Tzavara, E.T., Bymaster, F.P., Davis, R.J., Wade, M.R., Perry, K.W., Wess, J., McKinzie, D.L., Felder, C., Nomikos, G.G., 2004. M4 muscarinic receptors regulate the dynamics of cholinergic and dopaminergic neurotransmission: relevance to the pathophysiology and treatment of related CNS pathologies. *FASEB Journal* 18 (12), 1410-1412.
- Yohn, S.E., Conn, P.J., 2018. Positive allosteric modulation of M1 and M4 muscarinic receptors as potential therapeutic treatments for schizophrenia. *Neuropharmacology* 136, 438-448.
- Yousry, T.A., Schmid, U.D., Alkadhi, H., Schmidt, D., Peraud, A., Buettner, A., Winkler, P., 1997. Localization of the motor hand area to a knob on the precentral gyrus. A new landmark. *Brain* 120 (1), 141-157.
- Zavitsanou, K., Katsifis, A., Filomena, M., Xu-Feng, H., 2004. Investigation of M1/M4 muscarinic receptors in the anterior cingulate cortex in schizophrenia, bipolar disorder, and major depression disorder. *Neuropsychopharmacology* 29, 619-625.

A **B** **C** **D**

ACETYLCHOLINE

UNIQUE ALLOSTERIC
BINDING SITE

ORTHOSTERIC
BINDING SITE

UNIQUE ALLOSTERIC
BINDING SITE

Conflicts of Interest

The authors have no conflicts of interest.

Author Statement

The authors conceptualised, wrote and reviewed the manuscript that has been submitted.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Dean, B;Scarr, E

Title:

Muscarinic M1 and M4 receptors: Hypothesis driven drug development for schizophrenia

Date:

2020-06-01

Citation:

Dean, B. & Scarr, E. (2020). Muscarinic M1 and M4 receptors: Hypothesis driven drug development for schizophrenia. PSYCHIATRY RESEARCH, 288, <https://doi.org/10.1016/j.psychres.2020.112989>.

Persistent Link:

<http://hdl.handle.net/11343/241390>