- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19

2 Andrew D. King^{1,2}, Andy J. Pitman^{2,3}, Benjamin J. Henley^{1,2,4}, Anna M. Ukkola^{2,5}, Josephine
3 R. Brown^{1,2}

4

5

6
7

8
9

10
11

20

Corresponding author: Andrew King (andrew.king@unimelb.edu.au)

21 **Summary**

22 Much of Australia has been in severe drought since at least 2017. Here we link Australian
23 droughts to the absence of Pacific and Indian Ocean mode states which act as key drivers of
24 drought breaking rains. Predicting the impact of climate change on drought requires accurate
25 modelling of these modes of variability.

26 **Main**

27 Australia has an exceptionally variable climate¹ with periods of drought punctuated by heavy
28 rainfall events that can cause widespread flooding. This prevalence of floods and droughts is
29 likely to have been typical of Australia's long-term climate² and is relatively common in
30 instrumental observations over the past century or so. Droughts ranging from a year or two to
31 more than a decade in length (e.g., the Millennium Drought of 1997-2009) have tended to be
32 broken by widespread heavy rains (such as in 2010-2012, when rain also brought devastating
33 floods to Queensland) rather than simply a return to average rainfall conditions. Recently,
34 there has been considerable discussion around the causes of the ongoing drought in
35 southeastern Australia, including the possible influence of human-induced climate change.
36 Here we contend that the primary reason for the long-lasting dry conditions is a recent lack of
37 La Niña and negative Indian Ocean Dipole (IOD) events in the Pacific and Indian Oceans,
38 respectively.

39 **Unprecedented multi-year drought**

40 Large swaths of Australia, and in particular the Murray-Darling Basin (MDB) region of
41 southeastern Australia, have been in drought since 2017. The MDB is a crucial agricultural
42 region, sometimes described as the 'food bowl' of the country, and drought in this region has
43 consequences for food and cotton prices as well as cascading effects on the regional and
44 national economy. In addition, dry conditions can exacerbate risk of bushfires in terms of

45 how likely they are to occur and how difficult they are to control. Bushfires in eastern
46 Australia in spring 2019, which have continued into the summer, have been widespread and
47 intense due to the dryness of the fuel load in association with ongoing drought.

48 Drought may be analysed using a multitude of different methods and indices. In Australia, the
49 Bureau of Meteorology measures drought based on rainfall deficiencies across different
50 temporal scales (<http://www.bom.gov.au/climate/glossary/drought.shtml>). Here we focus on
51 rainfall, acknowledging that other variables influence the nature of drought.

52 The winter of 2016, which preceded the current drought, was particularly wet in southeast
53 Australia and ranked as the fourth wettest for the MDB since 1900, the start of the
54 instrumental record. This was associated with a negative IOD event, whereby anomalously
55 warm sea surface temperatures (SSTs) in the east Indian Ocean gave rise to dynamic and
56 thermodynamic conditions conducive to widespread rainfall³. Since then there have been
57 unusually dry conditions in the MDB with 12 consecutive seasons of below-average rainfall,
58 the longest such period since 1900 (Fig. 1a). Specifically, the occurrence of three consecutive
59 dry winters since 2017, (all in the lowest decile of observed values) is unprecedented since
60 1900. This has had severe impacts on crop harvests and water security, with several major
61 regional centres on the brink of running out of water.

62 Looking through the instrumental record we observe other periods of persistently dry
63 conditions, such as during the Federation, Second World War, and Millennium Droughts
64 (Fig. 1a). Palaeoclimate reconstructions suggest that other multi-year droughts have occurred
65 in previous centuries, but also that recent drying in southern Australia is unusual in a multi-
66 century context². Rainfall in the MDB is marked by very high variability, and there is no
67 statistically significant trend towards higher or lower annual rainfall totals since 1900⁴.

68 **Drivers of MDB rainfall variability**

69 Short-term rainfall variability in the MDB is related to the Southern Annular Mode (SAM),
70 the Sub-Tropical Ridge (STR) and the passing of individual weather systems (Fig. 1b). On
71 the interannual timescales relevant in particular to the ongoing MDB drought, ENSO and the
72 IOD are well-established drivers⁵. The IOD modulates MDB rainfall during austral winter
73 and early spring, while ENSO plays a strong role during austral spring^{6,7}. There are highly
74 significant correlations between sea surface temperature indices representing ENSO and the
75 IOD and seasonal MDB rainfall anomalies (Fig. 2a,b). It is also noteworthy that the
76 relationships between these climate modes and MDB rainfall are not always linear. On
77 average, rainfall totals in El Niño and ENSO-neutral seasons are similar, whereas La Niña
78 seasons are often substantially wetter (Fig. 2a). Whilst this was known previously⁸, it is
79 highly relevant in the context of drought – drought-breaking rainfall is considerably more
80 likely to occur during a La Niña season than either an El Niño or ENSO-neutral season. The
81 IOD-rainfall relationship is almost linear (Fig. 2b), but the probability of high rainfall totals is
82 substantially greater during negative IOD events than in either neutral or positive IOD
83 conditions for the MDB.

84 Without a La Niña or a negative IOD event, it is difficult for wet conditions to occur on a
85 large spatial scale across a season in the MDB or much of eastern Australia. This is
86 particularly clear in spring, when seven of the ten wettest seasons in the MDB since 1900
87 occurred during strong La Niña or negative IOD conditions, and the three remaining events
88 fall just short of being categorised as La Niña.

89 These results suggest that the Australian drought should be examined through a different lens.
90 Rather than considering only the causes of the dry conditions, analysis of the drought should
91 include a focus on the lack of wet conditions that break it.

92 **Dynamics that bring rain**

93 Considering the accumulated rainfall anomaly since a La Niña or negative IOD event has
94 occurred (Fig. 2c), we find that typically, the more time that has passed since a La Niña or
95 negative IOD event, the greater the rainfall deficits over the MDB. To the end of 2019, the
96 accumulated MDB rainfall deficit is over 450 mm; the greatest running deficit since a La
97 Niña or negative IOD event. In the instrumental record, when six months have passed since a
98 La Niña or negative IOD event, 67% of the time (32/48 cases) there was an accumulated
99 rainfall deficit in the MDB. Similarly, for events that were 12, 18 and 24 months after a La
100 Niña or negative IOD event, the MDB experienced an accumulated rainfall deficit 76%
101 (25/33), 81% (22/27) and 90% (19/21) of the time, respectively.

102 Major multi-year drought events are difficult to reliably contextualise using the instrumental
103 record since 1900. Nevertheless, the major Australian droughts of the last 100 years have
104 coincided with several of the longer-lasting periods when La Niña and negative IOD events
105 did not occur. The Second World War Drought from 1935-1945 includes two unusually long
106 periods when neither a La Niña nor a negative IOD event occurred from 1934-1938 and
107 1939-1942. Indeed, the 1938-39 La Niña brought temporary relief with wetter-than-average
108 conditions in the early part of 1939 in the MDB. The Millennium Drought from 1997-2009
109 included periods from 2001-2004 and 2004-2007 without a La Niña or negative IOD event
110 when rainfall deficits substantially worsened.

111 While there is a general pattern towards rainfall deficits accumulating between the La Niña
112 and negative IOD events that increase the probability of widespread heavy rainfall, there are
113 exceptions to this pattern. For example, two long time windows without either La Niña or
114 negative IOD events in the 20th century were associated with small rainfall surpluses in the
115 MDB during the early 1960s and mid-1990s (blue lines in Fig. 2c). The wettest months in
116 these outlier series occurred in summer and were associated with positive SAM conditions,
117 whereby the mid-latitude storm track is unusually far south and there are more frequent moist

easterly winds over southeast Australia related to atmospheric blocking in the Tasman Sea^{9,10}.

There are other occasions when positive SAM conditions fail to produce large-scale rainfall events in the MDB due to the seasonality of the SAM influence and its interaction with other climate modes⁶. Thus, the modes of variability do not provide a definitive guide to rainfall over the MDB due to natural variability and synoptic-scale weather systems. While the length of time since a La Niña or negative IOD event does not necessarily imply drought, it does increase the probability of drought due to the reduced chance of widespread rain.

Focus on interannual variability

Whilst Australia is no stranger to drought, the southeast of the continent is experiencing a particularly severe event at the time of writing. Droughts in Australia have wide-ranging impacts from large economic costs¹¹ to increased mental health disorders¹² and suicide rates¹³. While individual farms may experience some drought relief due to local storm events, southeast Australia is in urgent need of widespread heavy seasonal rainfall. Our analysis suggests that the probability of widespread drought-breaking rains is associated with La Niña or negative IOD events.

To understand the risks of human-induced climate change on the security of Australia's water supplies, its agriculture, the prevalence of droughts, and the intensity of bushfires will therefore require accurate projections of the future frequency of La Niña and negative IOD events in a warming world. These events may be increasing as the world warms^{14,15}, but there remains uncertainty in projections due to model deficiencies. In particular, climate models overstate the amplitude of the IOD¹⁶ and struggle with the spatial and temporal extent of La Niña¹⁷. This leaves Australia in a challenging position in terms of how to prioritise climate adaptation investment because it is unclear how the IOD and ENSO will change in the future and whether these changes will exacerbate drought conditions. Clearly, a continent as

vulnerable to drought as Australia has a vested interest in accelerating research that addresses the present and future role of the IOD and La Niña in explaining large-scale rainfall.

Acknowledgements

We thank the Bureau of Meteorology, the Bureau of Rural Sciences and CSIRO for providing the Australian Water Availability Project data. Several authors received funding through the Australian Research Council: A.D.K. (DE180100638), and A.J.P. and A.M.U. (CE170100023). B.J.H. received funding from an Australian Research Council Linkage Project (LP150100062), which was co-funded by Melbourne Water and the Victorian Department of Environment, Land, Water and Planning, and supported by the Australian Bureau of Meteorology.

Author contributions

A.D.K. conceived the study and performed the analysis. All authors contributed to the methodological design and the writing of the paper.

Additional Information

Supplementary information is available for this paper.

References

1. Nicholls, N., Drosowsky, W. & Lavery, B. Australian rainfall variability and change. *Weather* **52**, 66–72 (1997).
2. Freund, M., Henley, B. J., Karoly, D. J., Allen, K. J. & Baker, P. J. Multi-century cool- and warm-season rainfall reconstructions for Australia's major climatic regions. *Clim. Past* **13**, 1751–1770 (2017).
3. King, A. D. Natural Variability not Climate Change drove the Record Wet Winter in

- 164 Southeast Australia. *Bull. Am. Meteorol. Soc.* **99**, 54–59 (2017).
- 165 4. Ukkola, A. M., Roderick, M., Barker, A. W. & Pitman, A. Exploring the stationarity of
166 Australian temperature, precipitation and pan evaporation records over the last century.
167 *Environ. Res. Lett.* (2019). doi:10.1088/1748-9326/ab545c
- 168 5. Ummenhofer, C. C. *et al.* What causes southeast Australia’s worst droughts? *Geophys.*
169 *Res. Lett.* **36**, L04706 (2009).
- 170 6. Gallant, A. J. E., Kiem, A. S., Verdon-Kidd, D. C., Stone, R. C. & Karoly, D. J.
171 Understanding hydroclimate processes in the Murray-Darling Basin for natural
172 resources management. *Hydrol. Earth Syst. Sci.* **16**, 2049–2068 (2012).
- 173 7. Risbey, J. S., Pook, M. J., McIntosh, P. C., Wheeler, M. C. & Hendon, H. H. On the
174 Remote Drivers of Rainfall Variability in Australia. *Mon. Weather Rev.* **137**, 3233–
175 3253 (2009).
- 176 8. Power, S., Haylock, M., Colman, R. & Wang, X. The Predictability of Interdecadal
177 Changes in ENSO Activity and ENSO Teleconnections. *J. Clim.* **19**, 4755–4771
178 (2006).
- 179 9. Meneghini, B., Simmonds, I. & Smith, I. N. Association between Australian rainfall
180 and the Southern Annular Mode. *Int. J. Climatol.* **27**, 109–121 (2007).
- 181 10. Hendon, H. H., Lim, E.-P. & Nguyen, H. Seasonal Variations of Subtropical
182 Precipitation Associated with the Southern Annular Mode. *J. Clim.* **27**, 3446–3460
183 (2014).
- 184 11. van Dijk, A. I. J. M. *et al.* The Millennium Drought in southeast Australia (2001–
185 2009): Natural and human causes and implications for water resources, ecosystems,
186 economy, and society. *Water Resour. Res.* **49**, 1040–1057 (2013).

- 187 12. Austin, E. K. *et al.* Drought-related stress among farmers: findings from the
188 Australian Rural Mental Health Study. *Med. J. Aust.* **209**, 159–165 (2018).
- 189 13. Hanigan, I. C., Butler, C. D., Kokic, P. N. & Hutchinson, M. F. Suicide and drought in
190 New South Wales, Australia, 1970-2007. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 13950–5
191 (2012).
- 192 14. Cai, W. *et al.* Increased frequency of extreme La Niña events under greenhouse
193 warming. *Nat. Clim. Chang.* **5**, 132–137 (2015).
- 194 15. Zheng, X.-T. *et al.* Indian Ocean Dipole Response to Global Warming in the CMIP5
195 Multimodel Ensemble. *J. Clim.* **26**, 6067–6080 (2013).
- 196 16. Weller, E. & Cai, W. Realism of the Indian Ocean Dipole in CMIP5 Models: The
197 Implications for Climate Projections. *J. Clim.* **26**, 6649–6659 (2013).
- 198 17. Taschetto, A. S. *et al.* Cold Tongue and Warm Pool ENSO Events in CMIP5: Mean
199 State and Future Projections. *J. Clim.* **27**, 2861–2885 (2014).

Figures

Figure 1. The recent drought in context and the drivers of seasonal rainfall variability in the Murray-Darling Basin of Australia. a Observed seasonal rainfall departures as a percentage of the respective seasonal 1961-1990 climatology. The brown bars mark the times of significant multi-year drought defined following Freund *et al.*² The dark blue bars above

the graph indicate if the season in question is categorised as La Niña (LN) or negative IOD (nIOD). **b** Schematic showing the climate modes related to rainfall variability in the Murray-Darling Basin. Climate modes are coloured by the timescale on which they exhibit variability.

Figure 2. A lack of recent La Niña or negative IOD conditions is contributing to the ongoing drought in the Murray-Darling Basin. a. Scatter of September-November Niño-3.4 against concurrent MDB rainfall anomalies. **b.** Scatter of July-September Dipole Mode Index against concurrent MDB rainfall anomalies. **a,b.** Second-order polynomial best fits are plotted (grey; as in Power *et al.*⁸). Spearman rank correlation coefficients are shown and the most recent seasons of data are displayed (black crosses). **c.** Graph of accumulated rainfall anomalies during the time since a La Niña or negative IOD event has occurred. For series greater than three years in length the period from six months onwards is coloured and marked

223 by the end year. In the case of 2019, this is to the end of the year. The grey filled box shows
224 the percentage of occurrences for each time duration since a La Niña or negative IOD event
225 when the accumulated rainfall anomaly was in deficit.

MDB rainfall anomaly from seasonal average (%)

b

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

King, AD;Pitman, AJ;Henley, BJ;Ukkola, AM;Brown, JR

Title:

The role of climate variability in Australian drought

Date:

2020-02-24

Citation:

King, A. D., Pitman, A. J., Henley, B. J., Ukkola, A. M. & Brown, J. R. (2020). The role of climate variability in Australian drought. NATURE CLIMATE CHANGE, 10 (3), pp.177-179. <https://doi.org/10.1038/s41558-020-0718-z>.

Persistent Link:

<http://hdl.handle.net/11343/241631>