A general model of the thermal constraints on the world's most destructive locust,

Schistocerca gregaria

Running head: Thermoregulation in locusts

Author: Koutaro Ould Maeno^{1,2*}, Cyril Piou³, Michael R. Kearney⁴, Sidi Ould Ely^{2,6},

Sid' Ahmed Ould Mohamed², Mohamed El Hacen Jaavar², and Mohamed Abdallahi Ould

Babah Ebbe^{2,5}

¹ Japan International Research Center for Agricultural Sciences (JIRCAS), Livestock and

Environment Division, Ohwashi 1-1, Tsukuba, Ibaraki, 305-8686, Japan

² The Mauritanian National Desert Locust Centre: Centre National de Lutte

Antiacridienne (CNLA), Nouakchott, BP: 665, Mauritania

14 ³ CIRAD, UMR CBGP, INRA, IRD, Montpellier SupAgro, Univ. Montpellier. F-34398

15 Montpellier, France

16 ⁴ School of BioSciences, The University of Melbourne, Parkville, Australia

17 ⁵ Institut du Sahel (INSAH)/CILSS, BP 1530 Bamako, Mali (www.insah.org)

18 ⁶ Direction of Agriculture, Ministry of Rural Development, Nouakchott, Mauritania

19

20 Corresponding Author

21 Koutaro Ould Maeno

22 Japan International Research Center for Agricultural Sciences (JIRCAS), Livestock and

23 Environment Division, Ohwash 1-1, Tsukuba, Ibaraki 305-8686, Japan

24 Email address: kmaeno@affrc.go.jp

25 Tel./fax: +81-298-6622 +81-298-6621

26

27 **Abstract**

28 All terrestrial ectotherms are constrained to some degree by their thermal environment
29 and the extent that they can behaviorally buffer variable thermal conditions. New
30 biophysical modelling methods (NicheMapR) allow the calculation of the body
31 temperature of thermoregulating animals anywhere in the world from first principles, but
32 require detailed observational data for parameterization and testing. Here we describe the
33 thermoregulatory biology of marching bands of the desert locust, *Schistocerca gregaria*,
34 in the Sahara Desert of Mauritania where extreme heat and strong diurnal fluctuations are
35 a major constraint on activity and physiological processes. Using a thermal infrared
36 camera in the field, we showed that gregarious nymphs altered the microhabitats they
37 used, as well as postural thermoregulatory behaviors, to maintain relatively high body
38 temperature (nearly 40 °C). Field and laboratory experiments demonstrated that the
39 preferred body temperature accelerated digestive rates. Migratory bands frequently left

40 foraging sites with full guts before consuming all vegetation and moved to another habitat
41 before emptying their foregut. Thus, the repertoire for behavioral thermoregulation in the
42 desert locust strongly facilitates foraging and digestion rates which may accelerate
43 developmental rates and increase survival. We used our data to successfully parameterize
44 a general biophysical model of thermoregulatory behavior which could capture hourly
45 body temperature and activity at our remote site using globally available environmental
46 forcing data. This modelling approach provides a stronger basis for forecasting thermal
47 constraints on locust outbreaks under current and future climates.

48

49 *Keywords:* body temperature, desert locust, general biophysical model, thermal infrared
50 camera, NicheMapR, thermoregulatory behaviors

51

52 **Introduction**

53 In ecology there is an urgent need to develop better forecasting capacities so that we
54 can anticipate how species of conservation or economic importance will respond to
55 environmental change (Dietze 2017). Pest insects are one of the most economically
56 important groups of species requiring forecasting capacity. Any efforts to predict
57 environmental constraints on the behavior, distribution and abundance of terrestrial
58 ectotherms such as insects must adequately capture how environment and behavior
59 interact to determine body temperature, because virtually all biological processes are
60 temperature dependent (Josephson 1981, Bennett 1985, Angilletta 2009). This is a
61 significant challenge due the complex, nonlinear responses of heat exchange between
62 organisms and their microclimates, but it is possible to compute such responses from first
63 principles using the techniques of biophysical ecology (Porter et al. 1973, Gates 1980,
64 Kearney and Porter 2004, 2009, Buckley 2008).

Crucial to the development of temperature-explicit predictive models of organisms is the inclusion of behavioral responses. The body temperature (T_b) of ectotherms is largely determined by environmental temperatures; however, behavioral thermoregulation, such as shuttling between sun and shade and altering posture (Chappell and Whitman 1990, Heinrich 1993), often allows ectotherms to maintain physiologically suitable T_b even when thermal conditions in their habitats are subject to variation (Heinrich 1999). Indeed, behavioral thermoregulation may buffer many of the negative effects of extreme thermal conditions (Kearney et al. 2009, Huey et al. 2012) in parallel to plastic and evolutionary adjustments in physiological heat tolerance (Cloudsley-Thompson 1975, Heinrich 1993, Araújo et al. 2013, Hoffmann et al. 2013).

Although the computational tools now exist for behaviorally explicit modelling of the field body temperatures of terrestrial ectotherms (e.g. Kearney and Porter, 2019), they require detailed field data for model parameterization and testing. In this study we report

78 on an extensive field study of the field thermoregulatory behavior of one of the world's
79 most destructive insects, *Schistocerca gregaria* (Forskål, 1775), living in the Sahara
80 Desert - one of the world's most extreme thermal environments. We focus on how
81 thermoregulatory behavior in this species is likely to affect its ability to feed and avoid
82 predation and use our data to develop and test a general predictive model of its field body
83 temperature.

84 The desert locust, *S. gregaria*, inhabits arid and semi-arid areas, where desert air
85 temperatures cycle on a 24-hr period, potentially ranging from below freezing to over
86 40 °C in a single day (Symmons and Cressman 2001, Ward 2009, Maeno et al. 2016,
87 2019). The habitats of this species are patchily distributed and are frequented temporally
88 depending on the amount and timing of erratic rain (Tucker et al. 1985, Sword et al. 2010,
89 Showler 2019). Locusts exhibit density-dependent phase polyphenism. Specifically,
90 solitary individuals occur at low densities and are cryptic while gregarious ones occur

at high densities, are attracted to each other, and move long distances in groups (Pener and Simpson 2009, Cullen et al. 2017, Ayali 2019). This collective movement is called “marching” in nymphs and “swarming” in flying adults (Ellis and Ashall 1957, Uvarov 1977).

To our knowledge, the gregarious nymphs of *S. gregaria* are the only ectotherms that diurnally march on the hot ground of the Sahara Desert, travelling up to 1.5 km per day (Ellis and Ashall 1957). Large marching bands deplete food resources quickly, necessitating frequent migration to new food resources. Diurnal migration in the Sahara Desert by walking or jumping is likely risky and energetically expensive for a variety of reasons. First, operative temperatures in open ground within their habitat may become lethal at midday, necessitating behavioral, physiological, and morphological responses to avoid overheating. Second, marching on hot ground increases metabolic and water loss rates, requiring the locusts to refuel efficiently or conserve energy and water resources

during migration. Third, digestion and its nutritional outcome is extremely sensitive to body temperature (Miller et al. 2009b, Coggan et al. 2011, Clissold et al. 2013), enhancing the importance of thermoregulatory responses. Fourth, migratory bands are visually easy to detect by predators, which may lead to special anti-predatory strategies during migration. Predation risk also increases the importance of behavioral thermoregulation during cool mornings when anti-predatory behaviors, such as jumping, are constrained by low temperatures (Maeno et al. 2019).

Much work has been done on the adaptations of gregarious desert locust nymphs to their physical and predatory environments, especially from laboratory studies (Maeno et al. 2013a). However, detailed field studies remain limited, especially where multiple limiting factors are jointly considered, and predictive models are lacking. Thus, in this study we aimed to collect detailed field observations of wild migratory bands of gregarious nymphs in their natural habitat in the Sahara Desert of Mauritania to jointly

consider the hypotheses that: 1) the gregarious nymphs of *S. gregaria* have physiologically high heat tolerance and avoid overheating through certain behaviors associated with microhabitats, 2) they feed frequently, 3) they maintain relatively high T_b to accelerate digestive performance when heat is available, 4) predation and parasitism are low due to high temperatures, and 5) gregariousness (i.e., aggregations) function to increase T_b . We then used our data to develop a biophysical model of these responses to enable us to generalize the thermoregulatory response of *S. gregaria* through space and time and thereby enhance our predictive capacity of the thermal constraints on locust outbreaks.

Material and methods

Study area and climate

Mauritania, in West Africa, is an important area for desert locust outbreaks (Ould Babah

1997, Babah Ebbe 2010, Piou et al. 2017). The study site was 5 km × 5 km in size (19°23'N, 14°35'W) and was located near Akjoujt, northwestern Mauritania. The area is a vast plain with a variety of cover types, including dunes and playa. Sparse, low-growing desert annuals (grasses, herbs, vines, etc.) grew between the bushes and trees and plants were patchily distributed at the study site. *Calotropis procera* (Asclepiadaceae), *Acacia tortilis* (Fabaceae), *Maerua crassifolia* (Capparaceae), and *Boscia senegalensis* (Capparaceae) were frequently used as night-roosting plants (Maeno and Ould Babah Ebbe 2018). We conducted field surveys in November 2016. At this time, rain was scarce and the desert annuals were still mostly green, but began to dry out at the end of the study. Sunrise and sunset occurred at about 07:10 and 18:30 local time, respectively.

Study animals

We studied wild late instar (4th and 5th) nymphs of *S. gregaria*. The body length of the

5th instar nymphs was $43.04 \text{ mm} \pm 0.60$ ($n = 50$) for females and $40.66 \text{ mm} \pm 0.60$ ($n = 50$) for males. Near the survey site, group oviposition was observed approximately one month earlier (i.e., October), and most hatchlings had black bodies, which was typical coloration of the gregarious-phase state (Uvarov 1966, 1977). Almost all individuals had a yellow background with black patterns, and showed diurnal marching behavior, which was typical of the gregarious-phase state. Ground population densities ranged from 0 to more than 5,000 individuals/100 m². Hence, the locusts at our study were highly aggregated and patchily distributed, with dense locust aggregations forming in some areas. This type of distribution and behavior were indicative of the gregarious-phase (Uvarov 1977).

Thermoregulatory behavior

On November 10–19 and 24–27, 2016, at our study site, we surveyed T_b at various times

156 of the day. In the field, we observed that most nymphs within a group behaved
157 synchronously, but many individuals displayed different behaviors, including feeding,
158 marching and resting, as reported by Ellis and Ashall (1957). Probably, these behaviors
159 were caused by intraspecific differences in physiological states, especially hunger levels.

160 We distinguished 12 behaviors, including thermoregulatory behaviors, within the daily
161 behavior cycle classified according to Chappell and Whitman (1990), and measured the
162 associated *T_b* for each (Fig. 1). These behaviors were: 1) night-roosting, 2) basking on
163 night-roosting plants, 3) descending from night-roosting plants, 4) first walking on the
164 ground, 5) basking on the ground during the morning (locusts remained stationary on the
165 ground and their longitudinal body axis was perpendicular to the sun's rays, maximizing
166 the total body surface area exposed to direct solar radiation) (07:00–12:00), 6) feeding,
167 7) marching, 8) stiling on the ground (extending its legs and lift their body from the
168 ground and their body axis was parallel to the sun's rays), 9) remaining under shade, 10)

perching on plants, 11) basking on the ground during the afternoon (12:01–19:00), and 12) ascending into night-roosting plants. During the field survey, if any predation occurred, we recorded the predator and any potential predators.

Measurements of body temperature (T_b)

Obtaining T_b measurements of grasshoppers is technically difficult (Chappell and Whitman 1990). Direct measurements must be made quickly using fine-gauge thermocouples, because the small size and low heat capacity of grasshoppers lead to rapid changes in T_b when the animals are transferred to a different thermal environment for measurement. Free-living nymphs of *S. gregaria* are very sensitive to an approaching observer and can escape quickly (Maeno et al. 2013b). To overcome these problems, we adopted two techniques: First, preliminary trials showed that if the observer (always KOM) approached stationary resting locusts very slowly and silently, nymphs remained

182 in position without fleeing. Second, instead of fine-gauge thermocouples, we used
183 thermal infrared images, thus avoiding the need to capture or disturb the insects (Hunt et
184 al. 2011). To take thermal images as close to free-living nymphs as possible, two
185 additional methods were used, depending on whether locusts were stationary or moving.
186 When locusts were stationary, the observer walked very slowly towards the locust until
187 they were 2–3 m away. When migratory bands actively marched, the observer positioned
188 himself in the predicted direction of the front of the head (more than 20 m), and waited
189 for them to approach, to minimize disturbance. When they passed in front of the observer,
190 thermal images were taken. Thermal images were taken by a handheld thermal imaging
191 camera (FLIR E8, FLIR Systems, Inc.), with 320×240 pixels (9 Hz refresh-rate). This
192 camera measured infrared wavelengths at a range of 7.5–13 μm . All images were taken
193 from more than 50 cm away from the target insects and the emissivity of the camera was
194 set to 0.95 to view locusts. To compensate for low resolution images, high resolution

images were also taken using a digital camera (WG-5, Ricoh Imaging Company, Ltd) to confirm the location and posture of the locust. Each trial disturbed the locusts at the end of recording; thus, each group was recorded just once. Transects were not used in this study; instead, we randomly walked around the study area to find locusts, because gregarious nymphs actively march and frequently change direction and habitats. This procedure was repeated for 255 randomly chosen locusts over a period of 13 days. Locusts sometimes escaped before recordings were completed. In these instances, we only recorded the type of behavior and time, to examine the relationship between background temperature for substrate (T_s , see below) and occurrence of behaviors.

We used a thermo-hygro recorder (TR-72wf, ONDOTORI, Tokyo) to record air temperature and humidity in the shade at 50 cm above the ground. A thermo recorder probe (TR-71nw, ONDOTORI, Tokyo) was also placed above the sandy ground to record the surface temperature of the surface of sunlit soil every 10 min (Fig. 2a). To examine

the interaction between behavioral repertoires for each recording and these environmental factors, the recordings of air temperature and humidity by loggers within 5 mins were used. Also, to assess the range of T_s available to *S. gregaria* throughout the day, we measured ground temperature in shade of spreading shrubs and medium-sized trees, *C. procera*, from 09:00 to 18:00 on November 12, 2016.

Thermal imaging

FLIR Tools software (FLIR Systems, Inc.) was used to extract surface temperature data from thermal images using methods modified from Vogel et al. (2016), specifically extracting the temperature value of the pixel at either the center of aggregating locusts (at least five locusts) or as the one closest to the thorax of an individual as possible. For measurements of individuals based on pictures where nymphs could be visible, the mean values of three insects were used (T_b). To compare these values against ground surface

221 temperature, open ground within 2 m of the targeted locusts was also extracted. When
222 locusts roosted on trees at night, the surface of the tree within 50 cm of the locust was
223 extracted as the background temperature. Mean values of surface temperature within the
224 selected area were used (T_s).

225

226 **Feeding patterns**

227 Gregarious nymphs mainly foraged during the daytime (Kennedy 1939). To examine the
228 interaction between feeding patterns and thermoregulatory behavior, including night-
229 roosting (06:00–09:00), stilting (11:00–15:00), marching (11:00–17:30), and group-
230 basking (16:00–17:00), nymphs displaying one of these behaviors were dissected and the
231 state of their foreguts was recorded. The foreguts were grouped into five size categories:
232 0), empty, 1) little, 2) full, 3) extended, and 4) fully extended, based on methods modified
233 from Ellis (1951). The last three categories were considered to be actively digesting, while

the first two were considered to be mostly digested. Size 0 and 1 were almost similar in appearance, but the contents of the foregut was recorded based on dissection.

Effects of preferred body temperature (T_b) on digestive rate under field conditions

The food processing rate (i.e., rate of passage) is used as an indirect indicator of the overall digestive rate (Dorcas and Peterson 1997). To examine whether preferred T_b during the daytime accelerated the digestive rate, marching last instar nymphs collected just after feeding in the field were housed in a nylon mesh cage (length $\times$ width $\times$ height: 100 cm $\times$ 50 cm $\times$ 50 cm). These nymphs were given no food and were exposed to either direct sunlight or shade. Their foregut size was recorded by dissecting 10 females and 10 males every 30 mins from 13:00 till 15:00 on November 24, 2016. Foregut size was categorized according to the criteria established here. At the same time, we documented the presence of any visible parasite inside the body.

247

248 **Effects of preferred body temperature (T_b) on digestive rate under laboratory**
249 **conditions**

250 The effects of preferred T_b on the digestive rate was also examined under laboratory
251 conditions, based on methods modified from Baines et al. (1973). The *S. gregaria* nymphs
252 used in this study were issued from a line maintained by the CBGP (Montpellier, France).
253 Groups of approximately 100–200 individuals were placed in large cages ($40 \times 40 \times 50$
254 cm) at $32 \pm 1^\circ\text{C}$, with a 12:12 h light:dark photoperiod and 20–50% relative humidity.
255 They were fed wheat seedling and wheat bran *ad libitum*. Newly molted fifth-instar
256 nymphs were removed from the stock cages each day and were kept in a small cage (20
257 $\times 20 \times 35$ cm) in groups not exceeding 40 insects per cage. Insects were maintained in
258 the same rearing room. Only males that were 3 to 4 days old were used. The locusts were
259 kept for 24 h without food, so that the whole gut was almost empty. Then, food (wheat

seedling) was introduced and the locusts were watched until they stopped feeding. The remaining food was removed from the cages to prevent further feeding. Typically, the end of feeding was determined as the time when the locusts moved away from the food source and rested for a period by climbing the wall of the cages. After the test meal (within 25 min of introducing food), all insects were subjected to one of the three treatments. Specifically, they were placed in cages with (1) a heat lamp (100 W) at 32 °C, (2) no lamp at 32 °C, or (3) 20 °C with no lamp. In the first case, locusts were able to thermoregulate freely in the cage by moving closer to or further from the lamp. Locusts were killed at 20 °C after 1 h to examine their digestive rate. Locusts were dissected and only the foregut was weighed, because the midgut and hindgut does not significantly vary in migratory locusts, *Locusta migratoria* (Baines et al. 1973). As a control, the foregut of starved locusts (24 h) and just fed locusts were also examined. A total of 20 insects were examined for each treatment.

273

274 **Statistical analysis**

275 To evaluate the effects of 12 different behaviors on T_b and the effects of body temperature
276 on the weight of the foregut, Tukey–Kramer HSD tests were conducted. To determine
277 differences in temperature for T_s and T_b when they displayed either warming-up (group-
278 basking), cool-down (stilting, remaining in shade, and perching on plants), marching, or
279 feeding, we used the Wilcoxon rank-test. Statistical analyses were conducted using the
280 software package R, version 3.5.0 (R Development Core T. 2018) and JMP (SAS Institute,
281 Cary, NC, USA).

282

283 **Modelling microclimates, body temperature and thermoregulatory behaviors**

284 Biophysical modelling was done using the NicheMapR package (v. 2.0) which comprises
285 a microclimate model (Kearney and Porter 2017) and an ectotherm model (Kearney and

Porter 2019). The microclimate model was forced with the global gridded weather data obtained from the National Centers for Environmental Prediction (NCEP) reanalysis dataset through the RNCEP package (Kemp et al. 2012). These data are at 2° resolution of and were downscaled using the microclima package via the NicheMapR function ‘micro_ncep’, as described in detail in Kearney et al. (2020). All default parameters of the ‘micro_ncep’ function were used except that an organic cap was not simulated at the surface (parameter ‘cap’ = 0, i.e. bare sand) and the longwave downwelling radiation was computed as a function of screen temperature, relative humidity, shade and cloud cover rather than directly using the NCEP longwave radiation. Microclimates were simulated for the year 2016 at the site (longitude/latitude 19.38023, -14.58222) and the local height simulated was 1 cm to approximate the midpoint of a standing locust.

297

The parameters of the ectotherm model were obtained from the present study and

299 presented in table 2. By default the ectotherm model attempts to find a suitable
300 environment for activity by first altering posture (adpressing to the substrate and orienting
301 perpendicular to the sun's rays when cool, orienting parallel to the sun's rays when too
302 hot), then raising core temperature above the preferred temperature to as high as the
303 maximum foraging temperature, then seeking shade. To capture the fact that the locust
304 retreats to trees at night, we overwrote the local 1 cm height values of air temperature,
305 relative humidity and wind speed to that of the reference height (2 m) for nighttime hours,
306 and altered the sky temperature to the shaded value, in the microclimate inputs prior to
307 running the ectotherm model.

308

309 We compared the microclimatic information with the recorded temperature in the field as
310 well as the prediction of the model of body temperature and behaviors with the field
311 observations of November 2016. As the model predicted only basking and foraging

activities we simplified the field data to 3 categories: inactive (night roosting, basking on plants on early morning and descending), basking (morning and afternoon group basking) and foraging (other behaviors).

Once the ectotherm model was verified against field observation from Akjoujt we explored the prediction of this model for the full 2016 year for the study site as well as for a site in Morocco in the Draa valley near Fam el Hisn ($28^{\circ}39'N$, $8^{\circ}52'W$). The Draa valley is frequently hosting solitarious desert locust (some adults were observed at these coordinates in October 2016), and reproduction typically occurs in April to June (Piou et al. 2017); but no gregarious nymphs were observed since 2005 at this site.

Results

Daily thermal regime

Temperature and humidity varied greatly throughout each day in the Sahara Desert (Fig. 2). Following sunrise, air and ground temperature increased, while humidity decreased. Locusts experienced dynamic daily fluctuations in ambient temperature (mean, 26.7 °C; range, 14.3–38.7 °C) and ground temperature (mean, 28.7 °C; 11.5–56.4 °C). This wide range of thermal conditions exceeded the preferred temperature of species and low-temperature thresholds for certain behaviors (see Discussion). Relative humidity also varied greatly within a day (mean, 25.5 %; range, 6.6–60.3 %). Figure 3 shows an example of temperature variation over the course of a day in the field. When solar radiation was available, the range of available temperatures noticeably increased.

Relationship of daily cyclical behavior to T_b

Figure 4 and table 1 present an overview of the humidity, mean air and background temperature, and the surface body temperature of nymphs displaying various

thermoregulatory behaviors. Locusts altered their behavior throughout the day, marching and feeding in a cyclical manner. At night and dawn, when the temperature was coolest, locusts roosted in trees and larger bushes. Locust T_b was very close to T_s before dawn, suggesting no scope for thermoregulation (Figs. 1a & 4; Tukey–Kramer HSD tests, $P > 0.05$). Following sunrise, the locusts moved to the eastern side of the night-roosting plants (Fig. 1b), descended to the ground, and formed dense aggregations (Fig. 1c). Dense carpets of locusts on patches of bare ground (group-basking) were observed, with individuals warming their bodies via solar basking (Fig. 1b, c & d). Their T_b was higher than the T_s (Figs. 4 & 5a; Tukey–Kramer HSD tests, $P < 0.05$). They began to feed (Fig. 1e) and march, alternately, on relatively bare ground (Fig. 1f). Marching and feeding occurred at ground temperatures ranging between 21.2 °C and 54.1 °C throughout the day, while the T_b was slightly lower than the ground temperature (Fig. 4; Tukey–Kramer HSD tests, $P < 0.05$). At midday, when ground temperature exceeded 40 °C, the locusts often

351 stopped marching and remained on the ground, exhibiting stiling behavior (Fig. 1g & h),
352 and seeking shade in low bushes or on the ground in the shade of bushes (Fig. 1i). Some
353 locusts perched on plants (Figs. 1j & 5c, d). Their T_b were maintained at almost 40 °C,
354 within a set-point range that was cooler than the ground temperature (Figs. 4 & 5; Tukey–
355 Kramer HSD tests, $P < 0.05$). In the afternoon, as air and ground temperature fell, locusts
356 again began to march and forage cyclically. Near dusk, they basked in groups on the
357 ground, with T_b being maintained higher than T_s , as observed in the morning (Fig. 1k).
358 They also flew or climbed into trees and larger bushes for nocturnal roosting (Fig. 1l).
359 Thus, group-basking appeared to have a warming effect, while stiling, remaining in the
360 shade, and perching on plants served as a cooling mechanism.

361 To examine how the warm-up and cool-down behaviors influenced locust T_b , the T_b
362 of either warming up or cooling down were plotted against T_s (Fig. 6a). When locusts
363 displayed warm-up behavior, T_b exceeded T_s by 2.49 °C on average (Fig. 6b; Wilcoxon-

test, $z = 4.29$, $n = 95$, $P < 0.001$). In comparison, locusts maintained lower T_b when they displayed cooling down behaviors, with T_b being $8.0\text{ }^{\circ}\text{C}$ lower than T_s on average (Fig. 6b; Wilcoxon-test, $z = -8.55$, $n = 50$, $P < 0.001$). Before sunrise, T_b of night-roosting locusts was not regulated (Fig. 6b; Wilcoxon-test, $z = 0.825$, $n = 29$, $P = 0.41$).

No significant difference was found between T_b and T_s when locusts marched (Fig. 7; Wilcoxon-test, $z = -1.56$, $n = 53$, $P = 0.12$) and fed on grass (Fig. 7; Wilcoxon-test, $z = -0.81$, $n = 54$, $P = 0.42$), but their T_b remained below T_s when the background temperature exceeded $40\text{ }^{\circ}\text{C}$.

Environmental temperature and occurrence of thermoregulatory behavior

Within a day, warming behavior exhibited bimodal patterns, peaking in morning and around dusk (Fig. 8a). In comparison, cooling behavior peaked at midday (Fig. 8d). These patterns matched diel thermal patterns. For instance, warming behavior occurred at

377 relatively low temperatures, while cooling behavior mainly occurred at hot temperatures
378 (Figs. 8b, c, e & f). Marching and feeding occurred throughout the daytime, but the
379 frequency was relatively low during the early morning (low temperature) and at midday
380 (high temperature) (Fig. 9). Figure 10 shows the percentages of various thermoregulatory
381 behaviors within a day. At night, almost all locusts roosted on large trees or large bushes,
382 while various behaviors co-occurred during the daytime.

383

384 **Feeding patterns and digestive rates**

385 Some locusts fed on the leaves of trees in which they roosted at night, but most fed
386 on grass during daytime. The foregut of locusts roosting on night plants was almost empty
387 or small amounts of food were present in the morning (Fig. 11). In comparison, foregut
388 size was large, and contained a large amount of food during the daytime, irrespective of
389 their behaviors (Fig. 11).

When locusts were exposed to the sun or shade at midday, the digestive rate of locusts receiving solar radiation was faster than those in the shade (Fig. 12). Digestion was almost complete within 2 h in the sun. Digestion was also accelerated at preferred T_b under laboratory conditions, when measuring foregut weight instead of size (Fig. 13). When last instar male nymphs were allowed to thermoregulate in their preferred way, the weight of the foregut significantly reduced when compared to locusts kept at 20 °C (Fig. 13; Tukey–Kramer HSD tests, $P < 0.05$).

Modelling thermoregulatory behaviors

The microclimate model accurately predicted the observed temperatures (Fig. 14a, b) even though it was driven by down-scaled gridded data not directly measured on-site. Similarly, the ectotherm biophysical model predictions of body temperature corresponded closely to the measured ones (Fig. 14c). The relative mean squared deviation for surface

temperature, air temperature at 2m and body temperature were, respectively, 5.1, 4.6 and 3.5 °C. These were relatively small errors given that the daily values of these metrics usually fluctuated more than 20 °C between days and nights. The temporal predictions of activity of the ectotherm model also corresponded to the field observations (Fig. 14d). The contingency matrix of the predicted vs. observed behaviors (Table 3) gave an accuracy of 66.6% with only 2 false predictions of inactivity during actual foraging and 8 false predictions of foraging during actual inactivity, but 90 predictions of foraging while actually basking ($n = 305$). Hence, most of the inaccuracy of the model was not problematic as resided in the basking vs. foraging part of the confusion matrix.

Model simulations over the full 2016 year for the Akjoujt field site (Fig. 15a) showed that foraging activity could occur throughout year, with activity possible immediately after sunrise from mid-May to end of September. In comparison, the predictions for the Draa

valley site (Fig. 15b) inferred low-temperature constraints on morning foraging through most of the year, with numerous days of zero or limited activity due to cold weather in autumn, winter and spring despite sunny conditions.

Discussion

All organisms must maintain thermal homeostasis within their tolerance limits but how they achieve this, and the costs and benefits of thermoregulatory precision, will vary with the environment (Huey and Slatkin 1976). Avoidance of overheating through thermoregulatory behavior is essential for ectotherms living in hot environments with little shade, as in our study system. Our aim in this study was to obtain empirical data on the manner and extent to which *S. gregaria* thermoregulate in a desert environment, and then attempt to capture these responses with a mechanistic model integrating microclimatic calculations, heat budget calculations and behavioral algorithm.

429

430 **Thermoregulatory behavior and tolerance of *S. gregaria***

431 We observed that gregarious nymphs of *S. gregaria* avoided overheating as reported
432 previously in this species (Kennedy 1939; Ellis and Ashall 1957; Stower 1963; Culmsee
433 2002). We additionally found that they maintained relatively high T_b (nearly 40 °C) at
434 midday by displaying a variety of behaviors in response to wide-ranging available T_e
435 (from 28 °C to 56 °C). These field-estimated optimal body temperatures are consistent
436 with the laboratory-determined preferred ranges of 38 °C to 41 °C (Chapman 1965). They
437 also correspond to the preferred T_b of locusts and grasshoppers reported to range from
438 35 °C to 42 °C (Chappell and Whitman 1990, Miller et al. 2009, Coggan et al. 2011,
439 Clissold et al. 2013).

440 We inferred that the upper lethal temperature causing 50% mortality was 50.9°C,
441 which is similar to 50–51 °C tolerance documented for the desert-dwelling darkling beetle,

442 *Onymacris plana* (Edney 1971), and 55.1 °C for the Sahara Silver ant, *Cataglyphis bicolor*,
443 which is one of the most heat tolerant land animals alive (Wehner et al. 1992).

444 We frequently observed stiling as a cooling mechanism. This can be a potent
445 regulatory mechanism because of the strong vertical temperature gradient above the
446 ground surface during the day; temperatures can be up to 18 °C lower at just 1 cm above
447 than on the ground (Whitman 1987). However, stiling was not always cooling enough
448 and at these times locusts moved from open ground to cooler locations, such as vegetation
449 and shade. Our data hence show that *S. gregaria* could successfully maintain high but
450 tolerable temperatures in the hostile environment of the Sahara Desert.

451

452 **Thermoregulation, feeding and assimilation**

453 Enhanced thermoregulation is frequently associated with the processes of food
454 uptake and assimilation (Angilletta 2009). In general, ectotherms tend to elevate T_b

455 following feeding (see review, Angilletta 2009) and the preferred T_b is expected to have
456 evolved to match the temperature at which digestion is most efficient (Huey and Bennett
457 1987). Furthermore, the rate of energy gain at any temperature depends on the quantity
458 and quality of food. Because T_b is closely associated with performance, thermoregulatory
459 strategies have frequently coevolved with various life-history traits and spatiotemporal
460 habitats (Angilletta 2009).

461 Why do gregarious nymphs of *S. gregaria* maintain relatively high T_b ? To our
462 knowledge, few studies have documented for locusts in the field the relationship between
463 preferred T_b and spontaneous feeding, digestion, marching patterns, and anti-predatory
464 strategies (Ellis and Ashall 1957). Laboratory experiments showed that the development
465 and growth rates of grasshoppers are highly temperature dependent (Whitman 1988,
466 Chappell and Whitman 1990), including *S. gregaria* (Hamilton 1936, 1950, Symmons
467 et al. 1974), driven by the effects of T_b on food intake, digestion and mobilization rates

468 (Baines et al. 1973, Whitman, 1987, Harrison and Fewell 1995, Miller et al. 2009, Coggan
469 et al. 2011, Clissold et al. 2013). The present study showed that fed gregarious nymphs
470 digested food at the preferred T_b (nearly 40 °C) through behavioral thermoregulation,
471 even in the field. Locusts usually rest for 15 to 40 mins after feeding, to allow digestion
472 (Ellis 1951, Dkhili et al. 2019). Maintaining a high T_b accompanied with stationary
473 cooling down behaviors allows *S. gregaria* to digest food at the preferred T_b during post-
474 prandial resting when high T_e (>40 °C) is available. When T_e was below 40 °C in the
475 afternoon, locusts tended to bask in groups on the ground to raise T_b close to 40 °C.
476 Although the ecological significance of aggregation has remained unclear (Uvarov 1977),
477 this stationary group basking activity might facilitate heat gain by reducing surface-
478 volume ratios and convective heat exchange, creating a thicker boundary layer of still air
479 over the entire locust mass. This is important at our site because, under the cool nighttime
480 conditions there was little scope for behavioral thermoregulation and the digestion rate

481 was very slow.

482 The feeding patterns of gregarious locusts in the field are poorly understood.

483 Despite nymphs digesting most food in the foregut within 2 h during the daytime, we

484 observed that large amounts of food were retained in their foregut throughout the daytime,

485 supporting previous observations (Ellis and Ashall 1957), except when conditions were

486 extremely hot at midday or cool during the early morning (Kennedy 1939). The present

487 study did not monitor individual locusts to determine feeding patterns of meals' quantity

488 and inter-meal intervals. However, our analyses of foregut contents and general

489 observations show that gregarious nymphs frequently fed on small amounts of grass to

490 continuously fill their foregut. Similar feeding patterns (i.e., small meals over short

491 intervals) were observed for gregarious last instar nymphs of *L. migratoria* under

492 laboratory conditions at a constant temperature with *ad libitum* feeding (Simpson 1982,

493 Simpson and Raubenheimer 2000). And Dkhili et al. (2019) observed in the lab that

previously fed *S. gregaria* hoppers stopped their organized marches to feed for a short time when encountering food.

Foraging and migration behaviors

Migration appears to be an adaptive strategy to access sufficient food resources (Dingle 2014). If emigration occurs from habitats that are still favorable for use, it is considered obligatory (Farrow 1990). This may be the case in our population of *S. gregaria* because we frequently observed that marching bands left foraging sites before all plants were consumed. This foraging strategy has a risk of failing to reach favorable habitats. We observed an isolated migratory band in the middle of dunes where food resource was scanty at another site in 2015 (data not shown) which had apparently migrated over a large distance but failed to reach a food resource. However, at our study site, we seldom observed gregarious nymphs with empty foreguts during the daytime,

507 except for individuals just before molting. Thus, migration allows large numbers of
508 gregarious *S. gregaria* nymphs to access sufficient quantities of food constantly,
509 compensating for energy and water loss.

510

511 **Thermoregulation and predation risk**

512 In the wild, the thermoregulatory and foraging activity of prey ectotherms are
513 frequently constrained by predation and the risk of being parasitized because feeding and
514 basking in sunshine immediately exposes preys to predators (Huey and Slatkin 1976).
515 Therefore, many prey ectotherms must trade-off the conflicting needs of thermoregulation
516 and foraging with predation risk and thermal stress (Angilletta et al. 2002). Very high
517 thermal tolerance functions as a defense mechanism if it allows an organism to live where
518 predators cannot be active (Chappell 1983, Heinrich 1993). Although nymphs of *S.*
519 *gregaria* may face much higher losses from bird predation, bird activity tends to be

520 constrained during the hottest hours of the day (Mulli 2009). At our study site terrestrial
521 mammalian and reptile predators were rarely detected. Such predators include jerboas
522 (*Jaculus jaculus*), hedgehogs (*Atelerix albiyentris*), and terrestrial predatory insects. Also,
523 parasitic insects were not detected inside last instar gregarious nymphs in the current
524 study ($n = 1150$). Thus, high T_b might prevent locusts being parasitized or might kill the
525 parasites (Thomas and Blanford 2003). At night, when the escaping performance of
526 locusts was reduced, due to darkness and low temperature, they roosted on relatively high
527 branches of trees and bushes, which might protect them from ground predators (Maeno
528 and Ould Babah Ebbe 2018). This positioning is also favorable for basking in the early
529 morning sun (Whitman, 1988). At low temperature, conspicuous body coloration may
530 compensate low escaping performance as aposematic body coloration protects against
531 predators when nymphs consumed toxic plants such as, *Hyoscyamus muticus* (Sword
532 1999, Sword et al. 2000). Aggregation might also function as “selfish herds” (Sword et

al. 2005).

Predicting *S. gregaria* thermoregulatory behavior and activity constraints

We were able to use the biophysical modelling tools in NicheMapR to capture the thermal constraints on body temperature and foraging behavior in *S. gregaria* at our site.

There are two major challenges to such an endeavor – capturing the available microclimates and properly incorporating behavioral thermoregulation.

To predict microclimates, we use a recently developed approach for downscaling the NCEP historical global weather data (Kalnay et al. 1996) from ~200 km grids to the microclimate scale, via the integration of the microclima and NicheMapR packages (Kearney et al. 2020). Our comparisons of locally measured air temperatures with locally measured surface and air temperatures indicate that this was successful (Fig. 14a, b). The ectotherm model of NicheMapR, which incorporated postural and shade-seeking

thermoregulatory behavior, was then able to capture the observed body temperature and foraging patterns in *S. gregaria*.

The fact that we were able to accurately simulate the thermoregulatory behavior of *S. gregaria* has two implications. First, the simulation assumes that the organism behaves in a way to maximize its time foraging at its preferred temperature and only ceases foraging when it is not thermally suitable. The correspondence between the observations and predictions thus means that this species, at our site, is responding to its thermal environment in a manner to be expected if there were minimal costs to thermoregulation.

Second, because the model is a mechanistic one, we can use it to infer the thermoregulatory constraints and consequences at any other location or time. As an example, our simulation extended to the rest of the year at Akjoujt and for the whole year in Draa Waddy (Fig. 15) shows that the latter site would provide less foraging time, especially in the cooler months. More accurate estimates of body temperature and

foraging constraints can be used as input into forecasting models of locust outbreak (Piou et al. 2019) and movement patterns. This should help improve forecasting capacity and preventive control, both short-term in response to weather events and long-term in response to climate change (Meynard et al. 2017).

Funding

This work was supported by Grants-in-Aid for Scientific Research (KAKENHI grant No. 15K18808) from the Japan Society for the Promotion of Science. CP was funded by ANR JCJC PEPPER (ANR-18-CE32-0010-01).

Acknowledgements

We thank T. Tidjani and E. Mohamed and members of Centre National de Lutte Antiacridienne (CNLA) for providing assistance with the field surveys. We also thank L.

572 Nicolas and E. Christophe and members of Locust ecology and management research at
573 CBGP for helping in various aspects of the research. We greatly appreciate the
574 constructive and valuable comments on the manuscript from D. Cullen and one
575 anonymous reviewer.

576

577 **Literature Cited**

- 578 Angilletta, M. J. 2009. Thermal Adaptation: A theoretical and empirical synthesis.
579 Oxford University Press.
- 580 Angilletta, M. J., P. H. Niewiarowski, and C. A. Navas. 2002. The evolution of thermal
581 physiology in ectotherms. *Journal of Thermal Biology* 27:249–268.
- 582 Araújo, M. B., F. Ferri-Yáñez, F. Bozinovic, P. A. Marquet, F. Valladares, and S. L.
583 Chown. 2013. Heat freezes niche evolution. *Ecology Letters* 16:1206–1219.
- 584 Ayali, A. 2019. The puzzle of locust density-dependent phase polyphenism. *Current*
585 *Opinion in Insect Science* 35:41–47.
- 586 Babah Ebbe, M. A. 2010. Biogéographie du Criquet pèlerin en Mauritanie. Hermann,
587 Paris.
- 588 Baines, D. M., E. A. Bernays, and E. M. Leather. 1973. Movement of food through the
589 gut of fifth-instar males of *Locusta migratoria migratorioides* (R. & F.). *Acrida*

590 2:319–332.

591 Bennett, A. F. 1985. Temperature and muscle. *Journal of Experimental Biology*

592 344:333–344.

593 Buckley, L. B. 2008. Linking traits to energetics and population dynamics to predict

594 lizard ranges in changing environments. *American Naturalist* 171:1–19.

595 Chapman, R. F. 1965. The behaviour of nymphs of *Schistocerca gregaria* (Forskål)

596 (Orthoptera, Acrididae) in a temperature gradient, with special reference to

597 temperature preference. *Behaviour* 24:283–317.

598 Chappell, M. A. 1983. Thermal limitations to escape responses in desert grasshoppers.

599 *Animal Behaviour* 31:1088–1093.

600 Chappell, M. A., and D. W. Whitman. 1990. Grasshopper thermoregulation. In:

601 Chapman RF, Joern A (eds) *Biology of grasshoppers*. John Wiley, New York,

602 pp:143–172.

- 603 Clissold, F. J., N. Coggan, and S. J. Simpson. 2013. Insect herbivores can choose
604 microclimates to achieve nutritional homeostasis. *The Journal of Experimental*
605 *Biology* 216:2089–2096.
- 606 Cloudsley-Thompson, J. L. 1975. Adaptations of arthropoda to arid environments.
607 *Annual Review of Entomology* 20:261–283.
- 608 Coggan, N., F. J. Clissold, and S. J. Simpson. 2011. Locusts use dynamic
609 thermoregulatory behaviour to optimize nutritional outcomes. *Proceedings of the*
610 *Royal Society B* 278:2745–2752.
- 611 Cullen, D. A., A. J. Cease, A. V. Latchininsky, A. Ayali, K. Berry, J. Buhl, R. De
612 Keyser, B. Foquet, J. C. Hadrich, T. Matheson, S. R. Ott, M. A. Poot-Pech, B. E.
613 Robinson, J. M. Smith, H. Song, G. A. Sword, J. Vanden Broeck, R. Verdonck, H.
614 Verlinden, and S. M. Rogers. 2017. From molecules to management: mechanisms
615 and consequences of locust phase polyphenism. *Advances in Insect Physiology*

- 616 53:167–285.
- 617 Culmsee, H. 2002. The habitat functions of vegetation in relation to the behaviour of the
- 618 desert *Schistocerca gregaria* (Forskål)(Acrididae: Orthoptera). A study in
- 619 Mauritania (West Africa). *Phytocoenologia* 32:645–664.
- 620 Dietze, M. 2017. *Ecological Forecasting*. Princeton University Press.
- 621 Dingle, H. 2014. *Migration, The biology of life on the move*. Oxford Univ. Press,
- 622 London, U.K.
- 623 Dkhili, J., K. O. Maeno, L. Mina, I. Hassani, S. Ghaout, and C. Piou. 2019. Effects of
- 624 starvation and vegetation distribution on locust collective motion. *Journal of Insect*
- 625 *Behavior* 32:207–217.
- 626 Dorcas, M. E., and C. R. Peterson. 1997. The thermal biology of digestion in rubber
- 627 boas (*Charina bottae*): physiology, behavior, and environmental constraints.
- 628 *Physiological Zoology* 70:292–300.

- 629 Edney, E. B. 1971. The body temperature of tenebrionid beetles in the Namib desert of
630 Southern Africa. *Journal of Experimental Biology* 55:253–272.
- 631 Ellis, P. E. 1951. The marching behaviour of hoppers of the African migratory locust
632 (*Locusta migratoria migratorioides* R. & F.) in the laboratory. *Anti-Locust Bull*
633 7:1–46.
- 634 Ellis, P. E., and C. Ashall. 1957. Field studies on diurnal behaviour, movement and
635 aggregation in the desert locust (*Schistocerca gregaria* Forskål). *Anti-Locust*
636 *Bulletin* 25:1–103.
- 637 Farrow, R. A. 1990. Flight and migration in acridoids. In: Chapman RF, Joern A (eds)
638 *Biology of grasshoppers*. John Wiley, New York, pp.
- 639 Gates, D. M. 1980. *Biophysical Ecology*. Springer Verlag, New York.
- 640 Hamilton, A. G. 1936. The relation of humidity and temperature to the development of
641 three species of African locusts-*Locusta migratoria migratorioides* (R.& F.),

- 642 *Schistocerca gregaria* (FORSK.), and *Nomadacris septemfasciata* (SERV.).
- 643 Transactions of the Royal Entomological Society of London 85:1–60.
- 644 Hamilton, A. G. 1950. Further studies on the relation of humidity and temperature to the
- 645 development of two species of African locusts-*Locusta migratoria migratorioides*
- 646 (R. & F.) and *Schistocerca gregaria* (Forsk). Transactions of the Royal
- 647 Entomological Society of London 101:1–58.
- 648 Harrison, F., and H. Fewell. 1995. Thermal effects on feeding behavior and net energy
- 649 intake in a grasshopper intake energy experiencing large in body diurnal
- 650 fluctuations temperature. Physiological Zoology 68:453–473.
- 651 Heinrich, B. 1993. The hot-blooded insects. Harvard University Press, Cambridge,
- 652 Mass.
- 653 Heinrich, B. 1999. The thermal warriors. Harvard University Press, Cambridge, MA.
- 654 Hoffmann, A. A., S. L. Chown, and S. Clusella-Trullas. 2013. Upper thermal limits in

- 655 terrestrial ectotherms: how constrained are they ? *Functional Ecology*:934–949.
- 656 Huey, R. B., and A. F. Bennett. 1987. Phylogenetic studies of coadaptation: preferred
- 657 temperatures versus optimal performance temperatures of lizards. *Evolution*
- 658 41:1098–1115.
- 659 Huey, R. B., M. R. Kearney, A. Krockenberger, J. A. M. Holtum, M. Jess, and S. E.
- 660 Williams. 2012. Predicting organismal vulnerability to climate warming : roles of
- 661 behaviour , physiology and adaptation. *Proceedings of the Royal Society B*
- 662 367:1665–1679.
- 663 Huey, R. B., and M. Slatkin. 1976. Cost and benefits of lizard thermoregulation. *The*
- 664 Quarterly Review of Biology 51:363–384.
- 665 Hunt, V. L., G. D. Lock, S. G. Pickering, and A. K. Charnley. 2011. Application of
- 666 infrared thermography to the study of behavioural fever in the desert locust.
- 667 *Journal of Thermal Biology* 36:443–451.

- 668 Josephson, R. K. 1981. Temperature and the mechanical performance of insect muscle.
669 In: Heinrich B (ed) Insect thermoregulation. John Wiley, New York, pp:19–44.
- 670 Kalnay, E., M. Kanamitsu, R. Kistler, W. Collins, D. Deaven, L. Gandin, M. Iredell, S.
671 Saha, G. White, J. Woollen, Y. Zhu, M. Chelliah, W. Ebisuzaki, W. Higgins, J.
672 Janowiak, K. C. Mo, C. Ropelewski, J. Wang, A. Leetmaa, R. Reynolds, R. Jenne,
673 and D. Joseph. 1996. The NCEP/NCAR 40-year reanalysis project. Bulletin of the
674 American Meteorological Society 77:437–471.
- 675 Kearney, M., and W. P. Porter. 2004. Mapping the fundamental niche: physiology,
676 climate, and the distribution of a nocturnal lizard. Ecology 85:3119–3131.
- 677 Kearney, M., and W. Porter. 2009. Mechanistic niche modelling: combining
678 physiological and spatial data to predict species' ranges. Ecology Letters 12:334–
679 350.
- 680 Kearney, M. R., and W. P. Porter. 2017. NicheMapR – an R package for biophysical

- 681 modelling: the microclimate model. *Ecography* 40:664–674.
- 682 Kearney, M. R., and W. P. Porter. 2019. NicheMapR – an R package for biophysical
- 683 modelling: the ectotherm and Dynamic Energy Budget models. *Ecography* 43:85–
- 684 96.
- 685 Kearney, M., R. Shine, and W. P. Porter. 2009. The potential for behavioral
- 686 thermoregulation to buffer “cold-blooded” animals against climate warming.
- 687 *Proceedings of the National Academy of Sciences* 106:3835–3840.
- 688 Kearney, M. R., P. K. Gillingham, I. Bramer, J. P. Duffy, and I. M. D. Maclean. 2020.
- 689 A method for computing hourly, historical, terrain-corrected microclimate
- 690 anywhere on earth. *Methods in Ecology and Evolution* 11:38–43.
- 691 Kemp, M. U., E. Emiel van Loon, J. Shamoun-Baranes, and W. Bouten. 2012. RNCEP:
- 692 Global weather and climate data at your fingertips. *Methods in Ecology and*
- 693 *Evolution* 3:65–70.

- 694 Kennedy, J. S. 1939. The behaviour of the desert locust *Schistocerca gregaria*
695 (Forsk.)(Orthoptera) in an outbreak centre. Transactions of the Royal
696 Entomological Society of London 89:385–542.
- 697 Maeno, K. O., and M. A. Ould Babah Ebbe. 2018. Aggregation site choice by
698 gregarious nymphs of the desert locust, *Schistocerca gregaria*, in the Sahara Desert
699 of Mauritania. Insects 9:1–9.
- 700 Maeno, K. O., C. Piou, M. A. Ould Babah, and S. Nakamura. 2013a. Eggs and
701 hatchlings variations in desert locusts: phase related characteristics and starvation
702 tolerance. Frontiers in physiology 4:345.
- 703 Maeno, K. O., C. Piou, S. Ould Ely, M. A. Ould Babah, B. Péliissié, S. Ould Mohamed,
704 M. E. H. Jaavar, M. Etheimine, and S. Nakamura. 2013b. Plant size-dependent
705 escaping behavior of gregarious nymphs of the Desert locust, *Schistocerca*
706 *gregaria*. Journal of Insect Behavior 26:623–633.

- 707 Maeno, K. O., S. Ould Ely, S. Nakamura, K. Abdellaoui, S. Cissé, M. E. H. Jaavar, S.
 708 Ould Mohamed, M. Atheimine, and M. A. Ould Babah. 2016. Daily microhabitat
 709 shifting of solitarious-phase Desert locust adults: implications for meaningful
 710 population monitoring. *SpringerPlus* 5:107.
- 711 Maeno, K. O., S. Ould Ely, S. Ould Mohamed, M. E. H. Jaavar, S. Nakamura, and M.
 712 A. Ould Babah Ebbe. 2019. Defence tactics cycle with diel microhabitat choice
 713 and body temperature in the desert locust, *Schistocerca gregaria*. *Ethology*
 714 125:250–261.
- 715 Meynard, C. N., P.-E. Gay, M. Lecoq, A. Foucart, C. Piou, and M.-P. Chapuis. 2017.
 716 Climate-driven geographic distribution of the desert locust during recession
 717 periods: Subspecies' niche differentiation and relative risks under scenarios of
 718 climate change. *Global Change Biology* 23:4739–4749.
- 719 Miller, G. A., F. J. Clissold, D. Mayntz, and S. J. Simpson. 2009. Speed over efficiency:

- 720 locusts select body temperatures that favour growth rate over efficient nutrient
721 utilization. *Proceedings of the Royal Society B* 276:3581–3589.
- 722 Mullié, W. C. 2009. Chapter 14. Birds, locusts and grasshoppers. In: Zwarts L, Bijlsma
723 RG, van der Kamp J, Wymenga E, editors. *Living on the edge. Wetlands and birds*
724 *in a changing Sahel*. Zeist: KNNV Publishing.
- 725 Ould Babah, M. A. 1997. Strategy for controlling the desert locust in Mauritania. In:
726 Krall S, Peveling R, Ba Diallo D) (ed) *New strategies in locust control*. Page (B. D.
727 D. Krall S, Peveling R, Ed.). Birkhaeuser Verlag, Basel, Switzerland.
- 728 Pener, M. P., and S. J. Simpson. 2009. Locust phase polyphenism:an update. *Advances*
729 *in insect physiology* 36:1–286.
- 730 Piou, C., M. E. H. Jaavar Bacar, M. A. Ould Babah Ebbe, J. Chihrane, S. Ghaout, S.
731 Cisse, M. Lecoq, and T. Ben. 2017. Mapping the spatiotemporal distributions of
732 the Desert Locust in Mauritania and Morocco to improve preventive management.

- 733 Basic and Applied Ecology 25:37–47.
- 734 Piou, C., P. E. Gay, A. S. Benahi, M. A. O. Babah Ebbe, J. Chihrane, S. Ghaout, S.
- 735 Cisse, F. Diakite, M. Lazar, K. Cressman, O. Merlin, and M. J. Escorihuela. 2019.
- 736 Soil moisture from remote sensing to forecast desert locust presence. *Journal of*
- 737 *Applied Ecology* 56:966–975.
- 738 Porter, W. P., J. W. Mitchell, W. A. Beckman, and C. B. DeWitt. 1973. Behavioral
- 739 implications of mechanistic ecology - Thermal and behavioral modeling of desert
- 740 ectotherms and their microenvironment. *Oecologia* 13:1–54.
- 741 R Development Core T. 2018. R: A language and environment for statistical computing.
- 742 R Foundation for Statistical Computing, Vienna, Austria. <[http://www.R-](http://www.R-project.org)
- 743 [project.org](http://www.R-project.org)>.
- 744 Showler, A. T. 2019. Desert locust control: the effectiveness of proactive interventions
- 745 and the goal of outbreak prevention. *American Entomologist* 65:180–191.

- 746 Simpson, S. J. 1982. Patterns in feeding: a behavioural analysis using *Locusta*
747 *migratoria* nymphs. *Physiological Entomology* 7:325–336.
- 748 Simpson, S. J., and D. Raubenheimer. 2000. The hungry locusts. *Advances in the Study*
749 *of Behavior* 29:1–44.
- 750 Stower, W. J. 1963. Photographic techniques for the analysis of locust “hopper”
751 behaviour. *Animal Behaviour* 11:198–205.
- 752 Sword, G. A. 1999. Density-dependent warning coloration. *Nature* 397:217.
- 753 Sword, G. A., P. D. Lorch, and D. T. Gwynne. 2005. Migratory bands give crickets
754 protection. *Nature* 433:703.
- 755 Sword, G. A., S. J. Simpson, O. Taleb, M. El Hadi, and H. Wilps. 2000. Density-
756 dependent aposematism in the désert locust. *Proceedings of the Royal Society B*.
757 267:63–68
- 758 Sword, G. A., M. Lecoq, and S. J. Simpson. 2010. Phase polyphenism and preventative

- 759 locust management. *Journal of insect physiology* 56:949–57.
- 760 Symmons, P. M., and K. Cressman. 2001. Desert locust guidelines 1: Biology and
761 behavior. Rome: FAO.
- 762 Symmons, P. M., S. M. Green, R. A. Robertson, and K. G. Wardhaugh. 1974. The
763 production of distribution maps of the incubation and hopper development periods
764 of the desert locust *Schistocerca gregaria* (Forsk.) (Orthoptera, Acrididae).
765 *Bulletin of Entomological Research* 64:443–451.
- 766 Thomas, M. B., and S. Blanford. 2003. Thermal biology in insect-parasite interactions.
767 *Trends in Ecology & Evolution* 18:344–350.
- 768 Tucker, C. J., J. U. Hielkema, and J. Roffey. 1985. The potential of satellite remote
769 sensing of ecological conditions for survey and forecasting desert-locust activity.
770 *International Journal of Remote Sensing* 6:127–138.
- 771 Uvarov, B. P. 1966. Grasshoppers and Locusts. Vol. 1. Cambridge University. Press,

- 772 Cambridge, U.K.
- 773 Uvarov, B. P. 1977. Grasshoppers and Locusts. Vol. 2. Centre for Overseas Pest
774 Research, London.
- 775 Vogel, B., H. Wagner, J. Gmoser, A. Wörner, A. Löschberger, L. Peters, A. Frey, U.
776 Hofmann, and S. Frantz. 2016. Touch-free measurement of body temperature using
777 close-up thermography of the ocular surface. *MethodsX* 3:407–416.
- 778 Ward, D. 2009. The biology of deserts. Oxford University. Press Inc., Oxford UK, UK.
- 779 Wehner, R., A. C. Marsh, and S. Wehner. 1992. Desert ants on a thermal tightrope.
780 *Nature* 357:586–587.
- 781 Whitman, D. W. 1987. Thermoregulation and daily activity patterns in a black desert
782 grasshopper, *Taeniopoda eques*. *Animal Behaviour* 35:1814–1826.
- 783 Whitman, D. W. 1988. Function and evolution of thermoregulation in the desert
784 grasshopper *Taeniopoda eques*. *The Journal of Animal Ecology* 57:369–383.
- 785

Table 1. Maximum, minimum, and mean air temperature, humidity, ground temperature, background (ground or plants) temperature, and locust body temperature, as well as time when locusts were first and last seen over 13 days when locusts displayed 12 different behaviors within a day.

				Air temperature (°C) from			Humidity (RH) from			Ground temperature (°C) from			Background temperature (°C) from			Body temperature (°C) from				
		Timeof	Timeof	a data logger			a data logger			a data logger			an infrared thermal camera			an infrared thermal camera				
	Behavioral categories	the first seen	the last seen	Max	Min	Mean	Max	Min	Mean	Max	Min	Mean	Max	Min	Mean	Max	Min	Mean	n	
1	Night roosting	19:00	6:30	26.2	16.5	19.65 ± 0.43 e	52.3	14.9	26.51 ± 1.29 a	24.9	13.7	17.33 ± 0.95 f	26.5	15.1	18.91 ± 1.03 f	27	15.5	19.34 ± 0.72 e	29	
2	Basking on plant	6:54	10:49	24.8	22.1	22.72 ± 1.05 e	22.3	15.9	20.92 ± 3.10 abcd	29.9	20.5	22.74 ± 2.28 ef	23.6	16.4	21.20 ± 2.48 def	23.4	20.7	21.54 ± 1.73 e	5	
3	Descending from night-roosting plants	6:55	10:49	25.3	22.1	22.74 ± 1.05 e	22.2	16.8	21.12 ± 3.10 abc	26	21.1	22.08 ± 2.28 ef	26.2	18.4	20.46 ± 2.48 ef	26.6	21.9	23.86 ± 1.73 e	5	
4	First walking	7:15		22.3	22.3	22.3 e	22.3	22.3	22.3 ab	20.5	20.5	20.5 f	19.7	15.8	18.21 ± 2.09 f	22.8	18.5	20.54 ± 1.46 e	7	
5	Morning basking	7:23	11:59	31.2	24	28.26 ± 0.29 d	39.2	8.9	17.75 ± 0.86 b	39.7	26.7	33.68 ± 0.63 d	39.6	21.5	29.06 ± 0.69 d	39.4	25.6	31.83 ± 0.48 d	65	
6	Feeding	7:27	17:48	33.8	22.6	29.95 ± 0.32 c	38.9	6.8	10.36 ± 0.94 de	52.4	21.2	39.61 ± 0.69 c	51.9	18	36.3 ± 0.75 c	43.8	19.5	34.74 ± 0.53 c	54	
7	Marching	7:15	18:47	34.8	28.2	31.63 ± 0.32 b	21.9	6.8	8.52 ± 0.95 e	54.1	28.7	43.32 ± 0.70 b	51.9	25.8	41.50 ± 0.76 b	48.7	27.9	40.30 ± 0.53 a	53	
8	Stilting	9:47	16:32	33.6	33.3	33.51 ± 0.74 ab	9.1	8.8	8.89 ± 2.19 cde	46.4	45.3	46.07 ± 1.61 ab	55.5	43.8	46.27 ± 1.75 ab	41.1	39	39.86 ± 1.22 ab	10	
9	Shading	11:02	16:08	37.7	31.9	34.63 ± 0.60 a	15.7	8	11.08 ± 1.79 cde	54.5	39.5	50.97 ± 1.32 a	52.9	41.9	48.46 ± 1.43 a	43	35.4	39.36 ± 1.00 ab	15	
10	Perching on plants	11:02	17:20	37.7	31.9	34.29 ± 0.47 a	17.2	6.8	11.08 ± 1.39 cde	54.5	39.5	49.21 ± 1.02 a	55.5	41.9	47.51 ± 1.11 a	43.1	34.7	39.57 ± 0.77 a	25	
11	Afternoon basking	13:10	18:40	35.7	24.6	31.89 ± 0.43 b	43.9	7.1	18.74 ± 1.27 b	42.5	30.2	34.86 ± 0.93 d	38.7	27.2	33.72 ± 1.01 c	39.1	30	35.63 ± 0.71 bc	30	
12	Ascending to night-roosting plants	16:09	18:47	35.1	29.2	31.26 ± 0.88 abcd	18.8	8.7	11.76 ± 2.62 bcde	36.4	28.7	31.44 ± 1.93 de	34.7	25.7	29.71 ± 2.09 cde	35.8	29	32.30 ± 1.46 cd	7	

Air temperature, humidity, and ground temperature was recorded by a data logger.

Background substrate (ground or plants where locusts remained) temperature and locust body temperature was recorded by an infrared

792 thermal camera.

793 Different letters after each value indicate significant differences at $P < 0.05$ from Tukey-Kramer HSD

794

795 Table 2. Parameters used in the ectotherm model for *Schistocerca gregaria* hoppers.

Parameter	Description	Unit	Value	Reference/remarks
$CT_{\max}$	lethal maximum body temperature	°C	50	This study
$CT_{\min}$	lethal minimum body temperature	°C	1	Lab observations
T_{pref}	thermoregulation target body temperature	°C	40	This study
$T_F^{\max}$	maximum foraging body temperature	°C	43	This study
$T_F^{\min}$	minimum foraging body temperature	°C	25	This study
$T_B^{\min}$	minimum basking temperature	°C	15	This study

$T_{RB}^{\min}$	minimum temperature for movement to basking site	°C	15	This study
burrow	can the animal move underground?		false	
climb	can the animal climb?		true	
shade_seek	can the animal seek shade?		true	
shape	body shape		cylinder	
W_w	wet body mass	g	5	overestimated*
% wet	fraction of body surface acting as a free-water surface		0.1	estimated
%eyes	fraction of eyes surface acting as a free-water surface		0	estimated

minshades	daily minimum shade values	%	0	study sites are semi-desert areas with
maxshades	daily maximum shade values	%	90	vegetation

796 * but at the lower limit of the model computation capacity

797

798

799 Table 3. Confusion matrix of the observed vs. predicted behaviors of hoppers with the ectotherm-microclimate model.

		predicted behaviors		
		inactive	basking	foraging
observed behaviors	inactive	37	0	2
	basking	0	5	90
	foraging	8	0	163

800

801

802 **Figure legends**

803 Figure 1. Various behaviors of gregarious nymphs of *Schistocerca gregaria* from night

804 (a) to dusk over a 1-day period (l). (a) Night-roosting on tree, (b) nymphs moved to east

805 side of night-roosting plant and descended to the ground, (c) ground basking, (d) basking

806 posture: their body direction was perpendicular to the sun's rays at cooler temperatures

807 ($< 40^{\circ}\text{C}$), (e) feeding on low vegetation, (f) marching bands, (g) stilting on the hot ground,

808 (h) lateral view of stilting locust: such stilting locusts remained on the ground and lifted

809 their bodies to be distant from the ground surface, (i) locusts moved to the shade, (j)

810 locusts perched on dried *Calotropis procera* when ground temperature was intolerably

811 hot ($> 50^{\circ}\text{C}$), (k) ground-basking during the late afternoon, and (l) ascending to night-

812 roosting tree around dusk.

813

814 Figure 2. Diel fluctuations in mean air temperature ($\circ$), ground surface temperature in

815 sunlight (●) and humidity (△) for 15 days (November 2016) near Akjoujt, north-west
816 Mauritania. Error bars represent SE. Dark bar along the horizontal axis represents night-
817 time.

818

819 Figure 3. Example of diel fluctuation in mean ground surface temperature exposed to
820 sunlight (○) and under the shade of a spreading shrub or medium-sized tree, *Calotropis*
821 *procera* (●), from 09:00 to 18:00, and grass surface temperature of *Indigofera argentea*
822 of less than 10 cm height (▲) taken by an infrared thermal camera on Nov. 12, 2016 near
823 Akjoujt, north-west Mauritania. Error bars represent SE. Gray-colored area is available
824 temperature for locusts during the daytime in the field.

825

826

827 Figure 4. Relationships between various behaviors and mean air temperature (○),

828 background temperature (●), humidity (△), and locust body temperature (●) over 1-day.

829 Note, locusts kept their body temperature close to 40 °C when it was hot (> 40 °C) during

830 the daytime by displaying cooling-down behaviors. In comparison, locusts warmed-up

831 by basking at cool temperatures during the morning and afternoon. Different letters above

832 each bar indicate significant differences at $P < 0.05$ from Tukey-Kramer HSD for body

833 temperature (°C). Numbers in the figure represent the sample size.

834

835 Figure 5. Thermal images of locusts displaying group basking during the morning (09:04)

836 (a & b) or shading and perching on *Calotropis procera* at midday (14:05), taken by an

837 infrared thermal camera. Arrows in the photos indicate the positions of locusts that were

838 present.

839

840 Figure 6. (a) Relationships between background temperature (T_s) and locust body

temperature (T_b) when locusts displayed roosting on plants at night (●), warming-up behavior (●: morning and afternoon basking), and cooling-down behaviors (▲:stilting, shading, and perching on plants). (b) Comparison between T_s and T_b for each thermoregulatory behaviors. N.S., $P > 0.05$; ***, $P < 0.001$, represent significant differences from the Wilcoxon-test.

Figure 7. (a) Relationship between background temperature (T_s) and locust body temperature (T_b) when locusts actively marched (●) or fed on plants (●) during the daytime. (b) Comparison between T_s and T_b for each thermoregulatory behaviors. N.S., $P > 0.05$, represent significant differences from the Wilcoxon-test.

854 Figure 8. Frequency of gregarious nymphs displaying warming-up (group-basking, $n =$
 855 324) (a, b & c) or cooling-down behaviors (stilting, shading, and perching on plants, $n =$
 856 262) (d, e & f) at various times of the day (a & d), air temperatures ($^{\circ}\text{C}$)(b & e), and
 857 ground temperatures ($^{\circ}\text{C}$) (c & f). Basking behaviors were mainly displayed during
 858 morning and late afternoon, while cooling-down behaviors were displayed at midday
 859 when it was hottest (a & d). Basking behavior occurred at cool temperatures, while
 860 cooling down behavior occurred at hot temperatures (b, c, e & f).

861

862 Fig. 9. Frequency of gregarious nymphs displaying marching ($n = 335$) (a, b & c) or
 863 feeding ($n = 207$) (d, e & f) behavior at various times of the day (a & d), air temperatures
 864 ($^{\circ}\text{C}$) (b & e), and ground temperatures ($^{\circ}\text{C}$) (c & f). Both behaviors were observed at any
 865 time during the daytime and wide range of temperatures; however, most behaviors
 866 occurred after 09:00 when the temperature rose. Most marching occurred after feeding.

867

868 Fig. 10. Percentage of gregarious nymphs displaying various behaviors at various times
869 of the day. Field observations over 13 days were pooled. (Different locust groups, $n =$
870 1584). Note, locusts displayed various behaviors during a single daytime.

871

872 Fig. 11. Status of the foregut of gregarious nymphs that displayed night-roosting (06:00–
873 09:00), basking (16:00–17:00), stilting (11:00–15:00), and marching (11:00–17:30)
874 behavior. The foregut was grouped into five categories depending on size. Numbers above
875 the bars represent the sample size.

876

877 Fig. 12. Effects of preferred body temperature on the digestive rate in gregarious nymphs
878 on Nov. 24 2016. Just after feeding, locusts were exposed to either sun (a) or shade (b),
879 and the state of their foreguts was recorded every 30 min from 13:00 to 15:00. $n = 20$ (10

880 females and 10 males) each.

881

882 Fig. 13. Effects of preferred body temperature on digestive rate in gregarious nymphs

883 under laboratory conditions. Just after feeding, locusts were exposed to a heat lamp or no

884 heat lamp of 32 °C and 20 °C, and the weight of their foregut was measured after 1 h

885 exposure. $n = 20$ males each.

886

887 Fig. 14. Predicted surface temperature (a), air temperature at 2m (b), body temperature

888 (c) and behavior (d) following the microclimate and the biophysical ectotherm models.

889 The field observations are displayed in red.

890

891 Fig. 15. Predictions of hourly behaviors from the ectotherm model along the 2016 year

892 for a) the study site near Akjoujt in Mauritania (19°23'N, 14°35'W) and b) the Draa

893 Waddy near Fam el Hisn in Morocco ($28^{\circ}39'N$, $8^{\circ}52'W$). Color code: dark blue = night
894 time (inactivity), light blue = basking, orange = foraging, and white = inactivity during
895 light hours. The red rectangle in subplot (a) indicates the period of the field study in
896 Mauritania.

897

898

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8

Fig. 9

Fig. 10

Fig. 11

Fig. 12

Fig. 13

Fig. 14

1013

1014 Fig. 15

1015

1016

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Maeno, KO;Piou, C;Kearney, MR;Ould Ely, S;Ould Mohamed, S;Jaavar, MEH;Ould Babah Ebbe, MA

Title:

A general model of the thermal constraints on the world's most destructive locust, *Schistocerca gregaria*

Date:

2021-04-07

Citation:

Maeno, K. O., Piou, C., Kearney, M. R., Ould Ely, S., Ould Mohamed, S., Jaavar, M. E. H. & Ould Babah Ebbe, M. A. (2021). A general model of the thermal constraints on the world's most destructive locust, *Schistocerca gregaria*. *ECOLOGICAL APPLICATIONS*, 31 (4), <https://doi.org/10.1002/eap.2310>.

Persistent Link:

<http://hdl.handle.net/11343/275141>