Background-dependent *Wolbachia*-mediated insecticide resistance in
Laodelphax striatellus

Yongteng Li ¹, Xiangdong Liu ², Na Wang¹, Yueliang Zhang¹, Ary Hoffmann³ and
Huifang Guo ^{1*}

1. Institute of Plant Protection, Jiangsu Academy of Agricultural Sciences, Nanjing,
People's Republic of China

2. Department of Entomology, Nanjing Agricultural University, Nanjing, People's
Republic of China

3. School of Biosciences, Bio21 Institute, The University of Melbourne, Parkville, VIC
3010, Australia

*Address correspondence to Huifang Guo:

Institute of Plant Protection, Jiangsu Academy of Agricultural Sciences, N0.50,
Zhongling street, Nanjing, 210014, China.

E-mail addresses of all authors, in order of authorship:

liyongteng19880213@163.com, liuxd@njau.edu.cn, wangna1214104@aliyun.com,

moonjaas@126.com, ary@unimelb.edu.au, guohfjaas@163.com

Telephone: +86-25-84390891

Running title: *Wolbachia*-mediated resistance in planthoppers

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/1462-2920.14974](https://doi.org/10.1111/1462-2920.14974)

Originality-Significance Statement

Heritable symbionts can be a major evolutionary force in eukaryote–prokaryote interactions in invertebrates by influencing host reproduction. They can also protect hosts against natural enemies and pathogens, but their effects on insecticide resistance are unclear. Our work provides evidence that a widespread endosymbiont, *Wolbachia*, may protect planthoppers against a chemical insecticide in some genetic backgrounds. To our knowledge, this is the first time an increase of insecticide resistance has been linked to an endosymbiont infection and background effects, with implications for pest control.

Summary

While facultative endosymbionts are now known to protect insect hosts against pathogens and parasitoids, the effects of endosymbionts on insecticide resistance are still unclear. Here we show that *Wolbachia* are associated with increased resistance to the commonly used insecticide, buprofezin, in the small brown planthopper (*Laodelphax striatellus*) in some genetic backgrounds while having no effect in other backgrounds. In three *Wolbachia*-infected lines from experimental buprofezin-resistant strains and one line from a buprofezin-susceptible line established from Chuxiong, Yunnan province, China, susceptibility to buprofezin increased after removal of *Wolbachia*. An increase in susceptibility was also evident in a *Wolbachia*-infected line established from a field population in Rugao, Jiangsu province. However, no increase was evident in two field populations from Nanjing and Fengxian, Jiangsu province, China. When *Wolbachia* was introgressed into different genetic backgrounds, followed by *Wolbachia* removal, the data pointed to *Wolbachia* effects that depend on the nuclear background as well as on the *Wolbachia* strain. However, there was no relationship between *Wolbachia* density and the component of buprofezin resistance associated with

the symbiont. The results suggest that *Wolbachia* effects associated with chemical resistance are complex and unpredictable, but also that they can be substantial.

Keywords: buprofezin resistance, *Laodelphax striatellus*, *Wolbachia*

Introduction

Wolbachia are extremely common facultative symbionts notably found in insect pests, and well known for their effects on host reproduction involving cytoplasmic incompatibility, parthenogenesis, feminization or male killing (O'Neill *et al.*, 1997; Werren *et al.*, 2008). In recent years, effects of *Wolbachia* other than reproductive effects have started to receive increasing attention, particularly through data collected from flies and mosquitoes. These studies have shown that some *Wolbachia* strains protect against some RNA viruses (Hedges *et al.*, 2008; Teixeira *et al.*, 2008; Bian *et al.*, 2010; Walker *et al.*, 2011; Martinez *et al.*, 2014), and that some *Wolbachia* strains provide fitness benefits in terms of life history traits, nutrition and other effects (Weeks *et al.*, 2007; Moriyama *et al.*, 2015; Zug *et al.* 2015). However, *Wolbachia* can also have the opposite effects on hosts both in terms of protection and life history benefits; for instance, a greater susceptibility to viruses has been noted for *Wolbachia* in

Spodoptera exempta (Graham *et al.*, 2012), and *Wolbachia* can have a range of deleterious effects on the life history of their hosts (Fleury *et al.* 2000; Ross *et al.* 2019).

The relationship between insecticide resistance and symbiont presence has been investigated in some studies. In *Culex pipiens*, there was a higher *Wolbachia* load in mosquitoes carrying an organophosphate resistant gene than in susceptible individuals lacking the gene, but the presence of *Wolbachia* did not affect the strength of resistance (Berticat *et al.*, 2002; Duron *et al.*, 2006). And in *Aedes aegypti*, the wMel strain of *Wolbachia* had no impact on resistance to several insecticides, namely the organophosphate temephos, the insect growth regulator s-methoprene, the pyrethroid bifenthrin, and the entomopathogen *Bacillus thuringiensis* var. *israelensis* (Endersby and Hoffmann, 2012). However, another endosymbiont, *Rickettsia*, was associated with increased susceptibility to some insecticides in *Bemisia tabaci* (Ghanim and Kontsedalov, 2009), and a bacterial symbiont of the genus *Burkholderia* from soil, acquired by the bean bug *Riptortus pedestris*, degraded the pesticide fenitrothion and conferred resistance when present in the insect gut (Kikuchi *et al.*, 2012).

The small brown planthopper (SBPH), *Laodelphax striatellus* (Fallén), is a notorious pest of rice crops, which not only causes direct damage by feeding, but also acts as a vector of plant viruses such as the rice stripe virus (Kisimoto, 1967). Buprofezin has been one of the most used insecticides to control SBPH in China for many years, but its frequent application has resulted in resistance problems (Gao *et al.*, 2007; Wang *et al.*, 2008a). *Wolbachia* is a common symbiont in *L. striatellus* (Noda *et al.*, 2001, Li *et al.*, 2018) and we have found that susceptibility of a *Wolbachia*-infected line originating

from the buprofezin-resistant strain increased when *Wolbachia* was cleared (Li *et al.*, 2018, Liu and Guo, 2019). Here we tested whether *Wolbachia* in fact played a role in mediating resistance to buprofezin.

To test for resistance effects associated with *Wolbachia*, we established pairs of *Wolbachia*-infected and *Wolbachia*-cured lines with the same genetic background from insecticide-resistant and insecticide-susceptible strains and field populations, and compared buprofezin susceptibility of *Wolbachia*-infected and *Wolbachia*-cured lines. We then investigated the density of *Wolbachia* in all the *Wolbachia*-infected lines. Because effects of genetic background were evident in the results, a series of backcrosses was undertaken to separate host nuclear background and *Wolbachia* strain effects on variation in resistance.

Results

Wolbachia infection partly associates with buprofezin resistance

To analyse whether the *Wolbachia* strain from the Chuxiong population of *L. striatellus* could protect hosts from insecticide-induced mortality, buprofezin susceptibility of the three *L. striatellus* lines infected with *Wolbachia* established from the selected strain (CX-R1, CX-R2 and CX-R3) were characterized, along with one *L. striatellus* line from the experimental buprofezin-susceptible strain (CX-S) and their paired lines without *Wolbachia* infection. In all three paired lines from the selected strain with the same genetic background from Chuxiong, there was a significant increase in susceptibility to 200 mg/L buprofezin after *Wolbachia* were removed from the *Wolbachia*-infected line (R1: $\chi^2=4.86$, $P=0.027$, Fig. 1A1 ; R2: $\chi^2=11.307$, $P=0.001$,

Fig. 1A2; R3: $\chi^2=6.009$, $P=0.014$, Fig. 1A3), whereas in paired lines from the control strain, there was no difference in susceptibility to 200 mg/L buprofezin after the removal of *Wolbachia* (S: $\chi^2=1.69$, $P=0.194$; Fig. 1A4) .

We next investigated resistance across *Wolbachia*-infected lines derived from field populations. SPBH with and without *Wolbachia* were challenged by spraying with a 1000 mg/L buprofezin suspension. The Rugao line (established from Rugao population of *L. striatellus*, RG line) infected with *Wolbachia* showed high survival, and the insects of the same genetic background without *Wolbachia* exhibited a significantly lower survival ($\chi^2=6.77$, $P=0.009$, Fig. 1B1), consistent with the above results for *Wolbachia*. However, protection was not observed in the Nanjing line (established from the Nanjing population of *L. striatellus*, NJ line) and Fengxian line (established from the Fengxian population of *L. striatellus*, FX line), with no significant difference in survival between the *Wolbachia*-infected and *Wolbachia*-cured lines (NJ: $\chi^2=0.241$, $P=0.632$, Fig. 1B2; FX: $\chi^2=3.468$, $P=0.063$, Fig. 1B3). Comparing the susceptibility among different field populations, we found that there was no significant differences in survival among *Wolbachia*-infected field populations ($F_{2,15} = 1.083$, $P=0.346$), but significant differences in survival among *Wolbachia*-cured field populations ($F_{2,15} = 4.905$, $P=0.022$).

Based on the survival rate, we further tested the interaction of host origin and infectious status in an overall analysis of data from selected populations or field populations. For selected populations from Chuxiong, there was no significant interaction, with buprofezin resistance only affected by main effects of *Wolbachia* or

host origin (Table 1). For field populations, both *Wolbachia* and host origin did not affect buprofezin resistance, but a significant interaction between *Wolbachia* and host origin was found (Table 1).

Table 1 Effects of *Wolbachia* infection on buprofezin resistance of *L. striatellus* in CX resistance-selected populations and field populations as assessed by survival. The degrees of freedom and the mean square are given along with the F-statistic and its associated probability (in bold when less than 0.05).

Factors	df	Mean square	F	P
CX selected populations				
<i>Wolbachia</i>	1	0.20	15.22	0.001
Host origin	2	0.25	20.09	0.002
<i>Wolbachia</i> * host origin	2	0.01	0.35	0.766
Deviation	30	0.02		
Field populations				
<i>Wolbachia</i>	1	0.02	1.79	0.191
Host origin	2	0.02	1.23	0.308
<i>Wolbachia</i> * host origin	2	0.05	3.81	0.034
Deviation	30	0.01		

Introgression suggests a role for *Wolbachia* variation and nuclear background variation in resistance

To investigate whether the variable *Wolbachia* effects described above reflect the nuclear background or uncharacterized *Wolbachia* differences, the infection from each line was introgressed into two different genetic backgrounds (Table 2). Comparisons of LC₅₀s showed that *Wolbachia* infections from Chuxiong and Rugao increased buprofezin resistance in their original hosts (CX: $t = 3.44$, $df = 6$, $P = 0.014$; RG: $t =$

4.23, $df = 6$, $P = 0.006$; Fig. 2A1, Fig. 2B1, Fig. S2A, Fig. S2B), and also when *Wolbachia* was introgressed into a Fengxian background (FX w^{CX} : $t = 9.31$, $df = 6$, $P < 0.001$; FX w^{RG} : $t = 2.47$, $df = 6$, $P = 0.049$; Fig. 2A2 and 2B2) even though native *Wolbachia* decreased resistance in this background (FX: $t = 2.65$, $df = 6$, $P = 0.038$; Fig. 2C1, Fig. S2C). Although *Wolbachia* infections from Nanjing did not affect resistance in their original hosts ($t = 0.86$, $df = 6$, $P = 0.418$; Fig. 2D1, Fig. S2D), they did when *Wolbachia* was introgressed into other host backgrounds (CX: $t = 2.74$, $df = 6$, $P = 0.034$; RG: $t = 5.28$, $df = 6$, $P = 0.002$; Fig. 2D2, Fig. 2D3, Fig. S2D) which suggested resistance mediated by native *Wolbachia*. In contrast, the *Wolbachia* infections from Fengxian decreased the LC_{50} of buprofezin in their original hosts (Fig. 2C1, Fig. S2C), but increased it following introgression of *Wolbachia* from Fengxian to CX and RG hosts (CX: $t = 2.75$, $df = 6$, $P = 0.034$; RG: $t = 12.55$, $df = 6$, $P < 0.001$; Fig. 2C2, Fig. 2C3, Fig. S2C).

To further test whether *Wolbachia* affected insecticide resistance differently on the genetic background of the hosts, we tested the interaction of genetic background and infection status in an overall analysis of data from each backcross comparison (Table 2). For *Wolbachia* from Chuxiong, no significant interaction was found (Table 3), with buprofezin resistance only affected by main effects of *Wolbachia* or genetic background (Fig. 2A). For *Wolbachia* from Rugao, a significant interaction between symbiont and genetic background was found (Fig. 2B, Table 3). For the other two comparisons, there were significant main effects as well as interaction effects (Table 3). Overall LC_{50} data from the parental strains are consistent with the mortality data, highlighting differences

in *Wolbachia* effects among strains, and the LC₅₀ data highlight that strain differences are both related to the *Wolbachia* (or other maternal effect) and also to the nuclear background of the strain (Table 3).

Table 2 Designation of lines in bioassay experiments

Expected nuclear genetic background	Infection status	<i>Wolbachia</i> source	Line designation
Chuxiong (S strain)	+	Chuxiong	CX-S _{w+}
Chuxiong (S strain)	-		CX-S _{w-}
Fengxian	+	Fengxian	FX _{w+}
Fengxian	-		FX _{w-}
Nanjing	+	Nanjing	NJ _{w+}
Nanjing	-		NJ _{w-}
Rugao	+	Rugao	RG _{w+}
Rugao	-		RG _{w-}
Fengxian 96.9%, Chuxiong (S strain) 3.1%	+	Chuxiong	FX _w ^{CX}
Nanjing 96.9%, Chuxiong (S strain) 3.1%	+	Chuxiong	NJ _w ^{CX}
Fengxian 96.9%, Rugao 3.1%	+	Rugao	FX _w ^{RG}
Nanjing 96.9%, Rugao 3.1%	+	Rugao	NJ _w ^{RG}
Chuxiong (S strain) 96.9%, Fengxian 3.1%	+	Fengxian	CX _w ^{FX}
Rugao 96.9%, Fengxian 3.1%	+	Fengxian	RG _w ^{FX}
Chuxiong (S strain) 96.9%, Nanjing 3.1%	+	Nanjing	CX _w ^{NJ}
Rugao 96.9%, Nanjing 3.1%	+	Nanjing	RG _w ^{NJ}

Table 3 Effects of *Wolbachia* infection on buprofezin resistance of *L. striatellus* in backcrosses as assessed by LC_{50s} (log transformed for analysis). The degrees of freedom and the mean square are given along with the F-statistic and its associated

probability (in bold when less than 0.05).

Factors	df	Mean square	F	P
CX <i>Wolbachia</i>				
<i>Wolbachia</i>	1	0.56	38.94	<0.001
Genetic background	2	2.41	168.13	<0.001
<i>Wolbachia</i> * genetic background	2	0.02	1.68	0.214
Deviation	18	0.01		
RG <i>Wolbachia</i>				
<i>Wolbachia</i>	1	0.04	11.57	0.001
Genetic background	2	0.27	1.70	0.209
<i>Wolbachia</i> * genetic background	2	0.30	13.01	<0.001
Deviation	18	0.02		
FX <i>Wolbachia</i>				
<i>Wolbachia</i>	1	0.63	32.55	<0.001
Genetic background	2	1.14	58.67	<0.001
<i>Wolbachia</i> * genetic background	2	0.47	24.25	<0.001
Deviation	18	0.02		
NJ <i>Wolbachia</i>				
<i>Wolbachia</i>	1	0.72	48.13	<0.001
Genetic background	2	0.91	60.99	<0.001
<i>Wolbachia</i> * genetic background	2	0.27	17.87	<0.001
Deviation	18	0.02		

Variation in *Wolbachia* density is not related to resistance

When *Wolbachia* from Chuxiong was introduced to FX and NJ populations of *L. striatellus*, line differences were evident ($F_{2,11} = 33.760$, $P < 0.0001$; Fig. 3A), with *Wolbachia* density significantly increasing in the NJ_w^{CX} line, but not changing in the FX_w^{CX} line. When *Wolbachia* from Rugao was introduced to FX and NJ populations, line differences were marginally non-significant ($F_{2,11} = 3.900$, $P = 0.06$; Fig. 3B), with

Wolbachia density significantly decreasing in the FX_w^{RG} line, and not being affected in the NJ_w^{RG} line. Line differences were also detected ($F_{2,11} = 19.794$, $P=0.001$; Fig. 3C) when *Wolbachia* from Fengxian was introduced to the CX and RG populations, with *Wolbachia* density increasing in the CX_w^{FX} line. Finally, there were line differences ($F_{2,11} = 35.710$, $P<0.0001$; Fig. 3D) when *Wolbachia* from Nanjing was introduced to the CX and RG populations of *L. striatellus*, with *Wolbachia* density decreasing in the CX_w^{NJ} line, and increasing in the RG_w^{NJ} line.

These differences in *Wolbachia* density did not associate clearly with changes in buprofezin resistance (Fig. 4); in several cases such as in the *Wolbachia* from FX in the FX, CX and RG backgrounds, there was no direct connection between density in the crosses and the extent to which *Wolbachia* increased resistance (Fig. 3C). *Wolbachia* increased resistance even when present at a relatively low density, and vice versa.

Discussion

While *Wolbachia* has been shown to provide insects with protection from some entomopathogens including viruses and fungi (Panteleev *et al.*, 2007; Teixeira *et al.*, 2008; Osborne *et al.*, 2009), this is one of the few cases where *Wolbachia* has been implicated in providing protection to a chemical insecticide. In mosquitoes, no obvious effects of *Wolbachia* on resistance were found (Endersby and Hoffmann, 2012). Results from other endosymbionts seem to point to greater susceptibility to insecticides rather than greater resistance, such as in whiteflies where the presence of *Rickettsia* increased susceptibility to four out of six insecticides (Kontsedalov *et al.*, 2008), and where double infections of *Rickettsia*–*Arsenophonus* and *Wolbachia*–*Arsenophonus* which

carried higher amounts of symbionts overall also showed increased susceptibility to three out of the six insecticides when compared to strains infected with *Arsenophonus* alone (Ghanim and Kontsedalov, 2009). Finally, in brown planthoppers a distinct strain of *Arsenophonus* decreased insecticide resistance (Pang *et al.*, 2018).

In our study we found that *Wolbachia* was associated with resistance to buprofezin in some lines of small brown planthopper, but also that this association was not always evident and depended on the combination of *Wolbachia* and nuclear backgrounds tested. While the pattern appeared clear in lines from Chuxiong regardless of their selection history, we only established this pattern in one of three *Wolbachia*-infected lines from Jiangsu province. This line difference had a complicated basis involving either the nuclear or *Wolbachia* backgrounds. We could not distinguish between *Wolbachia* strains based on a standard gene set (Fig. S1), but the different behavior of the *Wolbachia* strains may reflect other genomic differences that were not detected here and that likely will require a full genomic comparison of the *Wolbachia* which may reveal other DNA based differences (Chrostek and Teixeira, 2015). In addition, there might be an interaction between *Wolbachia* and other maternally inherited components, notably mitochondria, which hitchhike along with *Wolbachia* in populations as they spread (Hale and Hoffmann, 1990) and can interact with *Wolbachia* in influencing life history traits (Dean, 2006).

Buprofezin has been developed as an insecticide for more than 30 years (Nagata, 1986), and it has been intensively used against rice planthoppers in agricultural fields in China for more than 20 years. Buprofezin resistance in SBPH is serious in China in

recent years, and mechanisms have been attributed to evolutionary changes in pest insect genomes such as alteration of drug target sites, up-regulation of degrading enzymes, and enhancement of drug excretion (Gao *et al.*, 2007; Wang *et al.*, 2008a; Wang *et al.*, 2008b). A single P450 gene, CYP6CW1 plays a role in the resistance of *L. striatellus* to buprofezin (Zhang *et al.*, 2012). We have found that the RG population, which had the longest history of buprofezin exposure, showed significantly higher survival in *Wolbachia*-infected individuals, and it was also the only population to show an effect of the *Wolbachia* infection and P450 expression levels. However, we have not found this gene to be consistently upregulated in comparisons of cured and *Wolbachia* infected lines that expressed increased resistance versus no effect (Fig. S3). We also failed here to show an association between *Wolbachia* density and resistance effects, even though density has to some extent been previously associated with fitness-related phenomena in *Wolbachia* (Hoffmann *et al.*, 2015), including a correlation between the density of *Wolbachia* and the extent of antiviral protection in *Drosophila*: the three *Wolbachia* strains of wMel, wRi and wAu, which provide strong antiviral protection, were significantly more abundant than the strains of wHa and wNo which did not provide protection (Osborne *et al.*, 2009).

The mechanism involved in insecticide resistance related to *Wolbachia* remains unclear; it might be related to changes in metabolism, expression changes particularly in genes involved in detoxification, or *Wolbachia*-related triggering of immune responses in the insect hosts. The association between resistance and *Wolbachia* appears complex given the line differences in *Wolbachia* effects. Although these line differences may be

mediated by other symbionts, we did not find differences among lines for the presence of several symbiont groups previously described from planthoppers. *Wolbachia*-associated resistance adds yet another relevant factor when considering the distribution of *Wolbachia* within and across pest populations. Chemical insecticides are stressors that insect pests often face, and there are likely to be other cases where endosymbionts are associated with resistance. Such effects may need to be considered when introducing *Wolbachia* into natural populations for pest and disease control.

Conclusions

We found that *Wolbachia*, common facultative symbionts found in insect pests, can be associated with increased resistance to the widely used insecticide buprofezin in the small brown planthopper in some genetic backgrounds while having no effect in others. The data supported the notion that *Wolbachia* effects on insecticide resistance could depend on the nuclear background as well as on the *Wolbachia* strain or other maternal factors, which suggest that *Wolbachia* effects on chemical resistance are complex and unpredictable, but also that they can be substantial.

Methods

Small brown planthoppers and *Wolbachia*

Wolbachia-infected small brown planthoppers were obtained from field populations in Rugao (RG), Fengxian (FX), Nanjing (NJ), Jiangsu province, China and Chuxiong (CX), Yunnan province, China. The CX population had been collected from Yunnan in July 2001 (at that time, buprofezin was rarely used), and reared in the laboratory without contact with any insecticides. A buprofezin-resistant strain was developed from this

susceptible strain through 32 generations of selection (Zhang *et al.*, 2012). Both the buprofezin-resistant strain and buprofezin-susceptible strain from the CX population were used in experiments. The other three populations were collected in May, 2014. The insecticide exposure history was different among the three regions, buprofezin has been widely used in RG since 1990 and in NJ since 2001, while it is rarely used in FX. All populations were infected by *Wolbachia*.

All the strains and lines were reared at the same density on rice seedlings hosts under the same environmental conditions. For maintaining each line, about 2000 7-day-old seedlings were randomly assigned to lines in a cage (length 29.5cm×width 20cm×height 18cm), and there were 1800 *L. striatellus* individuals in each cage. At the time of the experiment, we set up multiple cages per line to rear the insects, so the experimental subjects were combined across more than one cage. Seedlings were replaced by new seedlings every week. The insects were reared at a constant temperature of 27 (±1) °C with a photoperiod of 14:10 h light: dark.

Preparation of *Wolbachia*-infected and *Wolbachia*-cured lines

Before the establishment of lines, three isofemale lines of *L. striatellus* with different levels of resistance from the buprofezin-selected strain were established. Each line was developed from a pair of newly emerged female and male adults randomly selected from the resistant strain to minimize variation in genetic background within each strain. The same procedure was followed to establish lines from one buprofezin-susceptible strain from the Yunnan population and the three field populations from Jiangsu.

In each line, the presence of all symbionts known to be potentially present in

Author Manuscript

planthoppers, including *Wolbachia*, *Arsenophonus*, *Cardinium hertigii*, *Acinetobacter*, *Chryseobacterium*, *Serratia*, *Arthrobacter* and *Spiroplasma*, was checked in adult insects as previously described (*Wolbachia*: Zhou et al., 1998; *Arsenophonus* Thao et al., 2004; *Cardinium*: Nakamura et al., 2009; *Acinetobacter*: Vanbroekhoven 2004; *Chryseobacterium*: Alonso, 2007; *Serratia* : Zhu 2008; *Arthrobacter* : Koch, 1994; *Spiroplasma*: Sanada-Morimura, 2013). Both the females and males used to establish lines were only infected with *Wolbachia*. Their offspring were used to identify the *Wolbachia* infection through molecular markers (see below).

To generate SBPH lines free of *Wolbachia*, each *Wolbachia*-infected planthopper line was treated with 0.2% tetracycline by oral ingestion (Noda et al., 2001). One hundred newly hatched SBPH were fed rice seedlings sprayed with a 0.2% tetracycline solution. The rice seedlings were treated with antibiotics every 3 days, and new seedlings were supplied at these times. SBPH were treated with antibiotics over the entire nymphal stage for two generations, and newly emerged SBPH were then paired. After egg production, the presence of *Wolbachia* was checked in adult insects. Only female and male offspring without *Wolbachia* were kept. Before bioassays were performed, insects were reared for more than five generations without exposure to antibiotics. The symbiont infection status of the *Wolbachia*-infected and *Wolbachia*-cured lines was reconfirmed prior to experiments by diagnostic PCR (as specified above) and quantitative PCR (as specified below).

A further eight massbred lines were created by introgressing *Wolbachia* infections from each of the three locations into the genetic background of the other locations, by mating

50+ virgin infected females with an equivalent number of *Wolbachia*-cured males, and then using repeated backcrossing for five generations to ensure that the nuclear background was (a predicted) 96.9% of the one from which insects were sourced for backcrossing (Kriesner *et al.*, 2016). The backcrosses were only done with males that were from a stock treated with antibiotics. Males only transfer the nuclear genome, not the endosymbionts or mitochondrial genomes. Thus the “native” components remain intact with respect to the *Wolbachia* and associated backgrounds. All mass-bred lines are listed in Table 2.

Evaluation of buprofezin resistance

Wolbachia-cured and *Wolbachia*-infected lines of *L. striatellus* were used to compare susceptibility to buprofezin (98%, Jiangsu Anpon Electrochemical CO., LTD, China) using the rice seedling dipping method (Wang *et al.*, 2018a). Before the bioassay, to validate the *Wolbachia* infection of the lines, at least 20 individuals were randomly collected from each tested line for diagnostic detection of *Wolbachia*. To test whether there was an effect of tetracycline treatment on gut bacteria of *L. striatellus*, 10 individuals were randomly collected from each tested line, and diagnostic detection of the gut bacteria including *Enterobacter* sp. and *Enterococcus* sp. was conducted using previously described methods (Xia *et al.*, 2018).

In an initial assay to compare susceptibility of lines to buprofezin, all paired *Wolbachia*-cured and *Wolbachia*-infected lines prior to introgression were treated with one diagnostic concentration of buprofezin to compare mortality. In a later assay, the LC_{50s} of all introgressed lines were estimated based on a series of 5-6 concentrations of

buprofezin. In the bioassay procedure, five rice seedlings were immersed in buprofezin solution for 10 s, before the plants were dried at 25 (± 1) °C. The seedlings were placed in a plastic cup, and about fifteen 3rd instar nymphs were added. For each line, nymphs treated with distilled water were used as controls. All experiments were carried out at 27 (± 1) °C and with a photoperiod of 14:10 h light: dark. Insect mortality was recorded every 24 h over a period of 144 h. For survivorship comparisons based on a single concentration (lines from Chuxiong: 200 mg/L, lines from other sites: 1000 mg/L), each line treated with buprofezin was replicated 6 times (6 cups), while each line treated with water was replicated 4 times (4 cups). For LC₅₀ comparisons, there were 4 replicates (4 cups) at each concentration and four LC₅₀ estimates associated with *Wolbachia* source were obtained from linear regressions of log concentration against mortality (expressed as probits).

Analysis of *Wolbachia* density

Real-time quantitative PCR was performed with an ABI StepOne Real-Time PCR System (Applied Biosystems Inc, Foster City, CA, USA) to measure the density of *Wolbachia*. For each line, a total of 30 3rd instar nymphs of *L. striatellus* were collected as one sample, and the DNA was extracted with a Wizard® Genomic DNA Purification Kit (Promega). The primers for the reaction were the sense primer (*Wsp*-F) 5'-ATGTA ACTCCAGAAATCAA ACTC-3' and the antisense primer (*Wsp*-R) 5'-GATACCAGCATCATCCTTAGC-3'. The 20 µL RT-PCR reaction system consisted of 10 µL SYBR® Premix Ex Taq (Tli RNaseH Plus) (2X) (Takara, Japan), 0.4 µL forward and 0.4µL reverse primers, 0.4 µL ROX Reference Dye, 2 µL DNA and

6.8 μ L ddH₂O. The RT-PCR conditions were 95°C for 30 s, followed by 40 cycles of 95°C for 5 s and 60°C for 31 s, and then 95°C for 15 s, 60°C for 1 min, and 95°C for 15 s. The standard curve of Real-time fluorescent quantitative PCR of the *Wolbachia* *wsp* gene was built to determinate *Wolbachia* *wsp* gene copy number as described previously for *L. striatellus* (Zhou et al., 2010). Note that no host gene was included but individuals were compared at the same life stage and had been reared under the same conditions. Three technical replicates were carried out for each sample, and four biological replicates were run for each line.

Statistics

All data were checked for normality and homogeneity of variances by Shapiro-Wilk tests before analysis, and LC₅₀s were log transformed to reduce differences in variance between treatments. Survival curves between *Wolbachia*-infected and *Wolbachia*-cured lines in the presence of buprofezin were compared using Kaplan-Meier analyses and log-rank tests. LC₅₀ of paired lines (*Wolbachia*-infected and *Wolbachia*-cured line with the same host nuclear background) were compared by t-tests, and LC₅₀ of all lines with the same *Wolbachia* source and paired *Wolbachia*-cured lines were compared by two-way ANOVA, with *Wolbachia* and genetic background as the factors in the analysis. Density among different lines was also compared with ANOVAs. Pearson's correlation was computed to test the association between the ratio of LC₅₀ values for infected individuals over uninfected individuals and *Wolbachia* density. IBM Statistics (SPSS 19.0) software was used for these statistical analyses.

Acknowledgements

We are grateful to Dongxiao Zhao for building the unrooted species tree and Zhichun Zhang for revising the figure. We thank Baohua Qiu and Zongquan Liu at the Plant Protection Station of Fengxian County, Jiangsu Province, China, for collecting the small brown planthopper, *Laodelphax striatellus*.

Funding

This research was supported by grants from the National Natural Science Foundation of China (31672027, 31972265) and the Independent Innovation Fund of Agricultural Science and Technology in Jiangsu province, China (cx(16)1001). AAH was partly supported by a Fellowship from the National Health and Medical Research Council.

Author contributions

HG conceived the experiments. HG and XL designed the experiments. YL, NW and YZ collected the data. HG, YL and AH analysed the data. HG, AH and YL wrote and reviewed the manuscript.

Competing interests

We have no competing interests.

Data accessibility statement

We agree to archive the data associated with this manuscript.

References

- Alonso, C., Warnecke, F., Amann, R. and Pernthaler, J. (2007) High local and global diversity of Flavobacteria in marine plankton. *Environ Microbiol* **9**: 1253-1266.
- Baldo, L., Dunning Hotopp, J.C., Jolley, K.A., Bordenstein, S.R., Biber, S.A., Choudhury, R.R., et al. (2006) Multilocus sequence typing system for the

- endosymbiont *Wolbachia pipientis*. Appl Environ Microbiol **72**:7098–7110.
- Berticat, C., Rousset, F., Raymond, M., Berthomieu, A. and Weill, M. (2002) High *Wolbachia* density in insecticide-resistant mosquitoes. Proc R Soc Lond B **269**: 1413–1416.
- Bian, G.W., Xu, Y., Lu, P., Xie, Y. and Xi, Z.Y. (2010) The endosymbiotic bacterium *Wolbachia* induces resistance to dengue virus in *Aedes aegypti*. PLoS Pathog doi:org/10.1371/journal.ppat.1000833.
- Chrostek, E. and Teixeira, L. (2015) Mutualism breakdown by amplification of *Wolbachia* genes. PLoS Biol doi:org/10.1371/journal.pbio.1002065.
- Dean, M.D. (2006) A *Wolbachia*-associated fitness benefit depends on genetic background in *Drosophila simulans*. Proc R Soc Lond B **273**: 1415–1420.
- Duron, O., Labbé, P., Berticat, C., Rousset, F., Guillot, S., Raymond, M. And Weill, M. (2006) High *Wolbachia* density correlates with cost of infection for insecticide resistant *Culex pipiens* mosquitoes. Evolution **60**: 303–314.
- Endersby, N.M. and Hoffmann, A.A. (2012) Effect of *Wolbachia* on insecticide susceptibility in lines of *Aedes aegypti*. Bull Entomol Res **103**: 269–277.
- Fleury, F., Vavre, F., Ris, N., Fouillet, P. and Boulétreau, M. (2000) Physiological cost induced by the maternally-transmitted endosymbiont *Wolbachia* in the *Drosophila* parasitoid *Leptopilina heterotoma*. Parasitology **121**: 493–500.
- Gao, B., Wu, J., Huang, S., Mu, L. and Han, Z. (2007) Insecticide resistance in field populations of *Laodelphax striatellus* Fallén (Homoptera: Delphacidae) in China and its possible mechanisms. Int J Pest Manag **54**: 13–19.

- Ghanim, M. and Kontsedalov, S. (2009) Susceptibility to insecticides in the Q biotype of *Bemisia tabaci* is correlated with bacterial symbiont densities. *Pest Manag Sci* **65**: 939–942.
- Graham, R.I., Grzywacz, D., Mushobozi, W.L. and Wilson, K. (2012) *Wolbachia* in a major African crop pest increases susceptibility to viral disease rather than protects. *Ecol Lett* **15**: 993–1000.
- Hale, L.R. and Hoffmann, A.A. (1990) Mitochondrial DNA polymorphism and cytoplasmic incompatibility in natural populations of *Drosophila simulans*. *Evolution* **44**: 1383–1386.
- Hedges, L.M., Brownlie, J.C., O'Neill, S.L. and Johnson, K.N. (2008) *Wolbachia* and virus protection in insects. *Science* **322**: 702.
- Hoffmann, A.A., Ross, P.A. and Rašić, G. (2015) *Wolbachia* strains for disease control: ecological and evolutionary considerations. *Evol Appl* **8**: 751–768.
- Li, Y.T, Liu, X.D. and Guo, H.F. (2018) Variations in endosymbiont infection between buprofezin-resistant and susceptible strains of *Laodelphax striatellus* (Fallén). *Current Microbiol* doi: 10.1007/s00284-018-1436-x.
- Liu, X. and Guo, H.F. (2019) Importance of endosymbionts *Wolbachia* and *Rickettsia* in insecticide resistance development. *Current Opinion in Insect Science* **33**: 84–90.
- Kikuchi, Y., Hayatsu, M., Hosokawa, T., Nagayama, A., Tago, K. and Fukatsu, T. (2012) Symbiont-mediated insecticide resistance. *Proc Natl Acad Sci USA* **109**: 8618–8622.
- Kisimoto, R. (1967) Genetic variation in the ability of a planthopper vector; *Laodelphax striatellus* (Fallén) to acquire the rice stripe virus. *Virology* **32**: 144–512.

- Koch, C., Rainey, F.A. and Stackebrandt, E. (1994) 16S rDNA studies on members of *Arthrobacter* and *Micrococcus*: An aid for their future taxonomic restructuring. FEMS Microbiol Lett **123**: 167–171.
- Kontsedalov, S., Zchori-Fein, E., Chiel, E., Gottlieb, Y., Inbar, M. and Ghanim, M. (2008) The presence of *Rickettsia* is associated with increased susceptibility of *Bemisia tabaci* (Homoptera: Aleyrodidae) to insecticides. Pest Manag Sci **64**: 789–792.
- Kriesner, P., Conner, W.R., Weeks, A.R., Turelli, M. and Hoffmann, A.A. (2016) Persistence of a *Wolbachia* infection frequency cline in *Drosophila melanogaster* and the possible role of reproductive dormancy. Evolution **70**: 979–997.
- Martinez, J., et al. (2014) Symbionts commonly provide broad spectrum resistance to viruses in insects: a comparative analysis of *Wolbachia* strains. PLoS Pathog. doi:org/10.1371/journal.ppat.1004369.
- Moriyama, M., Nikoh, N., Hosokawa, T. and Fukatsu, T. (2015) Riboflavin provisioning underlies *Wolbachia*'s fitness contribution to its insect host. mBio doi: 10.1128/mBio.01732-15.
- Nagata, T. (1986) Timing of buprofezin application for control of the brown planthopper, *Nilaparvata lugens* Stal (Homoptera: Delphacidae). Appl Entomol Zool **21**: 357–362.
- Nakamura, Y. et al. (2009) Prevalence of *Cardinium* bacteria in planthoppers and spider mites and taxonomic revision of “*Candidatus Cardinium hertigii*” based on detection of a new *Cardinium* group from biting midges. Appl Environ Microbiol **75**: 6757–6763.
- Noda, H., Koizumi, Y., Zhang, Q. and Deng, K. (2001) Infection density of *Wolbachia*

and incompatibility level in two planthopper species, *Laodelphax striatellus* and *Sogatella furcifera*. Insect Biochem Mol Biol **31**: 727–737.

O'Neill, S.L., Werren, J.H., Hoffmann, A.A. Influential passengers: inherited microorganisms and arthropod reproduction. UK: Oxford University Press. 1997.

Osborne, S.E., Leong, Y.S., O'Neill, S.L. and Johnson, K.N. (2009) Variation in antiviral protection mediated by different *Wolbachia* strains in *Drosophila simulans*. PLoS Pathog doi:org/10.1371/journal.ppat.1000656.

Pang, R., Chen, M., Yue, L., Xing, K., Li, T., Kang, K., Liang, Z., Yuan, L. and Zhang, W. (2018) A distinct strain of *Arsenophonus* symbiont decrease insecticide resistance in its insect host. PLoS Genetics **14**: e1007725.

Pantelev, D.I., Goriacheva, I.I., Andrianov, B.V., Reznik, N.L., Lazebnyĭ, O.E. and Kulikov, A.M. (2007) The endosymbiotic bacterium *Wolbachia* enhances the nonspecific resistance to insect pathogens and alters behavior of *Drosophila melanogaster*. Russ J Genet **43**: 1066–1069.

Ross, P.A., Turelli, M. and Hoffmann, A.A. (2019) Evolutionary ecology of *Wolbachia* releases for disease control. Ann Rev Genet **53**: 93-116.

Sanada-Morimura, S., Matsumura, M. and Noda, H. (2013) Male killing caused by a *Spiroplasma* dymbiont in the small brown planthopper, *Laodelphax striatellus*. J Hered doi:10.1093/jhered/est052.

Teixeira, L., Ferreira, Á. and Ashburner, M. (2008) The bacterial symbiont *Wolbachia* induces resistance to RNA viral infections in *Drosophila melanogaster*. PLoS Biol doi:org/10.1371/journal.pbio.1000002.

- Thao, M.L. and Baumann, P. (2004) Evidence for multiple acquisition of *Arsenophonus* by whitefly species (Sternorrhyncha: Aleyrodidae). *Curr Microbiol* **48**: 140–144.
- Vanbroekhoven, K., Ryngaert, A., Wattiau, P., De Mot, R. and Springael, D. (2004) *Acinetobacter* diversity in environmental samples assessed by 16S rRNA gene PCR–DGGE fingerprinting. *FEMS Microbiol Ecol* **50**: 3750.
- Walker, T., et al. (2011) The *wMel* *Wolbachia* strain blocks dengue and invades *Aedes aegypti* populations. *Nature* **476**: 450–453.
- Wang, L.H., Fang, J.C. and Liu, B.S. (2008b) Relative toxicity of insecticides to *Laodelphax striatellus* (Fallén) (Homoptera: Delphacidae) and the resistance of field populations from different areas of East China. *Acta Entomol Sin* **51**: 930–937.
- Wang, Y., Gao, C., Xu, Z., Zhu, Y.C., Zhang, J., Li, W., Dai, D., Lin, Y., Zhou, W. and Shen, J. (2008a) Buprofezin susceptibility survey, resistance selection and preliminary determination of the resistance mechanism in *Nilaparvata lugens* (Homoptera: Delphacidae). *Pest Manag Sci* **64**: 1050–1056.
- Weeks, A.R., Turelli, M., Harcombe, W.R., Reynolds, K.T. and Hoffmann, A.A. (2007) From parasite to mutualist: rapid evolution of *Wolbachia* in natural populations of *Drosophila*. *PLoS Biol* doi:org/10.1371/journal.pbio.0050114.
- Werren, J.H., Baldo, L. and Clark, M.E. (2008) *Wolbachia*: master manipulators of invertebrate biology. *Nat Rev Microbiol* **6**: 741–751.
- Xia, X., Sun, B., Gurr, G.M., Vasseur, L., Xue, M. and You, M. (2018) Gut microbiota mediate insecticide resistance in the diamondback moth, *Plutella xylostella* (L.). *Frontiers Microbiol* doi: 10.3389/fmicb.2018.0002.

- Zhang, Y.L., Guo, H.F., Yang, Q., Li, S., Wang, L.H., Zhang, G.F. and Fang, J.C. (2012) Overexpression of a P450 gene (*CYP6CW1*) in buprofezin-resistant *Laodelphax striatellus* (Fallén). *Pestic Biochem Phys* **104**: 277–282.
- Zhou, L.L., Zhang, K.J., Song, Z.W. and Hong, X.Y. (2010) Relationship of WO phage and *Wolbachia* infection in *Laodelphax striatellus* (Fallen) (Hemiptera: Delphacidae). *Acta Entomol Sin* **53**: 978–984.
- Zhou, W.G., Rousset, F. and O'Neill, S. (1998) Phylogeny and PCR-based classification of *Wolbachia* strains using *wsp* gene sequences. *Proc R Soc Lond B* **265**: 509–515.
- Zhu, H., Sun, S.J. and Dang, H.Y. (2008) PCR detection of *Serratia* spp. using primers targeting *pfs* and *luxS* genes involved in AI-2-dependent quorum sensing. *Curr Microbiol* **57**: 326–330.
- Zug, R. and Hammerstein, P. (2015) Bad guys turned nice? A critical assessment of *Wolbachia* mutualisms in arthropod hosts. *Biol Rev Camb Philos Soc* **90**: 89–111.

Fig. 1 Survivorship of *Wolbachia*-infected and *Wolbachia*-cured *L. striatellus* lines exposed to buprofezin (black - buprofezin; gray - water). Error bars represent standard error of the mean calculated from six (chemical) or four (water control) replicates of 15 nymphs scored for survival at each concentration. Black lines refer to survival of *L. striatellus* exposed to buprofezin, gray lines refer to survival of *L. striatellus* exposed to water. (A1-A4) Comparison of the survival of four *L. striatellus* lines from selected and control strains exposed to 200 mg/L buprofezin. (B1-B4) Comparison of the survival of lines derived from field populations from Nanjing, Fengxian and Rugao exposed to 1000 mg/L buprofezin. Log rank test on Kaplan-Meier curves show the survival of *Wolbachia*-infected and *Wolbachia*-cured *L. striatellus* is significantly different among paired R1 lines ($P<0.05$), R2 lines ($P<0.01$), R3 lines ($P<0.01$) and RG lines ($P<0.01$), and is not significantly different among paired S, NJ and FX lines (all $P>0.05$) (* $P<0.05$, ** $P<0.01$).

Fig. 2 Relationship between mortality (expressed as probits) and log concentration of buprofezin (mg/L) in *Wolbachia*-introgressed and *Wolbachia*-cured *L. striatellus* lines with different genetic backgrounds. Values are based on four replicates of 15 nymphs per concentration and LC50 values (probit = 5) were computed separately for each replicate. Lines represent linear regressions based on average data points. (A) FX and NJ lines introgressed with *Wolbachia* from Chuxiong. (B) FX and NJ lines introgressed with *Wolbachia* from Rugao. (C) CX and RG lines introgressed with *Wolbachia* from Fengxian. (D) CX and RG lines introgressed with *Wolbachia* from Nanjing (* $P<0.05$, ** $P<0.01$, *** $P<0.001$).

Fig. 3 Comparison of *Wolbachia* density in the *Wolbachia*-introgressed and original lines of *L. striatellus*. (A) CX line and FX, NJ introgressed lines. (B) RG and FX, NJ introgressed lines. (C) FX and CX, RG introgressed lines. (D) NJ and CX, RG introgressed lines. The *Wolbachia* density was expressed as copy number of *wsp* gene per individual.

Fig. 4 Relationship between density and protection of *Wolbachia* in *L. striatellus*. No significant relationship was found between *Wolbachia* density and the ratio of LC₅₀ values for infected individuals over uninfected individuals.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Li, Y;Liu, X;Wang, N;Zhang, Y;Hoffmann, AA;Guo, H

Title:

Background-dependent Wolbachia-mediated insecticide resistance in *Laodelphax striatellus*

Date:

2020-07-01

Citation:

Li, Y., Liu, X., Wang, N., Zhang, Y., Hoffmann, A. A. & Guo, H. (2020).
Background-dependent Wolbachia-mediated insecticide resistance in *Laodelphax striatellus*. ENVIRONMENTAL MICROBIOLOGY, 22 (7), pp.2653-2663. <https://doi.org/10.1111/1462-2920.14974>.

Persistent Link:

<http://hdl.handle.net/11343/275523>