DR. RAHUL BARMANRAY (Orcid ID : 0000-0002-1433-2239)

DR. MERVYN KYI (Orcid ID : 0000-0002-4341-8250)

Article type : Letters to the editor (comment)

Article begins on page three of this document.

Title	Routine glucose assessment in the emergency department for detecting unrecognised diabetes: a cluster randomised trial
--------------	--

Authors:

	Title	First name	Mid inits	Last name	Postnom (eg, PhD) [3 only for publication]	Position1	Address1	Position2	Address2	Tel	Email
1	Assoc. Prof.	Spiros		Fourlanos	MBBS, FRACP, PhD	Director, Dept Diabetes & Endocrinology	1	Clinical Associate Professor	2	03 9342 7365	spiros.fourlanos@mh.org.au
2	Dr.	Rahul		Barmanray	MBBS(Hons), BMedSci	Endocrinology Advanced Trainee	1		2	03 9342 7000	rbarmanray@gmail.com
3	Dr.	Mervyn		Kyi	MBBS(Hons), FRACP, PhD, BMedSci	Endocrinologist	1		2	03 9342 7365	mervyn.kyi@mh.org.au
4											
5											

Number of corresponding author:	2
Number of alternative corresponding author:	

Addresses:

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/MJA2.50550](#)

This article is protected by copyright. All rights reserved

	Institution		City	State	Post Code	
1	Royal Melbourne Hospital		Melbourne	VIC	3050	
2	University of Melbourne		Melbourne	VIC	3050	
3						
4						
5						

Postal address of first corresponding author (if different from the institutional address given above)	
--	--

Primary Keywords [Office use only]	Endocrine system diseases; Health occupations
Secondary keywords [Office use only]	Diabetes mellitus, type 2; Hospital medicine
Notes:	

Article details (press ctrl – 9 to enter details):

Article type	Letter
Blurb	
Pullquote	

Office use

<i>Ms. Number</i>	mja19.01200
<i>Medical editor</i>	Christine Gee
<i>Medical editor email</i>	cgee@mja.com.au
<i>Structural editor</i>	Laura Teruel
<i>Structural editor email</i>	lteruel@mja.com.au
<i>Section/Category</i>	Letters to the editor (comment)
<i>Strapheading</i>	Letter
<i>Substrap</i>	

Wiley – file data:

Filename for copyediting	bar_mja19.01200_ms.docx
Accompanying graphics	None
Stock images	None
Appendices	None

Office use – history:

Event	Date
Original submission received	26/11/2019

Event	Date
Accept	16/01/2020

Proof sent to author	
Proof returned by author	
Published (date format xx/xx/xx)	
Issue	
Vol	
DOI	
Journal	
Original article DOI (for response)	

Routine glucose assessment in the emergency department for detecting unrecognised diabetes: a cluster randomised trial

TO THE EDITOR: We congratulate Cheung and colleagues¹ on their large cluster randomised trial of routine blood glucose and automated glycated haemoglobin (HbA_{1c}) testing in emergency departments. This trial reaffirmed the high prevalence of unrecognised diabetes in patients presenting to the emergency department, while demonstrating the feasibility of algorithmic detection. However, the rate of documented follow-up plans in patients with suspected or newly diagnosed diabetes was low and did not benefit from the trial intervention. Cheung and colleagues¹ and Hare and Shaw,² in their accompanying editorial, suggest that this may relate to diabetes services already operating at full capacity or to overburdened staff documenting abbreviated plans at discharge. The trial highlights the difficulty in improving outcomes when multiple non-integrated health professionals manage a condition and, hence, the importance of continuity of care.

The RAPIDS trial³ was an early intervention model of care consisting of integrated continuous acute diabetes care provided by a dedicated, proactive specialist inpatient diabetes team (IDT). The intervention involved an IDT using a networked blood glucose meter system to remotely identify inpatients with diabetes (known and newly diagnosed) to directly manage these patients, compared with usual care, where diabetes management was mostly provided by parent unit teams.³ This trial showed that direct diabetes management by a dedicated IDT improved glycaemia and decreased the rate of hospital-acquired infections.

During the RAPIDS trial, in patients with newly discovered hyperglycaemia (random capillary glucose > 11.1 mmol/L without known diabetes), treatment and follow-up plans were documented in 11/34 patients (33%) with usual care, comparable to findings by Cheung et al. However, with the IDT intervention, 22/28 patients (79%) had treatment and follow-up plans. Similarly, in patients with newly diagnosed diabetes (HbA_{1c} ≥ 6.5%), diabetes treatment was commenced in 8/17 patients (47%) with usual care, and in 11/12 patients (92%) with IDT intervention³ (unpublished data). It is likely that the presence of an IDT at one of the control hospitals in the trial by Cheung and colleagues contributed significantly to the improved plan documentation in that arm.

We thus echo the editorial and professional society voices asserting the importance of resourcing clinical services for diabetes in Australian hospitals.⁴ Establishing IDTs in our hospitals will enable excellent diabetes care despite the increasing prevalence of this disease in Australia.

Competing interests: No relevant disclosures.

Author details

Spiros Furlanos^{1,2}

Rahul Barmanray^{1,2}

Mervyn Kyi^{1,2}

1 Royal Melbourne Hospital, Melbourne, VIC.

2 University of Melbourne, Melbourne, VIC.

rahul.barmarney@mh.org.au

doi: 10.5694/mja19.01200

References

- 1 Cheung NW, Campbell LV, Fulcher GR, et al. Routine glucose assessment in the emergency department for detecting unrecognised diabetes: a cluster randomised trial. *Med J Aust* 2019; 211: 454-459. <https://www.mja.com.au/journal/2019/211/10/routine-glucose-assessment-emergency-department-detecting-unrecognised-diabetes>
- 2 Hare MJL, Shaw JE. Inpatient diabetes care requires adequate support, not just HbA_{1c} screening. *Med J Aust* 2019; 211: 452-453. <https://www.mja.com.au/journal/2019/211/10/inpatient-diabetes-care-requires-adequate-support-not-just-hba1c-screening>
- 3 Kyi M, Colman PG, Wraight PR, et al. Early intervention for diabetes in medical and surgical inpatients decreases hyperglycemia and hospital-acquired infections: a cluster randomized trial. *Diabetes Care* 2019; 42: 832-840.
- 4 Cheung NW, Chippis D, Cornelius S, et al. Australian Diabetes Society guidelines for routine glucose control in hospital. Australian Diabetes Society, 2012. <https://diabetessociety.com.au/documents/ADSGuidelinesforRoutineGlucoseControlinHospitalFinal2012.pdf> (viewed Nov 2019).

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Fourlanos, S;Barmanray, R;Kyi, M

Title:

Routine glucose assessment in the emergency department for detecting unrecognised diabetes: a cluster randomised trial

Date:

2020-07-01

Citation:

Fourlanos, S., Barmanray, R. & Kyi, M. (2020). Routine glucose assessment in the emergency department for detecting unrecognised diabetes: a cluster randomised trial. MEDICAL JOURNAL OF AUSTRALIA, 213 (2), <https://doi.org/10.5694/mja2.50550>.

Persistent Link:

<http://hdl.handle.net/11343/275541>