THE survival of auditory neurones depends on the continued supply of trophic factors. Early postnatal spiral ganglion cells (SGC) in a dissociated cell culture were used as a model of auditory innervation to test the trophic factors leukaemia inhibitory factor (LIF) and neurotrophin-3 (NT-3) for their ability, individually or in combination, to promote neuronal survival. The findings suggest that LIF supports neuronal survival in a concentration-dependent manner. Moreover LIF potentiated NT-3-mediated spiral ganglion neuronal survival in a synergistic fashion.

Key words: Auditory neurones; Cytokine; Neuronal degeneration; Neurotrophic effects; Neurotrophin; Pharmacological therapy

LIF potentiates the NT-3-mediated survival of spiral ganglia neurones *in vitro*

P. L. Marzella,^{CA} G. M. Clark,
R. K. Shepherd, P. F. Bartlett¹
and T. J. Kilpatrick¹

Department of Otolaryngology, University of Melbourne, Royal Victorian Eye and Ear Hospital, 32 Gisborne Street, East Melbourne Australia 3002; ¹Neuroimmunology Laboratory, The Walter and Eliza Hall Institute of Medical Research, Parkville Australia 3052

^{CA}Corresponding Author

Introduction

Degeneration of spiral ganglion cells (SGC) is one of the most common histopathological correlates of sensorineural hearing loss.¹ Hair cells within the cochlea produce and supply trophic factors:² when cochlear hair cells are lost as a result of trauma, disease or ageing, there is a secondary wave of SGC degeneration,³ suggesting that the continued supply of growth factors is responsible for maintaining neuronal integrity.

Brain-derived neurotrophic factor (BDNF), neurotrophin-3 (NT-3) and neurotrophin 4/5 (NT-4/5) are members of a family of growth factors reported to exert activity on SGC, providing protection against cisplatin toxicity and promoting neuronal survival.⁴ In addition, BDNF and NT-3 have been reported to play a role in the development and patterning of developing auditory neurones.^{5,6} Findings in mutant mice null for NT-3 report a significant loss of SGC.⁶ Moreover, studies of neurotrophin receptors (trkB and trkC), revealed that spiral ganglia neurones in trkB^{-/-};trkC^{-/-} double mutant mice were reduced by 61% compared with wild type.⁷

The cytokine and growth factor ciliary-derived neurotrophic factor (CNTF) has also been reported to have trophic effects on SGC.^{8,9} CNTF was found to increase the survival of auditory neurones in dissociated cell cultures.⁹ In addition, there is

evidence to suggest that NTs and cytokines act additively upon SGC. Indeed, a combination of CNTF and BDNF was found to be more effective in promoting the survival of neurones in dissociated cell cultures of 5-day-old rat than either factor individually.⁹ By contrast, in rat spiral ganglion explant treatment with NT-3 gave maximal survival; this remained unchanged when either BDNF or CNTF was added.⁸

Leukaemia inhibitory factor (LIF) is a cytokine structurally homologous to CNTF.¹⁰ Furthermore, the receptors for LIF and CNTF also share two identical receptor components, which are believed to induce their overlapping biological activity.¹⁰ Because LIF exhibits activity on several neuronal classes,^{10–13,15} and CNTF has demonstrated activity on SGC^{8,9} it is reasonable to assume that LIF would also have trophic activity on SGC. In addition, a limiting factor to the heterodimerisation of the CNTF receptor complex is the CNTF α -receptor (CNTFR α) component, which is restricted in its distribution.¹⁰ By contrast, LIF only requires the receptor components gp130 and LIFR β for signal transduction. The latter two receptor components are widely distributed,¹⁰ suggesting that LIF may have a greater number of targets (cells or receptors) than CNTF. The distribution of CNTFR α in the cochlea is unknown, but if its distribution in this organ is restricted in comparison to gp130 and LIFR β it is possible that LIF may have greater potency upon SGC and surrounding cells than CNTF.

We therefore tested whether LIF could potentiate neuronal survival in the mammalian cochlea. In addition, we assessed whether NT-3 and LIF acted in concert in either an additive or synergistic manner to potentiate cochlear neuronal survival.

Materials and Methods

Culture of auditory neurones: Five-day-old (P5) rat pups (Wistar) were cooled then rapidly decapitated, the heads were rinsed in 70% alcohol and under aseptic conditions the skull was opened longitudinally. The temporal bone was identified and the bony capsule of the inner ear removed and placed in HEPES buffered Eagle's media (HEM). The cochleae were dissected, and after careful removal of the organ of Corti the remaining gelatinous modiolus was digested in HEM, containing 0.025% trypsin (Sigma) and 0.001% DNase (Sigma) (30 min at 37°C). Trypsinisation was terminated by the addition of 1 ml fetal bovine serum (FBS). Digested cochleae were centrifuged ($1500 \times g$ for 5 min at room temperature [RT]) and the pellet was gently triturated in HEM containing 0.001% DNase. The SGC suspension was again centrifuged ($1500 \times g$ for 5 min at RT) and the pellet resuspended in Dulbecco modified Eagle's medium (DMEM) with N1 supplement and glucose adjusted to a final concentration of 4.5 g/l.

The dissociated SGC suspension was pre-plated in 35 mm tissue culture plates (Falcon; 30 min at 37°C, 10% CO₂). The supernatant of the enriched SGC was collected and a cell count was performed on 10 µl of SGC suspension. The SGC suspension was then plated in 96-well culture plates (Falcon), previously coated with poly-L-lysine (20 µg/ml) and laminin (0.01 mg/ml) at a density of 1×10^4 cells/well, containing an average of 518 ± 6 neurones. Cell cultures were maintained at 37°C in 10% CO₂ for 3 days in 200 µl serum free media (DMEM + N1 supplement + 4.5 g/l glucose). Growth factors were added to the medium at time of plating at a range of concentrations. To determine the initial seeded number of SGC/well a separate plate was incubated at 37°C in 10% CO₂ for 4 h, and the cells were then fixed and immunostained.

Immunocytochemistry: Immunostaining used the avidin-biotin complex (ABC) method (Vectastain kit). The cultures were fixed in 100% methanol (30 min) then rinsed in phosphate buffered saline (PBS). To prevent non-specific binding the cultures were pretreated with 2% FBS in PBS (30 min at RT) then incubated with the primary antibody: anti-200 kDa neurofilament (α NF 200 rabbit Ab 1:20 dilution, RT for 30 min). The cells were rinsed with 2% FBS in PBS, then reacted with the biotinylated secondary

antibody (anti-rabbit in 2% FBS) and subsequently with ABC reagents according to the manufacturer's protocol. Following this step the cultures were stained for 5 min at RT with diaminobenzidine (DAB; 1 tablet in 10 ml PBS with 30 µl H₂O₂). Surviving neurones were defined as cells whose somas immunostained with the α NF 200 antibody and possessed neuritic projections with a length equal to at least three times the width of the neuronal soma. To measure neuronal survival all the immunostained neurones in each well were counted. Experiments were repeated four times in which there were four wells for each given concentration of NTs or combination thereof. The data were analysed for significance using ANOVA on Ranks and Student-Newman-Keuls method for multiple comparison. The care and use of the animals reported on in this study were approved by the Animal Experimentation Ethics Committee of the RVEEH and performed in accordance with the principles of the NHMRC of Australia.

Results

Dissociated SGC were plated at 10^4 cells/well with a mean ($\pm$ s.e.m.) seed number of 518 ± 6 neurones per well. After 3 days *in vitro* the average number of viable cells (or cells with projections) in the untreated wells had decreased to 11 ± 2 cells per well, representing a survival rate of 2.1% compared with the number of neurones present 4 h after plating (seed). The addition of LIF promoted the survival of SGC in a concentration-dependent manner, with a significant increase in neuronal survival detected at concentrations as low as 0.1 ng/ml compared with the untreated wells (Fig. 1). Maximum neuronal survival using LIF alone was at 10 ng/ml (54 ± 5 cells/well). At higher concentrations (100 ng/ml), there was a significant decrease in the survival of SGC ($p < 0.05$; Student-Newman-Keuls pairwise comparison).

When used concurrently with NT-3, LIF retained a concentration-dependent survival promoting efficacy, with a maximal survival promoting effect at a concentration of 10 ng/ml/50 ng/ml. In addition, LIF and NT-3 showed significant survival-promoting activity at concentrations as low as 0.1 ng/ml/0.1 ng/ml (LIF/NT-3). When combined, the survival-promoting action of LIF and NT-3 appeared to be synergistic (Fig. 1), since the average number of neurones surviving in the presence of both factors was 1.6-fold greater than the aggregate of the maximal increases recorded when using each factor alone.

Discussion

This study indicates that LIF individually or in combination with NT-3 can promote survival of SGC

FIG. 1. Effects of leukaemia inhibitory factor (LIF) and neutrophin-3 (NT-3) on the survival of P5 rat spiral ganglia after 72 h *in vitro*. The total number of neurofilament antibody-stained spiral ganglia neurones counted in each well (left y-axis) or as a percentage of survival (right y-axis) is expressed as a function of the concentration of neurotrophic factor or combination of factors (x-axis). Each condition is represented by four culture wells repeated in four experiments, $n = 16$ for each data bar. Results are expressed as mean $\pm$ s.e.m. Seed number of neurones 518 ± 6 . ^a $p < 0.001$ between treatment condition and control wells; ^b $p < 0.001$ between 100 ng/ml and 10 ng/ml LIF.

in culture. At present the role of LIF in the auditory system or under what conditions it may be expressed is not known. LIF's known pleiotropic activity across other classes of neurones^{10-13,15} suggests that it may also have various roles within the auditory system.

Previously, [¹²⁵I]LIF binding sites have been detected in both neural crest- and placode-derived sensory ganglia.¹² Because SGC derive from the otic placode, while Schwann cells and supporting cells are of neural crest origin,¹⁴ it is possible that during embryogenesis these cells could also express [¹²⁵I]LIF binding sites, suggesting that LIF may have a developmental role in the cochlea. LIF may also have an action on SGC in response to injury: this hypothesis is supported by another study reporting increased retrograde transport in axotomized neurones.¹⁶ At present it is also unknown whether LIF may also have a target-derived neurotrophic role (i.e. whether it is present in hair cells). Although a recent study shows that the CNTF family of factors and their respective receptors are found in the postnatal cochlea¹⁷ we await the results of further *in situ* studies to elucidate whether LIF may also have a target-derived neurotrophic role within the adult cochlea.

A previous study reported that NT-3 treatment resulted in maximal survival of rat SGC, which remained unaffected by the concomitant addition of

either BDNF or CNTF.⁸ By contrast, we report that LIF can potentiate the NT-3-induced survival of SGC by up to 1.6-fold. It is unclear why LIF has greater efficacy than CNTF or BDNF in potentiating NT-3-induced survival of SGC. However, the widespread distribution of LIF receptor components across other classes of neurones¹⁰ suggests that LIF receptor may be broadly expressed on SGC and/or on surrounding cells. This broad distribution of the LIF receptor components may underpin the ability of LIF to potentiate NT-3-induced survival of SGC.

Furthermore, it is undetermined whether the potentiating effect of LIF upon NT-3-induced survival is occurring on the same neurones. The converging second messenger pathways for the neural cytokines and NTs,¹⁰ and the synergistic action demonstrated in these studies suggest that LIF and NT-3 may indeed be acting on the same SGC to potentiate each other's survival effects. Conversely, in dissociated cell cultures other non-neuronal cells (i.e. fibroblast and glial cells) proliferate, and therefore, it is possible that LIF may act indirectly via these cells as well as exert direct action on the SGC themselves. In addition, studies with antibodies directed at the trkB and trkC receptors for the NTs revealed that *in vitro* some populations of SGC are NT-independent.⁴ Thus the population of surviving neurones may include SGCs targeted by both factors, SGCs supported by only one of the factors and SGCs indirectly affected by LIF and NT-3. Double labelling studies may elucidate the site(s) of LIF-receptor complexes and thus help determine the site(s) of action.

Conclusion

We report that the cytokine LIF displays trophic activity and improves the NT-3-induced neuronal survival in dissociated SGC cultures. These data support the theory that NTs and cytokines act concomitantly on inner ear innervation in the early post-natal period, when the complex pattern of adult innervation is being established. *In situ* hybridization studies will determine transcription levels of the LIF receptor components within the auditory system, and thus elucidate where LIF may be acting and whether there are developmental periods of critical expression. Moreover, they will help determine if there is a relationship between non-neuronal cell injury and neuronal repair.

In addition, the results suggest that a combination of several trophic factors may provide a better approach than single factor delivery when developing pharmacological therapies for auditory neurone repair.

References

1. Otte J, Schuknecht HF and Kerr AG. *Laryngoscope* **88**, 1231–1246 (1978).
2. Lefebvre PP, Weber T, Rigo JM *et al.* *Hear Res* **58**, 185–192 (1992).
3. Spoendlin H. *Ann Otol Rhinol Laryngol* **93** (Suppl. **112**), 76–82 (1984).
4. Zheng JL, Stewart RR and Gao WQ. *J Neurosci* **15**, 5079–5087 (1995).
5. Staecker H, Van De Water TR, Lefebvre PP *et al.* *Dev Brain Res* **92**, 49–60 (1996).
6. Fariñas I, Jones KR, Backus C *et al.* *Nature* **369**, 658–661 (1994).
7. Minichello L, Piehl F, Vazquez E *et al.* *Development* **121**, 4067–4075 (1995).
8. Staecker H, Liu W, Hartnick C *et al.* *NeuroReport* **6**, 1533–1537 (1995).
9. Hartnick CJ, Staecker H, Malgrange B *et al.* *J Neurobiol* **30**, 246–254 (1996).
10. Ip NY and Yancopoulos GD. *Annu Rev Neurosci* **19**, 491–515 (1996).
11. Murphy M, Reid K, Brown MA *et al.* *Development* **117**, 1173–1182 (1993).
12. Qui LQ, Bernd P and Fukada K. *Dev Biol* **163**, 516–520 (1994).
13. Hilton DJ and Gough NM. *J Cell Biochem* **46**, 21–26 (1991).
14. Whitehead M. Development of the cochlea. In Altschuler RA, Hoffman DW and Bobbin RP, eds. *Neurobiology of Hearing*, New York: Raven Press, 1986: 191–211.
15. Kwon YW, Abbondanzo SJ, Stewart CL *et al.* *J Neurobiol* **28**, 35–50 (1995).
16. Hendry IA, Murphy M, Hilton DJ *et al.* *J Neurosci* **12**, 3247–3434 (1992).
17. Malgrange B, Lefebvre PP, Rogister B, *et al.* Abstracts of the Association for Research in Otolaryngology. XVIIIth Midwinter Research Meeting Abs# 18 (1997).

ACKNOWLEDGMENTS: P.L.M. is funded by the Ian Potter Foundation. G.M.C. is funded by the Human Communication Research Centre of the Department of Otolaryngology, University of Melbourne. R.K.S. is a recipient of a Senior Research fellowship from The Garnett Passe and Rodney Williams Foundation. P.F.B. is funded by NH MRC. T.J.K. is a recipient of a Senior Medical Research fellowship from The Sylvia and Charles Viertel Foundation.

Received 6 February 1997;
accepted 6 March 1997

General Summary

A combination of neuronal growth factors promoted the survival of auditory nerve cells *in vitro* in a synergistic manner. The neuronal cytokine leukaemia inhibitory factor (LIF) and the neurotrophin-3 (NT-3) were tested for their ability, individually or in combination, to promote auditory nerve cells survival. The findings suggest that treatment with LIF supports auditory nerve cell survival in a concentration-dependent manner, and potentiates NT-3-mediated auditory nerve cells survival in a synergistic fashion. The results suggest that a combination of several nerve growth factors may provide a better approach than single factor therapy when developing pharmacological therapies for the treatment of deafness.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Marzella, P. L.;Clark, Graeme M.;Shepherd, R. K.;Bartlett, P. F.;Kilpatrick, T. J.

Title:

LIF potentiates the NT-3-mediated survival of spiral ganglia neurones in vitro

Date:

1997

Citation:

Marzella, P. L., Clark, G. M., Shepherd, R. K., Bartlett, P. F., & Kilpatrick, T. J. (1997). LIF potentiates the NT-3-mediated survival of spiral ganglia neurones in vitro. *NeuroReport*, 8, 1641-1644.

Persistent Link:

<http://hdl.handle.net/11343/27559>