DR CLARE SLANEY (Orcid ID : 0000-0002-6986-6115)

Received Date : 06-Feb-2020

Revised Date : 14-Mar-2020

Accepted Date : 24-Mar-2020

Article type : Viewpoint

Cross-talk between tumors at anatomically distinct sites

Amanda J. Oliver^{1,2}, Phillip K. Darcy^{1,2}, Joseph A. Trapani^{1,2}, Michael H. Kershaw^{1,2} and Clare Y. Slaney^{1,2,*}.

Affiliations:

¹Cancer Immunology Program, Peter MacCallum Cancer Center, Melbourne, Victoria, Australia.

²Sir Peter MacCallum Department of Oncology, University of Melbourne, Parkville, Victoria, Australia.

***Correspondence:** Clare Slaney, clare.slaney@petermac.org

Key words: tumor cross-talk, abscopal effect, immunotherapy, tumor microenvironment, seed and soil hypothesis

Conflict of interest statement: The authors declare no conflict of interest.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/FEBS.15316](https://doi.org/10.1111/FEBS.15316)

This article is protected by copyright. All rights reserved

Acknowledgements: This work was supported by grants from the National Health and Medical Research Council (NHMRC) of Australia (1103352 and 1176935) the National Breast Cancer Foundation (NBCF) of Australia (IIRS-18-064 and IIRS-20-073) and Susan G. Komen Breast Cancer Foundation (16376637). PKD was supported by a Principal Research Fellowship from the NHMRC. CYS was supported by a Postdoctoral Fellowship from the NBCF. JAT is a Rosie Lew Fellow of the Peter MacCallum Cancer Foundation.

Running title: Cross-talk between metastatic tumors

List of abbreviations:

TME – Tumor microenvironment

PLR – paraneoplastic leukemoid reaction

GM-CSF/G-CSF – granulocyte-macrophage/granulocyte colony stimulating factor

IL – interleukin

TNF – Tumor necrosis factor

MDSC – myeloid derived suppressor cell

VEGF – vascular endothelial growth factor

DC – dendritic cell

EV – extracellular vesicle

EGFR – epithelial growth factor receptor

TCR – T cell receptor

NK – natural killer

Treg – regulatory T cell

FLT3L – FMS-like tyrosine kinase receptor ligand

SC - subcutaneous

Abstract

Cancer tissue is not homogenous, and individual metastases at different anatomical locations can differ from the primary tumor and from one another in both their morphology and cellular composition, even within an individual patient. Tumors are composed of cancer cells and a range of

other cell types, which, together with a variety of secreted molecules, collectively comprise the tumor microenvironment. Cells of the tumor microenvironment can communicate with each other and with distant tissues in a form of molecular cross-talk to influence their growth and function. Cross-talk between cancer cells and local immune cells is well described, and can lead to the induction of local immunosuppression. Recently, it has become apparent that tumors located remotely from each other, can engage in cross-talk that can influence their responsiveness to various therapies, including immunotherapy. In this article, we review studies that describe how tumors systemically communicate with distant tissues through motile cells, extracellular vesicles and secreted molecules that can affect their function. In addition, we summarize evidence from mouse studies and the clinic that indicate an ability of some tumors to influence the progression and therapeutic responses of other tumors in different anatomical locations.

Introduction

Tumors in humans often spread to multiple different sites within one patient. All tissues of the body are connected through lymphatic and vascular systems, and systemic immune responses against tumors have been shown to impact distal tissues and tumors. Tumor cells and their associated cells, collectively known as the tumor microenvironment (TME), in one location can have a significant influence on the TME of distal sites. Various factors secreted from tumors, such as growth factors and tumor-derived extracellular vesicles (EVs), have been observed in the circulation (**Figure 1A and B**). Immune cell populations in peripheral blood and lymphoid organs are influenced by tumor cell derived factors, leading to systemic changes within the host (**Figure 1D**).

The impact of tumors on non-cancerous distal tissues is largely demonstrated in studies of the pre-metastatic niche, where secreted factors from a primary tumor create an environment within the pre-metastatic tissue conducive to tumor cell growth (**Figure 1C**). In addition, the term “abscopal effect” that was first used by Mole in 1953 in radiotherapy describes a phenomenon where local treatment of one tumor leads to regression of a distant tumor [1]. This suggests it is possible that tumors in different metastatic sites can influence each other during tumor development, progression and therapy. However, very few studies have investigated how tumors in different metastatic sites might impact each other. In this article, we review studies that consider the existing evidence on the concept of cross-talk between tumors in different anatomical locations (**Figure 1E**).

Systemic changes mediated by tumor cells

Circulating factors from tumors

Tumor cells secrete various molecules including cytokines and growth factors that maintain the local microenvironment. These tumor molecules can also be released into the circulation where they can influence distant tissues. A clinical example of this comes from paraneoplastic leukemoid reaction (PLR) which describes a variety of hematological clinical disorders due to a non-hematolymphoid tumor. PLR has been reported in patients with a variety of cancers, including lung cancers, melanoma, mesothelioma, head and neck cancer and cancers of other origins [2]. The most commonly secreted cytokine by these tumors include granulocyte colony stimulating factor (G-CSF), granulocyte-macrophage colony stimulating factor (GM-CSF), IL-1, IL-6 and TNF- α . In fact, GM-CSF is expressed by a number of tumor types constitutively, with one study reporting 23 of 75 human tumor lines producing GM-CSF [3]. These cytokines stimulate bone marrow myelopoiesis and inhibit myeloid cell differentiation that leads to accumulation of immature myeloid cells, which are called myeloid-derived suppressor cells (MDSCs) that suppress effector cell responses.

Another common tumor-secreted protein which has systemic effects is vascular-endothelial growth factor (VEGF). Increased plasma levels of VEGF in cancer patients correlates with poorer prognosis [4] and is thought to impact on dendritic cell (DC) differentiation [5]. Furthermore, arginase activity is higher in peripheral blood of gastric cancer patients and could potentially be involved in suppressing functions of circulating immune cells [6]. In each of these examples, tumor growth can result in an altered serum proteome which, can impact on distal tissues and alter the normal physiology of the host. Of course, many other proteins are secreted by tumor cells, but have no function in malignancy or transformation of distal tissues. Examples include elevated serum AFP (hepatocellular cancer), HCG (choriocarcinoma), PSA (prostate cancer), CEA (colorectal cancer), CA 125 (ovarian cancer) and CA 15-3 (breast cancer). Many of these proteins have been used for decades as biomarkers of tumor size indicating both tumor progression and response to therapy [7].

Extracellular vesicles

EVs are small membrane-encapsulated particles, derived from cells, which contain DNA, RNA and proteins representative of the host cells. EVs describe both exosomes and microvesicles which can mediate communication between cells, through their content, to modulate both the local and distal microenvironments. Exosomes are released from the cell

by fusion of multivesicular endosomes with the plasma membrane whereas microvesicles are formed by budding off the plasma membrane [8]. Tumor-derived or tumor-associated EVs can exchange molecules from malignant or non-malignant cells within the TME to other cells throughout the body [9]. This communication can impact on signaling between the tumor and host, and can influence many aspects of tumorigenesis. Expression of ligands such as PD-L1, which can engage with receptors expressed by T cells to impair their function, have been identified on EVs. High levels of EV PD-L1 associate with clinical non-responders to anti-PD-1 therapy in metastatic melanoma [10].

EVs can be found in the blood and lymph through which they travel to distant tissues. A number of studies have demonstrated the influence of EVs on the pre-metastatic niche. Tumor-derived EVs have been shown to reprogram fibroblasts in pre-metastatic sites to generate an acidic microenvironment inhibitory for anti-tumor immune responses [11-13]. Epidermal growth factor receptor (EGFR)-expressing EVs secreted by gastric cancer cells enhance liver metastasis through induction of hepatocyte growth factor signaling in liver stromal cells [14]. Melanoma-derived EVs have been shown to induce vascular leakiness, allowing for increased accessibility for tumor cells into pre-metastatic sites [15]. In addition, tumor-derived EVs can also suppress monocyte maturation, leading to increased MDSCs, effectively increasing systemic tumor-mediated immunosuppression [16]. Thus, tumor derived EVs represent a form of communication from tumors to distal tissues that may potentially impact on distal TMEs and immune responses in other tumors.

Circulating cells

Various studies have focused on changes in the peripheral blood of cancer patients and pre-clinical cancer models. One study generated a novel system to match interactions across tissues to link pathways within primary tumors with changes in the systemic response in breast cancer patients [17]. This study revealed that expression of genes in circulating blood cells are tightly linked to the genes in the primary tumors, and such gene expression patterns vary with breast cancer subtypes. Therefore, the RNA profiles of tumors can be indicative of systemic immune responses. A separate study of breast cancer patients identified changes in the peripheral blood immune cell profile between breast patients with or without lymph node metastasis [18].

Indication of a systemic T cell response against cancer is evident from studies where T cells from peripheral blood of cancer patients can destroy tumor cells *in vitro* [19]. In addition, the presence of MDSCs in peripheral blood is associated with increased disease burden and negative prognosis in multiple solid tumors, presumably by infiltrating tumor sites, pre-metastatic niches and with their potential to inhibit circulating T cells [20, 21]. In ovarian cancer patients, the diversity in T cell receptor (TCR) clones within the circulation decreased during disease progression [22]. The diversity of peripheral TCR clones is decreased in lung and ovarian cancer patients compared to healthy controls, indicating that the presence of negatively impacts on T cells systemically[22, 23]. Intriguingly, during treatment with chemotherapy, the circulating T cell receptor (TCR) clonality is altered in various cancers, indicating a potential enhanced immune response that can be identified systemically in these cases [24, 25]. In breast cancer patients, increased diversity of the circulating TCR repertoire of CD8⁺ T cells was associated with increased response to chemotherapy[25].Furthermore, a study in melanoma found that the combinatorial diversity evenness in the TCR repertoire pretreatment can be predictive of checkpoint blockade efficacy [26].

A recent study investigated systemic immune responses in experimental mouse models. Spitzer *et al.* found multiple changes within lymphoid tissue and within the circulation of mice receiving effective immunotherapies during the priming and rejection phase of tumors [27]. In peripheral blood, there were increases in NK cells, DCs, activated B cells, naïve CD4⁺ T cells, CD90^{hi} T-helper cells and CD8⁺ T cells expressing Ly6C and CD44, observed during the priming phase. During the rejection phase, there were substantial decreases in immune cell frequencies in the circulation likely denoting migration of immune cells into the tissues. However, many cell populations in the circulation at this time retained expression of Ki67, indicating potential ongoing expansion and maintenance of these cells.

Thus, changes in the TME and immunity against a tumor is not just confined to the tumor, and/or the tumor-draining lymph nodes, but can cause systemic immune dysfunction. In this light, a study of osteosarcoma found that T cells in the spleen of tumor bearing mice showed an exhaustion status [28]. In addition, there was an increase in the percent populations of Tregs and MDSCs in the spleens of the tumor bearing mice. In this model, treatment with anti-PD-L1 decreased T cell exhaustion and increased monocyte maturation. These data suggest that cancer induced systemic immune dysregulation should be considered and targeted during cancer treatment.

Seed and soil hypothesis

In 1889 Stephen Paget proposed the seed and soil hypothesis, where metastatic spread depends on cross-talk between the cancer cells (the ‘seed’) and the organ microenvironment (the ‘soil’)[29]. This hypothesis has been built upon with clinical observations and murine models of cancer in which certain tumor types tend to metastasize to specific organs independent of vascularization, rate of blood flow or the number of tumor cells delivered.

Experimental data to support this hypothesis first came from a study of B16 murine melanoma cells which grow in the lungs when injected intravenously. It had been assumed that the explanation for this is trivial – that the melanoma nodules develop in the lungs because they lodge in the first capillary bed they encounter. However, B16 cells also grew in lung tissue, but not in renal tissue, that had been implanted intra-muscularly demonstrating that the tissue microenvironment is important in determining the site of metastasis [30]. Comparison of B16 and K-1735 melanoma cell lines injected into the internal carotid artery revealed differences in brain metastasis patterns, as they produced only meningeal or parenchymal growths, respectively [31]. This indicates different regions and microenvironments, even within the same organ, may support the seeding and growth of different tumor cell types.

Furthermore, establishment of a ‘pre-metastatic niche’ has now been identified as a precursor for metastasis. The pre-metastatic niche is defined as aberrant immune cells and extracellular matrix that initiate the transformation of a healthy microenvironment to one supportive of tumor cell growth. The first-described pre-metastatic niche found a role for VEGFR1⁺ hematopoietic progenitor cells that formed cellular clusters in pre-metastatic sites before tumor cell arrival [32]. Blocking VEGFR1 function or removing these cells ablated the cellular clusters and decreased tumor metastasis. Various proteins such as fibronectin, collagen IV, tenascin, periostin and certain chemokines and cytokines can be aberrantly expressed by tumor-influenced stromal cells in distal tissues [33-37]. EVs from pancreatic cancer can establish pre-metastatic niches in the liver and promote liver metastasis [38]. Tumor derived EVs can also be taken up by Kupffer cells of the liver, which produce TGFβ and fibronectin to recruit bone marrow derived cells (BMDCs) that support metastasis [39].

Cells attracted to the pre-metastatic niche include MDSCs, neutrophils, macrophages and regulatory T cells (Tregs) [33, 36, 40-45]. For example, a murine study of breast cancer demonstrated that granulocytic MDSCs were important for establishing a pre-metastatic niche [46]. Thus, it is well established that signals from the primary tumor can influence distal tissues and impact healthy organ microenvironments. It could be expected that tumors in different primary and metastatic sites can also influence each other in a similar manner.

Abscopal effects of localized radiation

A well-documented example of one tumor impacting on a distal tumor in metastatic cancer patients is the abscopal effect of local radiation - in rare cases, localized radiation therapy of one tumor is followed by regression of distant untreated tumor/s in an immune-dependent manner. The abscopal effect has been observed in melanoma [47], renal cell [48], breast [49], hepatocellular [50], non-small cell lung and thymic cancers [51]. Between 1969 and 2014, there were an estimated 46 case reports of the abscopal effect from radiation therapy [52].

The involvement of the immune system in abscopal responses has been demonstrated in mouse models. In a mouse fibrosarcoma model, the dose needed for 50% tumor control in T cell-deficient mice was 64.5 Gray compared with 30 Gray in immunocompetent mice, demonstrating the key role of the immune system [53]. In a bilateral mouse model of breast cancer, irradiation in one flank along with systemic delivery of FMS-like tyrosine kinase receptor 3 ligand (FLT3L) resulted in a T cell dependent delay in growth of both irradiated and non-irradiated tumors [54]. Another study using the Lewis lung carcinoma model observed control of lung metastases when footpad tumors were irradiated and the mice received systemic Flt3L [55].

Since the initial observations of abscopal responses, combined immunotherapy and radiation has increased the reports of abscopal responses. In a clinical trial combining radiotherapy with GM-CSF in patients with solid cancer, abscopal responses were observed in 30% of patients [51]. In melanoma, α PD-1 or α CTLA4/ α PD-L1 in combination with radiotherapy also increased abscopal responses [56, 57].

Mechanistic studies on combined radio-immunotherapy have shown that cross-presentation by BATF3-dependent DCs, CD8 T cells and type I interferons are required for abscopal responses [58-61]. Radiation therapy is thought to enhance the release of tumor associated

antigens and promote local promotion of inflammatory responses, especially stimulation of the innate immune system, which consequently leads to CD8⁺ T cell priming by BATF3-dependent DCs and T cell circulation and homing to tumor sites [62, 63]. In summary, tumors in distal sites can influence each other, in the context of radiation therapy, whereby tumor destruction by radiation leads to enhanced anti-tumor immune responses that become systemic.

Abscopal responses to other localized therapies have been observed when they are combined with PD-1/PD-L1 inhibitors [27]. In the MMTV-PyMT mouse model, which produce spontaneous tumors in multiple mammary glands, only those tumors that were directly injected with a tumor-binding antibody therapy regressed. However, when combined with systemic anti-PD-L1, intra-tumoral injection resulted in regression of distal tumors, even though anti-PD-L1 alone did not bring about systemic immune responses. Another study, which utilized vascular-targeted photodynamic therapy in primary murine Renca kidney tumors, resulted in systemic immune responses against lung metastases, when administered alongside anti-PD-1/PD-L1 [64].

Cross-talk between tumors in different anatomical sites

Evidence from experimental models

As discussed so far, complex interactions occur between the tumor and host, and these interactions can have systemic consequences. Thus, it is reasonable to hypothesize that metastatic tumors in different anatomical sites are not completely independent entities and that connectivity between tumors could impact on tumor growth and response to therapy. The limited number of studies that have investigated this hypothesis to date are discussed in detail in the following section.

Studies into how tumors can affect each other's growth and response to immunity and therapy go back many years. In 1905, a study by Paul Ehrlich found that engrafted tumors exhibit a negative effect on each other. In this study, earlier implanted tumors inhibited the growth of tumors injected later and the inhibitory effect increased with the size of the first tumor [65]. Studies in anti-angiogenic therapies have observed a decrease in the growth of metastatic tumors by angiogenesis inhibitors, only when the primary tumor is present. The serum and urine of tumor bearing mice contained angiostatin which inhibits endothelial cell

proliferation and was determined to be a systemic mediator of the effects in metastatic tumors in this study [66].

Another study used mathematical modelling to investigate dynamic behavior and interactions between tumor and immune cells at various metastatic sites, with or without therapy [67]. Simulation of T cell recruitment in lung and breast tumors was assessed in isolation, and upon metastatic seeding after 200 days, aimed at analyzing T cell redistribution and immune cell population dynamics between the two sites, and to model the effect of surgery or radiation. This mathematical modelling revealed that a proportion of the T cells activated in the breast were recruited to the metastatic site. A larger fraction of circulating immune cells traffic through the pulmonary circulation and therefore have a greater chance of extravasation into the lung. The authors hypothesized that this would result in increased primary tumor growth, given the relocation of tumor reactive T cells to the lungs and estimated a 30% reduction in lung tumor size when a secondary breast tumor is seeded. Thus, this study provides a simulation and mathematical model for interconnectivity between existing tumor sites, which is based on assumptions from redistribution of T cells and systemic immunity in various tumor studies and other mathematical models of tumor growth.

A study by our laboratory was the first to demonstrate the influence of one tumor on immunotherapy responses to tumor located in a distant anatomical site [68]. Using the Renca kidney cancer model, in which subcutaneous (SC) tumors were responsive and orthotopic kidney tumors resistant to a triple antibody (trimAb) therapy, this study assessed therapy responses in mice simultaneously implanted with SC and kidney tumors. When mice had only contralateral SC tumors, responses to trimAb therapy were sustained. However, when mice had one SC tumor and one kidney tumor, growth of the SC tumor was accelerated with therapy, indicating that the Renca kidney tumor was inhibiting the response of the SC tumor.

In exploring a possible mechanism, we determined that M2 macrophages, which are commonly found in Renca kidney tumors increased in number in SC tumors when a kidney tumor was also present. The increase in M2 macrophages was accompanied by a decrease in T and NK cell numbers and cytotoxicity within the SC tumors. Furthermore, CCL2 blockade or systemic macrophage depletion using Clodrolip improved therapy responses in SC tumors when kidney tumors were present. As there were no soluble factors identified to mediate the suppressive effects, we hypothesized that M2 macrophages generated within the suppressive

kidney tumor could traffic to the SC tumor, where they suppressed immune responses and decreased the therapeutic response.

A more recent study investigated the efficacy of dual checkpoint blockade (α PD-1/ α CTLA4) on melanoma brain metastases [69]. Taggart *et al* found that an extracranial tumor (SC, in their model) had to be present for α PD-1/ α CTLA4 to be efficacious against intra-cranial tumors. Mice implanted only with intra-cranial B16 melanoma cells responded poorly to α PD-1/ α CTLA4, whereas those bearing both intra-cranial and SC tumors had enhanced survival and decreased growth of both extra- and intra- cranial tumors. When an extracranial tumor was present, intra-cranial tumors had enhanced infiltration of immune cells and particularly, CD8⁺ T cells, bone marrow-derived macrophages and brain microglia. The beneficial effect of an extracranial tumor was abolished when mice were depleted of CD8⁺ T cells and NK cells. Furthermore, CD44⁺CD62L⁻ effector CD8⁺ T cells were increased in the peripheral blood of mice with extracranial disease and transfer of these cells from spleens of α PD-1/ α CTLA4 treated mice demonstrated enhanced homing into intracranial tumors compared with the same cell population from spleens of treatment-naïve mice. The authors also demonstrated that this enhanced homing may be due to enhanced T cell trafficking determinants on intracranial blood vessels when extracranial tumors were present.

Interestingly, further evidence in support of immune cross-talk between tumors can be found in studies utilizing neoadjuvant therapy [70]. In this approach, mice bearing both primary breast tumors and lung metastases were treated with immunotherapy prior to surgical removal of the primary. Mice undergoing neoadjuvant therapy survived longer, with decreased primary recurrence and progression of lung metastases, compared with mice undergoing surgery followed by adjuvant immunotherapy. The presence of the primary tumor during immunotherapy enabled a more pronounced tumor-specific T cell response that subsequently impacted on tumor relapse and metastatic formation.

Evidence from human cancers

There is increasing evidence for tumors in different metastatic sites influencing each other in human cancers. Evaluation of 371 patients with metastatic melanoma treated with high-dose IL-2 revealed that cutaneous and subcutaneous melanoma metastases were more responsive than visceral metastases [71]. Interestingly, patients with both cutaneous/subcutaneous and visceral metastases had an overall response rate of 17%, whereas patients with only

cutaneous/subcutaneous metastases had on overall response rate of 50%, supporting the notion of visceral metastases having a negative impact on therapy response.

Data presented at the 2017 American Society of Clinical Oncology meeting reported that metastasis to the liver in melanoma patients was associated with reduced α PD-1 response [72]. The authors recapitulated these findings in the B16 melanoma mouse model, in that mice with liver and SC tumors were unable to achieve SC tumor clearance upon treatment with α PD-1, whereas 30% of mice with SC tumors alone achieved tumor rejection. Interestingly, the presence of liver metastases from an unrelated MC38 tumor line did not impact on SC tumor rejection, suggesting a tumor- or antigen-specific process. Although neither study controlled for the possibility that visceral/liver metastases signify more aggressive or progressive disease, these studies suggest that in humans, metastatic tumors situated in specific organs can have an influence on the overall response to therapy.

Recent results from human trials of immune checkpoint blockade in the neoadjuvant setting support the observations in mice described above [73, 74]. Patients with melanoma or glioblastoma undergoing neoadjuvant checkpoint blockade had better outcomes than patients receiving adjuvant checkpoint blockade, with increased time to progression and increased overall survival. Enhanced systemic immune responses were observed in some patients receiving neoadjuvant therapy, supporting the interpretation that, despite a larger tumor burden, the presence of the primary tumor during checkpoint blockade led to an increased T cell response against the remaining unresectable malignant disease.

Concluding remarks

Given the complexity of cancers, with metastases to various organs, and the ability of tissues to communicate across very distinct anatomical sites, it is not surprising that tumors might also affect each other's progression and responses to therapy. Tumors at different sites can differ in their immune composition, and the immune system is made up of a diverse variety of motile cells and secreted molecules that have the potential to move throughout the body. In addition, other cell types within tumors, including malignant cells, can also produce a variety of molecules and particles, many of which presumably remain to be identified, that may influence other tissues and tumors remotely.

In this review, we have described studies supporting the concept of cross-talk between tumors, which can impact on the growth of metastases in various sites in the body and their response to various therapies, including immunotherapy (**Figure 1**). In some instances, cross-talk can clearly lead to a reduced response to therapy at certain sites, but there are also instances where tumors can become more responsive to therapy. With further advancement of this field, it may be possible to identify tumors that, based on their histological origin and anatomical site, might have a positive or a negative impact on the response to therapy. In at least some limited instances, it may be feasible to consider excising tumors at certain specific sites, or subjecting them to localized radiotherapy, in order to enhance the patient's overall response and prolong survival.

Figure Legend

Figure 1: Mechanisms of cross-talk of tumors with distal tumors and tissues. The presence of a tumor in one location results in systemic changes and can impact on distal tumors in other locations. **A)** Tumor-derived EVs and tumor-secreted proteins from malignant and non-malignant cells within the tumor microenvironment (TME), along with cells from the TME, can be released into the circulatory and lymphatic systems. **B)** The consequent changes in the molecular and cellular composition systemically have been detected in cancer patients compared to healthy individuals, and patients before, during and after treatment with various anti-cancer therapies. This highlights that events happening within the TME have an impact on the entire host. **C)** and **D)** Distant healthy tissues are impacted by secreted factors from the primary tumor which can disrupt the extracellular matrix and promote local immunosuppression in order to create the pre-metastatic niche (C) and impact on the composition and production of immune cells within lymphoid organs (D). For example, GM-CSF secreted by tumor cells can increase production of myeloid derived suppressor cells (MDSCs) from the bone marrow. **E)** The impact of one tumor on other distal tumors has recently been investigated in a limited number of studies. It seems that tumors can exhibit both positive and negative effects on each other, especially in the context of immunotherapy responses. For example, murine studies indicate that the presence of a primary tumor can enhance the generation of tumor specific T cell responses in metastatic tumors upon treatment with immunotherapies. Other studies suggest that immunosuppressive cells, such as M2 macrophages, that are generated in one tumor can traffic to, infiltrate and exhibit local immunosuppression in distal tumors thus dampening immunotherapy responses.

References

1. Mole, R. H. (1953) Whole body irradiation; radiobiology or medicine?, *Br J Radiol.* **26**, 234-41.
2. Abukhiran, I. A., Jasser, J. & Syrbu, S. (2020) Paraneoplastic leukemoid reactions induced by cytokine-secreting tumours, *J Clin Pathol.*
3. Bronte, V., Chappell, D. B., Apolloni, E., Cabrelle, A., Wang, M., Hwu, P. & Restifo, N. P. (1999) Unopposed production of granulocyte-macrophage colony-stimulating factor by tumors inhibits CD8+ T cell responses by dysregulating antigen-presenting cell maturation, *J Immunol.* **162**, 5728-37.
4. Toi, M., Kondo, S., Suzuki, H., Yamamoto, Y., Inada, K., Imazawa, T., Taniguchi, T. & Tominaga, T. (1996) Quantitative analysis of vascular endothelial growth factor in primary breast cancer, *Cancer.* **77**, 1101-6.
5. Gabrilovich, D. I., Chen, H. L., Girgis, K. R., Cunningham, H. T., Meny, G. M., Nadaf, S., Kavanaugh, D. & Carbone, D. P. (1996) Production of vascular endothelial growth factor by human tumors inhibits the functional maturation of dendritic cells, *Nat Med.* **2**, 1096-103.
6. Wu, C. W., Chi, C. W., Lin, E. C., Lui, W. Y., P'Eng F, K. & Wang, S. R. (1994) Serum arginase level in patients with gastric cancer, *J Clin Gastroenterol.* **18**, 84-5.
7. Duffy, M. J. (2001) Clinical uses of tumor markers: a critical review, *Crit Rev Clin Lab Sci.* **38**, 225-62.
8. Raposo, G. & Stoorvogel, W. (2013) Extracellular vesicles: exosomes, microvesicles, and friends, *J Cell Biol.* **200**, 373-83.
9. Tung, K. H., Ernstoff, M. S., Allen, C. & Shu, S. (2019) A Review of Exosomes and their Role in The Tumor Microenvironment and Host-Tumor "Macroenvironment", *J Immunol Sci.* **3**, 4-8.
10. Chen, G., Huang, A. C., Zhang, W., Zhang, G., Wu, M., Xu, W., Yu, Z., Yang, J., Wang, B., Sun, H., Xia, H., Man, Q., Zhong, W., Antelo, L. F., Wu, B., Xiong, X., Liu, X., Guan, L., Li, T., Liu, S., Yang, R., Lu, Y., Dong, L., McGettigan, S., Somasundaram, R., Radhakrishnan, R., Mills, G., Lu, Y., Kim, J., Chen, Y. H., Dong, H., Zhao, Y., Karakousis, G. C., Mitchell, T. C., Schuchter, L. M., Herlyn, M., Wherry, E. J., Xu, X. & Guo, W. (2018) Exosomal PD-L1 contributes to immunosuppression and is associated with anti-PD-1 response, *Nature.* **560**, 382-386.
11. Shu, S., Yang, Y., Allen, C. L., Maguire, O., Minderman, H., Sen, A., Ciesielski, M. J., Collins, K. A., Bush, P. J., Singh, P., Wang, X., Morgan, M., Qu, J., Bankert, R. B., Whiteside, T. L., Wu, Y. & Ernstoff, M. S. (2018) Metabolic reprogramming of stromal fibroblasts by melanoma exosome microRNA favours a pre-metastatic microenvironment, *Sci Rep.* **8**, 12905.

12. Asgharzadeh, M. R., Barar, J., Pourseif, M. M., Eskandani, M., Jafari Niya, M., Mashayekhi, M. R. & Omid, Y. (2017) Molecular machineries of pH dysregulation in tumor microenvironment: potential targets for cancer therapy, *Bioimpacts*. **7**, 115-133.
13. Bellone, M., Calcinotto, A., Filipazzi, P., De Milito, A., Fais, S. & Rivoltini, L. (2013) The acidity of the tumor microenvironment is a mechanism of immune escape that can be overcome by proton pump inhibitors, *Oncoimmunology*. **2**, e22058.
14. Zhang, H., Deng, T., Liu, R., Bai, M., Zhou, L., Wang, X., Li, S., Wang, X., Yang, H., Li, J., Ning, T., Huang, D., Li, H., Zhang, L., Ying, G. & Ba, Y. (2017) Exosome-delivered EGFR regulates liver microenvironment to promote gastric cancer liver metastasis, *Nat Commun*. **8**, 15016.
15. Peinado, H., Aleckovic, M., Lavotshkin, S., Matei, I., Costa-Silva, B., Moreno-Bueno, G., Hergueta-Redondo, M., Williams, C., Garcia-Santos, G., Ghajar, C., Nitadori-Hoshino, A., Hoffman, C., Badal, K., Garcia, B. A., Callahan, M. K., Yuan, J., Martins, V. R., Skog, J., Kaplan, R. N., Brady, M. S., Wolchok, J. D., Chapman, P. B., Kang, Y., Bromberg, J. & Lyden, D. (2012) Melanoma exosomes educate bone marrow progenitor cells toward a pro-metastatic phenotype through MET, *Nat Med*. **18**, 883-91.
16. Domenis, R., Cesselli, D., Toffoletto, B., Bourkoura, E., Caponnetto, F., Manini, I., Beltrami, A. P., Ius, T., Skrap, M., Di Loreto, C. & Gri, G. (2017) Systemic T Cells Immunosuppression of Glioma Stem Cell-Derived Exosomes Is Mediated by Monocytic Myeloid-Derived Suppressor Cells, *PLoS One*. **12**, e0169932.
17. Dumeaux, V., Fjukstad, B., Fjosne, H. E., Frantzen, J. O., Holmen, M. M., Rodegerdts, E., Schlichting, E., Borresen-Dale, A. L., Bongo, L. A., Lund, E. & Hallett, M. (2017) Interactions between the tumor and the blood systemic response of breast cancer patients, *PLoS Comput Biol*. **13**, e1005680.
18. Zuckerman, N. S., Yu, H., Simons, D. L., Bhattacharya, N., Carcamo-Cavazos, V., Yan, N., Dirbas, F. M., Johnson, D. L., Schwartz, E. J. & Lee, P. P. (2013) Altered local and systemic immune profiles underlie lymph node metastasis in breast cancer patients, *Int J Cancer*. **132**, 2537-47.
19. Slingluff, C. L., Jr., Cox, A. L., Stover, J. M., Jr., Moore, M. M., Hunt, D. F. & Engelhard, V. H. (1994) Cytotoxic T-lymphocyte response to autologous human squamous cell cancer of the lung: epitope reconstitution with peptides extracted from HLA-Aw68, *Cancer Res*. **54**, 2731-7.
20. Veglia, F., Perego, M. & Gabrilovich, D. (2018) Myeloid-derived suppressor cells coming of age, *Nat Immunol*. **19**, 108-119.
21. Zhang, S., Ma, X., Zhu, C., Liu, L., Wang, G. & Yuan, X. (2016) The Role of Myeloid-Derived Suppressor Cells in Patients with Solid Tumors: A Meta-Analysis, *PLoS One*. **11**, e0164514.

22. Cui, J. H., Lin, K. R., Yuan, S. H., Jin, Y. B., Chen, X. P., Su, X. K., Jiang, J., Pan, Y. M., Mao, S. L., Mao, X. F. & Luo, W. (2018) TCR Repertoire as a Novel Indicator for Immune Monitoring and Prognosis Assessment of Patients With Cervical Cancer, *Front Immunol.* **9**, 2729.
23. Liang, N., Chen, S., Yi, Y., Guan, Y.-F., Xia, X. & Yi, X. (2019) Characteristics of T-cell receptor repertoire between lung cancer patients and healthy people, *Journal of Clinical Oncology.* **37**, e20728-e20728.
24. Zhang, C., Palashati, H., Tan, Q., Ku, W., Miao, Y., Xiong, H. & Lu, Z. (2018) Immediate and substantial evolution of T-cell repertoire in peripheral blood and tumor microenvironment of patients with esophageal squamous cell carcinoma treated with preoperative chemotherapy, *Carcinogenesis.* **39**, 1389-1398.
25. Lin, K. R., Pang, D. M., Jin, Y. B., Hu, Q., Pan, Y. M., Cui, J. H., Chen, X. P., Lin, Y. X., Mao, X. F., Duan, H. B. & Luo, W. (2018) Circulating CD8(+) T-cell repertoires reveal the biological characteristics of tumors and clinical responses to chemotherapy in breast cancer patients, *Cancer Immunol Immunother.* **67**, 1743-1752.
26. Hogan, S. A., Courtier, A., Cheng, P. F., Jaberg-Bentele, N. F., Goldinger, S. M., Manuel, M., Perez, S., Plantier, N., Mouret, J. F., Nguyen-Kim, T. D. L., Raaijmakers, M. I. G., Kvistborg, P., Pasqual, N., Haanen, J., Dummer, R. & Levesque, M. P. (2019) Peripheral Blood TCR Repertoire Profiling May Facilitate Patient Stratification for Immunotherapy against Melanoma, *Cancer Immunol Res.* **7**, 77-85.
27. Spitzer, M. H., Carmi, Y., Reticker-Flynn, N. E., Kwek, S. S., Madhireddy, D., Martins, M. M., Gherardini, P. F., Prestwood, T. R., Chabon, J., Bendall, S. C., Fong, L., Nolan, G. P. & Engleman, E. G. (2017) Systemic Immunity Is Required for Effective Cancer Immunotherapy, *Cell.* **168**, 487-502 e15.
28. Markel, J. E., Noore, J., Emery, E. J., Bobnar, H. J., Kleinerman, E. S. & Lindsey, B. A. (2018) Using the Spleen as an In Vivo Systemic Immune Barometer Alongside Osteosarcoma Disease Progression and Immunotherapy with alpha-PD-L1, *Sarcoma.* **2018**, 8694397.
29. Paget, S. (1889) The distribution of secondary growths in cancer of the breast. 1889, *Cancer Metastasis Rev.* **8**, 98-101.
30. Hart, I. R. & Fidler, I. J. (1980) Role of organ selectivity in the determination of metastatic patterns of B16 melanoma, *Cancer Res.* **40**, 2281-7.
31. Schackert, G. & Fidler, I. J. (1988) Site-specific metastasis of mouse melanomas and a fibrosarcoma in the brain or meninges of syngeneic animals, *Cancer Res.* **48**, 3478-84.
32. Kaplan, R. N., Riba, R. D., Zacharoulis, S., Bramley, A. H., Vincent, L., Costa, C., MacDonald, D. D., Jin, D. K., Shido, K., Kerns, S. A., Zhu, Z., Hicklin, D., Wu, Y., Port, J. L., Altorki, N., Port, E. R., Ruggero,

521 D., Shmelkov, S. V., Jensen, K. K., Rafii, S. & Lyden, D. (2005) VEGFR1-positive haematopoietic bone
522 marrow progenitors initiate the pre-metastatic niche, *Nature*. **438**, 820-7.

523 33. Liu, Y., Gu, Y., Han, Y., Zhang, Q., Jiang, Z., Zhang, X., Huang, B., Xu, X., Zheng, J. & Cao, X. (2016)
524 Tumor Exosomal RNAs Promote Lung Pre-metastatic Niche Formation by Activating Alveolar
525 Epithelial TLR3 to Recruit Neutrophils, *Cancer Cell*. **30**, 243-256.

526 34. Oskarsson, T. (2013) Extracellular matrix components in breast cancer progression and
527 metastasis, *Breast*. **22 Suppl 2**, S66-72.

528 35. Barkan, D., Green, J. E. & Chambers, A. F. (2010) Extracellular matrix: a gatekeeper in the
529 transition from dormancy to metastatic growth, *Eur J Cancer*. **46**, 1181-8.

530 36. Erler, J. T., Bennewith, K. L., Cox, T. R., Lang, G., Bird, D., Koong, A., Le, Q. T. & Giaccia, A. J.
531 (2009) Hypoxia-induced lysyl oxidase is a critical mediator of bone marrow cell recruitment to form
532 the premetastatic niche, *Cancer Cell*. **15**, 35-44.

533 37. Hiratsuka, S., Watanabe, A., Aburatani, H. & Maru, Y. (2006) Tumour-mediated upregulation of
534 chemoattractants and recruitment of myeloid cells predetermines lung metastasis, *Nat Cell Biol*. **8**,
535 1369-75.

536 38. Costa-Silva, B., Aiello, N. M., Ocean, A. J., Singh, S., Zhang, H., Thakur, B. K., Becker, A., Hoshino,
537 A., Mark, M. T., Molina, H., Xiang, J., Zhang, T., Theilen, T. M., Garcia-Santos, G., Williams, C., Ararso,
538 Y., Huang, Y., Rodrigues, G., Shen, T. L., Labori, K. J., Lothe, I. M., Kure, E. H., Hernandez, J., Dousot,
539 A., Ebbesen, S. H., Grandgenett, P. M., Hollingsworth, M. A., Jain, M., Mallya, K., Batra, S. K.,
540 Jarnagin, W. R., Schwartz, R. E., Matei, I., Peinado, H., Stanger, B. Z., Bromberg, J. & Lyden, D. (2015)
541 Pancreatic cancer exosomes initiate pre-metastatic niche formation in the liver, *Nat Cell Biol*. **17**,
542 816-26.

543 39. Hoshino, A., Costa-Silva, B., Shen, T. L., Rodrigues, G., Hashimoto, A., Tesic Mark, M., Molina, H.,
544 Kohsaka, S., Di Giannatale, A., Ceder, S., Singh, S., Williams, C., Soplop, N., Uryu, K., Pharmed, L., King,
545 T., Bojmar, L., Davies, A. E., Ararso, Y., Zhang, T., Zhang, H., Hernandez, J., Weiss, J. M., Dumont-Cole,
546 V. D., Kramer, K., Wexler, L. H., Narendran, A., Schwartz, G. K., Healey, J. H., Sandstrom, P., Labori, K.
547 J., Kure, E. H., Grandgenett, P. M., Hollingsworth, M. A., de Sousa, M., Kaur, S., Jain, M., Mallya, K.,
548 Batra, S. K., Jarnagin, W. R., Brady, M. S., Fodstad, O., Muller, V., Pantel, K., Minn, A. J., Bissell, M. J.,
549 Garcia, B. A., Kang, Y., Rajasekhar, V. K., Ghajar, C. M., Matei, I., Peinado, H., Bromberg, J. & Lyden,
550 D. (2015) Tumour exosome integrins determine organotropic metastasis, *Nature*. **527**, 329-35.

551 40. Giles, A. J., Reid, C. M., Evans, J. D., Murgai, M., Vicioso, Y., Highfill, S. L., Kasai, M., Vahdat, L.,
552 Mackall, C. L., Lyden, D., Wexler, L. & Kaplan, R. N. (2016) Activation of Hematopoietic
553 Stem/Progenitor Cells Promotes Immunosuppression Within the Pre-metastatic Niche, *Cancer Res*.
554 **76**, 1335-47.

41. Chafe, S. C., Lou, Y., Sceneay, J., Vallejo, M., Hamilton, M. J., McDonald, P. C., Bennewith, K. L., Moller, A. & Dedhar, S. (2015) Carbonic anhydrase IX promotes myeloid-derived suppressor cell mobilization and establishment of a metastatic niche by stimulating G-CSF production, *Cancer Res.* **75**, 996-1008.
42. Sceneay, J., Chow, M. T., Chen, A., Halse, H. M., Wong, C. S., Andrews, D. M., Sloan, E. K., Parker, B. S., Bowtell, D. D., Smyth, M. J. & Moller, A. (2012) Primary tumor hypoxia recruits CD11b⁺/Ly6C^{med}/Ly6G⁺ immune suppressor cells and compromises NK cell cytotoxicity in the premetastatic niche, *Cancer Res.* **72**, 3906-11.
43. Qian, B. Z., Li, J., Zhang, H., Kitamura, T., Zhang, J., Campion, L. R., Kaiser, E. A., Snyder, L. A. & Pollard, J. W. (2011) CCL2 recruits inflammatory monocytes to facilitate breast-tumour metastasis, *Nature.* **475**, 222-5.
44. Kowanetz, M., Wu, X., Lee, J., Tan, M., Hagenbeek, T., Qu, X., Yu, L., Ross, J., Korsisaari, N., Cao, T., Bou-Reslan, H., Kallop, D., Weimer, R., Ludlam, M. J., Kaminker, J. S., Modrusan, Z., van Bruggen, N., Peale, F. V., Carano, R., Meng, Y. G. & Ferrara, N. (2010) Granulocyte-colony stimulating factor promotes lung metastasis through mobilization of Ly6G⁺Ly6C⁺ granulocytes, *Proc Natl Acad Sci U S A.* **107**, 21248-55.
45. Olkhanud, P. B., Baatar, D., Bodogai, M., Hakim, F., Gress, R., Anderson, R. L., Deng, J., Xu, M., Briest, S. & Biragyn, A. (2009) Breast cancer lung metastasis requires expression of chemokine receptor CCR4 and regulatory T cells, *Cancer Res.* **69**, 5996-6004.
46. Ouzounova, M., Lee, E., Piranlioglu, R., El Andaloussi, A., Kolhe, R., Demirci, M. F., Marasco, D., Asm, I., Chadli, A., Hassan, K. A., Thangaraju, M., Zhou, G., Arbab, A. S., Cowell, J. K. & Korkaya, H. (2017) Monocytic and granulocytic myeloid derived suppressor cells differentially regulate spatiotemporal tumour plasticity during metastatic cascade, *Nat Commun.* **8**, 14979.
47. Postow, M. A., Callahan, M. K., Barker, C. A., Yamada, Y., Yuan, J., Kitano, S., Mu, Z., Rasalan, T., Adamow, M., Ritter, E., Sedrak, C., Jungbluth, A. A., Chua, R., Yang, A. S., Roman, R. A., Rosner, S., Benson, B., Allison, J. P., Lesokhin, A. M., Gnjatic, S. & Wolchok, J. D. (2012) Immunologic correlates of the abscopal effect in a patient with melanoma, *N Engl J Med.* **366**, 925-31.
48. Wersall, P. J., Blomgren, H., Pisa, P., Lax, I., Kalkner, K. M. & Svedman, C. (2006) Regression of non-irradiated metastases after extracranial stereotactic radiotherapy in metastatic renal cell carcinoma, *Acta Oncol.* **45**, 493-7.
49. Hu, Z. I., McArthur, H. L. & Ho, A. Y. (2017) The Abscopal Effect of Radiation Therapy: What Is It and How Can We Use It in Breast Cancer?, *Curr Breast Cancer Rep.* **9**, 45-51.

587 50. Ohba, K., Omagari, K., Nakamura, T., Ikuno, N., Saeki, S., Matsuo, I., Kinoshita, H., Masuda, J.,
588 Hazama, H., Sakamoto, I. & Kohno, S. (1998) Abscopal regression of hepatocellular carcinoma after
589 radiotherapy for bone metastasis, *Gut*. **43**, 575-7.

590 51. Golden, E. B., Chhabra, A., Chachoua, A., Adams, S., Donach, M., Fenton-Kerimian, M., Friedman,
591 K., Ponzo, F., Babb, J. S., Goldberg, J., Demaria, S. & Formenti, S. C. (2015) Local radiotherapy and
592 granulocyte-macrophage colony-stimulating factor to generate abscopal responses in patients with
593 metastatic solid tumours: a proof-of-principle trial, *Lancet Oncol*. **16**, 795-803.

594 52. Abuodeh, Y., Venkat, P. & Kim, S. (2016) Systematic review of case reports on the abscopal
595 effect, *Curr Probl Cancer*. **40**, 25-37.

596 53. Stone, H. B., Peters, L. J. & Milas, L. (1979) Effect of host immune capability on radiocurability
597 and subsequent transplantability of a murine fibrosarcoma, *J Natl Cancer Inst*. **63**, 1229-35.

598 54. Demaria, S., Ng, B., Devitt, M. L., Babb, J. S., Kawashima, N., Liebes, L. & Formenti, S. C. (2004)
599 Ionizing radiation inhibition of distant untreated tumors (abscopal effect) is immune mediated, *Int J*
600 *Radiat Oncol Biol Phys*. **58**, 862-70.

601 55. Chakravarty, P. K., Alfieri, A., Thomas, E. K., Beri, V., Tanaka, K. E., Vikram, B. & Guha, C. (1999)
602 Flt3-ligand administration after radiation therapy prolongs survival in a murine model of metastatic
603 lung cancer, *Cancer Res*. **59**, 6028-32.

604 56. Ribeiro Gomes, J., Schmerling, R. A., Haddad, C. K., Racy, D. J., Ferrigno, R., Gil, E., Zanuncio, P. &
605 Buzaid, A. C. (2016) Analysis of the Abscopal Effect With Anti-PD1 Therapy in Patients With
606 Metastatic Solid Tumors, *J Immunother*. **39**, 367-372.

607 57. Twyman-Saint Victor, C., Rech, A. J., Maity, A., Rengan, R., Pauken, K. E., Stelekati, E., Benci, J. L.,
608 Xu, B., Dada, H., Odorizzi, P. M., Herati, R. S., Mansfield, K. D., Patsch, D., Amaravadi, R. K.,
609 Schuchter, L. M., Ishwaran, H., Mick, R., Pryma, D. A., Xu, X., Feldman, M. D., Gangadhar, T. C., Hahn,
610 S. M., Wherry, E. J., Vonderheide, R. H. & Minn, A. J. (2015) Radiation and dual checkpoint blockade
611 activate non-redundant immune mechanisms in cancer, *Nature*. **520**, 373-7.

612 58. Grass, G. D., Krishna, N. & Kim, S. (2016) The immune mechanisms of abscopal effect in radiation
613 therapy, *Curr Probl Cancer*. **40**, 10-24.

614 59. Vatner, R. E., Cooper, B. T., Vanpouille-Box, C., Demaria, S. & Formenti, S. C. (2014)
615 Combinations of immunotherapy and radiation in cancer therapy, *Front Oncol*. **4**, 325.

616 60. Mittal, D., Gubin, M. M., Schreiber, R. D. & Smyth, M. J. (2014) New insights into cancer
617 immunoediting and its three component phases--elimination, equilibrium and escape, *Curr Opin*
618 *Immunol*. **27**, 16-25.

619 61. Lynch, T. J., Bondarenko, I., Luft, A., Serwatowski, P., Barlesi, F., Chacko, R., Sebastian, M., Neal,
620 J., Lu, H., Cuillerot, J. M. & Reck, M. (2012) Ipilimumab in combination with paclitaxel and

carboplatin as first-line treatment in stage IIIB/IV non-small-cell lung cancer: results from a randomized, double-blind, multicenter phase II study, *J Clin Oncol.* **30**, 2046-54.

62. Rodriguez-Ruiz, M. E., Vanpouille-Box, C., Melero, I., Formenti, S. C. & Demaria, S. (2018) Immunological Mechanisms Responsible for Radiation-Induced Abscopal Effect, *Trends Immunol.* **39**, 644-655.

63. Ngwa, W., Irabor, O. C., Schoenfeld, J. D., Hesser, J., Demaria, S. & Formenti, S. C. (2018) Using immunotherapy to boost the abscopal effect, *Nat Rev Cancer.* **18**, 313-322.

64. O'Shaughnessy, M. J., Murray, K. S., La Rosa, S. P., Budhu, S., Merghoub, T., Somma, A., Monette, S., Kim, K., Corradi, R. B., Scherz, A. & Coleman, J. A. (2018) Systemic Antitumor Immunity by PD-1/PD-L1 Inhibition Is Potentiated by Vascular-Targeted Photodynamic Therapy of Primary Tumors, *Clin Cancer Res.* **24**, 592-599.

65. Ehrlich, P. & Apolant, H. (1905) *Beobachtungen über maligne Mäusetumoren*, L. Schumacher.

66. O'Reilly, M. S., Holmgren, L., Shing, Y., Chen, C., Rosenthal, R. A., Moses, M., Lane, W. S., Cao, Y., Sage, E. H. & Folkman, J. (1994) Angiostatin: a novel angiogenesis inhibitor that mediates the suppression of metastases by a Lewis lung carcinoma, *Cell.* **79**, 315-28.

67. Walker, R., Poleszczuk, J., Pilon-Thomas, S., Kim, S., Anderson, A., Czerniecki, B. J., Harrison, L. B., Moros, E. G. & Enderling, H. (2018) Immune interconnectivity of anatomically distant tumors as a potential mediator of systemic responses to local therapy, *Sci Rep.* **8**, 9474.

68. Devaud, C., John, L. B., Westwood, J. A., Yong, C. S., Beavis, P. A., Schwendener, R. A., Darcy, P. K. & Kershaw, M. H. (2015) Cross-talk between tumors can affect responses to therapy, *Oncoimmunology.* **4**, e975572.

69. Taggart, D., Andreou, T., Scott, K. J., Williams, J., Rippaus, N., Brownlie, R. J., Ilett, E. J., Salmond, R. J., Melcher, A. & Loriger, M. (2018) Anti-PD-1/anti-CTLA-4 efficacy in melanoma brain metastases depends on extracranial disease and augmentation of CD8(+) T cell trafficking, *Proc Natl Acad Sci U S A.* **115**, E1540-E1549.

70. Liu, J., Blake, S. J., Yong, M. C., Harjunpaa, H., Ngiow, S. F., Takeda, K., Young, A., O'Donnell, J. S., Allen, S., Smyth, M. J. & Teng, M. W. (2016) Improved Efficacy of Neoadjuvant Compared to Adjuvant Immunotherapy to Eradicate Metastatic Disease, *Cancer Discov.* **6**, 1382-1399.

71. Chang, E. & Rosenberg, S. A. (2001) Patients with melanoma metastases at cutaneous and subcutaneous sites are highly susceptible to interleukin-2-based therapy, *J Immunother.* **24**, 88-90.

72. Lee, J. C.-C., Tsai, K. K., Algazi, A. P., Rosenblum, M., Bluestone, J. & Daud, A. (2017) Relationship between liver metastases and PD-1 blockade in melanoma in, American Society of Clinical Oncology,

73. Amaria, R. N., Reddy, S. M., Tawbi, H. A., Davies, M. A., Ross, M. I., Glitza, I. C., Cormier, J. N., Lewis, C., Hwu, W. J., Hanna, E., Diab, A., Wong, M. K., Royal, R., Gross, N., Weber, R., Lai, S. Y.,

Ehlers, R., Blando, J., Milton, D. R., Woodman, S., Kageyama, R., Wells, D. K., Hwu, P., Patel, S. P., Lucci, A., Hessel, A., Lee, J. E., Gershenwald, J., Simpson, L., Burton, E. M., Posada, L., Haydu, L., Wang, L., Zhang, S., Lazar, A. J., Hudgens, C. W., Gopalakrishnan, V., Reuben, A., Andrews, M. C., Spencer, C. N., Prieto, V., Sharma, P., Allison, J., Tetzlaff, M. T. & Wargo, J. A. (2018) Neoadjuvant immune checkpoint blockade in high-risk resectable melanoma, *Nat Med.* **24**, 1649-1654.

74. Cloughesy, T. F., Mochizuki, A. Y., Orpilla, J. R., Hugo, W., Lee, A. H., Davidson, T. B., Wang, A. C., Ellingson, B. M., Rytlewski, J. A., Sanders, C. M., Kawaguchi, E. S., Du, L., Li, G., Yong, W. H., Gaffey, S. C., Cohen, A. L., Mellinghoff, I. K., Lee, E. Q., Reardon, D. A., O'Brien, B. J., Butowski, N. A., Nghiemphu, P. L., Clarke, J. L., Arrillaga-Romany, I. C., Colman, H., Kaley, T. J., de Groot, J. F., Liao, L. M., Wen, P. Y. & Prins, R. M. (2019) Neoadjuvant anti-PD-1 immunotherapy promotes a survival benefit with intratumoral and systemic immune responses in recurrent glioblastoma, *Nat Med.* **25**, 477-486.

febs_15316_f1.tif

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Oliver, AJ;Darcy, PK;Trapani, JA;Kershaw, MH;Slaney, CY

Title:

Cross-talk between tumors at anatomically distinct sites

Date:

2020-04-21

Citation:

Oliver, A. J., Darcy, P. K., Trapani, J. A., Kershaw, M. H. & Slaney, C. Y. (2020). Cross-talk between tumors at anatomically distinct sites. FEBS JOURNAL, 288 (1), pp.81-90. <https://doi.org/10.1111/febs.15316>.

Persistent Link:

<http://hdl.handle.net/11343/275671>