Heads Up: Children may be less affected than adults by novel coronavirus (COVID-19)

Author: Kevat, A¹

1. *Sydney Children's Hospital, Sydney, Australia*

Corresponding Author:

Dr Ajay Kevat

Respiratory Staff Specialist

Sydney Children's Hospital

High Street, Randwick, NSW 2031

Australia

Phone: 02 9382 1477

Fax: 02 9382 1787

Email: ajay.kevat@health.nsw.gov.au

Keywords: infant, newborn

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1111/jpc.14876](https://doi.org/10.1111/jpc.14876)

Heads Up

Children may be less affected than adults by novel coronavirus (COVID-19)

COVID-19 is a new strain of coronavirus that has infected people from many countries around the world. Initial reports suggest that children are relatively spared by this virus in comparison to adults. The February 2020 World Health Organization-China Joint Mission on Coronavirus Disease (1) found that only 2.4% of cases were in those less than nineteen years of age. Furthermore, disease severity was less in infected children compared to the total infected population; only 2.5% of children developed severe disease (compared to 13.8% overall) and 0.2% of children developed critical disease (compared to 6.1% overall). Severe disease was defined as dyspnoea, tachypnoea, hypoxia or infiltrates affecting >50% of the lung fields within 48 hours, and critical disease was defined as respiratory failure, septic shock and/or multi-organ failure. Although reported case numbers are quite small, young infants seem to have relatively low rates of being severely affected; one study of nine infected infants found none required intensive care or had significant complications (2). Despite this, certain paediatric populations such as extremely prematurely-born babies are likely to be particularly vulnerable, with one death confirmed in this subgroup (3). Further evidence is likely to emerge with time on just how children as a cohort globally will fare with this epidemic illness.

References

(1) World Health Organization-China Joint Mission on Coronavirus Disease. *WHO*; 2020
Feb 16-24; p12

(2) Wei M et al. *JAMA*; 2020 Feb 14 (online); doi: 10.1001/jama.2020.2131

(3) Zhu H et al. *Transl Pediatr* 2020; 9(1): 51-60 doi: 10.21037/tp.2020.02.06

Reviewer: Dr Ajay Kevat, ajay.kevat@health.nsw.gov.au, Respiratory Staff Specialist,
Sydney Children's Hospital, Australia

Heads Up: Children may be less affected than adults by novel coronavirus (COVID-19)

Author: Kevat, A¹

1. *Sydney Children's Hospital, Sydney, Australia*

Corresponding Author:

Dr Ajay Kevat

Respiratory Staff Specialist

Sydney Children's Hospital

High Street, Randwick, NSW 2031

Australia

Phone: 02 9382 1477

Fax: 02 9382 1787

Email: ajay.kevat@health.nsw.gov.au

Keywords: infant, newborn

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

McMahon, J;Harvey, A;Reid, SM;May, T;Antolovich, G

Title:

Anxiety in children and adolescents with cerebral palsy

Date:

2020-05-15

Citation:

McMahon, J., Harvey, A., Reid, S. M., May, T. & Antolovich, G. (2020). Anxiety in children and adolescents with cerebral palsy. JOURNAL OF PAEDIATRICS AND CHILD HEALTH, 56 (8), pp.1194-1200. <https://doi.org/10.1111/jpc.14879>.

Persistent Link:

<http://hdl.handle.net/11343/275756>