ASSOC. PROF. LEON J WORTH (Orcid ID : 0000-0003-4304-6845)

PROF. DEBORAH ANNE WILLIAMSON (Orcid ID : 0000-0001-7363-6665)

Article type : Lessons from practice

Article begins on page three of this document.

Title	<i>Candida auris</i> in an Australian health care facility: importance of screening high risk patients
-------	--

Authors:

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/MJA2.50612](https://doi.org/10.1002/MJA2.50612)

This article is protected by copyright. All rights reserved

	Title	First name	Mid inits	Last name	Postnom (eg, PhD)	Position1	Address1	Position2	Address2	Tel	Email
1	Asso c. Prof.	Leon	J	Worth	MBBS, FRACP, PhD, Grad Dip Epi	Infectious diseases physician	1		2		leon.worth@mh.org.au
2	Asso c. Prof.	Simon	J	Harrison	MBBS, MRCP (UK), PhD, FRCPath (UK), FRACP		3			03 9656 1076	simon.harrison@petermac.org
3	Dr.	Michael		Dickinson	MBBS (Hons), D.Med Sci, FRACP, FRCPA	Haematologist	3				michael.dickinson@petermac.org
4	Dr.	Annaliese		van Diemen			4				Annaliese.VanDiemen@dhhs.vic.gov.au
5	Ms.	Jennifer		Breen	RN	Infectious diseases physician	3				jennifer.breen@petermac.org
6	Ms.	Susan		Harper	RN	Infectious diseases physician	3				susan.harper@petermac.org
7	Asso c. Prof.	Caroline		Marshall	MBBS, FRACP, PhD, Grad Dip Clin Epi	Principal Research Fellow	5	Physician	6	03 9342 8897	caroline.marshall@mh.org.au
8	Prof.	Deborah	A	Williamson	MB ChB, PhD, BSc (Hons), MRCP, FRCPA	Deputy Director	7				deborah.williamson@unimelb.edu.au
9	Prof.	Karin	A	Thursky	MBBS, FRACP, MD	Research manager	2		3		karin.thursky@petermac.org
10	Prof.	Monica	A	Slavin	MBBS, FRACP, MD	Head, Infectious Diseases	2		3	03 9656 1707	monica.slavin@petermac.org

Number of corresponding author:	1
Number of alternative corresponding author:	

Addresses:

	Institution	Postal address	City	State	Post Code	Nation
1	Victorian Healthcare Associated Infection Surveillance Coordinating Centre	Doherty Institute	Melbourne	VIC	3000	
2	National Centre for Infections in Cancer	Grattan Street	Melbourne	VIC	3000	
3	Peter MacCallum Cancer Centre	Grattan Street	Melbourne	VIC	3000	
4	Victorian Department of Health and Human Services	Lonsdale Street	Melbourne	VIC	3000	
5	Victorian Infectious Diseases Service, Royal Melbourne Hospital	RMH	Melbourne	VIC	3050	7
6	University of Melbourne	Royal Melbourne Hospital	Melbourne	VIC	3050	
7	Microbiological Diagnostic Unit Public Health Laboratory, Doherty Institute	792 Elizabeth Street	Melbourne	VIC	3000	

Postal address of first corresponding author (if different from the institutional address given above)	
--	--

Primary Keywords [Office use only]	Infectious diseases; Pharmaceutical preparations; Environment and public health; Medical genetics
Secondary keywords [Office use only]	Infectious diseases; Infection control; Drug resistance, microbial; Public health; Sequence analysis
Notes:	

Article details (press ctrl – 9 to enter details):

Article type	Lessons from Practice
Blurb	Timely screening and control is required in high risk settings
Pullquote	

Office use

<i>Ms. Number</i>	mja19.00271. R4
<i>Medical editor</i>	Wendy Morgan
<i>Medical editor email</i>	wmorgan@mja.com.au
<i>Structural editor</i>	Graeme Prince
<i>Structural editor email</i>	gprince@mja.com.au
<i>Section/Category</i>	Medical education/Lessons from practice
<i>Strapheading</i>	Medical education
<i>Substrap</i>	Lessons from practice

Wiley – file data:

Filename for copyediting	wor_mja19.00271_ms
Accompanying graphics	
Stock images	
Appendices	

Office use – history:

Event	Date
Original submission received	22/03/2019

Event	Date
Accept	06/12/2019

Proof sent to author	
Proof returned by author	
Published (date format xx/xx/xx)	15/05/20
Issue	11
Vol	212
DOI	10.5694/mja19.00271
Journal	The Medical Journal of Australia
Original article DOI (for response)	

Candida auris in an Australian health care facility: importance of screening high risk patients

Clinical record

A 70-year-old man with multiple myeloma was admitted to our hospital in 2018, having been hospitalised 10 months previously in the United Kingdom. Following admission to our facility, routine collection of clinical specimens was performed in the setting of an episode of febrile neutropenia. *Candida auris* was isolated in a urine specimen collected in the presence of an indwelling urinary catheter, without accompanying pyuria.

Screening of ward contacts ($n = 73$) was subsequently performed by collection of composite axilla and groin skin swabs, together with swabbing of possible clinical sites of infection (eg, wounds, catheter sites). Swabs were plated onto *Candida* chromogenic agar and incubated aerobically for 48 hours at 35°C. Any colonies not typical for *C. albicans* or *C. tropicalis* were identified using matrix-assisted laser desorption ionisation time-of-flight (MALDI-TOF) mass spectrometry. The routine regimen of daily cleaning and disinfection of rooms with 1000 ppm sodium hypochlorite solution was continued. Enhanced infection control measures, including contact precautions and single-room isolation were instituted. A multidisciplinary taskforce coordinated screening, laboratory and prevention strategies. Review of laboratory reports for the preceding 12 months confirmed this to be the first documented *C. auris* isolate at our facility.

One ward contact, a 38-year-old man with diffuse large B cell lymphoma, was identified as colonised with *C. auris*. The organism was detected in a urine specimen collected in the presence of a long term indwelling urinary catheter. This patient had been admitted to a health care facility in the United Arab Emirates, before direct transfer to our facility about 3 months earlier.

Colonised patients had been located in a common ward for 19 days, each in a single room with dedicated bathroom and patient care equipment. They had also been managed on an outlying ward for brief periods (3 and 2 days, respectively) separated in time by 2 days. Neither patient developed clinical features of urinary tract or disseminated *C. auris* infection and antifungal therapy was not administered.

Isolates were confirmed as *C. auris* by MALDI-TOF mass spectrometry (each with score of 1.75). Antifungal susceptibility testing by broth microdilution demonstrated isolates were resistant to fluconazole (minimum inhibitory concentration [MIC] > 256 mg/L) and susceptible to caspofungin (MIC, 0.25 mg/L) and anidulafungin (MIC, 0.12 mg/L for Patient 1 and 0.25 mg/L for Patient 2). To investigate relatedness of isolates, whole genome sequencing and bioinformatics analysis were performed. Phylogeographic analysis demonstrated that both were related globally to those contained

in the India–Pakistan clade. The median pairwise single nucleotide polymorphism distance between the two isolates was 167, suggesting that while these isolates were related, it was not possible to confirm whether transmission had occurred.

Discussion

Candida auris is an emerging, drug-resistant yeast, responsible for hospital outbreaks internationally.¹ First recognised as a new species of *Candida* in 2009, cases have been reported in over 30 countries, including the United Kingdom and United Arab Emirates.^{1,2} In outbreak settings, bloodstream, urinary tract and deep tissue infections have been reported, in addition to colonisation. The majority of isolates are fluconazole resistant,³ with variable resistance to amphotericin B and the echinocandin class of antifungal agents. Infection is associated with a crude mortality of 30%.³ Key differences between *Candida albicans* (the most frequently identified *Candida* species in Australia) and *C. auris* are summarised in the Box.

Risks for *C. auris* acquisition include admission to a high dependency unit, presence of invasive medical devices, underlying immunocompromise or chronic disease and receipt of antibiotic or antifungal agents.⁴ One case of *C. auris* invasive disease has previously been reported in Australia,⁵ but to our knowledge the two cases identified at our facility represent the first possible transmission of *C. auris* in Australia.

Identification of *C. auris* is challenging, with potential misidentification by routine biochemical methods. If *C. auris* is included in the reference profile database, MALDI-TOF mass spectrometry may be used to confirm diagnosis. DNA sequencing also provides confirmation, together with data regarding origins and potential transmission in health care settings.³

Collection of bilateral axilla and groin skin swabs as a combined screening specimen is recommended for optimal yield.⁶ European and United States guidelines recommend screening of all room contacts of patients with *C. auris*.^{6,7} Screening of additional patients (eg, whole ward) is necessary where more than one case is identified. Targeted surveillance of patients who have recently had at least one overnight stay in an overseas facility is also recommended, especially if from a country reporting *C. auris* cases.^{6,7} Our experience highlights the importance of this strategy.

Clinicians should be aware of risks for *C. auris* acquisition, including overseas health care encounters. In high risk settings, and where a case of *C. auris* infection has been identified, timely screening of patients is required to ensure that appropriate control measures are instituted.

Lessons from practice

- *Candida auris* is an emerging drug-resistant yeast, now reported in Australian health care facilities.

- In contrast to *C. albicans*, which is commonly isolated in community and health care settings, *C. auris* is generally only identified in high risk hospitalised populations. Risks for acquisition include intensive care or high dependency unit admission, presence of invasive medical devices, underlying immunocompromise or chronic disease, and receipt of broad spectrum antibiotics or antifungal agents.
 - Strict infection control measures, including contact precautions and isolation, are required to reduce risks of transmission. Screening for colonisation is an important element of infection control strategies, and a composite skin swab of axilla and groin is recommended.
 - Timely detection requires laboratory identification. MALDI-TOF mass spectrometry may be used for confirmation, and whole genome sequencing may provide additional information on possible transmission events.
 - Health care facilities must ensure processes are implemented for screening of patients who have received health care in overseas hospitals.
-

Acknowledgements: We thank Dr Sarah Kidd for her assistance.

Competing interests: No relevant disclosures.

Provenance: Not commissioned; externally peer reviewed.

Author details

Leon J Worth^{1,2}

Simon J Harrison³

Michael Dickinson³

Annaliese van Diemen⁴

Jennifer Breen³

Susan Harper³

Caroline Marshall^{5,6}

Deborah A Williamson⁷

Karin A Thursky^{2,3}

Monica A Slavin^{2,3}

1 Victorian Healthcare Associated Infection Surveillance Coordinating Centre, Melbourne, VIC.

2 National Centre for Infections in Cancer, Melbourne, VIC.

3 Peter MacCallum Cancer Centre, Melbourne, VIC.

4 Victorian Department of Health and Human Services, Melbourne, VIC.

5 Victorian Infectious Diseases Service, Royal Melbourne Hospital, Melbourne, VIC, 7.

6 University of Melbourne, Melbourne, VIC.

7 Microbiological Diagnostic Unit Public Health Laboratory, Doherty Institute, Melbourne, VIC.

leon.worth@mh.org.au

doi: 10.5694/mja19.00271

References

- 1 Eyre DW, Sheppard AE, Madder H, et al. A *Candida auris* outbreak and its control in an intensive care setting. *N Engl J Med* 2018; 379: 1322-1331.

- 2 Chow NA, Muñoz JF, Gade L, et al. Tracing the evolutionary history and global expansion of *Candida auris* using population genomic analyses. *mBio* 2020; 11. e03364-19.
- 3 Osei Sekyere J. *Candida auris*: a systematic review and meta-analysis of current updates on an emerging multidrug-resistant pathogen. *Microbiologyopen* 2018; 7: e00578.
- 4 Ruiz-Gaitan A, Moret AM, Tasiias-Pitarch M, et al. An outbreak due to *Candida auris* with prolonged colonisation and candidaemia in a tertiary care European hospital. *Mycoses* 2018; 61: 498-505.
- 5 Heath CH, Dyer JR, Pang S, et al. *Candida auris* sternal osteomyelitis in a man from Kenya visiting Australia, 2015. *Emerg Infect Dis* 2019; 25: 192-194.
- 6 Centers for Disease Control and Prevention. Infection prevention and control for *Candida auris*. Atlanta, GA: CDC, 2018. <https://www.cdc.gov/fungal/candida-auris/c-auris-infection-control.html> (viewed Oct 2018).
- 7 Public Health England. Guidance for the laboratory investigation, management and infection prevention and control for cases of *Candida auris*. August 2017 v2.0. London: PHE, 2017. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/637685/Updated_Candida_auris_Guidance_v2.pdf (viewed Oct 2018).

[Box]

Comparison of clinical and epidemiological characteristics of *Candida albicans* and *Candida auris*

	<i>Candida albicans</i>	<i>Candida auris</i>
Colonisation	Colonisation of patients in community and health care settings is common; a commensal of skin and gut of immunocompetent and immunocompromised hosts	Colonisation of patients associated only with hospital outbreaks or transmission, also identified in environment and equipment in hospital outbreak settings
Infection	Infection most frequently at mucosal sites (eg, oropharyngeal, vulvovaginal); bloodstream and urinary tract infections less frequent	Bloodstream, urinary tract and wound infections reported
Risks for infection	ICU or HDU admission, invasive medical devices, major abdominal surgery, solid tumours, haematological malignancies, broad spectrum antibiotics	ICU or HDU admission, invasive medical devices, underlying immunocompromise or chronic disease (eg, diabetes, chronic lung disease, renal failure, cardiovascular disease, or malignancy), broad spectrum antibiotics or antifungal agents
Geographical distribution	Ubiquitous, community and health care settings	Reported only in health care settings, expanding global distribution
Laboratory identification	Culture using selective chromogenic media	Culture together with MALDI-TOF or DNA sequencing
Antifungal resistance	Generally susceptible to fluconazole	Resistance to fluconazole is likely*

HDU = high dependency unit; ICU = intensive care unit; MALDI-TOF = matrix-assisted laser desorption/ionisation time-of-flight mass spectrometer. * Note: agreed fluconazole minimum inhibitory concentration breakpoints for *C. auris* have not been established

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Worth, LJ;Harrison, SJ;Dickinson, M;van Diemen, A;Breen, J;Harper, S;Marshall, C;Williamson, DA;Thursky, KA;Slavin, MA

Title:

Candida auris in an Australian health care facility: importance of screening high risk patients

Date:

2020-05-24

Citation:

Worth, L. J., Harrison, S. J., Dickinson, M., van Diemen, A., Breen, J., Harper, S., Marshall, C., Williamson, D. A., Thursky, K. A. & Slavin, M. A. (2020). Candida auris in an Australian health care facility: importance of screening high risk patients. MEDICAL JOURNAL OF AUSTRALIA, 212 (11), pp.510-+. <https://doi.org/10.5694/mja2.50612>.

Persistent Link:

<http://hdl.handle.net/11343/275780>