Title Page

Title: Wilson and Homer reply to Papadakis

Author names:

DR ALYCE N WILSON, Public Health Registrar, Burnet Institute, Melbourne*

Lecturer, School of Population and Global Health, Faculty of Medicine, Dentistry and Health Sciences, The University of Melbourne

Adjunct lecturer, Department of Food, Nutrition and Dietetics, Monash University, Melbourne

PROFESSOR CAROLINE SE HOMER, Co-Program Director Maternal and Child Health, Burnet Institute, Melbourne

Distinguished Visiting Professor of Midwifery, Faculty of Health, University of Technology Sydney

Adjunct Professor, School of Public Health and Preventive Medicine, Monash University, Melbourne

Honorary (Professorial Fellow), School of Population and Global Health, Faculty of Medicine, Dentistry and Health Sciences, The University of Melbourne

caroline.homer@burnet.edu.au

*Corresponding author:

alyce.wilson@burnet.edu.au

85 Commercial Rd, Prahran VIC 3181

+61 3 9282 2111

Conflict of interest:

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/AJO.13164](https://doi.org/10.1111/AJO.13164)

This article is protected by copyright. All rights reserved

The authors do not have any conflicts of interest to declare.

Acknowledgements

The authors were commissioned to undertake this review by the Australian Commission for Safety and Quality in Health Care and have permission to publish this paper.

DR. ALYCE N WILSON (Orcid ID : 0000-0001-9229-789X)

Article type : Letter to the Editor

Letter to the Editor

Dear Editor

Thank you for your letter regarding consideration of an additional peripartum parameter, which could contribute to improving clinical outcomes and your description of the Anterior Ninety-degree Elevation Forceps (ANEF) technique.¹ Myrikas and Papadakis in their 2018 paper describe the ANEF approach and their experience. They present their cumulative data in 360 consecutive women where the Neville–Barnes ANEF approach was used over the past 4.5 years on women having their first baby. In this population, the rate of severe perineal trauma was 1.9% (n=7).¹ As we are not specialist obstetricians we will not make a comment about the ANEF approach per se but will leave that to our obstetric colleagues.

We did not include this paper or the work of these authors as ours was a review of recently published systematic reviews (within the last 5 years).

Nonetheless, we completely agree that good communication and clinical skills are critical to improving prevention and management of third- and fourth-degree tears and welcome research in this field, especially in an Australian and New Zealand context. Communication – between the woman and the clinicians, between the clinicians and her partner and support people, between the obstetrician and the midwives, and within the discipline teams - are all crucial to ensure teamwork and patient safety remain woman centred and focussed.

Kind regards,

Alyce Wilson and Caroline Homer

1. Myrikas S, Papadakis K. Anterior non-episiotomy or natural forceps delivery: refining the technique and improving communication as a way of reducing obstetric anal sphincter injuries in instrumental deliveries. *Journal of Pelvic, Obstetric and Gynaecological Physiotherapy*, 2018; 122: 50–55.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Wilson, AN;Homer, CSE

Title:

Wilson and Homer reply to Papadakis

Date:

2020-06-01

Citation:

Wilson, A. N. & Homer, C. S. E. (2020). Wilson and Homer reply to Papadakis. AUSTRALIAN & NEW ZEALAND JOURNAL OF OBSTETRICS & GYNAECOLOGY, 60 (3), pp.E7-E7. <https://doi.org/10.1111/ajo.13164>.

Persistent Link:

<http://hdl.handle.net/11343/275809>