Susceptibility of the bird cherry-oat aphid, *Rhopalosiphum padi* (Hemiptera: Aphididae), to four insecticides

Paul A Umina^{1,2,*}, Jenny Reidy-Crofts³, Marielle Babineau¹, James L Maino¹ and Owain R Edwards³

¹cesar, 293 Royal Parade, Parkville, Victoria 3052, Australia

² School of BioSciences, The University of Melbourne, Victoria 3010, Australia

³ CSIRO Land & Water, Floreat, Western Australia 6014, Australia

* Corresponding author: pumina@unimelb.edu.au ph.: +613 9349 4723

Running title: Tolerance to insecticides in oat aphids

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/aen.12490](https://doi.org/10.1111/aen.12490)

Abstract

The bird-cherry oat aphid (*Rhopalosiphumpadi*) is a global pest, attacking most cereal crops including barley, wheat, oats and triticale. The aphids cause yield losses through direct feeding damage and the transmission of plant viruses. In Australia, feeding injury can reduce cereal yields by 6%, with the damage caused by aphid-vectored viruses reducing the yield of cereal crops by up to 30%. Aphid control in these crops is achieved almost exclusively with insecticides, and there is growing concern surrounding insecticide resistance evolution in multiple aphid species. In this study, nine field populations of *R. Padi* were collected from localities representing the major grain growing regions of Australia. Toxicity data against four insecticides (dimethoate, alpha-cypermethrin, pirimicarb, imidacloprid) was generated for each aphid population. This revealed little differences in population responses for three of these insecticides. For alpha-cypermethrin, a widely used insecticide in Australia, there were significant differences between several populations. These data will be important for future monitoring of insecticide responses of *R. Padi* and highlight the impending pest management challenges growers could encounter in Australia.

Key words: toxicity data, oat aphid, insecticides, resistance monitoring

Introduction

The bird cherry-oat aphid, *Rhopalosiphum padi* (Linnaeus, 1758), is one of the most important cereal pests worldwide. While it can be found feeding on ~20 plant families, it is a pest mainly on Poaceae (particularly wheat, barley, corn, oats and triticale) and to a lesser extent Rosaceae crops (Blackman and Eastop 2000). In Australia, *R. Padi* is present throughout all major grain growing regions and can cause damage through both direct feeding and by transmitting more than a dozen plant viruses, including Barley yellow dwarf virus (BYDV) (Lucio-Zavala *et al.* 2001; Bailey 2007; Edwards *et al.* 2008; Schliephake *et al.* 2013). Both types of damage significantly reduce the quality and yield of cereal crops. Direct feeding by *R. Padi* and other cereal aphids can reduce yield by 6%, while damage from aphid-vectored viruses can reduce yield by up to 30% in cereal crops such as barley, oats and wheat (Valenzuela & Hoffmann 2015).

In Australian cereals, insecticides are the most commonly used management tool to control *R. padi*, with growers largely relying on foliar applications of organophosphates, synthetic pyrethroids and carbamates, as well as neonicotinoids applied as a seed treatment prior to sowing (Edwards *et al.* 2008; GRDC 2010; Kirkland *et al.* 2018; APVMA 2020). Synthetic pyrethroids and organophosphates are estimated to account for >85% of insecticide use in cereal crops in Australia (Umina *et al.* 2019). Pyrethroids in particular are widely used prophylactically in cereals to reduce the transmission of aphid-vectored diseases including BYDV (Edwards *et al.* 2008). Neonicotinoids, predominantly imidacloprid, have become commonplace in many cereal-growing regions, particularly in areas where virus risk is deemed high and/or Russian wheat aphid, *Diuraphis noxia* (Mordvilko, 1913), is present (Yazdani *et al.* 2018; Umina *et al.* 2019). Pirimicarb, a carbamate insecticide that is highly effective against aphids (Xiao *et al.* 2015), is also registered and used (although to a lesser extent) against *R. padi* and other cereal aphids (APVMA 2020; Umina *et al.* 2019).

Rhopalosiphum padi is thought to have been introduced to Australia more than 100 years ago and is now widespread and very abundant across the country (Valenzuela *et al.* 2010; Macfadyen & Kriticos, 2012). In many parts of the world, *R. padi* is heteroecious holocyclic, whereby its lifecycle alternates between a sexual and asexual phase on a primary and a secondary host, respectively (Halkett *et al.* 2006). Although males are present in Australia, the primary host essential for mating is rare and population genetic studies have shown *R. padi* to be anholocyclic, with parthenogenetic females reproducing only by viviparity. There is also evidence of a very low proportion of androcyclic clones

(where males and viviparous females are produced) (Hughes *et al.* 1964; Simon *et al.* 1996; Delmotte *et al.* 2003; Valenzuela *et al.* 2010). In addition to cereal crops, grasses play a critical role in the lifecycle and biology of *R. padi*. Perennial grass pastures in particular are thought to be an important refuge for *R. padi* in some parts of Australia. de Barro *et al.* (1992) demonstrated irrigated pastures are capable of producing very large numbers of alates during late autumn, placing nearby cereal crops at risk from infestation of migrating aphids; importantly this coincides with the early establishment stage of winter cereals. Annual grasses are also considered important as they have the potential to link aphids in perennial grass pastures with cereal crops and/or by allowing low numbers of *R. padi* to persist between cropping seasons over summer (de Barro & Wallwork 1992). These grasses (and volunteer cereals) often persist during summer months interspersed within fields, in headlands and along roadside verges.

There is some conjecture regarding insecticide resistance in *R. padi*. There are currently no documented cases of field-evolved resistance captured in the Arthropod Pesticide Resistance Database, a global resource that captures insecticide resistance cases from 1914 to present (Mota-Sanchez & Wise 2020). However, there are multiple publications that claim low-level resistance is evolving in China (Cheng *et al.* 2004; Zhang *et al.* 2016; Zuo *et al.* 2016). There are also several studies that have selected for high-level resistance through repeated insecticide exposure in the laboratory (e.g. Chen *et al.* 2007; Wang *et al.* 2016; Wang *et al.* 2018), demonstrating the evolutionary capacity for resistance to evolve in *R. padi*. Furthermore, a recent analysis conducted on Australian grain pests which considered the evolutionary potential of resistance, targeted and off-target insecticide pressure, and the availability of pest refuges, identified *R. padi* as a species at risk of evolving resistance within Australia (McDonald *et al.* 2019).

An insecticide study conducted four years ago found that Australian *R. Padi* populations had relatively low sensitivity across a range of insecticides commonly applied to grain crops (Clouston *et al.* 2016). Importantly, Clouston *et al.* (2016) identified *R. padi* as one of two species (the other being *Aphis craccivora* Koch, 1854) that should be the focus of continued monitoring for insecticide resistance to strengthen future risk-mitigation tactics. The generation of regular toxicity data for high risk pests is paramount if growers are to implement proactive and long-term insecticide management strategies (Umina *et al.* 2019). Here, we conduct a comprehensive study to establish current insecticide toxicity data for field populations of *R. Padi* collected from all major grain growing regions of Australia.

Materials and Methods

Aphid collections and culturing

Between January and December 2017, *R.padi* were collected from agricultural regions of New South Wales (NSW), Victoria (Vic.), South Australia (SA) and Western Australia (WA). At each location, global positioning system (GPS) coordinates and plant host were recorded. Aphids at each location were sampled from multiple plants to increase the likelihood of collecting multiple clonal types. Aphids were collected from wheat, barley, oats or grass weeds (Table 1). As with most invasive aphid populations (Figuerola *et al.* 2018), *R. padi* in Australia has limited clonal variability and populations tend to be characterised by a small number of dominant clonal lineages, which show no clear geographic or host-related distributions (Valenzuela *et al.* 2010).

Field populations were isolated in the laboratory to ensure the absence of parasitoid wasps, using the method described in Umina *et al.* (2014), except we used oat (*Avenasativa*) leaf sections instead of sprouting radish. Individual aphid populations were then maintained in 55mm petri dishes containing 10 g/L agar inserted with excised *A. sativa* leaf sections. Petri dishes were kept in a Controlled Temperature (CT) cabinet set at 10°C, with a L14:D10h photoperiod; these conditions are ideal for maintaining healthy aphid populations but minimising the number of generations that occur in the laboratory. Aphids were transferred to new petri dishes every 14 days.

To produce sufficient numbers for bioassays, aphids were transferred to wheat plants (*Triticumaestivum*) grown in potting mix in 900mL pots. Both oat and wheat are recognised as highly suitable host plants for *R. padi*, which is considered to be polyphagous on C3 grasses and crops. Both species are routinely used as *R. Padi* hosts in laboratory rearing and insecticide assays. Additionally, these hosts do not differentially affect insecticide efficacy (Araya & Foster 1987). Pots containing wheat plants were placed inside insect-proof cages (45cm H, 28cm W, 38cm L) and kept under natural light at approximately 20°C until aphid numbers were sufficient (typically 21 days). Under these unconstrained growth conditions, there is unlikely to be strong selection against tolerant (or resistant) aphid lineages. We have not observed any phenotypic or genetic changes over time in colonies of other aphid species maintained this way and repeatedly assayed as known resistant or susceptible controls (e.g. de Little & Umina 2017; Umina *et al.* 2014).

Insecticides tested

Aphid populations were screened against insecticides representing four chemical modes of action: neonicotinoids (Group 4A) represented by imidacloprid, carbamates (Group 1B) represented by pirimicarb, organophosphates (Group 1A) represented by dimethoate, and synthetic pyrethroids (Group 3A) represented by alpha-cypermethrin (Table 2). A stock solution of the recommended field rate of each chemical product was prepared in water, and serially diluted to represent five concentrations ranging from 1 to 10^{-4} times the field rate. The recommended field rates were based on recommendations for aphid control, assuming an application in 100L water/ha. These equate to 525 mg a.i./L imidacloprid, 5000mg a.i./L pirimicarb, 2000mg a.i./L dimethoate and 125 mg a.i./L alpha-cypermethrin.

Bioassay procedures

The laboratory bioassays undertaken closely followed those described by Umina *et al.* (2014), Clouston *et al.* (2016) and the Insecticide Resistance Action Committee (IRAC) Susceptibility Test Method No. 019 (IRAC 2016). The plant host of *T.aestivum* was used to emulate insecticide contact in a leaf dip bioassay for each chemical product. For imidacloprid, a single excised *T.aestivum* leaf section (~25mm in length) was submerged for 10s in 0.0525 mg a.i./L solution, placed abaxial side up on 10 g/L agar in a 35mm petri dish and air dried. This was performed a further four times, resulting in five replicate petri dishes, each containing a single treated leaf. This was then repeated for each of the remaining imidacloprid solutions (moving sequentially from the lowest concentrations to the field rate). Five untreated control replicates were established in the same manner, except water was used in place of the chemical solution. Once dry, apterous aphids from a single population were transferred to each petri dish using a fine-haired paintbrush. Between eight and 10 adult aphids were added to each dish. Lids were then placed on the petri dish plates and the dishes inverted to simulate natural aphid feeding on the underside of leaves. These were placed at 20°C with a L14:D10h photoperiod for 48 h, after which aphids were scored visually as dead (not moving over 5s period), incapacitated (inhibited movement), or alive (moving freely). The above procedure was repeated for the remaining insecticides, with three populations tested simultaneously on the same day. In all cases, insecticide stock solutions were made fresh on

each bioassay day, and these were then serially diluted to generate fresh chemical concentrations for each bioassay.

Bioassays were performed for pirimicarb, dimethoate and alpha-cypermethrin using the same procedure described above, except excised *T. aestivum* leaf sections were submerged in each chemical solution for 1 s each (not 10 s).

Statistical analysis

As incapacitated individuals invariably die and therefore do not contribute to the next generation, we pooled incapacitated individuals with dead individuals for data analysis across all bioassays. Aphid mortality in each bioassay Y was modelled as a binomial response variable $Y \sim \text{Binomial}(p, n)$, where p is the probability of mortality, and n is the number of aphids in each replicate. The probability of mortality was related to linear covariates with a logistic link function $\text{logit}(p) = a_i + b_i \ln(x)$, where x is chemical dose (mg a.i./L) and a_i and b_i are intercepts and slope coefficients respectively for each population i (Robertson *et al.* 2017; Venables & Ripley 2002). Variance was robustly estimated by allowing for dispersion (McCullagh & Nelder 1989).

Each chemical product was analysed separately, with differences in population effects (intercepts) assessed using an analysis of variance (ANOVA). They were then examined in pairwise comparisons using Tukey's Honest Significant Difference (HSD) method at the 0.05 significance level, which corrects for Type I errors when performing multiple hypotheses tests (Hsu 1996). For each bioassay, LC_{50} values (with 95% confidence intervals) and slope coefficient estimates of dose-response slopes were computed. To test differences in mortality responses between products, an additional model was fitted that included chemical product as a model factor, which was similarly contrasted using Tukey's HSD method.

All analyses were performed using the R language for statistical computing (R Core Team 2018).

Results

χ^2 tests indicated the models adequately fitted the data at the 0.05 significance level across all populations for each insecticide tested. Importantly, the mortality of the untreated control

aphids was very low across all bioassays; there was no mortality in 32 of the bioassays, 2% mortality in three bioassays and 4% mortality in one bioassay.

The dose-responses of *R. padi* were significantly different between the four chemical products ($\chi^2 = 74.7$, $df = 3$, $p < 0.001$). Imidacloprid and alpha-cypermethrin were the most toxic chemical products to *R. padi*, but were not significantly different to one another ($p = 0.97$), with average LC_{50} values ($\pm$ s.e.) across all nine populations of $8.72 (\pm 1.86)$ mg a.i./L and $9.94 (\pm 2.56)$ mg a.i./L, respectively. Dimethoate (34.42 ± 3.04 mg a.i./L) was significantly less toxic when compared with imidacloprid ($p < 0.001$) and alpha-cypermethrin ($p < 0.001$), while pirimicarb (103.99 ± 14.71 mg a.i./L) was significantly less toxic than imidacloprid ($p < 0.001$), alpha-cypermethrin ($p < 0.001$), and dimethoate ($p = 0.005$). When averaged across populations, the slopes for alpha-cypermethrin (0.94 ± 0.07) than slopes for imidacloprid (1.23 ± 0.09), dimethoate (1.52 ± 0.17) and pirimicarb (3.73 ± 1.19), however, only differences between alpha-cypermethrin and pirimicarb were statistically significant ($p = 0.019$).

Imidacloprid

There was no significant difference in the response of the nine populations to imidacloprid ($\chi^2 = 13.33$, $df = 8$, $p = 0.10$), indicating all populations have a similar sensitivity to this chemical product. Further, there were no differences when comparing LC_{50} values and 95% confidence intervals (Table 3). The estimated LC_{50} values for imidacloprid across the nine populations ranged from 3.77 to 21.81 mg a.i./L. Unique regression slopes for populations exposed to imidacloprid did not significantly improve the model ($\chi^2 = 2.87$, $df = 8$, $p = 0.94$).

For eight of the nine populations, the imidacloprid field rate (525 mg a.i./L) was adequate in suppressing 100% of aphid individuals. Five aphids (from a total of 44) from the ADE population were still alive at the field rate after 48 h exposure.

Pirimicarb

There was no significant difference in responses of the nine populations to pirimicarb ($\chi^2 = 1.17$, $df = 8$, $p = 0.99$), indicating all populations have a similar sensitivity to this chemical product. There were also no differences when comparing LC_{50} values and 95% confidence

intervals (Table 4). The estimated LC_{50} values across all populations ranged from 38.19 to 172.08 mg a.i./L.

Unique regression slopes for populations exposed to pirimicarb did not significantly improve the model ($\chi^2 = 1.69$, $df = 8$, $p = 0.99$). However, regression slope coefficients could not be estimated for two populations: ADE and BKV, due to the partial-separation of the response variable (e.g. 96% of individuals from ADE survived at 50 mg a.i./L whereas 0% survived at the subsequent dose of 500 mg a.i./L). This indicates that the biological response of *R. padi* to pirimicarb varies and the doses we tested were appropriate for some, but not all, populations. Regardless, the removal of ADE and BKV from the analysis did not affect the conclusion for the remaining seven populations (data not shown).

For all nine populations, the highest rate of pirimicarb (5000 mg a.i./L) was adequate in suppressing 100% of individuals after 48 h.

Dimethoate

There was no significant difference in the response of the nine populations to dimethoate ($\chi^2 = 2.01$, $df = 8$, $p = 0.98$) and there were also no differences when comparing LC_{50} values and 95% confidence intervals (Table 5). This indicates dimethoate sensitivity was similar across all populations. Similarly, unique regression slopes for each population exposed to dimethoate did not significantly improve the model ($\chi^2 = 2.15$, $df = 8$, $p = 0.97$).

For all nine populations, the dimethoate field rate (2000 mg a.i./L) was more than adequate in suppressing 100% of individuals after 48 h. LC_{50} values across all populations ranged from 14.68 to 48.20 mg a.i./L (Table 5).

Alpha-cypermethrin

Noticeable differences in dose-response curves were detected between populations exposed to alpha-cypermethrin, which were found to be statistically significant ($\chi^2 = 23.92$, $df = 8$, $p = 0.002$). These differences are further supported by non-overlapping 95% confidence intervals of LC_{50} values (Table 6). Aphids from KAT were significantly more sensitive to alpha-cypermethrin than aphids from TMW ($p = 0.006$), MNJ ($p = 0.03$) and PIC ($p = 0.02$). Unique regression slopes for populations exposed to alpha-cypermethrin did not significantly improve the model ($\chi^2 = 3.16$, $df = 8$, $p = 0.92$).

LC₅₀ values across all nine populations ranged from 0.6 to 24.57 mg a.i./L (Table 6). The field rate of alpha-cypermethrin (125 mg a.i./L) controlled 98%, 93% and 91% of aphids from ADE, MNJ and TMW, respectively. For the remaining six populations, the field rate was adequate in suppressing 100% of individuals after 48 h (Table 6).

Discussion

Our results provide toxicity data for nine populations of *R. padi* against four insecticides widely used for aphid control in cereal crops within Australia. It is worth noting that the mortality responses quantified in our study represent variation between individual aphids and populations taken at a point in time, while other sources of variation may exist, e.g. the same populations exposed to different seasonal conditions may behave differently (see Robertson *et al.* 2017). Monitoring insecticide susceptibility is an important part of pest management strategies, particularly for species with a history of insecticide resistance. Insecticide toxicity data not only provides an indication of the level of insecticide susceptibility in the field, but also allows researchers to accurately assess if population responses to insecticide products are increasing or decreasing through time (Roush & Miller 1986). Resistance to neonicotinoids, carbamates and organophosphates has been selected under laboratory conditions (Chen *et al.* 2007; Wang *et al.* 2018), demonstrating *R. padi* could evolve field resistance if selection pressures were sufficiently strong, which highlights the value of the toxicity data we have generated.

Recently, *R. Padi* sourced from populations across Australia were shown to have lower insecticide sensitivity compared with several other grain aphids (Clouston *et al.* 2016). Further, at two locations in Queensland and South Australia, some *R. padi* populations were observed to survive field rates of alpha-cypermethrin when tested using laboratory bioassays (Clouston *et al.* 2016). Our study provides a broader survey of insecticide responses among *R. padi* populations and identified differences in sensitivity levels to alpha-cypermethrin, but not to other insecticides. Interestingly, out of the four insecticides tested, the estimated slopes from the dose-response curves were lowest for alpha-cypermethrin. Typically, a lower slope value indicates a more heterogeneous response to the insecticide within a population (Chilcuit & Tabashnik 1995), and thus a greater likelihood that the population will evolve more quickly to insecticide selection pressure.

Clouston *et al.* (2016) estimated LC₅₀ values for alpha-cypermethrin in the range of 15.36 – 72.10 mg a.i./L for *R. padi*, with an average LC₅₀ of 33.83 (± 8.14) mg a.i./L across six populations. This is considerably higher than the average LC₅₀ value calculated from the nine populations we tested of 9.94 (± 2.56) mg a.i./L, which followed the same bioassay methodology and tested the same formulated chemical product. At a finer level, two populations from our study (one each from WA and Vic.) were collected in close proximity (<20km) to populations from the Clouston *et al.* (2016) study, allowing a provisional comparison of how sensitivity levels may have shifted. In the four years since the Clouston *et al.* (2016) study, the LC₅₀ value for alpha-cypermethrin has dropped from 24.88 to 9.90 mg a.i./L at the two Vic. populations (Pimpinio and Natimuk) and from 30.84 to 0.60 mg a.i./L for the two WA populations (Katanning and Woodanilling). These shifts in sensitivity through time may be a consequence of changes in insecticide selection pressure. This could be linked to a reduced risk of early-season BYDV infections (vectored by *R. padi*), which is known to be influenced by changing rainfall patterns (Thackray *et al.* 2009). Like many other countries, Australia has also seen a considerable shift in the adoption of neonicotinoid seed treatments, which has reduced the reliance on foliar applications of synthetic pyrethroids and other broad-spectrum insecticides for early season aphid/virus control (Bass *et al.* 2015; Attwood & Paisley 2019; Umina *et al.* 2019).

In our study, the several *R. padi* populations were not fully controlled by alpha-cypermethrin applied at the recommended field rate of 125 mg a.i./L. However, insecticide rates applied in laboratory bioassays do not necessarily indicate that these populations are resistant under field conditions; laboratory conditions provide a standardised environment while field populations are typically under more stress, which can impact both survival and chemical responses (Cahill *et al.* 1996; Iwasaki *et al.* 2018; Heye *et al.* 2019). Furthermore, exposure to insecticides when applied in the field will be considerably different to what aphids experience when tested through laboratory bioassays. For example, aphid mortality was assessed 48 h after chemical exposure in our experiments. However, products like alpha-cypermethrin can persist in the environment for many weeks and thus the exposure period for aphids is likely to be substantially longer in the field. For the remaining insecticides, our findings are similar to those reported by Clouston *et al.* (2016). We found little variation in *R. Padi* sensitivity to dimethoate, pirimicarb and imidacloprid among populations, and the overall LC₅₀ values are comparable between studies. The imidacloprid results, in particular,

were very similar between those we found (8.72 ± 1.86 mg a.i./L) and those estimated by Clouston *et al.* (2016) (9.56 ± 1.73 mg a.i./L).

There was considerable variation in responses of *R. padi* to alpha-cypermethrin, with reduced sensitivity to this chemical product identified in three populations. Aphids from MNJ, PIC and TMW had LC₅₀ values 26-, 27- and 41-fold (respectively) greater than aphids from the most sensitive population, KAT. Despite this, it appears there has been an overall increase in sensitivity to alpha-cypermethrin across Australia, which may reflect changes in insecticide use. For the other products, we found a relatively consistent level of insecticide toxicity across *R. padi* field populations. The toxicity data we have generated are now available for comparison to those obtained in any future monitoring activities. Importantly, they have been established before field resistance has evolved, allowing for insecticide resistance management strategies to be developed and implemented before resistance frequencies become too high (see Roush 1998; Sudo *et al.* 2018). Field surveillance can now be prioritised to focus on alpha-cypermethrin and other pyrethroid chemicals, and in regions of Australia where toxicity responses in aphid populations are approaching levels that may impact the efficacy of chemical applications in the field.

Acknowledgements

The authors thank Siobhan de Little, Annie Johnson, Lisa Kirkland and Kathy Overton for technical assistance. Thanks to the Grains Research and Development Corporation for financial support, and those growers, advisors and researchers who aided in aphid collections.

References

- APVMA. 2020. Australian Pesticides and Veterinary Medicines Authority Public Chemical Registration Information System Search. Available from: <https://portal.apvma.gov.au/pubcris> [Accessed 28 January 2020].
- Araya IE& FosterJE. 1987. Control of *Rhopalosiphumpadi* (Homoptera: Aphididae) in selected wheat and oat cultivars with seed systemic insecticides in the greenhouse. *Journal of Economic Entomology* **80**, 1272–1277.
- Atwood D & Paisley-Jones C. 2017. US EPA - Pesticides industry sales and usage 2008-2012. United States Environmental Protection Agency, Washington, USA.
- Bailey PY. 2007. Pests of field crops and pastures: identification and control. CSIRO Publishing, Melbourne, Australia.
- Bass C, Denholm I, Williamson MS, Nauen R. 2015. The global status of insect resistance to neonicotinoid insecticides. *Pesticide Biochemistry and Physiology* **121**, 78–87.
- Blackman RL&Eastop VF. 2000. Aphids on the world's crops. John Wiley & Sons Ltd. West Sussex, UK.
- Cahill M, Denholm I, Ross G, Gorman K, Johnston D. 1996. Relationship between bioassay data and the simulated field performance of insecticides against susceptible and resistant adult *Bemisiatabaci* (Homoptera: Aleyrodidae). *Bulletin of Entomological Research* **86**, 109–116.
- Chen M, Han Z, Qiao X, Qu M. 2007. Mutations in acetylcholinesterase genes of *Rhopalosiphumpadi* resistant to organophosphate and carbamate insecticides. *Genome* **50**, 172–179.
- Cheng HZ, Zheng SY, Guo YY, Zhao SZ, Wang P. 2004. The resistance of wheat aphids to several insecticides. *Henan Agricultural Science* **6**, 50–53.
- Chilcuit CF & Tabashnik BE. 1995. Evolution of pesticide resistance and slope of the concentration-mortality line: are they related? *Journal of Economic Entomology* **88**, 11–20.
- Clouston A, Edwards O, Umina P. 2016. An insecticide baseline study of Australian broadacre aphids. *Crop Pasture Science* **67**, 236–244.
- de BarroPJ&Wallwork H. 1992. The role of annual grasses in the phenology of *Rhopalosiphumpadi* in the low rainfall belt of South Australia. *Annals of Applied Biology*, **121** 455–467.

389

390 de Barro PJ, Maelzer DA, Wallwork H. 1992. The role of refuge areas in the phenology
 391 of *Rhopalosiphum padi* in low rainfall cropping areas of South Australia. *Annals of Applied*
 392 *Biology*, **121** 521-535.

393 de Little SC & Umina P. 2017. Susceptibility of Australian *Myzus persicae* (Hemiptera:
 394 Aphididae) to three recently registered insecticides: spirotetramat, cyantraniliprole, and
 395 sulfoxaflor. *Journal of Economic Entomology* **110**, 1764-1769.

396 Delmotte F, Sabater-Muñoz B, Prunier-Leterme N, Latorre A, Sunnucks P, Rispe C, Simon J-
 397 C. 2003. Phylogenetic evidence for hybrid origins of asexual lineages in an aphid species.
 398 *Evolution* **57**, 1291–1303.

399 Edwards OR, Franzmann B, Thackray D, Micic S. 2008. Insecticide resistance and
 400 implications for future aphid management in Australian grains and pastures: a review.
 401 *Australian Journal of Experimental Agriculture* **48**, 1523–1530.

402 Figueroa CC, Fuentes-Contreras E, Molina-Montenegro MA, Ramírez CC. 2018.
 403 Biological and genetic features of introduced aphid populations in agroecosystems,
 404 *Current Opinion in Insect Science* **26**, 63-68.

405 GRDC. 2010. Cereal aphids factsheet. Grains Research and Development Corporation.
 406 Available from: www.grdc.com.au/GRDC-FS-CerealAphids [Accessed 28 January 2020].

407 Halkett P, Kindlmann M, Plantegenest M, Sunnucks P, Simon JC. 2006. Temporal
 408 differentiation and spatial coexistence of sexual and facultative asexual lineages of an aphid
 409 species at mating sites. *Journal of Evolutionary Biology* **19**, 809-815.

410 Heye K, Graumnitz S, Rybicki Met al. 2019. Laboratory-to-field extrapolation: increase in
 411 carbamazepine toxicity in a higher tier, multiple-stress experiment. *Ecotoxicology and*
 412 *Environmental Safety* DOI: 10.1016/j.ecoenv.2019.109481.

413 Hsu JC. 1996. Multiple comparisons. Chapman & Hall, London, UK.

414 Hughes RD, Casmir M, O'Loughlin GT, Martyn EJ. 1964. A survey of aphids flying over
 415 eastern Australia in 1961. *Australian Journal of Zoology* **12**, 174-200.

416 IRAC. 2016. Insecticide Resistance Action Committee susceptibility test method no. 019.
 417 Available from: [https://irac-online.org/content/uploads/Method_019-](https://irac-online.org/content/uploads/Method_019-v3.4_15Dec16_aphid.pdf)
 418 [v3.4_15Dec16_aphid.pdf](https://irac-online.org/content/uploads/Method_019-v3.4_15Dec16_aphid.pdf) [Accessed 15 April 2018].

419 Iwasaki Y, Schmidt TS, Clements WH. 2018. Quantifying differences in response of aquatic
 420 insects to trace metal exposure in field studies and short-term stream mesocosm experiments.
 421 *Environmental Science and Technology* **52**, 4378–4384.

422 Kirkland LS, Pirtle EI, Umina PA. 2018. Responses of the Russian wheat aphid
 423 (*Diuraphis noxia*) and bird cherry oat aphid (*Rhopalosiphum padi*) to insecticide seed
 424 treatments in wheat. *Crop and Pasture Science* **69**, 966–973.

425 Lucio-Zavaleta E, Smith DM, Gray SM. 2001. Variation in transmission efficiency among
 426 Barley yellow dwarf virus-RMV isolates and clones of the normally inefficient aphid vector
 427 *Rhopalosiphum padi*. *Phytopathology* **91**, 792–796.

428 McCullagh P & Nelder JA. 1989. Generalized linear models. Chapman and Hall, London, UK.

429 Macfadyen S & Kriticos DJ. 2012. Modelling the geographical range of a species with variable
 430 life-history. *PLoS One*, 7(7):e40313.

431 McDonald G, Umina P, Lye J, Maino J, Perry K, Baker G. 2019. Insecticide resistance in the
 432 southern region: current status, future risk and best management practices. ISBN: 978-1-
 433 921779-84-8. Grains Research and Development Corporation, Kingston, Australia.

434 Mota-Sanchez D & Wise JC. 2020. The Arthropod Pesticide Resistance Database. Michigan
 435 State University. <http://www.pesticideresistance.org> [Accessed 12 Jan 2020].

436 R Core Team. 2018. R: A language and environment for statistical computing. R Foundation
 437 for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>.

438 Robertson JL, Jones MM, Olguin E & Alberts B. 2017. Bioassays with arthropods, 3rd
 439 Edition, CRC Press, Boca Raton, USA.

440 Roush RT. 1998. Two-toxin strategies for management of insecticidal transgenic crops: can
 441 pyramiding succeed where pesticide mixtures have not? *Philosophical Transactions of the*
 442 *Royal Society B: Biological Sciences* **353**, 1777–1786.

443 Roush RT & Miller GL. 1986. Considerations for design of insecticide resistance monitoring
 444 programs. *Journal of Economic Entomology* **79**, 293–298.

445 Schliephake E, Habekuss A, Scholz M, Ordon F. 2013. Barley yellow dwarf virus
 446 transmission and feeding behaviour of *Rhopalosiphum padi* *Hordeum bulbosum* clones.
 447 *Entomologia Experimentalis et Applicata* **146**, 347–356.

- 448 Simon J-C, Martinez-Torres D, Latorre A, Moya A, Hebert PDN. 1996. Molecular
449 characterization of cyclic and obligate parthenogens in the aphid *Rhopalosiphumpadi* (L.).
450 *Proceedings of the Royal Society of London Series B***263**, 481–486.
- 451 Sudo M, Takahashi D, Andow DA *et al.*2018. Optimal management strategy of insecticide
452 resistance under various insect life histories: heterogeneous timing of selection and interpatch
453 dispersal. *Evolutionary Applications***11**,271-283.
- 454 Thackray DJ, Diggle AJ, Jones RAC. 2009. BYDV predictor: a simulation model to predict
455 aphid arrival, epidemics of *Barley yellow dwarf virus* and yield losses in wheat crops in a
456 Mediterranean type environment. *Plant Pathology***58**, 186–202.
- 457 Umina PA, Edwards O, Carson P, Van Rooyen A, Anderson A. 2014. High levels of
458 resistance to carbamate and pyrethroid chemicals widespread in Australian *Myzus persicae*
459 (Hemiptera: Aphididae) populations. *Journal of Economic Entomology***107**, 1626–1638.
- 460 Umina PA, McDonald G, Maino J, Edwards O, Hoffmann AA. 2019. Escalating insecticide
461 resistance in Australian grain pests: contributing factors, industry trends and management
462 opportunities. *Pest Management Science***75**,1494–1506.
- 463 Valenzuela I, Ridland PM, Weeks AR, Hoffmann AA. 2010. Patterns of genetic variation and
464 host adaptation in an invasive population of *Rhopalosiphumpadi* (Hemiptera:
465 Aphididae).*Annals of the Entomological Society of America***103**, 886–897.
- 466 Valenzuela I & Hoffmann AA. 2015. Effects of aphid feeding and associated virus injury on
467 grain crops in Australia. *Austral Entomology***54**, 292–305.
- 468 Venables WN & Ripley BD. 2002. Modern applied statistics with S. Springer, New York.
469 USA.
- 470 Wang K, Peng X, Zuo Y, Li Y, Chen M. 2016. Molecular cloning, expression pattern and
471 polymorphisms of NADPH-Cytochrome P450 reductase in the bird cherry-oat aphid
472 *Rhopalosiphumpadi* (L.). *PLoS ONE***11**, e0154633.
- 473 Wang K, Zhang M, Huang Y, Yang Z, Su S, Chen M. 2018. Characterisation of imidacloprid
474 resistance in the bird cherry-oat aphid, *Rhopalosiphumpadi*, a serious pest on wheat crops:
475 *Pest Management Science***74**, 1457–1465.
- 476 Xiao D, Yang T, Desneux N, Han P, Gao X. 2015. Assessment of sublethal and
477 transgenerational effects of pirimicarb on the wheat aphids *Rhopalosiphumpadi* and
478 *Sitobionavenae*. *PLoS ONE* **10**, e0128936.

- 479 Yang Q, Umina PA, Rasic Get *al.* 2020. Origin of resistance to pyrethroids in the redlegged
480 earth mite (*Halotydeus destructor*) in Australia: repeated local evolution and migration. *Pest*
481 *Management Science***76**, 509–519.
- 482 Yazdani M, Baker G, DeGraaf Het *al.* 2018. First detection of Russian wheat aphid
483 *Diuraphis noxia* Kurdjumov (Hemiptera: Aphididae) in Australia: a major threat to cereal
484 production. *Austral Entomology* **57**, 410–417.
- 485 Zhang M, Qiao X, Li Y, Fang B, Zuo Y, Chen M. 2016. Cloning of eight *Rhopalosiphum*
486 (Hemiptera: Aphididae) nAChR subunit genes and mutation detection of the $\beta 1$ subunit in
487 field samples from China. *Pesticide Biochemistry and Physiology***132**, 89–95.
- 488 Zuo Y, Wang K, Zhang M, Peng X, Piñero JC, Chen M. 2016. Regional susceptibilities of
489 *Rhopalosiphum* (Hemiptera: Aphididae) to ten insecticides. *Florida Entomologist***99**,
490 269–275.

Table 1. Sampling information of Australian *R.padi* populations used in study.

Location	State	Code	Date collected	Host plant	Latitude	Longitude
Tamworth	NSW	TMW	January 2017	<i>Triticumaestivum</i>	-31.18	150.91
Picola	Vic	PIC	March 2017	<i>Triticumaestivum</i>	-35.99	145.09
Dookie	Vic	DOK	June 2017	<i>Triticumaestivum</i>	-36.39	145.69
Pimpinio	Vic	PIM	June 2017	<i>Hordeum vulgare</i>	-36.62	142.03
Back Valley	SA	BKV	September 2017	Unknown	-35.52	138.51
Adelaide	SA	ADE	December 2017	Grass weeds	-34.97	138.63
Manjimup	WA	MNJ	August 2017	<i>Avenasativa</i>	-34.31	116.17
Capel	WA	CAP	August 2017	Grass weeds	-33.58	115.66
Katanning	WA	KAT	August 2017	<i>Avenasativa</i>	-33.70	117.57

Table 2. Insecticides tested in this study against *R.padi*.

Trade Name	Active ingredient	Chemical group	Formulation	Supplier
Nuprid [®] 200SC insecticide	imidacloprid	Neonicotinoid	200 g/L SC	NuFarm Australia
Pirimor [®] WGaphicide	pirimicarb	Carbamate	500 g/L WG	Syngenta Australia
Dimethoate 400SC [®]	dimethoate	Organophosphate	400 g/L SC	NuFarm Australia
Astound [®] Duo insecticide	alpha- cypermethrin	Synthetic pyrethroid	100 g/L EC	NuFarm Australia

SC = suspension concentrate, EC = emulsifiable concentrate, WG = water-dispersible granule.

Table 3. Toxicity data for *R. padi* after exposure to imidacloprid for 48 h.

Population	No. aphids tested	Regression slope coefficient ($\pm$ SE)	LC ₅₀ (mg a.i./L)	Lower-upper 95% CIs (mg a.i./L)
TMW ^a	259	1.39 (0.65)	3.77	1.22 – 11.65
PIC ^a	263	0.78 (0.29)	5.57	1.21 – 25.68
DOK ^a	258	1.31 (0.58)	8.06	2.46 – 26.37
PIM ^a	266	1.15 (0.47)	6.81	1.97 – 23.44
BKV ^a	266	1.54 (0.75)	7.40	2.65 – 20.68
ADE ^a	255	0.86 (0.34)	21.81	5.00 – 95.16
MNJ ^a	254	1.35 (0.63)	4.20	1.29 – 13.66
CAP ^a	259	1.56 (0.70)	12.83	4.19 – 39.31
KAT ^a	257	1.16 (0.49)	8.02	2.29 – 28.08

Different letters indicate significant differences between populations ($p < 0.05$).

Table 4. Toxicity data for *R.padi* after exposure to pirimicarb for 48 h.

Population	No. aphids tested	Regression slope coefficient ($\pm$ SE)	LC ₅₀ (mg a.i./L)	Lower-upper 95% CIs (mg a.i./L)
TMW ^a	266	1.42 (0.83)	134.50	27.78 – 651.26
PIC ^a	265	1.43 (0.87)	98.67	20.35 – 478.42
DOK ^a	266	2.55 (1.97)	113.01	25.33 – 504.14
PIM ^a	269	1.69 (1.03)	139.52	30.89 – 630.17
BKV ^a	261	9.66 (7702)	51.93	NC
ADE ^a	270	10.32 (4909)	67.59	NC
MNJ ^a	258	1.99 (1.80)	38.19	10.73 – 135.82
CAP ^a	266	1.93 (1.21)	172.08	39.10 – 757.16
KAT ^a	263	2.65 (2.02)	120.47	25.82 – 562.02

Different letters indicate significant differences between populations ($p < 0.05$). NC indicates 95% confidence intervals could not be computed.

Table 5. Toxicity data for *R.padi* after exposure to dimethoate for 48 h.

Population	No. aphids tested	Regression slope coefficient ($\pm$ SE)	LC ₅₀ (mg a.i./L)	Lower-upper 95% CIs (mg a.i./L)
TMW ^a	270	1.82 (1.52)	33.59	7.15 – 157.70
PIC ^a	272	1.26 (0.84)	40.36	6.00 – 271.12
DOK ^a	278	2.37 (2.17)	39.53	8.36 – 186.90
PIM ^a	268	1.54 (1.08)	48.20	8.31 – 279.45
BKV ^a	262	1.42 (1.05)	31.65	5.34 – 187.39
ADE ^a	261	2.09 (2.10)	30.77	6.86 – 137.93
MNJ ^a	258	1.36 (0.97)	36.49	5.96 – 223.29
CAP ^a	277	1.11 (0.70)	34.50	4.86 – 244.57
KAT ^a	266	0.73 (0.43)	14.68	1.24 – 173.70

Different letters indicate significant differences between populations ($p < 0.05$).

Table 6. Toxicity data for *R.padi* after exposure to alpha-cypermethrin for 48 h.

Population ¹	No. aphids tested	Regression slope coefficient ($\pm$ SE)	LC ₅₀ (mg a.i./L)	Lower-upper 95% CIs (mg a.i./L)	Mortality at registered rate ²	RF ³
TMW ^a	260	1.16 (0.50)	24.57	7.51 – 80.47	91%	41.0
PIC ^a	285	1.31 (0.54)	16.22	5.74 – 45.80	100%	27.0
DOK ^{ab}	273	1.06 (0.38)	3.84	1.16 – 12.68	100%	-
PIM ^{ab}	253	0.64 (0.24)	9.90	1.88 – 52.01	100%	-
BKV ^{ab}	257	0.92 (0.33)	2.34	0.63 – 8.61	100%	-
ADE ^{ab}	279	0.66 (0.23)	7.34	1.51 – 35.64	98%	-
MNJ ^a	265	0.95 (0.37)	15.34	4.08 – 57.68	93%	25.6
CAP ^{ab}	261	0.98 (0.37)	9.31	2.67 – 32.39	100%	-
KAT ^b	247	0.85 (0.31)	0.60	0.14 – 2.43	100%	-

¹Different letters indicate significant differences between the populations ($p < 0.05$).

²Registered field rate equates to 125 mg a.i./L alpha-cypermethrin.

³Resistance Factors (RF) were calculated by dividing the population LC_{50} value by 0.60 (the lowest LC_{50} value estimated from KAT).

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Umina, PA;Reidy-Crofts, J;Babineau, M;Maino, JL;Edwards, OR

Title:

Susceptibility of the bird cherry-oat aphid, *Rhopalosiphum padi* (Hemiptera: Aphididae), to four insecticides

Date:

2020-08-05

Citation:

Umina, P. A., Reidy-Crofts, J., Babineau, M., Maino, J. L. & Edwards, O. R. (2020). Susceptibility of the bird cherry-oat aphid, *Rhopalosiphum padi* (Hemiptera: Aphididae), to four insecticides. *AUSTRAL ENTOMOLOGY*, 59 (4), pp.838-844. <https://doi.org/10.1111/aen.12490>.

Persistent Link:

<http://hdl.handle.net/11343/276109>