DR. FATHIMA FERIAL ISMAIL (Orcid ID : 0000-0002-9515-7996)

PROF. RODNEY SINCLAIR (Orcid ID : 0000-0001-6751-1428)

Article type : Correspondence - online only

TITLE: Reversibility of alopecia caused by chronic cutaneous lupus erythematosus

AUTHORS:

Fathima Ferial Ismail, MBBS, BMedSci (corresponding author)

Sinclair Dermatology, Victoria, Australia

2 Wellington Parade, East Melbourne, Victoria, Australia

Email: f.ferial.ismail@gmail.com

Telephone: +61431345524

Nekma Meah, MBChB, MRCP (UK)

Sinclair Dermatology, Victoria, Australia

Rodney Sinclair, MBBS, MD, FACD

Sinclair Dermatology, Victoria, Australia

University of Melbourne, Victoria, Australia

Epworth Healthcare, Victoria Australia

Key words: cicatricial alopecia, chronic cutaneous lupus erythematosus, discoid lupus erythematosus, lupus profundus

Figures: 1

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/IJD.15148](#)

This article is protected by copyright. All rights reserved

Conflicts of interest: RS has served as a consultant or paid speaker for, or participated in clinical trials sponsored by, Leo Pharma, Amgen, Novartis, Merck & Co., Celgene, Coherus Biosciences, Janssen, Regeneron, MedImmune, GlaxoSmithKline, Cutanea, Samson Clinical, Boehringer Ingelheim, Pfizer, MSD, Oncobiologics, Roche, Eli Lilly and Bayer.

Dear Editor,

Chronic cutaneous lupus erythematosus (CCLE) is a common cause of cicatricial alopecia (CA) which represents almost 2% of presentations to specialist hair clinics¹. CA from CCLE is 7 times more common in women. CA is generally permanent and irreversible², although hair regrowth has been noted in some patients with CA secondary to CCLE³⁻⁵. We present 3 cases of CCLE where prompt treatment led to significant hair regrowth.

A 29-year-old female presented with a 2-month history of multiple circular patches of CA. She had a past history of Raynaud's syndrome. Dermoscopy revealed loss of follicular ostia, telangiectasia and significant post-inflammatory hyperpigmentation consistent with a cicatricial alopecia. Histology showed a CA with loss of terminal follicles, mid-dermal fibrosis, perifollicular, perivascular and interface lymphocytic infiltrate, necrotic keratinocytes, pigment incontinence and epidermal atrophy. Indirect immunofluorescence was positive for IgM in the basement membrane. Anti-nuclear antibody (ANA) and anti-double-stranded-DNA (dsDNA) were negative, however anti-PM/SCL was positive. The clinical and histological findings supported a diagnosis of discoid lupus erythematosus (DLE). Treatment was started with prednisolone 20mg daily and minoxidil 0.45mg daily. Hair regrowth was apparent at 2 weeks. Hydroxychloroquine 200mg twice daily was introduced and the patches of cicatricial alopecia were injected with triamcinolone acetonide 5mg/ml. On review at 6 weeks, there was marked hair regrowth in all patches (Fig. 1).

A 29-year-old male with no significant past medical history presented with a 2-month history of a solitary patch of CA with central atrophy and hyperpigmentation. Histology revealed a CA with prominent perifollicular lymphocytic inflammation and increased stromal mucin. There were twenty-four vellus follicles, but no terminal follicles were present. A weakly positive ANA (1/80) was present, with negative anti-dsDNA and extractable nuclear antigens (ENA). Clinical and histological features were consistent with DLE. He was treated with intralesional triamcinolone acetonide 5 mg/ml. Review at 4 weeks confirmed marked hair regrowth and reduced erythema. Hydroxychloroquine 200mg daily was commenced and the lesion was re-injected with triamcinolone 5mg/ml. Five months later he had near-complete remission (Fig. 1).

A 38-year-old female presented with a 3-year history of a solitary, 10 x 10cm, slowly enlarging, patch of CA. Past medical history included iron deficiency anaemia. Dermoscopy revealed areas of loss of follicular ostia and yellow dots. Histology showed a CA with associated basal vacuolar changes in the epidermis, thickening in the basement membrane, a marked increase in dermal mucin, a deep and superficial perivascular chronic inflammatory infiltrate with interface changes. Indirect immunofluorescence was non-specific with perivascular staining for fibrinogen, C3 and C1q. ANA and anti-dsDNA were negative, and anti-PM/SCL was low positive. Clinical and histological features were consistent with lupus profundus. She was treated with intralesional triamcinolone acetonide 5mg/ml. At week 6 weeks, hair regrowth was evident and the size of the alopecic patch had reduced to 7 x 6cm (Fig. 1).

CA associated with CCLE is generally considered irreversible. This case series highlights the potential reversibility of CA with prompt treatment.

FIGURE LEGEND

Figure 1

(a) Case 1 baseline photograph; (b) Case 1 after 6 weeks of treatment; (c) Case 2 baseline photograph; (d) Case 2 after 5 months of treatment; (e) Case 3 baseline photograph; (f) Case 3 after 6 weeks of treatment

REFERENCES

1. Vañó-Galván S, Saceda-Corralo D, Blume-Peytavi U, et al. Frequency of the Types of Alopecia at Twenty-Two Specialist Hair Clinics: A Multicenter Study. *Skin Appendage Disorders*. 2019;5:309-15.
2. Garza-Mayers AC, McClurkin M, Smith GP. Review of treatment for discoid lupus erythematosus. *Dermatol Ther*. 2016 Jul;29(4):274-83.
3. Milam EC, Ramachandran S, Franks AG Jr. Treatment of Scarring Alopecia in Discoid Variant of Chronic Cutaneous Lupus Erythematosus With Tacrolimus Lotion, 0.3. *JAMA Dermatol*. 2015 Oct;151(10):1113-6.
4. Hamilton T, Otberg N, Wu WY, et al. Successful hair re-growth with multimodal treatment of early cicatricial alopecia in discoid lupus erythematosus. *Acta Derm Venerol*. 2009;89(4):417-8.
5. Igari S, Yamamoto T. Dramatic effect of hydroxychloroquine on lupus alopecia. *J Dermatol*. 2018 Feb;45(2):194-7.

ijd_15148_f1.jpg

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Ismail, FF;Meah, N;Sinclair, R

Title:

Reversibility of alopecia caused by chronic cutaneous lupus erythematosus

Date:

2020-08-17

Citation:

Ismail, F. F., Meah, N. & Sinclair, R. (2020). Reversibility of alopecia caused by chronic cutaneous lupus erythematosus. INTERNATIONAL JOURNAL OF DERMATOLOGY, 59 (12), pp.E437-E438. <https://doi.org/10.1111/ijd.15148>.

Persistent Link:

<http://hdl.handle.net/11343/276156>