Bose Sucharita (Orcid ID: 0000-0003-4617-0577)
Sowdhamini Ramanathan (Orcid ID: 0000-0002-6642-2367)

Quaternary variations in the structural assembly of *N*-acetylglucosamine-6-phosphate deacetylase from *Pasteurella multocida*

Running Title: Quaternary structure of PmNagA

Lavanyaa Manjunath^{1, a,b}, David Coombes^c, James Davies^c, Mugdha Dhurandhar^d, Vikas R Tiwari^d, Renwick C.J. Dobson^{c,e}, R Sowdhamini^d, Ramaswamy S^{a,f} and Sucharita Bose^a

^a Institute for Stem Cell Science and Regenerative Medicine, NCBS, GKV Campus, Bellary Main Road, Bangalore, Karnataka, 560065, India.

^b Manpal Academy of Higher Education, Tiger Circle, Manpal, Karnataka, India.

^c Biomolecular Interaction Centre and School of Biological Sciences, University of Canterbury, PO Box4800, Christchurch 8140, New Zealand

^d National Centre for Biological Sciences, GKV Campus, Bellary Road, Bangalore, Karnataka, 560065, India.

^e Bio21 Molecular Science and Biotechnology Institute, Department of Biochemistry and Molecular Biology, University of Melbourne, Parkville, Victoria 3010, Australia

^f Department of Biological Sciences and Department of Biomedical Engineering, Purdue University, West Lafayette IN. 47907. USA

Correspondence email: sucharitab@instem.res.in

Acknowledgements

The research work was supported by DBT-Indo Swedish Grant (BT/IN/SWEDEN/06/SR/2017-18), DBT-B-life grant (BT/PR5081/INF/156/2012) NCBS X-ray facility grant (BT/PR12422/MED/31/287/214). We acknowledge the NCBS X-ray facility and thank Dr. Vinod Nayak, the X-ray facility manager for help in freezing crystals. We thank Dr. KR Vinothkumar for preparing the cryo-EM samples and data collection. We thank Dr. Kanaga Vijayan for his illustration work in Figure 1B. We thank Dr. Sathya Srinivasachari for the PmNagB experiments. We thank ESRF Access Program of RCB (supported by the Grant No. BT/INF/22/SP22660/2017 of the Department of Biotechnology, Ministry of Science and Technology. RCJD acknowledges the following for funding support, in part: 1) the New Zealand Royal Society Marsden Fund (contract UOC1506); 2) a Ministry of Business,

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/pro.25998

¹ present address: Department of Biology, University of York, Wentworth Way, Heslington, York, YO10 5DD, United Kingdom

Innovation and Employment Smart Ideas grant (contract UOCX1706); and 3) the Biomolecular Interactions Centre (University of Canterbury).

Abstract: *N*-acetylglucosamine 6-phosphate deacetylase (NagA) catalyzes the conversion of *N*-acetylglucosamine-6-phosphate to glucosamine-6-phosphate in amino sugar catabolism. This conversion is an essential step in the catabolism of sialic acid in several pathogenic bacteria, including *Pasteurella multocida*, and thus NagA is identified as a potential drug target. Here, we report the unique structural features of NagA from *P. multocida* (PmNagA) resolved to 1.95 Å. PmNagA displays an altered quaternary architecture with unique interface interactions compared to its close homolog, the *Escherichia coli* NagA (EcNagA). We confirmed that the altered quaternary structure is not a crystallographic artefact using single particle electron cryo-microscopy. Analysis of the determined crystal structure reveals a set of hot-spot residues involved in novel interactions at the dimer-dimer interface. PmNagA binds to one Zn²⁺ ion in the active site and demonstrates kinetic parameters comparable to other bacterial homologs. Kinetic studies reveal that at high substrate concentrations (~10-fold the K_M), the tetrameric PmNagA displays hysteresis similar to its distant neighbor, the dimeric *Staphylococcus aureus* NagA (SaNagA). Our findings provide key information on structural and functional properties of NagA in *P. multocida* that could be utilized to design novel antibacterials.

Keywords: Quaternary variations, *N*-acetylglucosamine-6-phosphate deacetylase, metal binding, amino sugar metabolism, sialic acid catabolism

1. Introduction

Pathogenic and commensal bacteria utilize host-derived nutrients as a source of energy and metabolic precursors to survive in the host environment. Several bacterial pathogens, including *Pasteurella multocida*, *Clostridium perfringens*, *Escherichia coli*, *Haemophilus influenzae*, and *Bacteroides fragilis* catabolize sialic acids, a structurally diverse group of nine-carbon keto sugars¹⁻³. Bacterial sialic acid catabolism begins with the import⁴⁻⁶ and conversion of sialic acid by the nanAKE enzymes into *N*-acetylglucosamine-6-phosphate (GlcNAc-6P)⁷⁻¹², a substrate for *N*-acetylglucosamine-6-phosphate deacetylase (NagA)^{13,14}. NagA belongs to the metal-dependent amidohydrolase family of proteins of the urease superfamily^{15,16} and deacetylates *N*-acetylglucosamine-6-phosphate (GlcNAc-6P) to form glucosamine-6-phosphate (GlcN-6P) and acetate¹⁷ (Figure 1A). The product, GlcN-6P is a branch metabolite and is either channeled into glycolysis upon conversion to fructose-6-phosphate by glucosamine-6-phosphate deaminase (NagB)¹⁸ or into peptidoglycan and lipopolysaccharide synthesis¹⁹ upon conversion into UDP-GlcNAc by phosphoglucosamine mutase (GlmM) and *N*-acetylglucosamine-1-phosphate-uridyl transferase (GlmU)²⁰. Thus, NagA, the enzyme that catalyzes the conversion of GlcNAc-6P to GlcN-6P occupies a central position at the crossroads of several biochemical pathways in bacteria (Figure 1B).

The structure of NagA has been studied from various bacteria including *E. coli* (EcNagA)²¹, *Bacillus subtilis* (BsNagA)³, *Thermotoga maritima* (TmNagA), *Vibrio cholerae* (VcNagA) and *Mycobacterium smegmatis* (MsNagA)²². All NagA homologs display the same structural framework of a distorted TIM barrel with (α/β)₈ topology and a small β -barrel domain. However, they differ in quaternary structure and the number of metal centres in the active site. NagA enzymes bind to various divalent metal ions including Fe²⁺ and Zn²⁺, and can either have a binuclear (BsNagA and MsNagA)

^{3,22} or a mononuclear metal centre (EcNagA, VcNagA and TmNagA). The minimal functional unit of NagA, as seen in BsNagA, TmNagA and MsNagA, is a dimer with the active site located at the dimer interface ^{3,22,23}. EcNagA ²¹ and VcNagA display a compact tetrameric structure assembled as dimer of dimers with D2 (222) symmetry.

NagA has been demonstrated to be critical for bacterial survival. Gene deletion studies of *nagA* in *Gluconacetobacter xylinus* shows that Δ nagA mutants are unable to survive in media containing *N*-acetylglucosamine as the sole carbon source ¹⁴. In *Listeria monocytogenes*, *nagA* mutants showed cell division defects with septal malformation, thinner cell-wall with sensitivity to antibiotics and reduced peptidoglycan and teichoic acid content ²⁴. The growth defects and the changes in cell morphology could not be rescued upon supplementing with GlcNAc. Similarly, *Staphylococcus aureus nagA* mutants showed reduced growth rates and increased sensitivity to cell wall inhibitors ¹⁹. These data suggest that targeting enzymes responsible for maintaining the intracellular pool of GlcN-6P may prove beneficial for designing novel antibiotics. Thus, NagA enzyme has gained attention as a potential antibacterial target ¹⁴.

Pasteurella multocida, a Gram-negative coccobacillus is a commensal and opportunistic pathogen present in populations of livestock, poultry and domestic animals like cats and dogs ². *P. multocida* causes chronic infections in both animals and humans resulting in systemic pasteurellosis, pneumonia, rhinitis, and septicemia. *P. multocida* can form biofilms ²⁵ and infections in humans are caused mostly due to bites and scratches by animals, making it a considerable threat to human and animal health. *In silico* analyses show that *P. multocida* has the sialic acid catabolism genes including the *nagAB* operon ^{2,26}, and is highly similar in sialobiology to *H. influenzae* ²⁷. Inhibition of sialic acid uptake in *H. influenzae* have adverse effects on the survival in biofilms and in serum ²⁷. Since *P. multocida* follows same sialobiology as *H. influenzae* ²⁷, inhibiting enzymes involved in sialic acid catabolism may also negatively affect the survival of *P. multocida*.

In this context, we pursued structure-activity relationship studies on NagA enzyme from *P. multocida* in order to lay groundwork for future structure-based drug design research. In this study we have characterized the unique structural and functional properties of this enzyme, especially the variations in the quaternary architecture as compared to its close homologs, the *Escherichia coli* NagA (EcNagA) and *Vibrio cholerae* NagA (VcNagA). Although PmNagA displays kinetic parameters similar to the other bacterial homologs, it also shows moderate hysteresis as seen in the distantly related *Staphylococcus aureus* NagA (SaNagA) ²⁸. Our structural data suggests that despite

PmNagA retaining the same quaternary state (a tetramer) as seen in EcNagA and VcNagA, the interactions at the dimer-dimer interface vary significantly. Experiments combining size-exclusion chromatography along with single-particle cryo electron microscopy suggest that PmNagA retains this altered interface and quaternary architecture in solution as well. Using bioinformatics, we have identified hot-spot residues present at the dimer-dimer interface that contribute significantly towards the altered architecture. Consurf analysis suggests that these hot-spot residues present at the surface are not conserved in the NagA enzyme family, especially in EcNagA and VcNagA, and have a faster mutation rate as compared to the slowly evolving core. Our study identifies PmNagA as a variant among the tetrameric homologs with a unique quaternary architecture and provides insight into how homo-oligomers evolved in the NagA enzyme family.

2. Materials and methods

2.1. Macromolecule production

The *P. multocida nagA* gene was synthesized and cloned into the pET300NT/DEST vector²⁹. Recombinant PmNagA was overexpressed in *E. coli* BL21 DE3 cells grown in Luria Bertani media supplemented with ampicillin (100 $\mu\text{g}\cdot\text{mL}^{-1}$). Cells were grown at 310 K until mid-log phase and protein overexpression was induced for 5 hours at 293 K by addition of 100 μM IPTG. Cells were pelleted by centrifugation at 6000g for 20 minutes at 277 K, subsequently resuspended in the lysis Buffer (70 mM Tris pH 8.0, 300 mM NaCl, 6% glycerol, 10 mM β -mercaptoethanol (βME) and one tablet of Roche cocktail protease inhibitor tablet per 100mL of lysis buffer) and lysed at 277 K using Emulsiflex-C5 (Avestin) at 15kpsi pressure. The protein was purified by immobilized metal affinity chromatography. Supernatant obtained after centrifugation (15000 g, 45 min, 277 K) was loaded onto a pre-equilibrated Econo Column containing 10 mL of Nickel-NTA agarose beads. The beads were washed with 20 column volumes (CV) of Buffer 1 (50 mM Tris pH 8.0, 300 mM NaCl, 6% glycerol, 10 mM βME), 5 CV of Buffer 2 (Buffer 1 + 50 mM Imidazole), 5 CV of Buffer 3 (Buffer 1 + 100 mM imidazole) and finally eluted with 3 CV of Buffer 4 (Buffer 1 + 300mM imidazole). The protein was concentrated (to 8 $\text{mg}\cdot\text{mL}^{-1}$) using Amicon 30K MCWO filter and injected onto a S200 preparative grade Size Exclusion Column equilibrated with 50 mM Tris pH 8.0, 50 mM NaCl, 500 μM ZnCl_2 and 10 mM βME .

2.2. Crystallization

Hanging drop vapor diffusion crystallization trials were carried out by mixing 0.2 μL of PmNagA (3-6 mg mL^{-1}) with 0.2 μL of commercial screens (PEG Ion 1 & 2 (Qiagen), Crystal Screen 1 & 2 (Emerald Biosystems) and Wizard Classic 1 & 2 (Rigaku) using Mosquito, the nanolitre robot from TTP Labtech. PmNagA crystals appeared in 3-4 days in the presence of 0.1 M HEPES sodium salt pH 7.5, 0.8 M sodium phosphate at 291 K.

2.3. Data collection and processing

The crystals were cryoprotected with reservoir buffer containing 20% ethylene glycol before flash cooling in liquid nitrogen. The single crystal diffraction data were collected at ID-29 beamline at ESRF, Grenoble. Data processing and reduction were performed using XDS³⁰ and AIMLESS³¹. The best PmNagA crystal diffracted to 1.95 Å.

2.4. Structure solution and refinement

Phases were obtained by molecular replacement using PHASER^{31,32} with a polyalanine model of *E. coli* NagA (PDB entry: 2p50)²³, which shares 54% sequence identity with PmNagA. The initial model was built with Autobuild³³, and manual building was executed in Coot³⁴ with iterative refinement cycles using Phenix_refine³⁵. Waters were added using ARP/WARP^{36,37} from the CCP4 suite³⁸. The PmNagA structure has been submitted to RCSB with a pdb code 6JKU. The data processing and refinement statistics are shown in Table I.

2.5. PISA, Dimplot and PPCheck for interface analysis

Dimer and tetramer interface of EcNagA (2p50), VcNagA (3iv8) and PmNagA (6jku) were analyzed using PISA³⁹ in the ccp4i2⁴⁰ suite. Dimplot⁴¹ was used for pictorial depiction of the monomer-monomer interface in PmNagA. PPCheck⁴² was used to identify hotspot residues at the dimer-dimer interface in PmNagA.

2.6. FoldX for assessing the role of the hotspot residues on thermodynamic stability of the PmNagA enzyme

The hotspot residues in PmNagA were computationally mutated to alanine to ascertain their energetic contribution towards the stability of the enzyme. The changes in stability was predicted by the Stability and AnalyzeComplex functions in FoldX⁴³. The ΔG prediction by FoldX is defined as the difference in free energy between the unfolded and the folded state of the protein ($\Delta G = G_{\text{folded}} - G_{\text{unfolded}}$). Thus, by measuring the difference of unfolding Gibbs free energy ($\Delta\Delta G$) between the mutant and the wild type (WT) PmNagA enzyme ($\Delta\Delta G = \Delta G_{\text{mutant}} - \Delta G_{\text{WT}}$), one can evaluate how the mutation affects the stability. The predicted $\Delta\Delta G$ value is positive if the mutation is destabilizing and negative when it is stabilizing⁴⁴. Based on the following $\Delta\Delta G$ values, we have considered a mutation to have a destabilizing ($\Delta\Delta G > 0.76 \text{ kcal/mol}$), neutralizing ($-0.76 \text{ kcal/mol} < \Delta\Delta G \leq +0.76 \text{ kcal/mol}$) or stabilizing ($\Delta\Delta G < -0.76 \text{ kcal/mol}$) effect.

Furthermore, the FoldX AnalyzeComplex command was used to determine the impact of the mutations on the interaction energy at the dimer-dimer interface of PmNagA involving chains A and C, and B and D, respectively. In order to predict the free energy of binding of a complex XY, FoldX computes the Gibbs free energies of the complex (ΔG_{XY}) and of the individual molecules X and Y alone. The interaction energy is then given by $\Delta G_{\text{interaction}} = \Delta G_{\text{XY}} - (\Delta G_{\text{X}} + \Delta G_{\text{Y}})$. While negative interaction energy suggests better binding, positive energy indicates no binding⁴⁵.

2.7. Multiple Sequence Alignment and Phylogenetic tree construction.

A multiple sequence alignment of PmNagA was performed with NagA sequences from various bacterial species and human NagA using MultAlign⁴⁶ and Clustal Omega⁴⁷. The sequence alignment in Figure 3C was generated using the ESPript3 server⁴⁸. A phylogenetic tree based on the maximum likelihood method was constructed using MEGA X⁴⁹ with several bacterial and human NagA protein sequences.

2.8. X-ray absorption spectra at Proxima-1 at Soleil

We validated the identity of the metal present in PmNagA by measuring the X-ray absorption spectra (XAS) of protein crystals. XAS scans were taken at PROXIMA-1 beamline at the SOLEIL synchrotron source at 107K. XAS is characterized by a sharp increase in absorption at specific energy, which is characteristic of the absorbing element. The spectrum derived from PmNagA crystals contains X-ray absorption lines characteristic of Zn with the peak at ~9660 eV, which corresponds to the K_{α} absorption edge of zinc.

2.9. Cryo-electron microscopy

Three μL of PmNagA in the size exclusion buffer ($0.5 \text{ mg}\cdot\text{mL}^{-1}$) was applied to the Quantifoil holey carbon grids (R 1.2/1.3) with blotting and freezing accomplished on a Vitrobot mark IV at 18°C and 100% RH. The data were collected with Titan Krios and Falcon 3 detector in counting mode at $1.07 \text{ \AA}/\text{pixel}$ sampling with images exposed for 60 seconds, with a total accumulated dose of $\sim 27 \text{ e}/\text{\AA}^2$ and dose fractionated into 25 frames, with each frame having a dose of $\sim 0.9 \text{ e}$. Unblur⁵⁰ was used for alignment and further processing was done in Relion 2.0⁵¹. The summed images were subsequently used for automated particle picking with Gautomatch, with template derived from previous data collection, and CTF was estimated with Gctf⁵². A total of 42,511 particles were extracted from 980 micrographs with a box size of 180 pixels and subjected to two rounds of 2D classification.

2.10. Kinetic assay

Kinetics analysis was carried out by measuring hydrolysis of the GlcNAc-6P carboxamide group at 215 nm ^{28,53}, using a Cary 100 Bio UV-Vis spectrophotometer (Agilent Technologies). An extinction coefficient for the GlcNAc-6-P carboxamide group was determined to be $352 \text{ M}^{-1} \text{ cm}^{-1}$ by measurement of absorbance before and after enzyme-catalyzed hydrolysis. Enzymatic activity was measured by initial velocity measurements performed with a final concentration of $2 \text{ }\mu\text{g}\cdot\text{mL}^{-1}$ NagA in 50 mM Tris-HCl pH 8.0 and varying amounts of GlcNAc-6P (Carbosynth UK), 1 mL total volume, with triplicate measurements undertaken and reproducible within 10% error. Control samples were tested with varied pH, temperature and enzyme concentration to ensure that the assay condition was near optimal and enzyme degradation during the assay was minimal. Initial rate data were then fitted to the Michaelis-Menten model using GRAPHPAD PRISM 6.0 software (GraphPad Software, La Jolla California USA, www.graphpad.com).

3. Results

3.1. PmNagA enzyme demonstrates kinetic hysteresis

PmNagA catalyzes the hydrolysis of the *N*-acetylglucosamine-6-phosphate carboxamide group to form glucosamine-6-phosphate. The carboxamide group absorbs strongly at 215 nm , and the hydrolysis can be easily followed by a decrease in absorbance in a UV spectrophotometer²⁸. The rate is directly proportional to the enzyme concentration, confirming that PmNagA has the predicted

deacetylase activity. Kinetic analysis demonstrates that PmNagA completes hydrolysis of GlcNAc-6P with kinetic parameters of $K_M = 0.13 \pm 0.004$ mM and $k_{cat} = 84 \pm 0.6$ s⁻¹, similar to other characterized NagA enzymes (Table II).

Kinetic characterization of EcNagA reports substrate inhibition¹³, which is visible at 2 mM GlcNAc-6P⁵³. Although closely related to EcNagA, PmNagA does not display substrate inhibition at high substrate concentrations (Figure 2A), and displays Michaelis-Menten model kinetics ($R^2 = 0.995$) more consistent with the distantly related MsNagA and SaNagA^{22,28}. Additionally, catalysis exhibits a hysteretic lag-phase (Figure 2B), as reported in *S. aureus* NagA (SaNagA)²⁸. Hysteresis is a slow increase in activity of an enzyme over time in response to fast changes in ligand concentration and is thought to be a way by which enzymes modulate their function⁵⁴. As seen in Figure 2B, the lag in enzyme activity becomes more pronounced at higher substrate concentrations. Here, we have taken the steep portion of the rate curve as the initial rate for the Michaelis-Menten curve fitting (Figure 2A), as it is expected to represent the steady-state⁵⁵. Kinetic characterization of SaNagA reveals a significant hysteretic lag phase at substrate concentrations above 0.5 mM²⁸. PmNagA experiences a smaller lag, where the rate at 0-10 seconds is 20% slower than the fastest rate and was only visible for substrate concentrations above 1 mM (Figure 2B). In summary, PmNagA does not display traditional substrate inhibition reported for the closely related EcNagA, but does display hysteresis, which is similar to the distantly related SaNagA.

3.2. Structure Solution

The structure of PmNagA was subsequently determined to 1.95Å resolution to see if the unique kinetic properties can be explained structurally. The enzyme crystallized in the spacegroup $P2_12_12_1$ with a tetrameric (dimer of dimers) assembly in the asymmetric unit. The final model includes residues -6 to 378 built in subunit A and B, -1 to 378 in subunit C and 0 to 378 in subunit D. The residues of the N terminal affinity tag (residues -6 to 0) are partially resolved in all the chains. Based on the presence of an unambiguous Fo-Fc density at 3σ and XAS data, one zinc ion was modeled in the active site of each subunit. The final refined structure had a R_{work}/R_{free} of 0.16/0.20, showed good geometry, with a low number of bond angle and bond length outliers. The data processing and structure refinement statistics are presented in Table I.

3.3. Overall Structure of PmNagA

PmNagA exists as a dimer of dimers with dihedral or D2 symmetry in the asymmetric unit. The PmNagA monomer is composed of two distinct domains as observed in previously published NagA structures^{3,21–23}. Domain I (residues 53–350) is a distorted (α/β)₈ TIM barrel, consisting of eight alternating α helices and β strands, and includes the zinc-bound catalytic site. Domain II is a β barrel structure composed of eight β strands and formed by residues from both the N-(1–52) and C-terminus (351–378) (Figure 3A). Structural superposition of PmNagA monomer on EcNagA (PDB: 2p50), BsNagA (PDB: 2vhl), TmNagA (PDB: 1o12) and MsNagA (PDB: 6fv3) resulted in an r.m.s.d ranging from 0.9Å to 1.52Å, which suggests that the overall tertiary structure of NagA is similar across these species.

A comparison of the β -barrel domain reveals varied architecture between NagA homologs. In BsNagA and TmNagA, strands β A and β B are undivided thereby forming a U-shaped β -barrel structure³. Whereas in EcNagA, strands β A and β B are divided into β A and β A1, and β B and β B1 respectively, thereby forming a β -sandwich structure. In this context, the β domain of PmNagA resembles the β sandwich structure of EcNagA²¹ (Figure 3A).

Dimerization is crucial for NagA enzyme activity, as the substrate interacts with residues from both subunits. Similar to other NagA enzymes^{3,22,23}, the α 6 and α 7 helices of both PmNagA subunits (Domain I) pack against each other, engage in multiple hydrophobic, salt bridge and hydrogen bond interactions, and bury a surface area of 3534 Å². Additionally, the β 6- α 7 loop of one monomer extends like an arm and interacts with its symmetry equivalent from the opposite monomer (Figure 3B). Substrate interaction has been characterized in MsNagA, where the guanidinium group of Arg-219, donated by the opposite subunit interacts with the phosphate moiety of GlcNAc-6P. Comparably, the guanidinium group of Arg-225 in PmNagA is positioned similarly to interact with the backbone oxygen of His-248 and a phosphate molecule present in the crystallization buffer.

Since the active site is present at the monomer-monomer interface with a large buried surface area, this interface is designated as the "catalytic interface" or "tight" interface. The catalytic interface is generally well conserved between NagA homologs. Comparison of the interfaces shows that although the number of polar interactions varies, several interacting residues are shared between homologs. Gly-224 carbonyl oxygen serves as an anchor point by forming hydrogen bonds with the carboxyl group of Asp-250 and carboxamide group of Asn-253 from opposite subunit. Additionally, an inter-subunit salt bridge interaction involving Arg-255 and Asp-235 is observed. These interactions

are conserved in EcNagA and VcNagA. Several water-mediated interactions connecting residues from opposite monomers are also observed. The carbonyl oxygen of Ile-221 of one subunit interacts with the main chain nitrogen of Ile-221 of the other subunit via a water molecule. Similarly, two water molecules connect Ser-220 with Lys-236 from the opposite subunits. Hydrophobic residues that constitute up to 50% of the interface residues also stabilize the dimer. Met-227 buries a surface area of 59 Å² and packs against Gly-224 from the opposite monomer (Figure S2). Multiple sequence alignment with various bacterial homologs shows that the residues present at the dimer interface are well conserved across bacterial species (Figure 3C).

3.4. Metal binding site and active site in PmNagA

Metal-dependent amidohydrolases have a mono or binuclear metal binding site and the role of the metal in the hydrolysis reaction varies with the enzyme and the substrate being hydrolyzed⁵⁶. BsNagA and MsNagA have binuclear metal binding sites^{3,22}, whereas EcNagA²³ and VcNagA are mononuclear. In most cases, this transition from a binuclear to a mononuclear arrangement is accompanied by the mutation of the HXH motif to a QXN motif and the presence of a histidine residue (His-143 in EcNagA), which is responsible for polarization of the carbonyl center of the substrate, GlcNAc-6P. PmNagA lacks the HXH motif required for binding the second metal ion at the M α site and is expected to have only one metal binding site (M β) (Figure S1). The presence of zinc in the PmNagA crystal was confirmed by the characteristic X-ray absorption spectra at 9.65 KeV at the Proxima I beamline in Soleil (Figure S4A). The metal bound PmNagA structure shows that zinc forms a tetrahedral complex at M β site, which coordinates with the metal ligating residues, Glu-128, His-192, His-213, and a phosphate molecule (Figure S4B & C). PmNagA was crystallized in a high salt condition with phosphate as part of the buffer and is anchored to the protein by direct hydrogen bonding to the carboxyl group of Asp-270. This phosphate from the buffer does not bind at the same place as the phosphate from GlcNAc-6P and the respective binding sites are in fact ~8.5Å apart. Analysis of the M α site reveals PmNagA does not bind a second metal ion. Instead, a water molecule occupies that position and interacts with several protein residues including Asn-61, Glu-128 and Asp-270 (Figure S4D). This contrasts with the binuclear metal binding as seen in MsNagA, wherein a zinc at M α site coordinates with His-65, Glu-122, Asp-267 and a water molecule shared with the M β zinc²².

3.5. Structure of PmNagA reveals an alternative tetrameric architecture

NagA homologs exist as either dimers or tetramers^{3,21,22}, with the dimer as the minimal functional unit. BsNagA, TmNagA and MsNagA exist as dimers both in solution and crystal (Figure 4A)^{3,22}. SaNagA is a dimer in solution²⁸. The theoretical molecular weight of the PmNagA monomer protein is 42.8 kDa. Size exclusion chromatography data suggest that PmNagA is likely to be a tetramer in solution since the protein ($8 \text{ mg} \cdot \text{mL}^{-1}$) eluted as a single peak at an elution volume of $V_e = 61.8 \text{ ml}$, which corresponds to a molecular weight of $\sim 180 \text{ kDa}$ (Figure S3A). Additionally, the cryo-electron microscopy 2D class average data of PmNagA ($0.5 \text{ mg} \cdot \text{mL}^{-1}$) indicates presence of a tetrameric assembly in solution as well (Figure S3B). EcNagA²¹ and VcNagA display a compact tetrameric structure assembled as dimer of dimers with D2 (222) symmetry. In both these enzymes, the β -barrel domain is shown to be involved in tetramerization. Amino acid sequence comparison revealed that although PmNagA shares $> 55\%$ sequence identity in the TIM barrel domain with EcNagA and VcNagA, the β -barrel domain is only 38% identical. This prompted us to have a close look at the quaternary architecture of PmNagA.

In the asymmetric unit of the crystal, PmNagA organizes as a D2 tetramer or a "dimer of dimers" with all the four subunits related by three two-fold axes (Figure 4B). Subunit A and B forms one dimer species, and subunit C and D forms the other dimer species. The monomer-monomer interactions involve subunits A and B, and C and D respectively. The dimer-dimer interactions are observed between subunits A and C, and B and D respectively. This interface is stabilized by hydrophobic interactions as well as two direct and eight water-mediated hydrogen bonds between residues contributed by the β domain and the $(\alpha/\beta)_8$ domain of respective subunits. The carboxamide nitrogen of Asn-362 present in the βK - βL loop of subunit A hydrogen bonds to the main chain carbonyl oxygen of Gly-48 in the βD - βE loop of subunit C. This interface is symmetric wherein equivalent amino acids from both the subunits are involved in various interactions. Thus, Asn-362 in subunit C also interacts with Gly-48 in subunit A. Another notable feature of the tetramer is the abundance of water molecules at the interface (Figure 5A & B), which mediate hydrogen bond interactions between residues from the relevant subunits. Residues Tyr-10 and Tyr-17, present in the βA1 and βB strands of subunit A interacts with their symmetry equivalents and Asn-362 in subunit C via water molecules. (Figure 5B). Hydrophobic residues dominate the interface (56-60%) with interactions observed between Val-8, Tyr-10 and Tyr-17 (the βA1 - βB1 loop) of subunit A and Val-318 (α8 helix), Leu-325 (α9 helix) and Phe-364 (βK - βL) loop of subunit C. Due to the symmetric

nature of interface, residues Val-8, Tyr-10 and Tyr-17 from subunit C also pack against Val-318, Leu-325 and Phe-364 of subunit A (Figure 5C). The BD interface also shows similar interactions. Since the dimer-dimer interface area is significantly smaller than the catalytic interface area, this interface is designated as the "weak interface" (Figure 4B).

3.6. PmNagA lacks the ionic interface as observed in EcNagA

Although PmNagA retains the same oligomeric state (D2 tetramer) as seen in the close homologs, EcNagA (54% identity) and VcNagA (58% identity), the dimer-dimer interface varies significantly. We have chosen EcNagA for comparison purposes, as this enzyme has been well characterized and the tetrameric nature was reported both in solution and crystal form²¹. EcNagA tetramer is more compact with greater buried surface area (6136 Å²), and the dimer-dimer interface involves three subunits and consists of a hydrophobic (AD interface) and an ionic interface (AC interface). In contrast, PmNagA has a buried surface area of (5561 Å²) with an open structure, where the dimer-dimer interface involves two subunits interacting through the hydrophobic interface (AC interface) (Figure 4B & C). Figure S5A compares the dimer-dimer interface between EcNagA, VcNagA and PmNagA enzymes. The AC interface in PmNagA is analogous to the AD interface in EcNagA. While in both cases this interface involves residues from the N and C terminus of the protein, differences lie in the amino acid residues involved and also in the nature of interaction. In EcNagA, a combination of hydrophobic, salt bridge, hydrogen bond interactions stabilize the hydrophobic interface (AD interface). In PmNagA, the dimer-dimer interface is stabilized mainly by hydrophobic interactions and water-mediated hydrogen bond interactions. An inter-subunit salt bridge and hydrogen bond interactions between Arg-8 (subunit A) and Glu-322 (subunit D) present in EcNagA (Figure S6A), is not conserved in PmNagA, as valine replaces arginine and glutamine replaces the glutamate residue. Moreover, the hydrophobic residues present at the interface also differ between the homologs. In EcNagA, Phe-15 in the βB1 strand, Val-321, Gly-325, Ile-326, Ala-327 in the α8 and α9 helices of subunit A pack against their symmetry equivalents from subunit D (Figure S6A). Although conserved in PmNagA, Gly-322 and Ile-323 do not form the tetrameric interface. In PmNagA, the hydrophobic patch involves residues Val-8, Tyr-10 and Tyr-17, Val-318, Leu-325 and Phe-364 (Figure 5C).

The major difference in the quaternary structure between these homologs is the absence of the charged interface in PmNagA, which is otherwise present in EcNagA (AC interface). In EcNagA, this interface is mainly stabilized by nine salt bridge interactions. The hydrophobic residues are also

under-represented (14%) (Figure S5A and S6B). Inter-subunit salt bridge pairs include Lys-204 and Glu14, Arg-341 and Asp-240, His-13 and Asp-240. In PmNagA, a glutamine replaces Lys-204 and precludes such an interaction at that position. Although, His-13, Glu-14, Asp-240, Arg-341 are either conserved or semi-conserved in PmNagA (Figure S5B), they do not form an interface. Consequently, due to the absence of the second interface, a solvent channel is observed at the center of the tetrameric assembly in PmNagA, otherwise absent in EcNagA (Figure 4B and C).

3.7. Identification of hotspot residues at the dimer-dimer interface in PmNagA.

Studies have shown that only few of the interface residues, also known as hotspot residues contribute significantly towards the total binding energy of the protein complex formation and play a crucial role in the stability of the complex⁵⁷. Since experimental methods using alanine scanning mutation are expensive and time consuming, computational methods have been extensively used to identify such hotspot residues at the protein-protein interface⁵⁸. With PPCheck we identified such hotspot residues at the dimer-dimer interface of PmNagA and subsequently used multiple sequence alignment to compare those residues in other homologs. PPCheck identified residues Tyr-17, Gly-18, Gln-319, Asn-362 and Asp-363 present at the center of the interface as potential hotspot residues, which engages in several polar interactions with residues from the opposite subunit. However, of all the hotspot residues, only Asp-363 is conserved in EcNagA. In EcNagA, Tyr-17 and Gly-18 are replaced by Asp residues (Asp-17 and Asp18), Gln-319 is replaced by Glu-319, and a smaller proline residue replaces Asn-362. Consurf analysis⁵⁹ also suggests that Tyr-17, Gly-18 and Asn-362, which are identified as hotspot residues in PmNagA, are not conserved in the NagA enzyme family. Our analysis suggests that the distinct quaternary architecture of PmNagA is possibly a result of rapidly mutating surface residues, which led to the formation of altered interactions at the interface. However, this change in dimer-dimer interface does not seem to have an effect on the catalytic properties of the enzyme. *In-silico* mutation of the hotspot residues in PmNagA to their EcNagA counterparts resulted in an unfavorable stabilization energy ($-105.74 \text{ kJ} \cdot \text{mol}^{-1}$ vs. $20.37 \text{ kJ} \cdot \text{mol}^{-1}$) of the dimer-dimer interface suggesting that these residues play an important role in forming the interface.

Moreover, we used the Stability and AnalyseComplex functions in the bioinformatic tool, FoldX, to quantitatively assess the energetic impact of these hotspot residues towards the overall stability of PmNagA oligomer. The hotspot residues were mutated to alanine to probe the thermodynamic stability and the interface interactions of the mutant structures in comparison to that

of the wild-type structure (See Methods). All most all the hotspot residues mutations to alanine have a destabilizing effect. Tyr-17-Ala (Y17A) is the most destabilizing mutation ($\Delta\Delta G = 4.8 \text{ kcal/mol}$), followed by Gly-18-Ala (G18A), Asp-363-Ala (D363A), Asn-362-Ala (N362A). Gln-319-Ala (Q319A) has a neutralizing effect on the stability of the protein ($\Delta\Delta G = 0.16 \text{ kcal/mol}$). Also, the simultaneous substitution of all five hotspot residues to alanine predicted a highly destabilizing effect ($\Delta\Delta G = 10.40 \text{ kcal/mol}$) (Figure S8A).

Similarly, the interaction energy profile at the dimer-dimer interface (AC and BD) of PmNagA, as predicted by AnalyseComplex also suggests that most of the hotspot residues when mutated destabilize the interface. The maximum effect was observed when all the five residues were substituted simultaneously, followed by Y17A mutation (Figure S8B). As predicted, the mutations at the dimer-dimer interface do not disrupt the catalytic interface (AB and CD), which shows a favorable interaction energy of -19 kcal/mol .

3.8. Cryo-EM data analysis confirms that PmNagA conserves the alternative tetramer architecture in solution.

We further investigated whether this altered dimer-dimer interface seen in the PmNagA crystal structure is also preserved in solution. For this purpose, we obtained the Cryo-EM 2D class average data of PmNagA in solution and compared it with the 2D projections of PmNagA crystal structure at the same resolution. We hypothesized that if the altered tetrameric architecture were maintained in solution, then both the 2D class average data and the 2D projections would show near identical features and dimensions. We created a map using the coordinates from the PmNagA crystal structure and filtered it to $10 \text{ \AA}$. Further, 2D projections of this map were created using Relion⁵¹ that resulted in 140 such projections at random angles. 2D classes from cryo-EM data were compared with these projections generated from the crystal structure using EMAN2 e2classvsproj.py⁶⁰. Although very few orientations of the protein were obtained from 2D class averaging, all these orientations matched well with the 2D projections from the crystal structure (Figure 6). This suggests that the overall tetrameric architecture and the altered interface in PmNagA are conserved both in the crystal environment and in solution as well.

4. Discussion

Protein oligomerization is crucial to its structure and function, as mis-assembled proteins have been implicated in several diseases. Levy and others⁶¹ showed that protein pairs that share 50-60%

sequence identity have the same oligomeric structure in >80% of the cases. In about 20% of the cases, variations in the quaternary architecture arise due to the dynamic nature of protein-protein interactions⁶², which emerge in evolution at a faster pace than a new protein fold⁶³. In this study, we present the unique quaternary variations observed in the structural assembly of tetrameric PmNagA, a deacetylase enzyme belonging to metallohydrolase family. Even though the primary, tertiary and oligomeric states are well conserved between EcNagA and PmNagA, close homologs sharing >50% sequence identity, interactions at the dimer-dimer interface differ significantly. In EcNagA, the dimer-dimer interface involves three subunits and comprises of a larger hydrophobic interface and a smaller ionic interface. Although PmNagA has a similar hydrophobic interface, it lacks the ionic interface.

This altered interface does not seem to affect the overall function of PmNagA, as both EcNagA and PmNagA display comparable kinetic properties (k_{cat} , K_M , k_{cat}/K_M). However, unlike EcNagA, PmNagA does not exhibit substrate inhibition. Also, similar to the dimeric SaNagA, PmNagA shows kinetic hysteresis, which could be linked to altered conformational dynamics at the active site⁵⁴. This is not surprising since NagA occupies a central position at the crossroads of various metabolic pathways and its function may need to be tightly regulated for the survival of the bacteria.

A maximum-likelihood method based phylogenetic tree constructed with several bacterial and human NagA enzymes suggest divergence of this enzyme family into two groups, differing in the metal center and in quaternary structure. The close homologs, BsNagA and MsNagA cluster together, possess an HXH motif, bind to an additional metal ion at the $M\alpha$ site, and are also coincidentally C2 dimers. One exception is TmNagA, which possess the HXH motif but has been shown to require a single metal ion for catalysis²³. EcNagA, VcNagA and PmNagA represent the second group, which exhibit mononuclear metal binding site and are D2 homotetramers (Figure S7C). Our study reveals that PmNagA represents yet another sub-group in the homo-tetramer population that demonstrates an altered quaternary architecture with a square topology (Figure S7C).

Studies suggest that D2 dihedral complexes are ten times more abundant than cyclic complexes and often arise from C2 dimeric intermediates. Multiple evolutionary pathways lead to the assembly of such dihedral complexes from its cyclic intermediates⁶⁴. Self-complementary or symmetrical interactions are over-represented in dihedral complexes and mostly arise due to amino acid substitutions⁶⁴. Although the interface symmetry is conserved at the dimer-dimer interface, residues involved in tetramerization vary between EcNagA and PmNagA. A Consurf analysis⁵⁹ on PmNagA

was performed to analyze the evolutionary conservation of amino acid residues based on the phylogenetic relations between NagA homologs. The analysis suggests that several of the surface residues (Tyr-17, Gly-48, Asn-50, Gln-319, Asn-362 and Phe-364) that form the dimer-dimer interface in PmNagA are variable and hence are susceptible to mutations. In contrast, the residues in the protein core, as well as the catalytic interface are highly conserved (Figure S7A & B). Moreover, several interface residues (Tyr-17, Gly-18, Gln-319 and Asn-362) were identified as potential hotspot residues that play a dominant role in the tetramer formation and when computationally mutated to alanine have destabilizing effect on the PmNagA oligomer. Further analysis suggests that these residues are not conserved in the other tetrameric homologs, EcNagA and VcNagA. Thus, it is possible that the diverse D2 tetramers (in EcNagA, VcNagA and PmNagA) evolved from the C2 dimers (BsNagA, MsNagA and TmNagA) as nature chose multiple routes for the assembly, and random mutations (such as Tyr-17, Gly-18, Gln-319 and Asn-362) gathered over a period on the protein surface have led to the emergence of a variant interface.

Why this altered interface and quaternary state were selected and propagated by evolutionary processes is open for speculation. Studies suggest that a large number of these homo-oligomers assemble into higher-order heteromeric complexes to fulfill various structural and functional roles in cells⁶⁵. It has been speculated that *in vivo*, NagA and NagB may form complexes that enable substrate channeling and improved overall activity²⁸. Like NagA, NagB is also known to form different oligomeric complexes. In *E.coli*, while NagA is a tetramer, NagB is a hexamer and is allosterically regulated by GlcNac-6P^{65,66}. In *P. multocida*, NagA is a tetramer and a preliminary study from our laboratory shows that NagB is a dimer (data not shown). It is tempting to hypothesize that the altered interface in PmNagA allows it to form functional complexes with PmNagB and thereby have a role in regulating the intracellular pool of the intermediates GlcNac-6P and GlcN-6P. Our data suggest that in the NagA family, the conserved binding mode at the monomer-monomer or the catalytic interface observed in both close and distant homologs indicates an ancient evolutionary origin whereas the diverse binding mode at the dimer-dimer interface between close homologs is suggestive of rapid acquisition of novel interactions in evolution.

Accession number

The accession numbers are from the NCBI or Uniprot or Genbank database.

Thermotoga maritima (WP_004080848.1); *Pasteurella multocida* (WP_014326154.1); *Escherichia coli* (WP_000271153.1); *Bacillus subtilis* (WP_003243413.1); *Vibrio cholerae* (WP_000271133.1); *Mycobacterium smegmatis* (WP_011728170.1); *Listeria monocytogenes* 1 (WP_070254771.1);

Listeria monocytogenes 2 (NP_464481.1); *Homo sapiens* (NP_001317378.1); *Haemophilus influenzae* (WP_011271899.1); *Fusobacterium nucleatum* (WP_008798351.1); *Caviibacterium pharyngocola* (WP_100295540.1); *Haemophilus ducreyi* (WP_010945619.1); *Aggregatibacter segnis* (WP_109104193.1); *Citrobacter koseri* (WP_047458163.1); *Salmonella enterica* (WP_000271196.1); *Streptococcus pneumoniae* (WP_001134463.1); *Lysinibacillus sphaericus* (Q84F86.1) ; *Streptococcus pyogenes* (PZO95670.1); *Shigella sonnei* (WP_000271169.1); *Burkholderia pseudomallei* K96243 (WP_004195336.1); *Bdellovibrio* sp (PIQ61352.1)

The PDB structures of NagA homologs discussed in this manuscript have the following RCSB codes: *Pasteurella multocida* (6jku), *Bacillus subtilis* (2vhl), *Mycobacterium smegmatis* (6fv3, 6fv4), *Thermotoga maritima* (1o12), *Vibrio cholerae* (3egj, 3iv8), *Escherichia coli* (2p50, 1yrr, 1ymy).

References

- (1) Brigham, C.; Caughlan, R.; Gallegos, R.; Dallas, M. B.; Godoy, V. G.; Malamy, M. H.; Al, B. E. T.; Ateriol, J. B. Sialic Acid (N -Acetyl Neuraminic Acid) Utilization by *Bacteroides Fragilis* Requires a Novel N -Acetyl Mannosamine Epimerase □ †. *J. Bacteriol.* **2009**, *191* (11), 3629–3638.
- (2) Steenbergen, S. M.; Lichtensteiger, C. A.; Caughlan, R.; Garfinkle, J.; Fuller, T. E.; Vimr, E. R. Sialic Acid Metabolism and Systemic Pasteurellosis. *Infect. Immun.* **2005**, *73* (3), 1284–1294.
- (3) Vincent, F.; Yates, D.; Garman, E.; Davies, G. J.; Brannigan, J. A. The Three-Dimensional Structure of the N-Acetylglucosamine-6-Phosphate Deacetylase, NagA, from *Bacillus Subtilis*: A Member of the Urease Superfamily. *J. Biol. Chem.* **2004**, *279* (4), 2809–2816..
- (4) Setty, T. G.; Cho, C.; Govindappa, S.; Apicella, M. A.; Ramaswamy, S. Bacterial Periplasmic Sialic Acid-Binding Proteins Exhibit a Conserved Binding Site. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **2014**, *70* (7), 1801–1811.
- (5) Wahlgren, W. Y.; Dunevall, E.; North, R. A.; Paz, A.; Scalise, M.; Bisignano, P.; Bengtsson-Palme, J.; Goyal, P.; Claesson, E.; Caing-Carlsson, R.; Andersson, R.; Beis, K.; Nilsson, U. J.; Farewell, A.; Pochini, L.; Indiveri, C.; Grabe, M.; Dobson, R. C. J.; Abramson, J.; Ramaswamy, S.; Friemann, R. Substrate-Bound Outward-Open Structure of a Na⁺-Coupled Sialic Acid Symporter Reveals a New Na⁺ Site. *Nat. Commun.* **2018**, *9* (1).
- (6) North, R. A.; Wahlgren, W. Y.; Remus, D. M.; Scalise, M.; Kessans, S. A.; Dunevall, E.; Claesson, E.; da Costa, T. P. S.; Perugini, M. A.; Ramaswamy, S.; Allison, J. R.; Indiveri, C.; Friemann, R.; Dobson, R. C. J. The Sodium Sialic Acid Symporter from *Staphylococcus Aureus* Has Altered Substrate Specificity. *Front. Chem.* **2018**, *6* (JUL).
- (7) North, R. A.; Horne, C. R.; Davies, J. S.; Remus, D. M.; Muscroft-Taylor, A. C.; Goyal, P.; Wahlgren, W. Y.; Ramaswamy, S.; Friemann, R.; Dobson, R. C. J. “Just a Spoonful of Sugar..”: Import of Sialic Acid across Bacterial Cell Membranes. *Biophysical Reviews.* 2018, pp 219–227.

- (8) North, R. A.; Kessans, S. A.; Griffin, M. D. W.; Watson, A. J. A.; Fairbanks, A. J.; Dobson, R. C. J. Cloning, Expression, Purification, Crystallization and Preliminary X-Ray Diffraction Analysis of *N*-Acetylmannosamine-6-Phosphate 2-Epimerase from Methicillin-Resistant *Staphylococcus Aureus*. *Acta Crystallogr. Sect. F Struct. Biol. Commun.* **2014**, 70 (5), 650–655.
- (9) Coombes, D.; Davies, J. S.; Newton-Vesty, M. C.; Horne, C. R.; Setty, T. G.; Subramanian, R.; Moir, J. W. B.; Friemann, R.; Panjikar, S.; Griffin, M. D. W.; North, R. A.; Dobson, R. C. J. The Basis for Non-Canonical ROK Family Function in the *N*-Acetylmannosamine Kinase from the Pathogen *Staphylococcus Aureus*. *J. Biol. Chem.* **2020**,
- (10) North, R. A.; Watson, A. J. A.; Pearce, F. G.; Muscroft-Taylor, A. C.; Friemann, R.; Fairbanks, A. J.; Dobson, R. C. J. Structure and Inhibition of *N*-Acetylneuraminate Lyase from Methicillin-Resistant *Staphylococcus Aureus*. *FEBS Lett.* **2016**, 590 (23), 4414–4428.
- (11) Manjunath, L.; Guntupalli, S. R.; Currie, M. J.; Rachel, A.; Dobson, R. C. J.; Nayak, V.; Subramanian, R.; Manjunath, L.; Guntupalli, R.; Currie, M. J.; Rachel, A. Crystal Structures and Kinetic Analyses of *N*-Acetylmannosamine-6-Phosphate 2-Epimerases from *Fusobacterium Nucleatum* and *Vibrio Cholerae* Crystal Structures and Kinetic Analyses of *N*-Acetylmannosamine-6-Phosphate 2-Epimerases from *Fusobacterium Nucleatum*. *Acta Crystallogr. Sect. F* **2018**, F74, 431–440.
- (12) Kumar, J. P.; Rao, H.; Nayak, V.; Ramaswamy, S. Crystal Structures and Kinetics of *N*-Acetylneuraminate Lyase from *Fusobacterium Nucleatum*. *Acta Crystallogr. Sect. F Struct. Biol. Commun.* **2018**, 74, 725–732.
- (13) Souza, J. M.; Plumbridge, J. A.; Calcagno, M. L. *N*-Acetylglucosamine-6-Phosphate Deacetylase From *Escherichia Coli*: Purification and Molecular and Kinetic Characterization. *Arch. Biochem. Biophys.* **1997**, 340 (2), 338–346.
- (14) Yadav, V.; Panilaitis, B.; Shi, H.; Numata, K.; Lee, K.; Kaplan, D. L. *N*-Acetylglucosamine 6-Phosphate Deacetylase (NagA) Is Required for *N*-Acetyl Glucosamine Assimilation in *Gluconacetobacter Xylinus*. *PLoS One* **2011**, 6 (6), 1–11.
- (15) Seibert, C. M.; Raushel, F. M. Structural and Catalytic Diversity within the Amidohydrolase Superfamily. *Biochemistry* **2005**, 44 (17), 6383–6391.
- (16) Holm, L.; Sander, C. An Evolutionary Treasure: Unification of a Broad Set of Amidohydrolases Related to Urease. *Proteins Struct. Funct. Genet.* **1997**, 28 (1), 72–82.
- (17) Plumbridge, J. A. A Dominant Mutation in the Gene for the Nag Repressor. *J. Gen. Microbiol.* **1992**, 138, 1011–1017.
- (18) Calcagno, M.; Campos, P. J.; Mulliert, G.; Suástegui, J. Purification, Molecular and Kinetic Properties of Glucosamine-6-Phosphate Isomerase (Deaminase) from *Escherichia Coli*. *Biochim. Biophys. Acta (BBA)/Protein Struct. Mol.* **1984**, 787 (2), 165–173.
- (19) Komatsuzawa, H.; Fujiwara, T.; Nishi, H.; Yamada, S.; Ohara, M.; McCallum, N.; Berger-Bächi, B.; Sugai, M. The Gate Controlling Cell Wall Synthesis in *Staphylococcus Aureus*. *Mol. Microbiol.* **2004**, 53 (4), 1221–1231.
- (20) Barreteau, H.; Kovac, A.; Boniface, A.; Sova, M.; Gobec, S.; Blanot, D. Cytoplasmic

- Steps of Peptidoglycan Biosynthesis. *Fed. Eur. Microbiol. Soc.* **2008**, 32, 168–207.
- (21) Ferreira, F. M.; Mendoza-Hernandez, G.; Castañeda-Bueno, M.; Aparicio, R.; Fischer, H.; Calcagno, M. L.; Oliva, G. Structural Analysis of N-Acetylglucosamine-6-Phosphate Deacetylase Apoenzyme from *Escherichia Coli*. *J. Mol. Biol.* **2006**, 359 (2), 308–321.
- (22) Ahangar, M. S.; Furze, C. M.; Guy, C. S.; Cooper, C.; Maskew, K. S.; Cameron, A. D.; Fullam, E. Structural and Functional Determination of Homologs of the *Mycobacterium Tuberculosis* N -Acetylglucosamine-6-Phosphate Deacetylase (NagA). *J Biol Chem* **2018**, 293 (25), 9770–9783.
- (23) Hall, R. S.; Brown, S.; Fedorov, A. A.; Fedorov, E. V.; Xu, C.; Babbitt, P. C.; Almo, S. C.; Raushel, F. M. Structural Diversity within the Mononuclear and Binuclear Active Sites of N-Acetyl-D-Glucosamine-6-Phosphate Deacetylase. *Biochemistry* **2007**, 46 (27), 7953–7962..
- (24) Popowska, M.; Osińska, M.; Rzczkowska, M. N-Acetylglucosamine-6-Phosphate Deacetylase (NagA) of *Listeria Monocytogenes* EGD, an Essential Enzyme for the Metabolism and Recycling of Amino Sugars. *Arch. Microbiol.* **2012**, 194 (4), 255–268.
- (25) de Emery, B. D.; Furian, T. Q.; Pilatti, R. M.; Chitolina, G. Z.; Borges, K. A.; Salle, C. T. P.; Moraes, H. L. S. Evaluation of the Biofilm Formation Capacity of *Pasteurella Multocida* Strains Isolated from Cases of Fowl Cholera and Swine Lungs and Its Relationship with Pathogenicity. *Pesqui. Vet. Bras.* **2017**.
- (26) Almagro-Moreno, S.; Boyd, E. F. Insights into the Evolution of Sialic Acid Catabolism among Bacteria. *BMC Evol. Biol.* **2009**, 9 (1), 118.
- (27) Johnston, J. W.; Coussens, N. P.; Allen, S.; Houtman, J. C. D.; Turner, K. H.; Zaleski, A.; Ramaswamy, S.; Gibson, B. W.; Apicella, M. A. Characterization of the N-Acetyl-5-Neuraminic Acid-Binding Site of the Extracytoplasmic Solute Receptor (SiaP) of Nontypeable *Haemophilus Influenzae* Strain 2019. *J. Biol. Chem.* **2008**, 283 (2), 855–865..
- (28) Davies, J. S.; Coombes, D.; Horne, C. R.; Pearce, F. G.; Friemann, R.; North, R. A.; Dobson, R. C. J. Functional and Solution Structure Studies of Amino Sugar Deacetylase and Deaminase Enzymes from *Staphylococcus Aureus*. *FEBS Lett.* **2019**, 593 (1), 52–66..
- (29) Bairy, S.; Gopalan, L. N.; Setty, T. G.; Srinivasachari, S.; Manjunath, L.; Kumar, J. P.; Guntupalli, S. R.; Bose, S.; Nayak, V.; Ghosh, S.; Sathyanarayanan, N.; Caing-Carlsson, R.; Wahlgren, W. Y.; Friemann, R.; Ramaswamy, S.; Neerathilingam, M. Automation Aided Optimization of Cloning, Expression and Purification of Enzymes of the Bacterial Sialic Acid Catabolic and Sialylation Pathways Enzymes for Structural Studies. *Microb. Biotechnol.* **2018**, 11 (2), 420–428.
- (30) Kabsch, W. XDS. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **2010**, 66 (2), 125–132.
- (31) Evans, P. R.; Murshudov, G. N. How Good Are My Data and What Is the Resolution? *Acta Crystallogr. Sect. D Biol. Crystallogr.* **2013**.
- (32) Read, R. J. Pushing the Boundaries of Molecular Replacement with Maximum Likelihood. *Acta Crystallogr. - Sect. D Biol. Crystallogr.* **2001**, 57 (10), 1373–1382.
- (33) Terwilliger, T. C.; Grosse-Kunstleve, R. W.; Afonine, P. V.; Moriarty, N. W.; Zwart,

- P. H.; Hung, L. W.; Read, R. J.; Adams, P. D. Iterative Model Building, Structure Refinement and Density Modification with the PHENIX AutoBuild Wizard. In *Acta Crystallographica Section D: Biological Crystallography*; 2007; Vol. 64, pp 61–69.
- (34) Emsley, P.; Lohkamp, B.; Scott, W. G.; Cowtan, K. Features and Development of Coot. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **2010**, 66 (4), 486–501.
- (35) Adams, P. D.; Afonine, P. V.; Bunkóczi, G.; Chen, V. B.; Davis, I. W.; Echols, N.; Headd, J. J.; Hung, L.-W.; Kapral, G. J.; Grosse-Kunstleve, R. W.; McCoy, A. J.; Moriarty, N. W.; Oeffner, R.; Read, R. J.; Richardson, D. C.; Richardson, J. S.; Terwilliger, T. C.; Zwart, P. H. *PHENIX*: A Comprehensive Python-Based System for Macromolecular Structure Solution. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **2010**, 66 (2), 213–221.
- (36) Lamzin, B. Y. V. S.; Wilson, K. S.; Molecular, E.; Hamburg, D.-. Automated Refinement of Protein Models. *Acta Cryst* **1993**, 49, 129–147.
- (37) Murshudov, G. N.; Nicholls, R. A. Research Papers REFMAC 5 for the Refinement of Macromolecular Crystal Structures Research Papers. *Acta Crystallogr. Sect. D Struct. Biol.* **2011**, 67, 355–367. <https://doi.org/10.1107/S0907444911001314>.
- (38) Winn, M. D.; Ballard, C. C.; Cowtan, K. D.; Dodson, E. J.; Emsley, P.; Evans, P. R.; Keegan, R. M.; Krissinel, E. B.; Leslie, A. G. W.; McCoy, A.; McNicholas, S. J.; Murshudov, G. N.; Pannu, N. S.; Potterton, E. A.; Powell, H. R.; Read, R. J.; Vagin, A.; Wilson, K. S. Overview of the CCP4 Suite and Current Developments. *Acta Crystallographica Section D: Biological Crystallography*. 2011, pp 235–242.
- (39) Krissinel, E.; Henrick, K. Inference of Macromolecular Assemblies from Crystalline State. *J. Mol. Biol.* **2007**, 372 (3), 774–797.
- (40) Evans, P. R.; Jenkins, H. T.; Keegan, R.; Krissinel, E.; Stevenson, K.; Lebedev, A.; McNicholas, S. J.; Nicholls, R. A.; Noble, M.; Pannu, N. S.; Roth, C.; Sheldrick, G.; Skubak, P. CCP 4 i 2 : The New Graphical User Interface to the CCP 4 Program Suite Research Papers. *Acta Crystallogr. Sect. D Struct. Biol.* **2018**, No. August 2017, 68–84.
- (41) Laskowski, R. A.; Swindells, M. B. LigPlot+: Multiple Ligand-Protein Interaction Diagrams for Drug Discovery. *J. Chem. Inf. Model.* **2011**, 51 (10), 2778–2786.
- (42) Sukhwai, A.; Sowdhamini, R. PPcheck: A Webserver for the Quantitative Analysis of Protein-Protein Interfaces and Prediction of Residue Hotspots. *Bioinform. Biol. Insights* **2015**, 9, 141–151..
- (43) Schymkowitz, J.; Borg, J.; Stricher, F.; Nys, R.; Rousseau, F.; Serrano, L. The FoldX Web Server: An Online Force Field. *Nucleic Acids Res.* **2005**, 33 (SUPPL. 2).
- (44) Li, L.; Ye, Y.; Sang, P.; Yin, Y.; Hu, W.; Wang, J.; Zhang, C.; Li, D.; Wan, W.; Li, R.; Li, L.; Ma, L.; Xie, Y.; Meng, Z. Effect of R119G Mutation on Human P5CR1 Dynamic Property and Enzymatic Activity. *Biomed Res. Int.* **2017**, 2017.
- (45) Pierri, C. L.; Bossis, F.; Punzi, G.; De Grassi, A.; Cetrone, M.; Parisi, G.; Tricarico, D. Molecular Modeling of Antibodies for the Treatment of TNF α -Related Immunological Diseases. *Pharmacol. Res. Perspect.* **2016**, 4 (1).
- (46) Corpet, F. Multiple Sequence Alignment with Hierarchical Clustering. *Nucleic Acids Res.* **1988**, 16 (22), 10881–10890.
- (47) Sievers, F.; Higgins, D. G. Clustal Omega. *Curr. Protoc. Bioinforma.* **2014**, 2014,

- 3.13.1-3.13.16.
- (48) Gouet, P.; Courcelle, E.; Stuart, D. I.; Métoz, F. ESPript: Analysis of Multiple Sequence Alignments in PostScript. *Bioinformatics* **1999**, *15* (4), 305–308.
 - (49) Kumar, S.; Stecher, G.; Li, M.; Knyaz, C.; Tamura, K. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Mol. Biol. Evol.* **2018**, *35* (6), 1547–1549.
 - (50) Brilot, A. F.; Chen, J. Z.; Cheng, A.; Pan, J.; Harrison, S. C.; Potter, C. S.; Carragher, B.; Henderson, R.; Grigorieff, N. Beam-Induced Motion of Vitrified Specimen on Holey Carbon Film. *J. Struct. Biol.* **2012**, *177* (3), 630–637.
 - (51) Kimanius, D.; Forsberg, B. O.; Scheres, S. H.; Lindahl, E. Accelerated Cryo-EM Structure Determination with Parallelisation Using GPUs in RELION-2. *Elife* **2016**, *5*.
 - (52) Zhang, K. Gctf: Real-Time CTF Determination and Correction. *J. Struct. Biol.* **2016**, *193* (1), 1–12.
 - (53) Hall, R. S.; Xiang, F. D.; Xu, C.; Raushel, F. M. Activation via a Single Divalent Metal Ion †. *Biochemistry* **2007**, *46* (27), 7942–7952.
 - (54) Frieden, C. Kinetic Aspects of Regulation of Metabolic Processes. The Hysteretic Enzyme Concept. *J. Biol. Chem.* **1970**, *245* (21), 5788–5799.
 - (55) Masson, P. Time-Dependent Kinetic Complexities in Cholinesterase-Catalyzed Reactions. *Biochem.* **2012**, *77* (10), 1147–1161.
 - (56) Sugrue, E.; Fraser, N. J.; Hopkins, D. H.; Carr, P. D.; Khurana, J. L.; Oakeshott, J. G.; Scott, C.; Jackson, C. J. Evolutionary Expansion of the Amidohydrolase Superfamily in Bacteria in Response to the Synthetic Compounds Molinate and Diuron. *Appl. Environ. Microbiol.* **2015**, *81* (7), 2612–2624.
 - (57) Wang, H.; Liu, C.; Deng, L. Enhanced Prediction of Hot Spots at Protein-Protein Interfaces Using Extreme Gradient Boosting. *Sci. Rep.* **2018**, *8* (1).
 - (58) Grosdidier, S.; Fernández-Recio, J. Identification of Hot-Spot Residues in Protein-Protein Interactions by Computational Docking. *BMC Bioinformatics* **2008**, *9*.
 - (59) Ashkenazy, H.; Abadi, S.; Martz, E.; Chay, O.; Mayrose, I.; Pupko, T.; Ben-Tal, N. ConSurf 2016: An Improved Methodology to Estimate and Visualize Evolutionary Conservation in Macromolecules. *Nucleic Acids Res.* **2016**, *44* (W1), W344–W350.
 - (60) Tang, G.; Peng, L.; Baldwin, P. R.; Mann, D. S.; Jiang, W.; Rees, I.; Ludtke, S. J. EMAN2: An Extensible Image Processing Suite for Electron Microscopy. *J. Struct. Biol.* **2007**, *157* (1), 38–46.
 - (61) Levy, E. D.; Erba, E. B.; Robinson, C. V.; Teichmann, S. A. Assembly Reflects Evolution of Protein Complexes. *Nature* **2008**, *453* (7199), 1262–1265.
 - (62) Larion, M.; Moore, L. B.; Thompson, S. M.; Miller, B. G. Divergent Evolution of Function in the ROK Sugar Kinase Superfamily: Role of Enzyme Loops in Substrate Specificity. *Biochemistry* **2007**, *46* (47), 13564–13572.
 - (63) Perica, T.; Marsh, J. A.; Sousa, F. L.; Natan, E.; Colwell, L. J.; Ahnert, S. E.; Teichmann, S. A. The Emergence of Protein Complexes: Quaternary Structure, Dynamics and Allostery. In *Biochemical Society Transactions*; 2012; Vol. 40, pp 475–491.
 - (64) Venkatakrishnan, A. J.; Levy, E. D.; Teichmann, S. A. Homomeric Protein Complexes: Evolution and Assembly. *Biochemical Society Transactions*. 2010, pp

- 879–882.
- (65) Oliva, G.; Fontes, M. R.; Garratt, R. C.; Altamirano, M. M.; Calcagno, M. L.; Horjales, E. Structure and Catalytic Mechanism of Glucosamine 6-Phosphate Deaminase from *Escherichia Coli* at 2.1 Å Resolution. *Structure* **1995**, 3 (12), 1323–1332.
- (66) Álvarez-Añorve, L. I.; Gaugué, I.; Link, H.; Marcos-Viquez, J.; Díaz-Jiménez, D. M.; Zonszein, S.; Bustos-Jaimes, I.; Schmitz-Afonso, I.; Calcagno, M. L.; Plumbridge, J. Allosteric Activation of *Escherichia Coli* Glucosamine-6-Phosphate Deaminase (NagB) in Vivo Justified by Intracellular Amino Sugar Metabolite Concentrations. *J. Bacteriol.* **2016**, 198 (11), 1610–1620.
- (67) Yamano, N.; Higashida, N.; Endo, C. Purification and Characterization of N-Acetylglucosamine-6-Phosphate Deacetylase from a Psychrotrophic Marine Bacterium, *Alteromonas* Species. *Mar* **2000**, 57–64.

Figure Legends

Figure 1 The reaction catalyzed by *N*-acetylglucosamine-6-phosphate deacetylase (NagA) and its role in bacteria. **A.** The reaction catalyzed by *N*-acetylglucosamine-6-phosphate deacetylase results in two products, glucosamine-6-phosphate and acetate. Figure was made using CHEMDRAW (Cambridge Software). **B.** The central role of NagA in bacterial metabolic processes. NagA is

involved in the sialic acid catabolic pathway (black arrows), which feeds into glycolysis. NagA also provides the substrates to make the precursor (UDP-GlcNAc) of the cell wall and LOS biosynthesis pathways. LOS is represented by green wavy line.

Figure 2 Kinetic analysis of PmNagA. **A.** Initial rate data plotted by substrate concentration fitted to Michaelis-Menten equation. The fit had a high correlation ($R^2 = 0.995$), giving kinetic parameters of $K_M = 0.133$ mM (95% CI, 0.124-0.141 mM) and $k_{cat} = 83.6$ s⁻¹ (95% CI, 82.3-89.9 s⁻¹). The fit is from three independent experiments with the data points marked as open circles (O). **B.** Catalytic rate curves with 1, 2 and 4 mM *N*-acetylglucosamine-6-phosphate (solid line). A lag-phase occurs after initiation and rates between 0-10 sec are approximately 20% lower than the fastest rate (dashed line). The lag phase increases with substrate concentration, which is consistent with kinetic analysis from SaNagA¹⁸.

Figure 3 Monomer and Dimer architecture in PmNagA. **A.** Cartoon structure showing the two domains of NagA. Domain I forms a distorted α/β TIM barrel and Domain II is comprised of the β sandwich fold. The α helices and β strands are numbered as per the EcNagA nomenclature **B.** Surface representation of the dimer structure showing the metal (magenta sphere) in the active site. Each monomer is represented in pink and yellow respectively. **C.** Multiple sequence alignment shows sequence conservation of the bacterial homologs at the dimer interface. PmNagA, EcNagA, VcNagA, BsNagA, MsNagA, and TmNagA represent the bacterial enzymes. The residues that are found at the dimer interface are marked with a green rectangle. Fully conserved residues are marked red while semi-conserved residues are marked yellow. The figure has been generated in Esript3.

Figure 4 Quaternary structures of NagA homologs. The oligomeric organization of MsNagA (**A.**) PmNagA (**B.**) and EcNagA (**C.**) are shown in schematic and surface representation. The monomers are colored green, cyan, magenta and yellow. The monomer-monomer interactions are depicted by red line. The dimer-dimer interactions are represented by blue line. PmNagA is a loose tetramer with a solvent channel in the middle. The dimer-dimer interface shows square topology and involves only two subunits. EcNagA displays a compact tetramer wherein the dimer-dimer interface involves interactions between three subunits. The dimeric MsNagA is a C2 dimer whereas EcNagA and PmNagA can be described as D2 tetramers or dimer of dimers.

Figure 5 The dimer-dimer interface in PmNagA. **A.** This interface in PmNagA is shown in surface and stick representation. The water molecules present at the interface are shown as blue spheres. The interface residues as identified by PISA analysis are shown in stick representation. **B.**

The inter-subunit hydrogen bond interactions, both direct and water-mediated are shown. Residues Tyr-10, Tyr-17, Gly-48 and Asn-362 are shown in sticks representation. The hydrogen bonds are shown as red dashes and the distances are shown in Å. **C.** Hydrophobic interactions involve residues Val-8, Tyr-10, Tyr-17, Val-318, Leu-325 and Phe-364. A black ellipse represents the non-crystallographic 2-fold axis at the dimer-dimer interface.

Figure 6 PmNagA is a tetramer both in solution and in crystal form. Comparison of the 2D class averages generated from the Cryo-EM data with 2D projections generated from the crystal structure at same resolution show near identical features. The pixel size is 1.07 Å and box size is 180 pixels for both the 2D class averages and the projections. This suggests that the quaternary architecture observed in the crystal state is also preserved in the solution state.

A.

B.

PROT_25996_Figure_1.tif

PROT_25996_Figure_2.tif

PROT_25996_Figure_3.tif

PROT_25996_Figure_4.tif

PROT_25996_Figure_5.tif

PROT_25996_Figures_6.tif

Table I Crystallography Data processing and Refinement Statistics. Numbers in parenthesis refers to the highest resolution bin.

Data processing	PmNagA (6JKU)
Diffraction source	ID-29, ESRF
Wavelength (Å)	0.974
Temperature (K)	100K
Space group	P2 ₁ 2 ₁ 2 ₁
a, b, c (Å)	89.16, 139.55, 166.56
α , β , γ (°)	90, 90, 90
Mosaicity (°)	0.09
Resolution range (Å)	47.13 -1 .95 (2.02-1.95)
Total No. of reflections	657486 (33122)
No. of unique reflections	134675 (6848)
Completeness (%)	88.9(92.7)
Redundancy	4.9 (4.8)
$\langle I/\sigma(I) \rangle$	10 (1.9)
CC0.5	0.997(0.635)
Rmerge	0.112 (0.745)
Rpim.	0.09 (0.46)
Overall B factor from Wilson plot (Å ²)	24.36
Refinement	
Reflections used in refinement	134630 (13893)
Reflections used for R-free	6601 (669)
R-work	0.163 (0.2393)
R-free	0.20(0.2888)
Number of non-hydrogen atoms	12819
Macromolecules	11483
Ligands	183
Protein residues	1529
RMSD (bonds)	0.007
RMSD (angles)	1.21
Ramachandran favored (%)	95.40
Ramachandran outliers (%)	0.85
Clash score	2.50
Average B-factor	27.37
Macromolecule	26.46
Ligand	41.28
Solvent	34.23

Table II Table comparing the kinetic properties from different bacterial enzymes including PmNagA, EcNagA, VcNagA, MsNagA and SaNagA.

	PmNagA	EcNagA ^{13,53}	VcNagA ⁶⁷	MsNagA ²²	SaNagA ²⁸
k_{cat} (s ⁻¹)	84 ± 0.6	102 ± 2 (105)	0.97	91 ± 6	345 ± 6
K_M (mM)	0.13 ± 0.004	0.08 ± 0.01 (0.30 ± 0.02)	0.03	3.2 ± 0.4	0.16 ± 0.01
k_{cat}/K_M (M ⁻¹ ·s ⁻¹)	6.3×10 ⁵	1.3×10 ⁶ (3.5×10 ⁵)	3.2×10 ⁴	2.8×10 ⁴	2.2×10 ⁶

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Manjunath, L;Coombes, D;Davies, J;Dhurandhar, M;Tiwari, VR;Dobson, RCJ;Sowdhamini, R;Ramaswamy, S;Bose, S

Title:

Quaternary variations in the structural assembly of N-acetylglucosamine-6-phosphate deacetylase from *Pasteurella multocida*

Date:

2020-09-14

Citation:

Manjunath, L., Coombes, D., Davies, J., Dhurandhar, M., Tiwari, V. R., Dobson, R. C. J., Sowdhamini, R., Ramaswamy, S. & Bose, S. (2020). Quaternary variations in the structural assembly of N-acetylglucosamine-6-phosphate deacetylase from *Pasteurella multocida*. *PROTEINS-STRUCTURE FUNCTION AND BIOINFORMATICS*, 89 (1), pp.81-93. <https://doi.org/10.1002/prot.25996>.

Persistent Link:

<http://hdl.handle.net/11343/276299>