DR. COLLIN K CHIN (Orcid ID : 0000-0002-2819-4041)
DR. KATHARINE LOUISE LEWIS (Orcid ID : 0000-0003-0549-5877)
DR. PREETESH JAIN (Orcid ID : 0000-0003-2735-168X)
DR. JOHN FRANCIS SEYMOUR (Orcid ID : 0000-0003-2188-6835)
PROF. DAVID RITCHIE (Orcid ID : 0000-0001-7329-6605)
DR. CONSTANTINE TAM (Orcid ID : 0000-0002-9759-5017)
DR. NATHAN FOWLER (Orcid ID : 0000-0002-9494-1877)
DR. FELIPE SAMANIEGO (Orcid ID : 0000-0001-6931-4583)

Article type : Ordinary Papers

Article Type : Original Article

Title : Autologous stem cell transplantation for untreated transformed indolent B-cell lymphoma in first remission: an international, multi-centre propensity-score matched study

Author(s) : ^{1,2}C. K. Chin; ²K. J. Lim; ^{3,5}K. Lewis; ¹P. Jain; ⁴Y. Qing; ⁴L. Feng; ^{3,5}C. Y. Cheah; ^{2,6}J. F. Seymour; ^{2,6}D. Ritchie; ²K. Burbury; ^{2,6,7}C. S. Tam; ¹N. H. Fowler; ¹L. E. Fayad; ¹J. R. Westin; ¹S. S. Neelapu; ¹F. B. Hagemeister; ¹F. Samaniego; ¹C. R. Flowers; ¹L. J. Nastoupil; ^{2,6}M. J. Dickinson.

Affiliation(s) : ¹*Department of Lymphoma/Myeloma, University of Texas MD Anderson Cancer Center, Houston, Texas;* ²*Department of Clinical Haematology, Peter MacCallum Cancer Centre and Royal Melbourne Hospital, Melbourne, Australia;* ³*Department of Haematology, Sir Charles Gairdner Hospital, Nedlands, Australia;* ⁴*Department of Biostatistics, University of Texas MD Anderson Cancer Center, Houston, Texas;* ⁵*Medical School, University of Western Australia, Nedlands, Australia;* ⁶*The University of Melbourne, Melbourne, Australia;* ⁷*St Vincent's Hospital Melbourne, Melbourne, Australia.*

Corresponding Author

Dr Michael J. Dickinson

Department of Clinical Haematology

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/BJH.17072](https://doi.org/10.1111/BJH.17072)

This article is protected by copyright. All rights reserved

1 Peter MacCallum Cancer Centre & Royal Melbourne Hospital

2 305 Grattan Street

3 Melbourne, VIC 3000

4 **Email:** Michael.Dickinson@petermac.org

5 **Phone:** +61 3 8559 5000

6 **Version Number and Date** : Version 3.0, Wednesday 08 July 2020

7 **Target Journal** : British Journal of Haematology

8 **Word count (abstract)** : 200 (limit – 200 words)

9 **Word count (total)** : 2843 (limit – 3000 words)

10 **Figures/Tables** : 6 (3 tables, 3 figures)

11 **Supplementary Data** : 1 Table

12 **References** : 28 (limit – 60)

13
14 **Key words** (*maximum 5*)

15 Transformed lymphoma, autologous stem cell transplantation, non-Hodgkin

16 lymphoma, indolent lymphoma

17
18
19
20 **Abstract**

21
22 High dose chemotherapy (HDC) and autologous stem cell transplantation (ASCT) is
23 used as consolidation in first remission (CR1) in some centres for untreated,
24 transformed indolent B cell lymphoma (Tr-iNHL) but the evidence-base is weak. 319
25 patients with untreated Tr-iNHL meeting pre-specified transplant eligibility criteria
26 (age<75, LVEF≥45%, no severe lung disease, CR by PET or CT≥3 months after at
27 least standard RCHOP intensity front-line chemotherapy) were retrospectively
28 identified. Non-diffuse large B-cell lymphoma transformations were excluded.
29 283(89%) patients had follicular lymphoma, 30(9%) marginal zone lymphoma, 6(2%)
30 other subtypes. 49 patients underwent HDC/ASCT in CR1, and a 1:2 propensity-
31 score matched cohort of 98 patients based on age, stage, and HGBL-DH was
32 generated. After a median follow-up of 3.7 (range 0.1-18.3) years, ASCT was
33 associated with significantly superior PFS (HR 0.51,0.27-0.98;P=0.043) with a trend
34 towards inferior OS (HR 2.36;0.87-6.42;P=0.1) due to more deaths from progressive

1 disease (8%v4%). 40(41%) patients experienced relapse in the non-ASCT cohort–
2 15 underwent HDC/ASCT with 7(47%) ongoing CR; 10 CAR-T therapy with 6(60%)
3 ongoing CR; 3 allogeneic SCT with 2(67%) ongoing CR. Although ASCT in CR1
4 improves initial duration of disease control in untreated Tr-iNHL, the impact on OS is
5 less clear with effective salvage therapies in the CAR-T era.
6
7
8

1 Introduction

2 Large cell transformation of indolent B-cell lymphoma (Tr-iNHL) can be manifest at
3 initial presentation or occur over time and accounts for 13% of *de novo* presentations
4 of diffuse large B-cell lymphoma (DLBCL) (1, 2). Several retrospective studies
5 suggest that untreated Tr-iNHL may have similar outcomes to non-transformed
6 DLBCL in the rituximab era(3-6). The optimal treatment approach of untreated Tr-
7 iNHL is unknown, but is generally extrapolated from prospective clinical trials in *de*
8 *novo* DLBCL. Additionally, some centres implement high dose chemotherapy (HDC)
9 and autologous stem cell transplantation (ASCT) as consolidation in first remission
10 (CR1) in fit patients where the transformed disease is not encompassable in a single
11 loco-regional radiation field. This approach is based on the understanding that
12 transformation of low grade lymphoma has a poor prognosis, but is supported by
13 limited data(1, 7, 8). We performed an international, multi-centre, retrospective study
14 of transplant eligible patients with untreated Tr-iNHL to assess the impact of
15 **consolidation ASCT in CR1 v. no ASCT in CR1** on progression free survival (PFS)
16 and overall survival (OS). In addition, we assessed the impact of patient & disease
17 factors on outcomes. Finally, we describe patterns of relapse & subsequent salvage
18 therapies to determine the effect of upfront ASCT consolidation on treatment
19 sequencing and outcomes.

22 Patients and methods

23 Patients

24 Transplant eligible patients with biopsy proven Tr-iNHL were retrospectively
25 identified across three sites in Australia and United States diagnosed between
26 01/2000 to 01/2019. Cases were reviewed by hematopathologists at each academic
27 medical centre per routine clinical practice. Chronic lymphocytic leukaemia
28 (CLL/SLL), mantle cell lymphoma (MCL) as primary diseases and non-diffuse large
29 B-cell lymphoma (DLBCL) transformations were excluded. Composite disease was
30 defined as aggressive & indolent lymphoma diagnosed within the same tissue
31 sample. Sequential transformation was defined as a diagnosis of transformed
32 lymphoma following an antecedent diagnosis of indolent lymphoma. Patients who
33 received prior therapy for indolent lymphoma were excluded. Concordant bone
34 marrow involvement with Tr-iNHL was defined as $\geq 5\%$ involvement with large cell

lymphoma, and discordant bone marrow involvement defined as morphologic evidence of bone marrow involvement with indolent lymphoma only(9). Flow cytometry and immunohistochemistry studies were utilized to confirm a clonal B-cell population or aberrant immunophenotyped where there was unresolved suspicion of involvement by lymphoma. Criteria and methods for performance and interpretation of fluorescence *in situ* hybridization (FISH) for *MYC*, *BCL2* and *BCL6* rearrangements were per the policy of each centre.

Transplant eligibility was defined as: age <75 years; left ventricular ejection fraction (LVEF) $\geq 45\%$; no severe lung disease; standard dose rituximab, cyclophosphamide, doxorubicin, vincristine & prednisone (R-CHOP) chemotherapy or greater intensity front-line chemotherapy; complete remission (CR) by positron emission tomography (PET) or computerized tomography (CT) imaging ≥ 3 months after front-line chemotherapy. **Patients with clinical, radiological or histopathologic evidence of relapsed or refractory disease any time after completing front line chemotherapy were excluded. Patients receiving salvage chemotherapy due to lack of response or refractory disease were excluded. ASCT was performed as consolidation of complete remission achieved after front-line chemotherapy.** One Australian centre routinely employed HDC & ASCT in CR1 in transplant eligible patients, while two did not. Additionally, HDC & ASCT in CR1 was performed at the other two sites at the discretion of the treating physician for patients with high grade B-cell lymphoma with *MYC*, *BCL2* and/or *BCL6* rearrangements (HGBL-DH). This study was approved by the institutional review board of each participating centre.

Statistical Analysis

Landmark analysis was performed with time zero at 3 months after completion of front-line chemotherapy. Progression free survival (PFS) was defined as the interval from time zero to disease relapse, death from any cause or last follow-up. Overall survival (OS) was defined as the interval from time zero to death from any cause or last follow-up. Disease response by CT and/or PET was determined by the Revised Response Criteria for Malignant Lymphoma(10). The Chi-square test or Fisher's exact test was used to evaluate the association between two categorical variables. Wilcoxon rank sum test was used to evaluate the difference in a continuous variable

1 between patient groups. Propensity score matching (PSM) analysis using the
2 “greedy match” algorithm was used to match the baseline covariates to adjust for
3 potential selection bias(11). The Kaplan-Meier method was used for time-to-event
4 analysis including PFS and OS. **Univariable (UVA) and multivariable (MVA) Cox**
5 **proportional hazards models were fitted to the data after PSM to evaluate the**
6 **association between ASCT in CR1 and time-to-event outcomes.** Co-variables with
7 p values <0.2 from UVA were used to build the MVA model, and a backward model
8 selection method was applied. The analyses were performed using R version 3.6.3
9 (2020-02-29) and SAS 9.4 (SAS, Cary, NC).

12 **Results**

13 Patient Characteristics

14 319 transplant eligible patients were identified with the following underlying indolent
15 histologies: 283 (89%) patients with follicular lymphoma, 30 (9%) marginal zone
16 lymphoma, 5 (2%) mucosa-associated lymphoid tissue (MALT) lymphoma and 1
17 (0.3%) Waldenström macroglobulinaemia. 49 patients underwent HDC and ASCT in
18 CR1. A matched cohort of 98 patients based on age, stage, and HGBL-DH at
19 diagnosis was generated with a 1:2 ratio using PSM. The standardized differences of
20 less than 9% for age, sex, stage, ECOG PS and HGBL-DH suggested that these
21 variables were balanced between the two treatment groups after PSM. The
22 associations between ASCT in CR1 and other covariates were evaluated (Table 1).
23 Patient, disease and treatment characteristics were similar between patients treated
24 with ASCT in CR1 v **no ASCT in CR1**, with the following exceptions in the ASCT
25 cohort: 1) fewer patients with sequential transformations or having received
26 maintenance rituximab, 2) more patients with elevated LDH.

28 Outcomes

29 Reported outcomes and survival analysis were performed on the PSM cohort
30 (n=147). With a median follow-up of 3.7 (range 0.1-18.3) years (estimated from the
31 censored observations), the 4 year OS and PFS rates were 91% (95% CI: 0.86-0.97)
32 and 61% (95% CI: 0.527-0.706), respectively. **The median follow-up was similar in**
33 **patients receiving ASCT in CR1 (3.7 years; range 0.2-10.6) compared to**
34 **patients who did not receive ASCT in CR1 (3.5 years; range 0.1-18.3).**

By univariable analysis, no significant differences in 4-year PFS or OS rates were observed with FL v. non-FL indolent histology (PFS: 66% v. 61%; P=0.7; OS: 91% v. 90%; P=0.2), sequential v. non-sequential transformation (PFS: 75% v. 65%; P=0.4; OS: 91% v. 91%; P=0.9) or advanced [3 or 4] v. limited stage [1 or 2] disease (PFS: 66% v. 67%; P=0.4; OS: 90% v. 100%; P=0.7). The factors associated with inferior PFS and OS by UVA were HGBL-DH (PFS: HR 3.0, 95% CI 1.5-5.8, P=0.001; 4 year PFS 39% v. 70%; OS: HR 5.1, 95% CI 1.5-17.2, P=0.009; 4 year OS 79% v. 93%) and "high" IPI score [4 or 5] (PFS: HR 3.7, 95% CI 1.4-9.5, P=0.007; 4 year PFS 41% v. 75%; OS: HR 8.1, 95% CI 2.1-30.8, P=0.002; 4 year OS 60% v. 96%) (Table 2). The following factors were associated with inferior PFS without an impact on OS: B symptoms (PFS: HR 2.5, 95% CI 1.4-4.4, P=0.001; 4 year PFS 45% v. 75%; 4 year OS 89% v. 93%, P=0.5) and ≥ 2 extranodal sites (PFS: HR 2.1, 95% CI 1.2-3.7, P=0.006; 4 year PFS 53% v. 73%; 4-year OS 89% v. 92%, P=0.8). A trend toward inferior PFS was observed with concordant v. no BM involvement (HR 1.7, 95% CI 0.8-3.9, P=0.18; 4 year PFS 63% v. 71%), discordant v. no BM involvement (HR 1.8, 95% CI 1.0-3.2, P=0.05; 4 year PFS 62% v. 71%) and ECOG PS >1 v. ECOG PS 0 or 1 (HR 2.0, 95% CI 0.7-5.6, P=0.2; 4 year PFS 54% v. 67%). A trend towards improved survival outcomes was observed with composite lymphoma v. non-composite presentations (PFS: HR 0.6, 95% CI 0.3-1.1, P=0.1; 4 year PFS 69% v. 56%; OS: HR 0.4, 95% CI 0.1-1.3, P=0.1; 4 year OS 92% v. 87%), maintenance rituximab (PFS: HR 0.6, 95% CI 0.3-1.3, P=0.2; 4 year PFS 79% v. 63%; OS: HR 0.2, 95% CI 0.1-1.6, P=0.1; 4 year OS 100% v. 89%) and ASCT in CR1 (PFS: HR 0.6, 95% CI 0.3-1.0, P=0.07; 4 year PFS 77% v. 60%; OS: HR 2.4, 95% CI 0.9-6.4, P=0.09; 4 year OS 88% v. 93%) (Figure 1).

By MVA, ASCT in CR1 was independently associated with improved PFS (HR 0.5, 95% CI 0.27-0.98; P=0.043) (Table 3) with the adjustment of the other important covariates in the model. The factors associated with inferior PFS were HGBL-DH (HR 2.31, 95% CI 1.10-4.84; P=0.026) and B symptoms (HR 1.9, 95% CI 1.1-3.5; P=0.03). The overall p-value for the association between BM involvement v. not involved was 0.104 (discordant BM involvement v. not involved [HR 1.9, 95% CI 1.03-3.60; P=0.04]; concordant BM involvement v. not involved [HR 1.72, 95% CI

0.75-3.94; P=0.2]). Due to limited number of deaths, MVA was not performed for OS analysis.

Relapse & deaths

Fifty-one patients experienced lymphoma relapse with an estimated 4 year post-relapse survival rate of 68%. 16 (31%) patients experienced relapse with indolent lymphoma only (median PFS 1.4 years, 4-year post relapse OS 87%) and 35 (69%) with large cell or transformed lymphoma (median PFS 1.4 years, 4-year post relapse OS 74%, [Figure 2A](#)). Fewer relapses occurred among patients who underwent **ASCT in CR1 v. no ASCT in CR1** (22% v. 41%) due to fewer relapses with indolent lymphoma (4% v. 14%) [[Figure 3](#)]. 11 (22%) patients in the **ASCT in CR1 cohort** experienced relapse (median PFS 1.4 years, 4-year OS 73%). At last follow-up, 4 (36%) had died due to progressive disease (PD), 2 (18%) died due to secondary malignancy, 2 (18%) underwent CAR-T with ongoing CR, 3 (27%) continue on salvage therapy. 40 (41%) patients in the **no ASCT in CR1 cohort** experienced relapse (median PFS 1.2 years, 4-year OS 84%, [Figure 2B](#)) – at last follow-up, 4 (10%) died due to PD, 15 underwent HDC & ASCT with 7/15 [47%] ongoing CR; 10 underwent CAR-T therapy (5 relapse post ASCT, 4 refractory disease, 1 relapse post CR1) with 6/10 [60%] ongoing CR; 3 underwent allogeneic SCT (2 relapse post ASCT, 1 relapse post CR1) with 2/3 [67%] ongoing CR. The remaining patients experienced relapse with either iNHL or aNHL and continue on salvage therapy or investigational agents with ongoing response. There were 17 deaths across the entire cohort. The 4 year non-relapse mortality (NRM) rate for the entire cohort was 6%, with no significant difference between patients treated with **ASCT in CR1** (9%) v **no ASCT in CR1** (5%), P=0.5. Causes of NRM in the **ASCT in CR1 cohort** include – 1 (2%) due to sepsis, 1 (2%) Parkinson's disease, 2 (4%) secondary malignancy. Causes of NRM in the **no ASCT in CR1 cohort** include – 1 (1%) due to treatment-related toxicity, 1 (1%) secondary malignancy, 3 (3%) unknown causes unrelated to lymphoma.

Discussion

In this international, multi-centre retrospective analysis of untreated Tr-iNHL patients in the rituximab era, we observed improved initial duration of disease control with upfront **ASCT in CR1** largely due to fewer relapses with indolent disease. In

addition, NRM was not significantly increased in the **ASCT in CR1** cohort (9% v 5%, P=0.5) and there was no excess of secondary malignancies. Our findings are consistent with a retrospective analysis by the Canadian group of 105 patients with Tr-iNHL, 82% having received prior therapy of whom 50 patients proceeded with ASCT consolidation. Patients who underwent ASCT in the rituximab era demonstrated favourable outcomes with a 3-year PFS of 54% and OS of 69%(12). **Similar findings have been demonstrated in a single-centre, retrospective Norwegian study(13).** In a separate retrospective study, investigators at Washington University School of Medicine reported on 105 patients with transformed FL of whom 24 underwent ASCT as consolidation therapy. Although survival outcomes were superior in the ASCT v. non ASCT cohorts (5-year PFS rate: 42% v. 30%; 5-year OS rate: 74% v. 63%) this did not reach statistical significance(14). Of note, the ASCT cohort was heavily pretreated with 62% of patients receiving prior therapy for FL compared to 38% in the non ASCT cohort, possibly accounting for this difference. **A retrospective Danish study of 85 patients with Tr-iNHL of whom 54 received ASCT consolidation reported a PFS improvement in sequential transformations, but not composite/discordant Tr-iNHL. In this context, 76% of sequential transformations had received one or more therapies prior transformation thereby being more representative of a salvage ASCT rather than consolidation ASCT in CR1(15).** There were too few patients with sequential transformations who received ASCT in CR1 in our study for meaningful interpretation. By limiting our analysis to treatment naïve patients and using propensity score matching to reduce bias, our study provides robust evidence examining the impact of ASCT in CR1 in patients with Tr-iNHL. However, small numbers and a relatively short follow-up of **3.7 years** may limit the interpretation of these findings. With an estimated number needed to treat (NNT) of 6 to prevent a single progression event with upfront ASCT, the true survival benefit needs to be weighed carefully against the short- and long- term complications of high dose chemotherapy, financial costs and excellent survival outcomes at relapse, especially with indolent lymphoma.

The presence of *MYC* rearrangements in *de novo* DLBCL is associated with inferior survival outcomes(16, 17). In our study, HGBL-DH was identified in 16% of Tr-iNHL and was independently associated with inferior survival outcomes. A retrospective

1 study of 312 patients with newly diagnosed DLBCL of whom 63 had concurrent or
2 antecedent indolent lymphoma demonstrated similar findings of inferior survival
3 outcomes associated with HGBL-DH. Despite HGBL-DH demonstrating equivalently
4 poor outcomes in transformed and non-transformed DLBCL, a higher rate of HGBL-
5 DH was reported in the transformed v. non-transformed DLBCL cohorts (32% v.
6 11%)(3). This may be explained by the high prevalence of *BCL2* translocations in FL;
7 however, lack of systematic *MYC* FISH testing and selection bias associated with
8 retrospective analysis limits this finding, which and needs confirmation in a
9 prospective manner.

11 Discordant, not concordant bone marrow involvement with transformed lymphoma
12 was independently associated with inferior PFS in our study, but any interpretation is
13 limited by small numbers. Our findings are discrepant to a retrospective analysis of
14 795 patients of Tr-iNHL which demonstrated concordant bone marrow involvement
15 as an independent negative prognostic marker(18). More than half of patients who
16 presented with concordant bone marrow involvement progressed within the first year
17 with a dismal 3 year PFS of 37% and OS of 49%. Therefore, most patients
18 presenting with concordant BM involvement were unlikely to have achieved
19 adequate disease control or survived to be eligible for **ASCT in CR1**, thereby being
20 excluded from our study and explaining this discrepant finding.

22 In our study, approximately one third of all relapses occur with indolent disease and
23 the remainder with transformed lymphoma. Patients who underwent ASCT in CR1
24 experienced fewer relapses largely due to fewer indolent relapses. However,
25 patients in the **ASCT in CR1** cohort whose disease relapsed as large cell lymphoma
26 had inferior overall survival due to poorer responses to second line and later therapy.
27 This may be accounted for by geographical differences in the availability of
28 potentially curative CAR-T therapy which remains inaccessible in Australia resulting
29 in more progression-related deaths in the **ASCT in CR1** cohort. Therefore, the
30 observed clinical benefit of ASCT in CR1 is the prevention of indolent disease
31 relapse. Given the rapidly growing treatment options for indolent B cell lymphomas
32 including immunomodulatory agents (lenalidomide), monoclonal antibodies
33 (obinutuzumab), PI3K inhibitors (idelalisib, duvelisib, copanlisib) and other
34 investigational agents (tazemetostat, ibrutinib, venetoclax) the need for ASCT in CR1

is put to question(19-25). Sequencing ASCT as second line or later salvage therapy may spare patients exposure to a high-cost, potentially toxic therapy and associated long term sequelae. The question of whether ASCT may be potentially curative by preventing indolent disease relapse exists; however, longer follow up is required. Further studies are needed to identify whether a subset of patients may benefit from potentially curative, upfront ASCT in CR1 such as mutations in *TP53*, *CDKN2A/B*, *NOTCH1* and *MYC*(26-28).

Our study has limitations. We selected PSM to reduce treatment assignment bias by accounting for multiple baseline covariates by individual matching to patients in the ASCT in CR1 cohort. Despite the efforts made to reduce potential bias, we acknowledge that confounding may exist due to unmeasured patient factors such as co-morbidities and fitness, as well as pre-specified practices at each centre determining whether ASCT consolidation was administered. This limitation is inherent to the retrospective nature of our study and limitations of recorded data available which we have attempted to account for by determining pre-specified transplant eligibility criteria in the study inclusion criteria. Thirdly, although a trend towards inferior overall survival was demonstrated, this observation is limited by short follow-up, limited number of events and cross-over effect of patients receiving salvage ASCT at relapse.

Conclusion

Upfront ASCT in CR1 for patients with untreated Tr-iNHL in the rituximab era is associated with improved initial duration of disease control due to fewer relapses with indolent lymphoma. The lack of an overall survival benefit is likely due to highly effective salvage therapies which may achieve long term disease control. Further observational and translational studies are needed to identify which patient subgroups may benefit from potentially curative front-line therapy with ASCT in CR1.

Acknowledgements

1 The authors would like to acknowledge Mansoor Noorani (UT MD Anderson Cancer
2 Center) and Kerrie Stokes (Peter MacCallum Cancer Centre) for their contribution to
3 this study.

4
5 Portions of this article were presented at the 2020 ASCO Annual Meeting, May 29 –
6 31, 2020, Chicago, Illinois, United States.

7 8 **Authorship Contributions**

9 *CKC & MD*: Designed the study.

10 *CKC, KJL, KLL, PJ*: Collected and assembled the data.

11 *CKC, YQ & LF*: Analysed and interpreted the data.

12 All authors: Wrote the manuscript.

13 All authors gave final approval for the manuscript.

14 *CYC, JFS, DR, KB, CST, NF, LEF, JW, SSN, FBH, FS, LN, MD*: Provided study
15 materials and patients.

16 17 **Funding**

18 This work was supported in part by the Cancer Center Support Grant (CCSG) at UT
19 MD Anderson Cancer Center.

20 21 **Disclosures**

22 *CKC, KJL, PJ, YQ, LF, KB, LEF, FS, LEF, FBH*: No conflicts of interest.

23 *KLL*: honoraria: Roche; travel expenses: Janssen, Novartis.

24 *CYC*: consulting/advisory/honoraria – Roche, Janssen, MSD, Gilead, Ascentage
25 Pharma, Acerta, Loxo Oncology, TG therapeutics; research funding – Celgene,
26 Roche, Abbvie; travel expenses – Roche.

27 *JFS*: received research support from AbbVie, Celgene, Janssen and Roche; served
28 as consultant and advisory board member for AbbVie, Acerta Pharma, Janssen,
29 Roche, Sunesis Pharmaceuticals and Takeda; Speaker's Bureau for AbbVie,
30 Celgene and Roche; received Honoraria from AbbVie, Acerta Pharma, Janssen,
31 Roche, Sunesis Pharmaceuticals and Takeda; received travel expenses from
32 AbbVie and Roche.

DR: received research support from Amgen, Bristol-Myers Squibb, Takeda, Beigene and Imago. Served as consultant and advisory board member for Amgen and Pfizer. Received honoraria from Amgen, Novartis and Sanofi.

CT: received research support from Janssen-Cilag and AbbVie; served as consultant and advisory board member for Janseen, Loxo, Roche, BeiGene and AbbVie; honoraria from Janssen-Cilag, AbbVie, Novartis, Beigene and Pharmacyclics.

NF: received research support from Celgene, Roche, Janssen, TG Therapeutics and AbbVie; served as consultant and advisory board member for Celgene, Roche, Janssen, TG therapeutics and AbbVie.

JW: has received research support from Novartis, Celgene, Janssen, Kite/Gilead, Unum, Genentech, Curis and 47 Inc; served as consultant and advisory board member for Novartis, Celgene, Juno, Janssen, Kite/Gilead, MorphoSys, Genentech and Curis.

SSN: has received research support from Kite/Gilead, Collectis, Poseida, Merck, Acerta, Karus, BMS, Unum Therapeutics, Allogene, and Precision Biosciences; served as consultant and advisory board member for Kite/Gilead, Celgene, Novartis, Unum Therapeutics, Pfizer, Merck, Precision Biosciences, Cell Medica, Incyte, Allogene, Calibr, and Legend Biotech; has patents related to cell therapy.

CF: has received research support from AbbVie, Acerta Pharma, Celgene, Gilead Sciences, Infinity Pharmaceuticals, Janssen Pharmaceuticals, Millenium/Takeda, National Institutes of Health, Onyx Pharmaceuticals, Pharmacyclics and Spectrum Pharmaceuticals; served as consultant and advisory board member for Spectrum Pharmaceuticals, Celgene, Optum Rx, Seattle Genetics, Kite/Gilead and Bayer.

LN: has received research support from TG Therapeutics, Janssen, Genentech and Celgene; has served as consultant and advisory board member for TG Therapeutics, Novartis, Janssen, Spectrum Pharmaceuticals, Kite/Gilead, Genentech, Bayer and Celgene.

MD: has received research support from Celgene, GSK, Takeda and Novartis; has served as consultant or advisory board member for Roche, GSK, Janssen, Takeda Celgene, Gilead Sciences and Novartis; has served as a speaker for Roche, Novartis, Janssen, Takeda and Gilead Sciences.

	No ASCT (n=98)	ASCT (n=49)	Total (n=147)	P-value
Centre				

- Peter MacCallum Cancer Centre	0	39 (80%)	39 (27%)	-
- Sir Charles Gairdner Hospital	3 (3%)	1 (2%)	4 (3%)	
- UT MD Anderson Cancer Center	95 (97%)	9 (18%)	104 (71%)	
*Age (range)	56 (26-74)	56 (29-74)	56 (26-74)	0.6
*Sex				
- Male	57 (58%)	30 (61%)	87 (59%)	0.9
- Female	41 (42%)	19 (39%)	60 (41%)	
*Stage (Ann arbor)				
- 1 or 2	9 (9%)	5 (10%)	14 (10%)	1.0
- 3 or 4	89 (91%)	44 (90%)	133 (90%)	
*HGBL-DH				
- Yes	15 (15%)	9 (18%)	24 (16%)	0.6
- No / not tested	83 (85%)	40 (82%)	123 (84%)	
*ECOG PS				
- 0 or 1	90 (95%)	47 (96%)	137 (95%)	1.0
- ≥2	5 (5%)	2 (4%)	7 (5%)	
Indolent lymphoma subtype				
- Follicular lymphoma	91 (93%)	42 (86%)	133 (90%)	1.0
- Marginal zone lymphoma	5 (5%)	3 (6%)	8 (5%)	
- Waldenström Macroglobulinaemia	0	1 (2%)	1 (1%)	
- MALT lymphoma	2 (2%)	0	2 (1%)	
Composite lymphoma				
- Yes	79 (81%)	34 (69%)	113 (77%)	0.2
- No	19 (19%)	15 (31%)	34 (23%)	
Timing of transformation				
- At presentation	84 (86%)	48 (98%)	132 (90%)	0.02
- Sequential	14 (14%)	1 (2%)	15 (10%)	
LDH				
- Normal	53 (67%)	18 (38%)	71 (56%)	0.002
- Elevated	26 (33%)	30 (62%)	56 (44%)	
- Not recorded	19	1	20	
Extranodal sites				
- 0 or 1 sites	65 (68%)	31 (63%)	96 (66%)	0.7
- ≥2 sites	31 (32%)	18 (37%)	49 (34%)	
- Not recorded	2	0	2	
IPI				
- IPI 0 or 1	21 (27%)	12 (25%)	33 (26%)	0.9
- IPI 2 or 3	51 (66%)	32 (67%)	83 (66%)	
- IPI 4 or 5	5 (6%)	4 (8%)	9 (7%)	

- Not recorded	21	1	22	
B symptoms				
- Yes	26 (27%)	12 (24%)	38 (26%)	0.8
- No	69 (73%)	37 (76%)	106 (74%)	
- Not recorded	3	0	3	
Bulk (≥10cm)				
- Yes	12 (13%)	12 (24%)	24 (17%)	0.1
- No	81 (87%)	37 (76%)	118 (83%)	
- Not recorded	5	0	5	
Type of BM involvement				
- Discordant	23 (25%)	20 (42%)	43 (31%)	0.6
- Concordant	13 (14%)	8 (16%)	21 (15%)	
- Not involved	55 (60%)	20 (42%)	75 (54%)	
- Not assessed	7	1	8	
Front line chemotherapy				
- R-CHOP	74 (76%)	36 (73%)	110 (75%)	0.6
- Da-EPOCH-R	20 (20%)	10 (20%)	30 (20%)	
- RHCVD	2 (2%)	3 (6%)	3 (6%)	
- OCHOP + Lenalidomide	2 (2%)	0	0	
Conditioning chemotherapy				
- BEAM	-	10 (20%)	10 (20%)	-
- Cyclophosphamide, carmustine, Etoposide	-	36 (73%)	36 (73%)	
- Busulfan, Melphalan	-	1 (2%)	1 (2%)	
- Busulfan, Melphalan, Gemcitabine, SAHA	-	2 (4%)	2 (4%)	
Maintenance rituximab				
- No	77 (79%)	45 (92%)	122 (83%)	0.06
- Yes	21 (21%)	4 (8%)	25 (17%)	
Disease status at first relapse or last follow-up				
- Indolent lymphoma	14 (14%)	2 (4%)	16 (11%)	0.5
- Large cell/transformed lymphoma	26 (27%)	9 (18%)	35 (24%)	
- Remission	58 (59%)	38 (78%)	96 (65%)	

Table 1. Baseline clinical characteristics by ASCT post 1:2 ratio PSM.

(*) 1:2 ratio PSM performed using based on age, Ann-arbor stage, HGBL-DH status.

PSM – Propensity score matching; ASCT – Autologous stem cell transplantation; HGBL-DH – High grade B-cell lymphoma with MYC, BCL2 and/or BCL6 rearrangements; ECOG – Eastern Cooperative Oncology Group; PS – performance status; WM – Waldenström macroglobulinaemia; MALT – mucosa-associated lymphoid tissue; COM – composite lymphoma; DIS – discordant lymphoma; SEQ – sequential lymphoma; LDH – lactate dehydrogenase; IPI – International Prognostic Index; BM – bone marrow, aNHL – aggressive non-Hodgkin lymphoma; RCHOP – Rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone; DA-EPOCH-R – Dose adjusted etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin, rituximab; RHCVD – rituximab, hyperfractionated cyclophosphamide, vincristine, doxorubicin, dexamethasone alternating with methotrexate & cytarabine; O-CHOP – Obinutuzumab, cyclophosphamide, doxorubicin, vincristine, prednisone; BEAM – carmustine, etoposide, cytarabine, melphalan; SAHA – suberoylanilide hydroxamic acid.

	PFS analysis			OS analysis		
	HR	95% CI	P value	HR	95% CI	P value
Age > 60 years	1.1	0.6-2.0	0.8	1.1	0.4-3.2	0.8
Male gender	1.0	0.6-1.8	0.9	0.9	0.4-2.5	0.9
Composite v. non-composite transformation	0.6	0.3-1.1	0.1	0.4	0.1-1.3	0.1
Sequential v. non-sequential transformation	0.6	0.2-1.9	0.4	0.9	0.1-7.0	0.9
Indolent lymphoma histology – FL v non-FL histology	1.2	0.5-2.9	0.7	2.1	0.6-7.6	0.2
Aggressive lymphoma histology – HGBL-DH v non-HGBL-DH/not tested	3.0	1.5-5.8	0.001	5.1	1.5-17.2	0.009
IPI “high” (4/5) v IPI “low-intermediate” (0-3)	3.7	1.4-9.5	0.007	8.1	2.1-30.8	0.002
Advanced stage (3/4) v limited stage (1/2)	1.5	0.6-4.3	0.4	1.5	0.2-11.6	0.7

ECOG PS ≥ 2	2.0	0.7-5.6	0.2	3.9	0.9-17.4	0.08
Elevated LDH	1.2	0.6-2.3	0.6	2.1	0.7-6.7	0.2
Extranodal sites ≥ 2	2.1	1.2-3.7	0.006	1.1	0.4-3.0	0.8
Presence of B symptoms	2.5	1.4-4.4	0.001	1.5	0.5-4.3	0.5
Bulk ≥ 10 cm	0.7	0.3-1.7	0.5	0.7	0.2-3.1	0.6
BM involvement v. not involved (overall)	-	-	0.12	-	-	0.7
- Concordant BM involvement v not involved	1.9	0.8-4.1	0.14	1.7	0.5-6.5	0.4
- Discordant BM involvement v not involved	1.7	1.0-3.2	0.07	1.2	0.4-3.4	0.8
Upfront chemotherapy with R-CHOP v DA-EPOCH-R	0.7	0.3-1.6	0.5	0.6	0.1-4.4	0.6
Maintenance Rituximab v no maintenance	0.6	0.3-1.3	0.2	0.2	0.02-1.41	0.1
ASCT in CR1 v no ASCT	0.6	0.3-1.0	0.07	2.4	0.9-6.4	0.09

Table 2. Univariable analysis of patient, disease & treatment factors on PFS and OS.

PFS – progression free survival; OS – overall survival; HR – hazard ratio; CI – confidence interval; FL – follicular lymphoma; HGBL-DH – high grade B cell lymphoma with MYC, BCL2 and/or BCL6 rearrangements; IPI – International Prognostic Index; ECOG – Eastern Cooperative Oncology Group; PS – performance status; LDH – lactate dehydrogenase; RCHOP – Rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone; DA-EPOCH-R – Dose adjusted etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin, rituximab; ASCT – autologous stem cell transplantation; CR1 – first remission.

	PFS		
	HR	95% CI	P value
<u>ASCT in CR1</u> v. No ASCT	0.51	0.27-0.98	0.043
<u>HGBL-DH</u> v. non-HGBL-DH or not tested	2.31	1.10-4.84	0.026
<u>B symptoms</u> v. no B symptoms	1.92	1.05-3.50	0.034
<u>Concordant BM involvement</u> v. not involved	1.72	0.75-3.94	0.200
<u>Discordant BM involvement</u> v. not involved	1.93	1.03-3.60	0.040

Table 3. Multivariable analysis of patient, disease & treatment factors on PFS.

PFS – progression free survival; HR – hazard ratio; CI – confidence interval; HGBL-DH - high grade B cell lymphoma with MYC, BCL2 and/or BCL6 rearrangements; IPI – International Prognostic Index; ASCT – autologous stem cell transplantation; CR1 – first remission.

(A)

Number at risk						
ASCT in CR1	49	24	7	0	0	0
No ASCT in CR1	98	38	16	5	1	0

Number censored						
ASCT in CR1	0	15	29	36	36	36
No ASCT in CR1	0	25	42	52	55	56

----- No ASCT in CR1 — ASCT in CR1

(B)

Number at risk						
ASCT in CR1	49	24	5	0	0	0
No ASCT in CR1	98	45	21	7	2	0

Number censored						
ASCT in CR1	0	20	36	41	41	41
No ASCT in CR1	0	48	70	83	87	89

----- No ASCT in CR1 — ASCT in CR1

Figure 1. (A) PFS based on ASCT in CR1 v no ASCT; (B) OS based on ASCT in CR1 v no ASCT.

ASCT – autologous stem cell transplantation; CR1 – first remission; PFS – progression free survival; OS – overall survival.

Figure 2. (A) Post-relapse OS depending on histologic grade at first relapse; (B) Post-relapse OS of ASCT in CR1 v no ASCT.

ASCT – autologous stem cell transplantation; CR1 – first remission; OS – overall survival; PFS – progression free survival.

ASCT in CR1

Remission
(n=38, 78%)

Aggressive relapse
Indolent relapse

No ASCT

Remission
(n=58, 59%)

Figure 3. Patterns of relapse, death and most recent line of therapy at last follow-up stratified by ASCT in CR1 v. no ASCT.

ASCT – Autologous stem cell transplantation; CR1 – first remission; CAR-T – chimaeric antigen receptor T-cell therapy; CR – complete remission; PR – partial remission; SD – stable disease; PD – progressive disease; HDC – high dose chemotherapy; ST – salvage therapy; rel. – relapse.

References

1. Reddy N, Oluwole O, Greer JP, Goodman S, Engelhardt B, Jagasia MH, et al. Superior long-term outcome of patients with early transformation of non-Hodgkin lymphoma undergoing stem cell transplantation. *Clin Lymphoma Myeloma Leuk*. 2012;12(6):406-11.
2. Wang Y, Link BK, Witzig TE, Maurer MJ, Allmer C, King RL, et al. Impact of concurrent indolent lymphoma on the clinical outcome of newly diagnosed diffuse large B-cell lymphoma. *Blood*. 2019;134(16):1289-97.
3. Behdad A, Boddy CS, Fought AJ, Taxter T, Falkiewicz MK, Ayers E, et al. Survival outcomes of diffuse large B-cell lymphoma by association with concurrent or antecedent follicular lymphoma and double hit status. *Leuk Lymphoma*. 2019;60(13):3266-71.
4. Wagner-Johnston ND, Link BK, Byrtek M, Dawson KL, Hainsworth J, Flowers CR, et al. Outcomes of transformed follicular lymphoma in the modern era: a report from the National LymphoCare Study (NLCS). *Blood*. 2015;126(7):851-7.
5. Magnano L, Balague O, Dlouhy I, Rovira J, Karube K, Pinyol M, et al. Clinicobiological features and prognostic impact of diffuse large B-cell lymphoma component in the outcome of patients with previously untreated follicular lymphoma. *Ann Oncol*. 2017;28(11):2799-805.
6. Mendez M, Torrente M, Sanchez-Beato M, Gonzalez-Rincon J, Royuela A, Gomez-Codina J, et al. Transformed follicular lymphoma in the rituximab era: A report from the Spanish Lymphoma Oncology Group. *Hematol Oncol*. 2019;37(2):143-50.
7. Villa D, George A, Seymour JF, Toze CL, Crump M, Lee C, et al. Favorable outcomes from allogeneic and autologous stem cell transplantation for patients with transformed nonfollicular indolent lymphoma. *Biol Blood Marrow Transplant*. 2014;20(11):1813-8.
8. Ghesquieres H, Berger F, Felman P, Callet-Bauchu E, Bryon PA, Traverse-Glehen A, et al. Clinicopathologic characteristics and outcome of diffuse large B-cell lymphomas presenting with an associated low-grade component at diagnosis. *J Clin Oncol*. 2006;24(33):5234-41.
9. Campbell J, Seymour JF, Matthews J, Wolf M, Stone J, Juneja S. The prognostic impact of bone marrow involvement in patients with diffuse large cell

lymphoma varies according to the degree of infiltration and presence of discordant marrow involvement. *Eur J Haematol*. 2006;76(6):473-80.

10. Cheson BD, Pfistner B, Juweid ME, Gascoyne RD, Specht L, Horning SJ, et al. Revised response criteria for malignant lymphoma. *J Clin Oncol*. 2007;25(5):579-86.
11. Austin PC. A comparison of 12 algorithms for matching on the propensity score. *Stat Med*. 2014;33(6):1057-69.
12. Villa D, Crump M, Keating A, Panzarella T, Feng B, Kuruville J. Outcome of patients with transformed indolent non-Hodgkin lymphoma referred for autologous stem-cell transplantation. *Ann Oncol*. 2013;24(6):1603-9.
13. Blaker YN, Eide MB, Liestol K, Lauritzsen GF, Kolstad A, Fossa A, et al. High dose chemotherapy with autologous stem cell transplant for patients with transformed B-cell non-Hodgkin lymphoma in the rituximab era. *Leuk Lymphoma*. 2014;55(10):2319-27.
14. Jackson A, Liu J, Bartlett NL, Cashen A. Transformed Follicular Lymphoma: Is Early Autologous Stem Cell Transplantation Indicated in the Modern Era? *Ann Hematol Oncol*. 2014;1(1):1002.
15. Madsen C, Pedersen MB, Vase MO, Bendix K, Moller MB, Johansen P, et al. Outcome determinants for transformed indolent lymphomas treated with or without autologous stem-cell transplantation. *Ann Oncol*. 2015;26(2):393-9.
16. Rosenwald A, Sehn LH, Maucourt-Boulch D. Prognostic Significance of *MYC* Single, Double, Triple Hit and *MYC*-Translocation Partner Status in Diffuse Large B-Cell Lymphoma - a Study By the Lunenburg Lymphoma Biomarker Consortium (LLBC). *Blood*. 2018;132:344-.
17. Sewastianik T, Prochorec-Sobieszek M, Chapuy B, Juszczynski P. *MYC* deregulation in lymphoid tumors: molecular mechanisms, clinical consequences and therapeutic implications. *Biochim Biophys Acta*. 2014;1846(2):457-67.
18. Sehn LH, Scott DW, Chhanabhai M, Berry B, Ruskova A, Berkahn L, et al. Impact of concordant and discordant bone marrow involvement on outcome in diffuse large B-cell lymphoma treated with R-CHOP. *J Clin Oncol*. 2011;29(11):1452-7.
19. Wang M, Fowler N, Wagner-Bartak N, Feng L, Romaguera J, Neelapu SS, et al. Oral lenalidomide with rituximab in relapsed or refractory diffuse large cell,

follicular and transformed lymphoma: a phase II clinical trial. *Leukemia*.

2013;27(9):1902-9.

20. Dreyling M, Santoro A, Mollica L, Leppa S, Follows G, Lenz G, et al. Long-term safety and efficacy of the PI3K inhibitor copanlisib in patients with relapsed or refractory indolent lymphoma: 2-year follow-up of the CHRONOS-1 study. *Am J Hematol*. 2019.

21. Gopal AK, Kahl BS, Flowers CR, Martin P, Ansell SM, Abella-Dominicis E, et al. Idelalisib is effective in patients with high-risk follicular lymphoma and early relapse after initial chemoimmunotherapy. *Blood*. 2017;129(22):3037-9.

22. Salles GA, Morschhauser F, Solal-Celigny P, Thieblemont C, Lamy T, Tilly H, et al. Obinutuzumab (GA101) in patients with relapsed/refractory indolent non-Hodgkin lymphoma: results from the phase II GAUGUIN study. *J Clin Oncol*. 2013;31(23):2920-6.

23. Fowler N, Nastoupil L, Chin CK, Strati P, Hagemeister F, Fayad L, et al. A Phase I/II Study of Lenalidomide Plus Obinutuzumab in Relapsed Indolent Lymphoma. American Society of Hematology Annual Meeting; Orlando, FL: *Blood*; 2019.

24. Morschhauser F, Tilly H, Chaidos A, Phillips T, Ribrag V, Campbell P, et al. Phase 2 Multicenter Study of Tazemetostat, an EZH2 Inhibitor, in Patients with Relapsed or Refractory Follicular Lymphoma. American Society of Hematology Annual Meeting; Orlando, FL: *Blood*; 2019.

25. Cheah CY, Fowler NH. Novel agents for relapsed and refractory follicular lymphoma. *Best Pract Res Clin Haematol*. 2018;31(1):41-8.

26. Lo Coco F, Gaidano G, Louie DC, Offit K, Chaganti RS, Dalla-Favera R. p53 mutations are associated with histologic transformation of follicular lymphoma. *Blood*. 1993;82(8):2289-95.

27. Pasqualucci L, Khiabani H, Fangazio M, Vasishtha M, Messina M, Holmes AB, et al. Genetics of follicular lymphoma transformation. *Cell Rep*. 2014;6(1):130-40.

28. Fitzgibbon J, Iqbal S, Davies A, O'Shea D, Carlotti E, Chaplin T, et al. Genome-wide detection of recurring sites of uniparental disomy in follicular and transformed follicular lymphoma. *Leukemia*. 2007;21(7):1514-20.

Number at risk						
ASCT in CR1	49	24	7	0	0	0
No ASCT in CR1	98	55	16	5	1	0

Number censored						
ASCT in CR1	0	15	29	36	36	36
No ASCT in CR1	0	28	42	52	55	55

----- No ASCT in CR1 — ASCT in CR1

Number at risk						
ASCT in CR1	49	24	5	0	0	0
No ASCT in CR1	98	45	21	7	2	0

Number censored						
ASCT in CR1	0	20	36	41	41	41
No ASCT in CR1	0	48	70	83	87	89

----- No ASCT in CR1 — ASCT in CR1

bjh_17072_f1.png

bjh_17072_f2.png

bjh_17072_f3.png

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Chin, CK;Lim, KJ;Lewis, K;Jain, P;Qing, Y;Feng, L;Cheah, CY;Seymour, JF;Ritchie, D;Burbury, K;Tam, CS;Fowler, NH;Fayad, LE;Westin, JR;Neelapu, SS;Hagemeister, FB;Samaniego, F;Flowers, CR;Nastoupil, LJ;Dickinson, MJ

Title:

Autologous stem cell transplantation for untreated transformed indolent B-cell lymphoma in first remission: an international, multi-centre propensity-score-matched study

Date:

2020-10-16

Citation:

Chin, C. K., Lim, K. J., Lewis, K., Jain, P., Qing, Y., Feng, L., Cheah, C. Y., Seymour, J. F., Ritchie, D., Burbury, K., Tam, C. S., Fowler, N. H., Fayad, L. E., Westin, J. R., Neelapu, S. S., Hagemeister, F. B., Samaniego, F., Flowers, C. R., Nastoupil, L. J. & Dickinson, M. J. (2020). Autologous stem cell transplantation for untreated transformed indolent B-cell lymphoma in first remission: an international, multi-centre propensity-score-matched study. [Abstract]. BRITISH JOURNAL OF HAEMATOLOGY, 191 (5), pp.806-815. <https://doi.org/10.1111/bjh.17072>.

Persistent Link:

<http://hdl.handle.net/11343/276451>