DR. OFER MICHAEL GONEN (Orcid ID : 0000-0002-3833-9128)

Article type : Full length original research paper

7T quantitative magnetic resonance spectroscopy of glutamate, GABA and glutathione in the PCC/precuneus in patients with epilepsy

Ofer M. Gonen^{1,2,3}, Bradford A. Moffat², Patricia M. Desmond^{2,5}, Elaine Lui^{2,5}, Patrick Kwan^{1,2,3,4}, Terence J. O'Brien^{1,2,3,4}

1. Department of Neurology, The Royal Melbourne Hospital, VIC, Australia. 2. The Department of Medicine & Radiology, The University of Melbourne, VIC, Australia. 3. The Department of Neurology, The Alfred Hospital, VIC, Australia. 4. The Department of Neuroscience, Central Clinical School, Monash University, VIC, Australia. 5. The Department of Radiology, The Royal Melbourne Hospital, VIC, Australia

Corresponding Author:

Name: Dr. Ofer M. Gonen

Address: Neurology Department, The Royal Melbourne Hospital
300 Grattan St, Parkville 3050, VIC, Australia

Telephone: (+61)-3-93427722

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/EPI.16731](https://doi.org/10.1111/EPI.16731)

This article is protected by copyright. All rights reserved

Fax: (+61)-3-93428628

E-mail: Ofer.Gonen@mh.org.au

17 **ORCID Numbers:**

Ofer M. Gonen 0000-0002-3833-9128

Terence J. O'Brien 0000-0002-7198-8621

18 **Keywords:** Default Mode Network; epilepsy; MRS; precuneus; posterior cingulate cortex; glutathione.

19 **Word Count (abstract):** 290; **Word Count (excluding abstract, acknowledgements / statements, references, tables, figures):** 4061;

20 **References:** 49; **Figures:** 3; **Tables:** 3; **Supplementary Tables:** 5; **Total pages:** 37

21 **Abstract**

22
23 **Objective:** The posterior cingulate cortex (PCC)/precuneus is a key hub of the default mode network, whose function is known to be altered in
24 epilepsy. Glutamate and γ -amino butyric acid (GABA) are the main excitatory and inhibitory neurotransmitters in the CNS, respectively.
25 Glutathione (GSH) is the most important free radical scavenging compound in the brain. Quantification of these molecules by magnetic resonance
26 spectroscopy (MRS) up to 4T is limited by overlapping resonances from other molecules. In this study we used ultra-high-field (7T) MRS to
27 quantify their concentrations in patients with different epilepsy syndromes.

28 **Methods:** 19 patients with temporal lobe epilepsy (TLE) and 16 with idiopathic generalized epilepsy (IGE) underwent MRI scans using a 7T
29 research scanner. Single-voxel (8 cm³) MRS, located in the PCC/precuneus, was acquired via Stimulated Echo Acquisition Mode (STEAM). Their
30 results were compared to 10 healthy volunteers.

Results: Mean concentrations of glutamate, GABA, and the glutamate/GABA ratio did not differ between the IGE, TLE and healthy volunteer groups. The mean($\pm$ SD) concentration of GSH was 1.9($\pm$ 0.3) mM in healthy controls, 2.0($\pm$ 0.2) mM in patients with TLE, and 2.2($\pm$ 0.4) mM in patients with IGE. One-way ANOVA with post-hoc Tukey-Kramer test revealed a significant difference in the concentration of GSH between patients with IGE and controls ($P=0.03$). Short-term seizure-freedom in patients with epilepsy was predicted by an elevated concentration of glutamate in the PCC/precuneus ($P=0.01$). In patients with TLE the concentration of GABA declined with age ($P=0.03$).

Significance: Patients with IGE have higher concentrations of GSH in the PCC/precuneus than healthy controls. There is no difference in the concentrations of glutamate, GABA, or their ratio in the PCC/precuneus between patients with IGE, patients with TLE, and healthy controls. Measuring the concentration of glutamate in the PCC/precuneus may assist with predicting drug-response.

Key Points

- The concentrations of glutamate and GABA in the PCC/precuneus are similar in patients with epilepsy and healthy controls.
- The concentration of glutathione in the PCC/precuneus is higher in patients with idiopathic generalized epilepsy in comparison to healthy controls.
- The concentration of GABA in the PCC/precuneus decreases with age in temporal lobe epilepsy patients but not in those with idiopathic generalized epilepsy.
- Elevated concentration of glutamate in the PCC/precuneus may predict short-term seizure-freedom in patients with epilepsy.

51

52 1. INTRODUCTION

53

54 Glutamate and γ -amino butyric acid (GABA) are the major excitatory and inhibitory neurotransmitters in the human brain, respectively.^{1,2} Their
55 concentration in the brain is affected by various antiseizure medications (ASMs) and it is believed that their ratio may reflect the propensity of the
56 brain for hyperexcitable conditions such as epilepsy.³ Glutathione (GSH), the main free radical scavenger compound in the brain, is metabolically
57 related to glutamate and GABA and is involved in a wide range of neurological disorders.⁴ Previous studies revealed a reduction in the serum level
58 of GSH in patients taking certain ASMs and the activity of Glutathione Peroxidase in erythrocytes was found to be significantly lower in patients
59 with idiopathic generalized epilepsy (IGE) when compared to a control group.^{5,6}

60 Several studies have used magnetic resonance spectroscopy (MRS) to measure the concentrations of these metabolites in various brain regions in
61 patients with epilepsy.⁷⁻¹¹ However, the accuracy of MRS has been limited by overlapping resonances from other molecules at magnetic fields up
62 to 4T. In comparison, 7T MRI has enabled the acquisition of spectra with superior signal to noise ratios and chemical shift dispersion, thus
63 improving the quantification of these metabolites.¹²

64 Indeed, in recent years several research groups have started using 7T MRS to study epilepsy. For example, GABA concentration was measured in
65 the thalamus using 7T MRS and was found to be higher in two patients with well-controlled epilepsy as compared to six patients with poorly-
66 controlled epilepsy.¹³ In another study, levels of glutamate were acquired using a chemical exchange saturation transfer (CEST) technique at 7T
67 in patients with non-lesional temporal lobe epilepsy (TLE) and found to correctly lateralize the seizure focus.¹⁴ Our own group quantified glutamate
68 concentrations at 7T in and around gliomas using both CEST and stimulated echo acquisition mode (STEAM) and demonstrated positive
69 correlation between its level and parameters of tumor aggressiveness and epileptogenesis.¹⁵

The posterior cingulate cortex (PCC)/precuneus is the structural and functional core of the default mode network (DMN), the main resting-state network of the brain, which is known to be involved in many neuropsychiatric disorders including epilepsy.^{16,17} However, MRS studies of the DMN in patients with epilepsy are scarce and limited by sample size and magnetic field strength. For example, a multi-voxel study at 3T demonstrated elevation of glutamate in various cortical and sub-cortical regions in four patients with IGE. No such changes were discovered in the cingulate cortex or the precuneus.¹⁸ Notably, GABA and GSH were not measured in that study and we are unaware of any study measuring GSH concentrations via MRS in any brain region in patients with IGE. GSH was measured via MRS in patients with focal epilepsy and reductions in its concentration were measured in both hemispheres.¹¹

The purpose of this study was to measure concentrations of glutamate, GABA, and GSH in the PCC/precuneus in patients with focal and generalized epilepsy at 7T and to compare these results with healthy controls. The clinical usefulness of finding distinct metabolic profiles in different groups derives from the potential application of such findings both for diagnostic purposes and for their use as biomarkers in epilepsy management, if future studies find changes in such profiles in response to therapy. As DMN alterations have been previously demonstrated with resting-state fMRI in both focal and generalized epilepsy, we expected such changes to be reflected by altered metabolite levels in both patient groups.^{19–23} Our hypotheses were that both patient groups will demonstrate increased glutamate, reduced GABA, increased glutamate to GABA ratio and reduced GSH in comparison with healthy controls.

2. METHODS

2.1 Participants

Patients were recruited from the Comprehensive Epilepsy Programs of the Royal Melbourne and Alfred Hospitals in Melbourne, Australia. TLE and IGE were defined according to well-established ILAE criteria.²⁴ They were excluded if their epilepsy was related to a known brain lesion (other than mesial temporal sclerosis) or if they had contraindications for 7T MRI.

Nineteen right-handed patients (12 female) with temporal lobe epilepsy (TLE), ranging in age from 19-72 years (mean 41.9 ± 13.9 ; median 42), and 16 right-handed patients (12 female) with IGE, ranging in age from 20-70 years (mean 39.2 ± 15.3 ; median 33.5), were recruited. Of the TLE patients, nine had left-sided TLE, six had right-sided TLE, and four had bilateral seizure foci. The patients were matched with 10 healthy, right-handed, volunteers (six female). Their mean($\pm$ SD) age was 40.0 ± 14.4 years (median 36.5). Mean($\pm$ SD) duration of epilepsy was 18.3 ± 14.7 years in the TLE group and 24.6 ± 18.4 years in the IGE group.

The age of participants was similar across the three groups (one-way ANOVA - $F(2,41)=0.13$; $P=0.88$) and there was no statistically significant difference of epilepsy duration between the TLE and IGE groups (Kruskal-Wallis H test - $\chi^2(1)=1.26$; $P=0.26$). The number of medications was also similar between the patient groups ($P=0.17$). Characteristics of all 35 recruited patients are provided in the supplementary material.

Seizure occurrence in the preceding 3 months was ascertained as a measure of drug-responsiveness to minimize recall bias. In patients who had started drug treatment for less than 3 months before the scan drug-responsiveness was verified three months later. Accordingly, short-term seizure freedom was defined as lack of self-reported seizures in the 3 months preceding the scan (other than seizures induced by medication withdrawal as part of video-EEG monitoring). Short-term seizure-freedom was reported by 11/19 patients with TLE (10/18 of the patients in the final analysis) and by 6/16 patients with IGE. Information about drug-treatment was collected from all patients. The study was approved by the Human Research Ethics Committee of Melbourne Health. All participants provided written informed consent.

2.2 Sample size estimation

107 Sample size estimation was limited by the lack of similar 7T studies of the aforementioned metabolites in patients with epilepsy. Nevertheless,
108 considering the study of Pan et al, who measured the concentration of GABA scaled to N-acetyl aspartate (NAA) in the thalamus of healthy
109 controls, patients with well controlled IGE, and patients with poorly controlled IGE, the smallest effect size existed between the healthy controls
110 (0.053 ± 0.012 ; $n=8$) and the poorly controlled group (0.038 ± 0.009 ; $n=6$), amounting to Cohen's $f=0.69$.¹³ Assuming $\alpha=0.05$ and $\beta=0.80$, a three
111 group ANOVA study looking for a similar effect size would need a total sample size of 24 participants (calculated by G*Power version 3.1.9.2).

112 **2.3 Imaging protocol**

113

114 A 7T research scanner (Siemens Healthcare, Erlangen, Germany) with a 32-channel head-coil (Nova Medical Inc, Wilmington MA, USA) was
115 used. MRS scans were preceded by MP2RAGE (magnetization prepared 2 rapid gradient echoes), a three-dimensional T1-weighted sequence, for
116 localization purposes with 0.9 mm isotropic resolution. Other parameters were TR=4900 ms, TE=2.9 ms, TI1=700 ms, FA1=5°, TI2=2700 ms,
117 FA2=6°. Single-voxel ¹H MRS was acquired using STEAM with a 20 x 20 x 20 mm³ cubical midline voxel (volume of 8 mL). The anterior inferior
118 border of the voxel was just above the corpus callosum, its anterior superior border just dorsal to the marginal branch of the cingulate sulcus, and
119 its posterior border ventral to the parieto-occipital sulcus, in a similar manner to the study of Kantarci et al – see Figure 1.²⁵

120 Shimming was performed prior to MRS acquisition. The Siemens standard automated shimming was employed followed by manual fine tuning
121 of gradients to optimize the line width. The TR was 8500 ms except for the scan of control number 5, who reached a specific absorption rate limit
122 and therefore TR=9300 ms was used in his case. TE was 6 ms. In total, thirty-two averages were acquired for the water-suppressed sequence
123 (acquisition time of 4:49 minutes), and four averages were used for the unsuppressed sequence (acquisition time of 51 seconds). The first spectrum
124 from the healthy control scans was compared to the single spectrum obtained from the patients. A representative spectrum from each group (TLE,
125 IGE, and controls) appears in Figure 2.

126 **2.4 Quantitative MRS calculation**

127

128 Metabolite concentrations were quantified using LCModel version 6.3-0B with a 7-Tesla basis set using scaling to unsuppressed water and Eddy-
129 current correction.²⁶ The LCModel output included estimated concentrations (mM) and Cramér–Rao lower bounds (CRLB), which are estimated
130 standard deviations, expressed in percent of the estimated concentrations.

131 For partial volume effect correction, we created binary masks for each MRS voxel using a MATLAB script created by Mr. Bartosz Kossowski
132 from The Polish Academy of Sciences (<https://www.nitrc.org/projects/rda2nifti/>). MATLAB version R2017b was used (MathWorks, Natick, MA,
133 United States). The ANTs toolbox (<http://stnava.github.io/ANTs/>) version 2.3.1 was used to transform the dimensions of the masks for consistency
134 with MP2RAGE coordinates.²⁷ Skull-stripping and brain extraction of the MP2RAGE images was done using FMRIB’s Brain Extraction Tool.²⁸
135 Each brain was then segmented using FMRIB’s Automated Segmentation Tool (FAST) into partial volume maps for gray matter, white matter and
136 CSF.²⁹

137 Fractions of the partial volume maps within each MRS voxel were determined using the fslstats utility of the FSL suite version 5.0.10.³⁰ The NMR-
138 visible water concentration (mM) in the voxel was estimated by $(43300f_{GM} + 35880f_{WM} + 55556f_{CSF}) / (1 - f_{CSF})$, where f_{GM} , f_{WM} , and f_{CSF} are the
139 volume fractions of gray matter, white matter, and CSF in the voxel.³¹

140 **2.5 Statistical Analysis**

141

142 Continuous variables (age of the participants, duration of epilepsy, the concentrations of glutamate, GABA, GSH, and the ratio of glutamate to
143 GABA concentrations) were compared between healthy controls, patients with TLE, and patient with IGE. Shapiro-Wilk test checked the
144 assumption of normality of these variables and Bartlett’s test was used to check for homoscedasticity. One-way analysis of variance (ANOVA)
145 with post-hoc Tukey-Kramer test for multiple comparisons was used to compare the means between the three groups when data passed these tests.
146 In cases of normality without homoscedasticity Welch’s ANOVA was used. In cases data did not pass the normality test Kruskal-Wallis H test

147 was used, adjusted for ties. In addition, the discrete ordinal variable of the number of medications was also compared between the patient groups
148 using Kruskal-Wallis H test as well as the analysis of medication effects in subgroups due to very small sample sizes.

149 Mean concentrations of glutamate, GABA, the concentration ratio glutamate/GABA and GSH all followed a Gaussian distribution and passed the
150 Shapiro-Wilk normality test with $P>0.05$. Other than the concentration of glutamate, all other MRS results that were the focus of this study also
151 passed Bartlett's test of homoscedasticity and one-way ANOVA was used for comparison between the groups. The concentrations of glutamate
152 were compared with Welch's ANOVA. The age of participants passed the Shapiro-Wilk normality test and Bartlett's test of homoscedasticity and
153 was compared using one-way ANOVA across the three groups. Epilepsy duration, however, did not pass Shapiro-Wilk's test in the TLE group
154 ($P=0.05$) and therefore Kruskal-Wallis H test was used for comparison of epilepsy duration.

155 Due to collinearity between metabolites and an overall small sample size exact logistic regression was employed for an exploratory analysis of the
156 relationship between short-term seizure freedom and each of the metabolites this study focused on in the entire patient population. Multiple linear
157 regression was used to study the effects of age and duration of epilepsy on metabolite concentrations in the patient groups. Beta regression was
158 similarly used to study the effects of age and duration of epilepsy on the fraction of CSF in the region of interest as a measure of atrophy in all 35
159 patients. All statistical analyses were conducted with Stata 15.0 (StataCorp, College Station, TX, USA). Statistical significance was set at $\alpha=0.05$.

160 3. RESULTS

161
162 Metabolite concentrations and age of participants for patients and controls are summarized in Table 1. Results of multiple regression of metabolite
163 concentrations in the volume of interest with age and duration of epilepsy in the patient groups are summarized in Table 2. Exact logistic regressions
164 of short-term seizure-freedom with metabolite concentrations are summarized in Table 3. Tests of normality and homoscedasticity, composition
165 of the volume of interest, analysis of drug effects, and clinical information for all the patients are summarized in the supplementary tables.

166 3.1 Quality Control

167

168 Values of CRLB for the various metabolites and Full Width at Half Maximum (FWHM) of the spectra were obtained from the LCModel output.
169 Results with CRLB above 20% were unreliable and excluded from further analysis, resulting in the exclusion of a single GABA concentration
170 from the spectrum of one patient from the IGE group (due to CRLB=44%).

171 Maximal CRLB for glutamate, GABA, and GSH were 2%, 11%, and 6%, respectively, in the control group. The mean($\pm$ SD) FWHM of the spectra
172 was 0.034 ± 0.003 ppm (10.1 ± 0.9 Hz) in these subjects, confirming adequate quality.

173 In the TLE group, one patient's spectrum had poor quality with a FWHM of 0.099 ppm (29.4 Hz), which was above an established rejection cut-
174 off of 0.070 ppm.³² That spectrum was excluded from further analysis, resulting in maximal CRLB of 3%, 14%, and 6% for glutamate, GABA,
175 and GSH, respectively, and a mean($\pm$ SD) FWHM of 0.034 ± 0.005 ppm (10.1 ± 1.5 Hz) for the remaining patients with TLE.

176 In the IGE group, after the exclusion of the single result mentioned above maximal CRLB were 4%, 18%, and 8% for glutamate, GABA, and
177 GSH, respectively. Mean $\pm$ SD FWHM of the spectra in the IGE group were 0.036 ± 0.008 ppm (10.7 ± 2.4 Hz).

178 3.2 Voxel Composition

179

180 Beta regression of the fraction of CSF in the volume of interest with age and duration of epilepsy showed a significant increase in the fraction of
181 CSF with age in the TLE group ($P=0.02$) with a marginal effect of 0.002 per year (95% confidence interval 0.0003-0.003). No independent effect
182 of epilepsy duration was found on the fraction of CSF ($P=0.19$). In the IGE group, no effect of patient age or the duration of epilepsy on the
183 fraction of CSF was demonstrated ($P=0.69$ and $P=0.79$, respectively). Considering the possibility of a differential effect of seizure focus laterality
184 on this atrophic process in the TLE group, beta regression of the CSF fraction vs age was performed on patients with left-sided TLE ($n=9$) and

185 patients with right-sided TLE (n=6). For right-sided TLE the effect of age was not significant ($P=0.89$) whereas for left-sided TLE it was ($P=0.009$)
186 with a marginal effect of 0.002 per year (95% confidence interval 0.001-0.004).

187 **3.3 Glutamate**

188
189 Welch's ANOVA showed no significant difference in the concentration of glutamate between the three groups. Multiple regression demonstrated
190 no significant effect of age or the duration of epilepsy on the concentration of glutamate in the patient groups.

191 **3.4 GABA**

192
193 One-way ANOVA showed no statistically significant difference in the concentration of GABA between the three groups. Multiple regression of
194 the concentration of GABA with age and the duration of epilepsy showed a significant reduction in GABA concentration with age in the TLE
195 group ($P=0.03$) – see Figure 3B. The effect size, as measured by η^2 , is 0.28 (Cohens' $f=0.62$). No such effect was found in the IGE group and no
196 independent effect of epilepsy duration on GABA concentration was demonstrated in the patient groups.

197 **3.5 The Glutamate to GABA Concentration Ratio**

198
199 One-way ANOVA showed no significant difference in the ratio of glutamate to GABA concentration between the three groups. Multiple regression
200 demonstrated no significant effect of age or the duration of epilepsy on this ratio in the patient groups.

201 **3.6 GSH**

202

203 One-way ANOVA demonstrated a statistically significant difference for the concentration of GSH between the groups, $F(2,41)=3.68$, $P=0.03$.
204 Post-hoc Tukey-Kramer test for multiple comparisons revealed a statistically significant difference between the IGE and control groups ($P=0.03$).
205 The effect size, as measured by η^2 , is 0.15 (Cohen's $f=0.42$). The concentration was not significantly different between the IGE and TLE groups
206 and between patients with TLE and healthy controls.

208 **3.7 Other Metabolites**

209
210 Figure 3A shows mean concentration and standard deviations of all metabolites with consistent CRLB<20% that were acquired from all groups:
211 glutamate, GABA, GSH, glutamine, inositol, NAA, and total creatine (creatine + phosphocreatine). To facilitate comparison between studies,
212 concentrations of glutamate, GABA, and GSH normalized to total creatine appear in the supplementary data (Table 2S). Notably, the stability of
213 total creatine was checked among the patient groups and the controls once its normality and homoscedasticity were established. It was found not
214 to be different between the groups, $F(2,41)=2.79$, $P=0.07$.

215 **3.8 Effect of Metabolite Concentrations on Short-Term Seizure-Freedom**

216
217 Exact logistic regression performed separately on glutamate, GABA, and GSH, evaluated their effect on seizure-freedom in the preceding 3 months
218 before the scan in the entire patient population (16 of the 34 patients in the final analysis reported short-term seizure-freedom). While no effect of
219 GABA, the ratio glutamate/GABA, and GSH on short-term seizure-freedom was found, the concentration of glutamate was demonstrated to be a
220 predictor of short-term seizure-freedom (odd ratio 2.14; 95% confidence interval 1.17-4.46; $P=0.01$) – see Table 3. In order to test whether patients
221 who reported short-term seizure-freedom had indeed higher concentrations of glutamate in the PCC/precuneus, a t-test was performed after
222 confirmation of normality, comparing the concentration of glutamate between the groups. In patients who reported short-term seizure-freedom,

223 the concentration of glutamate (mean±SD) was 10.6±1.4 mM, whereas in patients who did not the concentration was 9.4±1.2 mM, resulting in a
224 one-tailed value of $P=0.005$.

225 4. DISCUSSION

226
227 The results of this study found a higher concentration of GSH on high field MRS in the PCC/precuneus in the IGE group in comparison with
228 controls. No such difference was found for other metabolites that were examined in this study, which is in contradiction with our hypotheses and
229 requires explanation. One possibility is that changes in excitatory and inhibitory neurotransmitters do exist in these types of epilepsy in comparison
230 with each other and with healthy controls but are located in the epileptogenic networks themselves and not in the DMN. Another possibility is that
231 no such changes exist.

232 However, we did find evidence that glutamate concentration in the PCC/precuneus may be a predictor of short-term seizure-freedom, possibly
233 related to less DMN deactivation in these patients as a reflection of a more benign clinical course. There is evidence that task-induced deactivation
234 in the PCC/precuneus correlates with the glutamate/GABA concentration ratio in this region.³³ The finding that glutamate, and not its ratio to
235 GABA, was a predictor of short-term seizure-freedom suggests that DMN deactivation and metabolite concentrations in the PCC/precuneus may
236 be related. Moreover, seizure-freedom in patients with MRI-negative TLE is known to be associated with ipsilateral elevation of glutamate
237 concentration, which is speculated to reflect increased propensity for seizures on a background of preservation of tissue integrity.³⁴ It is possible
238 that our results were underpinned by a similar mechanism.

239 Focusing on the age-related reduction in the concentration of GABA in the TLE group, we have also noted the increasing fractional component of
240 CSF with age in patients with TLE as a measure of atrophy, a finding driven by patients with left-sided epileptogenic foci, and not demonstrated
241 in patients with IGE. These findings could be related. Interestingly, in the worldwide ENIGMA project, MRI scans of 367 patients with IGE, 754
242 with mesial TLE with hippocampal sclerosis, 1026 with other forms of epilepsy, and 1727 matched healthy controls were compared. No changes

243 of cortical thickness were seen for the IGE group in the PCC and precuneus regions. In contrast, in TLE patients with hippocampal sclerosis the
244 ENIGMA study revealed cortical thickness reductions of the bilateral precuneal cortices with left, but not right, seizure foci.³⁵ Our results are in
245 line with this finding. It is possible, therefore, that the precuneus region undergoes atrophic changes in patients with left-sided TLE, affecting
246 GABAergic more than glutamatergic neurons.

247 An important question to consider is whether the concentrations of the studied metabolites were affected by drug treatment. Some ASMs such as
248 vigabatrin, gabapentin, lamotrigine, and topiramate are known increase brain GABA concentrations.^{36,37} The effects of carbamazepine and
249 valproate were also assessed in several small studies. Petroff and coworkers compared metabolite concentrations in the occipital cortex of 14
250 patients with epilepsy to 10 healthy controls using 2.1T MRS. They reported reduced GABA levels in 4/8 patients taking carbamazepine. Of the
251 five patients taking valproate, one had reduced GABA levels, two had increased glutamate levels, and one demonstrated reduced glutamate
252 concentration.³⁸ Garcia and coworkers used 3T MRS focusing on the parietal and occipital lobes to acquire the metabolic profile of patients with
253 epilepsy taking valproate and compared them to healthy controls. While lower levels of glutamate were seen in the spectra of the parietal lobes in
254 the patient group, the difference was not significant.³⁹ Finally, Doelken et al showed increased GABA levels in patients with focal epilepsy who
255 responded to levetiracetam.⁴⁰

256 The four most prevalent ASMs in our patient population were carbamazepine, valproate, levetiracetam, and lamotrigine. No effect of these
257 medications on the levels of glutamate and GABA was found except for a reduction in the concentration of glutamate in patients with TLE treated
258 with valproate (Cohen's $d=1.9$; see Table 4S in the supplementary data). However, only three patients in this group were treated with this
259 medication, having little effect on the overall findings, and making it difficult to draw conclusions.

260 Could the higher levels of GSH in the PCC/precuneus in patients with IGE be related to drug effects? Ono and coworkers have shown that total
261 plasma glutathione was reduced in epilepsy patients treated with phenytoin and carbamazepine alone or with multiple ASMs in comparison with

controls. No such results were seen in patients treated with valproic acid or phenobarbital monotherapy.⁵ A similar study demonstrated a reduction in erythrocyte levels of GSH in children treated with valproic acid or carbamazepine.⁴¹

Animal models have shed light on the effects other ASMs have on GSH levels. Topiramate was shown to increase levels of GSH in erythrocytes in a rodent model of epilepsy, and oxcarbazepine was shown to decrease the brain levels of GSH in another model in mice^{42,43} Lamotrigine was found to have both increasing and decreasing effects on brain GSH levels in rodent models of epilepsy and levetiracetam was shown to increase levels of GSH in the hippocampi of mice in a pilocarpine model of epilepsy.^{43–45} No patient in our study was treated with phenytoin or oxcarbazepine, and only 2 of our IGE patients were taking topiramate (and none in the TLE group). As for carbamazepine, valproate, levetiracetam and lamotrigine, no effect of these medications on the concentration of GSH was found (see Table 4S in the supplementary data). As discussed above, the number of medications was similar between the patient groups. To summarize, no medication effect seems to have been responsible for the higher levels of GSH in the IGE group.

A possible explanation for this finding could be a response to oxidative stress. Numerous animal models have demonstrated the involvement of reactive oxygen species in the pathogenesis of epilepsy.⁴⁶ Similar mechanisms were shown in a range of neurodegenerative and psychiatric conditions, such as Huntington's disease, Parkinson's disease, Alzheimer's disease, and schizophrenia, with a compensatory initial elevation of GSH concentration eventually being overwhelmed.^{46,47} GSH levels were shown to be reduced in the parietooccipital region of both hemispheres in patients with focal epilepsy in a previous MRS study.¹¹ However, we are unaware of a previous MRS study measuring GSH levels in patients with IGE. Notably, a 7T MRS study of metabolic changes in the hippocampi of rats after induction of status epilepticus by pilocarpine injection discovered gradual elevation of the GSH (scaled to creatine) over 7 days, remaining elevated in chronic epileptic rats.⁴⁸

The study's main limitation is its relatively small sample size. While our results are exploratory in nature, they do suggest elevated GSH concentrations in the PCC/precuneus of patients with IGE. If the difference in GSH concentration between patients with IGE and other groups is found to be consistent in larger cohorts, it may have some value in the management of epilepsy patients in the future. For example, the exact

282 mechanism by which the ketogenic diet, which is used for treatment of intractable seizures, helps control seizures is not fully explained.
283 Nevertheless, it was shown that a possible mechanism may be related to activation of pathways resulting in increased production of antioxidants
284 such as GSH.⁴⁹ Monitoring serum and brain levels of GSH may prove to be important in guiding such modes of therapy.

285 The higher level of glutamate in patients who reported short-term seizure-freedom is another exploratory finding. If corroborated in larger cohorts,
286 it may prove useful as part of a future model of drug-response prediction.

287 5. CONCLUSIONS

288
289 In conclusion, no difference was found in the concentrations of glutamate, GABA, or their ratio in the PCC/precuneus between patients with IGE,
290 patients with TLE, and healthy controls. An age-related decline in the concentration of GABA in patients with TLE was demonstrated. Patients
291 with IGE were shown to have higher concentrations of GSH in the PCC/precuneus than healthy controls. The concentration of glutamate in the
292 PCC/precuneus may predict short-term seizure-freedom.

296 ACKNOWLEDGMENTS

297
298 The authors acknowledge the facilities, scientific and technical assistance, and salary support of Bradford A. Moffat from the National Imaging
299 Facility, a National Collaborative Research Infrastructure Strategy (NCRIS) capability, at the Melbourne Brain Centre Imaging Unit, The
300 University of Melbourne. Ofer M. Gonen was supported by the Melbourne Research Scholarship (MRS), The University of Melbourne and The
301 Neuroscience Foundation, The Royal Melbourne Hospital. Patrick Kwan is supported by a Medical Research Future Fund Practitioner Fellowship.

302 We would also like to thank Mr. Bartosz Kossowski, Laboratory of Brain Imaging, Nencki Institute of Experimental Biology, Polish Academy of
303 Sciences, for a tool which creates MRS voxel masks aligned to imaging data (<https://www.nitrc.org/projects/rda2nifti/>).
304

305 **Ethical Publication Statement**

306

307 We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this report is consistent with those
308 guidelines.
309

310 **Disclosures:** Neither of the authors has any conflict of interest to disclose.

311 **REFERENCES**

312

- 313 1 . Benarroch EE. Glutamate transporters: Diversity, function, and involvement in neurologic disease. *Neurology* 2010.
314 doi:10.1212/WNL.0b013e3181cc89e3.
- 315 2 . Kocsis JD, Mattson RH. GABA levels in the brain: A target for new antiepileptic drugs. *Neuroscientist*. 1996.
316 doi:10.1177/107385849600200610.
- 317 3 . van Veenendaal TM, Ijff DM, Aldenkamp AP, Hofman PAM, Vlooswijk MCG, Rouhl RPW *et al*. Metabolic and functional MR
318 biomarkers of antiepileptic drug effectiveness: A review. *Neurosci Biobehav Rev* 2015; 59: 92–99.
- 319 4 . Rae CD, Williams SR. Glutathione in the human brain: Review of its roles and measurement by magnetic resonance spectroscopy. *Anal*
320 *Biochem* 2017; 529: 127–143.
- 321 5 . Ono H, Sakamoto A, Sakura N. Plasma total glutathione concentrations in epileptic patients taking anticonvulsants. *Clin Chim Acta* 2000;
322 298: 135–143.
- 323 6 . Prasad DKV, Satyanarayana U, Shaheen U, Surya Prabha T, Munshi A. Oxidative stress in the development of genetic generalised

- 324 epilepsy: An observational study in southern Indian population. *J Clin Diagnostic Res* 2017; 11: BC05–BC08.
- 325 7 . Simister RJ, McLean MA, Barker GJ, Duncan JS. A proton magnetic resonance spectroscopy study of metabolites in the occipital lobes in
326 epilepsy. *Epilepsia* 2003. doi:10.1046/j.1528-1157.2003.19102.x.
- 327 8 . Simister RJ, McLean MA, Barker GJ, Duncan JS. Proton MRS reveals frontal lobe metabolite abnormalities in idiopathic generalized
328 epilepsy. *Neurology* 2003; 61: 897–902.
- 329 9 . Simister RJ, McLean MA, Barker GJ, Duncan JS. Proton MR spectroscopy of metabolite concentrations in temporal lobe epilepsy and
330 effect of temporal lobe resection. *Epilepsy Res* 2009; 83: 168–176.
- 331 10 . Hattingen E, Lückcrath C, Pellikan S, Vronski D, Roth C, Knake S *et al.* Frontal and thalamic changes of GABA concentration indicate
332 dysfunction of thalamofrontal networks in juvenile myoclonic epilepsy. *Epilepsia* 2014. doi:10.1111/epi.12656.
- 333 11 . Mueller SG, Trabesinger AH, Boesiger P, Wieser HG. Brain glutathione levels in patients with epilepsy measured by in vivo 1H-MRS.
334 *Neurology* 2001; 57: 1422–1427.
- 335 12 . Tkáč I, Andersen P, Adriany G, Merkle H, Ugurbil K, Gruetter R. In vivo 1H NMR spectroscopy of the human brain at 7 T. *Magn Reson*
336 *Med* 2001; 46: 451–456.
- 337 13 . Pan JW, Duckrow RB, Spencer DD, Avdievich NI, Hetherington HP. Selective homonuclear polarization transfer for spectroscopic
338 imaging of GABA at 7T. *Magn Reson Med* 2013. doi:10.1002/mrm.24283.
- 339 14 . Davis KA, Nanga RPR, Das S, Chen SH, Hadar PN, Pollard JR *et al.* Glutamate imaging (GluCEST) lateralizes epileptic foci in
340 nonlesional temporal lobe epilepsy. *Sci Transl Med* 2015; 7: 1–8.
- 341 15 . Neal A, Moffat BA, Stein JM, Nanga RPR, Desmond P, Shinohara RT *et al.* Glutamate weighted imaging contrast in gliomas with 7 Tesla
342 magnetic resonance imaging. *NeuroImage Clin* 2019; 22: 101694.
- 343 16 . Hagmann P, Cammoun L, Gigandet X, Meuli R, Honey CJ, Van W vedeen J *et al.* Mapping the structural core of human cerebral cortex.
344 *PLoS Biol* 2008; 6: 1479–1493.
- 345 17 . Raichle ME. The Brain's Default Mode Network. *Annu Rev Neurosci* 2015; 38: 433–447.

- 346 18 . Doelken MT, Mennecke A, Stadlbauer A, Kecskeméti L, Kasper BS, Struffert T *et al.* Multi-voxel magnetic resonance spectroscopy at 3
347 T in patients with idiopathic generalised epilepsy. *Seizure* 2010. doi:10.1016/j.seizure.2010.07.005.
- 348 19 . Haheef Z, Lenartowicz A, Yeh HJ, Engel J, Stern JM. Effect of lateralized temporal lobe epilepsy on the default mode network. *Epilepsy*
349 *Behav* 2012. doi:10.1016/j.yebeh.2012.07.019.
- 350 20 . Kay BP, Difrancesco MW, Privitera MD, Gotman J, Holland SK, Szaflarski JP. Reduced default mode network connectivity in treatment-
351 resistant idiopathic generalized epilepsy. *Epilepsia* 2013; 54: 461–470.
- 352 21 . McGill ML, Devinsky O, Kelly C, Milham M, Castellanos FX, Quinn BT *et al.* Default mode network abnormalities in idiopathic
353 generalized epilepsy. *Epilepsy Behav* 2012; 23: 353–359.
- 354 22 . Zanão TA, Lopes TM, de Campos BM, Yasuda CL, Cendes F. Patterns of default mode network in temporal lobe epilepsy with and
355 without hippocampal sclerosis. *Epilepsy Behav* 2019. doi:10.1016/j.yebeh.2019.106523.
- 356 23 . Zhang Z, Lu G, Zhong Y, Tan Q, Liao W, Wang Z *et al.* Altered spontaneous neuronal activity of the default-mode network in mesial
357 temporal lobe epilepsy. *Brain Res* 2010; 1323: 152–160.
- 358 24 . Scheffer IE, Berkovic S, Capovilla G, Connolly MB, French J, Guilhoto L *et al.* ILAE classification of the epilepsies: Position paper of
359 the ILAE Commission for Classification and Terminology. *Epilepsia* 2017. doi:10.1111/epi.13709.
- 360 25 . Kantarci K, Weigand SD, Petersen RC, Boeve BF, Knopman DS, Gunter J *et al.* Longitudinal 1H MRS changes in mild cognitive
361 impairment and Alzheimer’s disease. *Neurobiol Aging* 2007. doi:10.1016/j.neurobiolaging.2006.06.018.
- 362 26 . Provencher SW. Estimation of metabolite concentrations from localized in vivo proton NMR spectra. *Magn Reson Med* 1993; 30: 672–
363 679.
- 364 27 . Avants BB, Tustison NJ, Song G, Cook PA, Klein A, Gee JC. A reproducible evaluation of ANTs similarity metric performance in brain
365 image registration. *Neuroimage* 2011. doi:10.1016/j.neuroimage.2010.09.025.
- 366 28 . Smith SM. Fast robust automated brain extraction. *Hum Brain Mapp* 2002. doi:10.1002/hbm.10062.
- 367 29 . Zhang Y, Brady M, Smith S. Segmentation of brain MR images through a hidden Markov random field model and the expectation-

- maximization algorithm. IEEE Trans Med Imaging 2001. doi:10.1109/42.906424.
- 30 . Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM. FSL - Review. Neuroimage 2012. doi:10.1016/j.neuroimage.2011.09.015.
- 31 . Lee H, Caparelli E, Li H, Mandal A, Smith SD, Zhang S *et al.* Computerized MRS voxel registration and partial volume effects in single voxel 1H-MRS. Magn Reson Imaging 2013; 31: 1197–1205.
- 32 . Kreis R. Issues of spectral quality in clinical 1H-magnetic resonance spectroscopy and a gallery of artifacts. NMR Biomed 2004; 17: 361–381.
- 33 . Gu H, Hu Y, Chen X, He Y, Yang Y. Regional excitation-inhibition balance predicts default-mode network deactivation via functional connectivity. Neuroimage 2019; 185: 388–397.
- 34 . Pimentel-Silva LR, Casseb RF, Cordeiro MM, Campos BAG, Alvim MKM, Rogerio F *et al.* Interactions between in vivo neuronal-glia markers, side of hippocampal sclerosis, and pharmacoresponse in temporal lobe epilepsy. Epilepsia 2020; 61: 1008–1018.
- 35 . Whelan CD, Altmann A, Botia JA, Jahanshad N, Hibar DP, Absil J *et al.* Structural brain abnormalities in the common epilepsies assessed in a worldwide ENIGMA study. Brain 2018. doi:10.1093/brain/awx341.
- 36 . Puts NAJ, Edden RAE. In vivo magnetic resonance spectroscopy of GABA: A methodological review. Prog Nucl Magn Reson Spectrosc 2012; 60: 29–41.
- 37 . Kuzniecky R, Ho S, Pan J, Martin R, Gilliam F, Faught E *et al.* Modulation of cerebral GABA by topiramate, lamotrigine, and gabapentin in healthy adults. Neurology 2002; 58: 368–372.
- 38 . Petroff OAC, Rothman DL, Behar KL, Hyder F, Mattson RH. Effects of valproate and other antiepileptic drugs on brain glutamate, glutamine, and GABA in patients with refractory complex partial seizures. Seizure 1999; 8: 120–127.
- 39 . Garcia M, Huppertz HJ, Ziyeh S, Buechert M, Schumacher M, Mader I. Valproate-induced metabolic changes in patients with epilepsy: Assessment with 1H-MRS. Epilepsia 2009; 50: 486–492.
- 40 . Doelken MT, Hammen T, Bogner W, Mennecke A, Stadlbauer A, Boettcher U *et al.* Alterations of intracerebral -aminobutyric acid

- 390 (GABA) levels by titration with levetiracetam in patients with focal epilepsies. *Epilepsia* 2010; 51: 1477–1482.
- 391 41 . Cengiz M, Yüksel A, Seven M. The effects of carbamazepine and valproic acid on the erythrocyte glutathione, glutathione peroxidase,
392 superoxide dismutase and serum lipid peroxidation in epileptic children. *Pharmacol Res* 2000. doi:10.1006/phrs.1999.0603.
- 393 42 . Naziroğlu M, Kutluhan S, Uğuz AC, Çelik Ö, Bal R, Butterworth PJ. Topiramate and vitamin e modulate the electroencephalographic
394 records, brain microsomal and blood antioxidant redox system in pentylenetetrazol-induced seizure of rats. *J Membr Biol* 2009.
395 doi:10.1007/s00232-009-9177-1.
- 396 43 . Agarwal NB, Agarwal NK, Mediratta PK, Sharma KK. Effect of lamotrigine, oxcarbazepine and topiramate on cognitive functions and
397 oxidative stress in PTZ-kindled mice. *Seizure* 2011. doi:10.1016/j.seizure.2010.12.006.
- 398 44 . Arora T, Mehta AK, Sharma KK, Mediratta PK, Banerjee BD, Garg GR *et al.* Effect of carbamazepine and lamotrigine on cognitive
399 function and oxidative stress in brain during chemical epileptogenesis in rats. *Basic Clin Pharmacol Toxicol* 2010. doi:10.1111/j.1742-
400 7843.2009.00499.x.
- 401 45 . Oliveira AA, Almeida JPC, Freitas RM, Nascimento VS, Aguiar LMV, Júnior HVN *et al.* Effects of levetiracetam in lipid peroxidation
402 level, nitrite-nitrate formation and antioxidant enzymatic activity in mice brain after pilocarpine-induced seizures. *Cell Mol Neurobiol*
403 2007. doi:10.1007/s10571-006-9132-y.
- 404 46 . Cárdenas-Rodríguez N, Coballase-Urrutia E, Pérez-Cruz C, Montesinos-Correa H, Rivera-Espinosa L, Sampieri A *et al.* Relevance of the
405 glutathione system in temporal lobe epilepsy: Evidence in human and experimental models. *Oxid Med Cell Longev* 2014; 2014.
406 doi:10.1155/2014/759293.
- 407 47 . Wood SJ, Berger GE, Wellard RM, Proffitt TM, McConchie M, Berk M *et al.* Medial temporal lobe glutathione concentration in first
408 episode psychosis: A 1H-MRS investigation. *Neurobiol Dis* 2009. doi:10.1016/j.nbd.2008.11.018.
- 409 48 . Filibian M, Frasca A, Maggioni D, Micotti E, Vezzani A, Ravizza T. In vivo imaging of glia activation using 1H-magnetic resonance
410 spectroscopy to detect putative biomarkers of tissue epileptogenicity. *Epilepsia* 2012; 53: 1907–1916.
- 411 49 . Milder J, Patel M. Modulation of oxidative stress and mitochondrial function by the ketogenic diet. *Epilepsy Res* 2012.

Figure Legends

Figure 1 – T₁-weighted sagittal MRI with superimposed MRS 8 cm³ voxel

Figure 2 – Representative spectra from each group (A - TLE; B - IGE; C – Controls)

Figure 3 – Results of metabolite quantification (A – Metabolic Panel of all groups with SD error bars; B - GABA vs age in patients with TLE)

Table Captions:

Table 1 – Comparison between the groups

Table 2 – Metabolites vs age and duration of epilepsy

Table 3 – Exact logistic regression of short-term seizure-freedom with metabolite concentrations

Supplementary Material Tables:

430 **Table 1S** – Tests of normality and homoscedasticity (*P*-values)

431 **Table 2S** – Metabolite concentrations normalized to total creatine (mean±SD)

432 **Table 3S** – Voxel segmentation

433 **Table 4S** – Analysis of drug effects (Kruskal-Wallis H test)

434 **Table 5S** – Clinical data

435 **Table 1 – Comparison between the groups**

	TLE	IGE	Controls	Statistic	<i>P</i>-value
Mean(±SD) age (years)	41.7(±14.3)	39.2(±15.3)	40.0(±14.4)	$F(2,41)=0.13$	0.88
Mean(±SD) glutamate (mM)	9.7(±1.0)	10.2(±1.8)	10.3(±0.9)	Welch's $F(2,24.1)$	0.24
Mean(±SD) GABA (mM)	1.5(±0.2)	1.6(±0.4)	1.5(±0.2)	$F(2,40)=0.45$	0.64
Mean(±SD) glutamate/GABA	6.4(±1.0)	6.6(±1.1)	7.0(±0.8)	$F(2,40)=0.95$	0.40
Mean(±SD) GSH (mM)	2.0(±0.2)	2.2(±0.4)	1.9(±0.3)	$F(2,41)=3.68$	0.03

436

437

438 **Table 2 – Metabolites vs age and duration of epilepsy**

Metabolite	<i>P</i> (model)	R^2	β (age)	<i>p</i> (age)	β (duration)	<i>P</i> (duration)
TLE						
Glutamate	0.10	0.256	-0.04	0.05	0.01	0.74
GABA	0.02	0.397	-0.01	0.03	0.00	0.56
Glutamate/GABA	0.18	0.208	0.02	0.31	0.02	0.36

GSH	0.18	0.206	0.01	0.10	0.00	0.91
IGE						
Glutamate	0.02	0.452	-0.07	0.18	-0.01	0.88
GABA	0.06	0.369	-0.02	0.09	0.00	0.66
Glutamate/GABA	0.22	0.226	0.07	0.09	-0.04	0.19
GSH	0.44	0.119	-0.01	0.43	0.00	0.81

Table 3 – Exact logistic regression of short-term seizure-freedom with metabolite concentrations

Metabolite	Odds Ratio	95% Confidence Interval	<i>P</i>-value ($\geq$ Model Score)
Glutamate	2.14	1.17 - 4.46	0.01
GABA	8.24	0.71 - 146.90	0.09
GSH	7.56	0.67 - 115.10	0.10
Glutamate / GABA	0.92	0.45 - 1.82	0.82

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Gonen, OM;Moffat, BA;Desmond, PM;Lui, E;Kwan, P;O'Brien, TJ

Title:

Seven-tesla quantitative magnetic resonance spectroscopy of glutamate, gamma-aminobutyric acid, and glutathione in the posterior cingulate cortex/precuneus in patients with epilepsy

Date:

2020-10-28

Citation:

Gonen, O. M., Moffat, B. A., Desmond, P. M., Lui, E., Kwan, P. & O'Brien, T. J. (2020). Seven-tesla quantitative magnetic resonance spectroscopy of glutamate, gamma-aminobutyric acid, and glutathione in the posterior cingulate cortex/precuneus in patients with epilepsy. *EPILEPSIA*, 61 (12), pp.2785-2794. <https://doi.org/10.1111/epi.16731>.

Persistent Link:

<http://hdl.handle.net/11343/276538>