Effective treatment of disseminated superficial actinic porokeratosis using a novel topical cholesterol/simvastatin combination cream

Running head: DSAP treated with cholesterol/statin

Word count: 561

Figures: 1

Key words: porokeratosis, treatment, disseminated superficial actinic porokeratosis, cholesterol, statin, simvastatin

Authors:

Rebekka Jerjen¹, MChD

Wei-Liang Koh^{1,2}, MBBS, MRCP

Rodney Sinclair¹, MBBS, MD, FACD

¹ Sinclair Dermatology, East Melbourne, Victoria 3002, Australia

² Department of Dermatology, Changi General Hospital, Singapore

Corresponding author

Dr Rebekka Jerjen

Sinclair Dermatology

2 Wellington Parade, East Melbourne, Victoria 3002, Australia.

ORCID: 0000-0002-7749-1069

Tel: +61 405013585

Fax: +61 3 9650 9944

E-mail: Rebekka.jerjen@gmail.com

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/AJD.13473](https://doi.org/10.1111/AJD.13473)

This article is protected by copyright. All rights reserved

COI: None to declare

Funding: None

Acknowledgments: The patient has provided consent for use of photographs in this publication

Author Manuscript

DR. REBEKKA BARBARA JERJEN (Orcid ID : 0000-0002-7749-1069)

PROF. RODNEY SINCLAIR (Orcid ID : 0000-0001-6751-1428)

Article type : Case Letter

Effective treatment of disseminated superficial actinic porokeratosis using a novel topical cholesterol/simvastatin combination cream

Disseminated superficial actinic porokeratosis (DSAP) is the most common subtype of porokeratosis. It is characterised by pink-brown papules or plaques with a ridge-like border affecting areas of sun-exposed skin.¹ Sporadic and autosomal dominant inheritance patterns have been described and pathogenic germline mutations identified in the mevalonate pathway genes.² DSAP has been reported to undergo malignant transformation, most commonly to squamous cell carcinoma (SCC) but also basal cell carcinoma (BCC).³ Current treatments include topical therapy such as imiquimod, 5-fluorouracil, retinoids, corticosteroids, vitamin-D analogues; systemic retinoids and physical therapies such as cryotherapy.⁴ A recent small case series described the successful treatment of porokeratosis using a combination of topical 2% cholesterol and 2% lovastatin.¹ Lovastatin is not readily available in Australia nor the UK. Herein we present two patients with DSAP treated with an off-label combination of 2% cholesterol and 2% simvastatin cream.

Case 1: A 63-year-old female presented for a routine skin check. Her past medical history included two previously excised BCCs and frontal fibrosing alopecia. On examination, she had scattered pink-brown papules with a raised ridge-like border on the legs, consistent with DSAP. After an informed discussion, she was commenced on a compounded, twice-daily, topical 2% cholesterol/2% simvastatin cream. On three-month review, there was subjective and objective improvement in the lesions to which she had applied the cream. There were no adverse effects.

Case 2: A 52-year-old female presented with a 15-year history of DSAP affecting her arms and legs. She reported similar lesions in her mother and grandmother. Previous ineffective treatments included cryotherapy and 5-fluoruracil cream. On examination, there were scattered pink-brown papules with a ridge-like border on her legs and forearms consistent with DSAP (Figure 1). Given these characteristic clinical findings, no biopsy was performed. The patient commenced treatment with topical 2% cholesterol/ 2% simvastatin cream. On three-month review, there was an objective and subjective reduction in the number of lesions, with a markedly fainter appearance (Figure 1). She did not report any adverse effect from the cream and is continuing treatment.

Loss of function mutations in genes in the mevalonate pathway result in a reduction in its end products, including cholesterol, as well as accumulation of toxic intermediate metabolites.¹ Cholesterol is required to maintain skin integrity as it forms a key component of the extracellular lipid matrix in the stratum corneum. Cholesterol deficiency leads to increased keratinocyte sensitivity to apoptosis.⁵ Premature apoptosis and dysregulated keratinocyte differentiation have been identified in porokeratosis.² Consequently, it has been hypothesised that a cream containing an HMG-CoA reductase inhibitor (statin), that would prevent toxic metabolite accumulation, together with cholesterol to replenish essential mevalonate pathway end-products might be effective in porokeratosis. This was recently demonstrated by Atzmony *et al.*, who showed clinical improvement in one patient with DSAP, two patients with porokeratosis palmaris et plantaris disseminata and two patients with linear porokeratosis treated with topical 2% cholesterol/2% lovastatin ointment.¹

To the best of our knowledge, our two patients with DSAP are the first cases in Australia treated with topical 2% cholesterol/2% simvastatin cream, a pathogenesis-directed approach. We demonstrate simvastatin can be substituted for lovastatin in the compounded topical formulation. Our study is limited by the lack of placebo control and further controlled studies are required to verify the efficacy and safety of the medication as well as clarify the role of other toxic metabolites and depleted end-products that may play a role in porokeratosis pathogenesis.

REFERENCES

1. Atzmony L, Lim YH, Hamilton C, et al. Topical cholesterol/lovastatin for the treatment of porokeratosis: A pathogenesis-directed therapy. *J Am Acad Dermatol*. 2020;82(1):123-131. doi:10.1016/j.jaad.2019.08.043
2. Wang J, Liu Y, Liu F, et al. Loss-of-function Mutation in PMVK Causes Autosomal Dominant Disseminated Superficial Porokeratosis. *Sci Rep*. 2016;6(April):1-9. doi:10.1038/srep24226
3. Sasson M, Krain AD. Porokeratosis and cutaneous malignancy: A review. *Dermatologic Surg*. 1996;22(4):339-342. doi:10.1111/j.1524-4725.1996.tb00327.x
4. Weidner T, Illing T, Miguel D, Elsner P. Treatment of Porokeratosis: A Systematic Review. *Am J Clin Dermatol*. 2017;18(4):435-449. doi:10.1007/s40257-017-0271-3
5. Calay D, Vind-Kezunovic D, Frankart A, Lambert S, Poumay Y, Gniadecki R. Inhibition of akt signaling by exclusion from lipid rafts in normal and transformed epidermal keratinocytes. *J Invest Dermatol*. 2010;130(4):1136-1145. doi:10.1038/jid.2009.415

Figure 1: Disseminated superficial actinic porokeratosis on the upper extremities of a 52-year-old female (a) left (b) right. Treatment response after twice daily application of topical cholesterol 2%/simvastatin 2% cream for three months (c) left (d) right

ajd_13473_f1a.jpg

ajd_13473_f1b.jpg

ajd_13473_f1c.jpg

ajd_13473_f1d.jpg

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Al Dhafiri, M;Boeisa, AN

Title:

Localised acquired hypertrichosis of the leg after a tibial lengthening

Date:

2020-09-11

Citation:

Al Dhafiri, M. & Boeisa, A. N. (2020). Localised acquired hypertrichosis of the leg after a tibial lengthening. AUSTRALASIAN JOURNAL OF DERMATOLOGY, 62 (1), <https://doi.org/10.1111/ajd.13460>.

Persistent Link:

<http://hdl.handle.net/11343/278538>