Title page

The Bairnsdale Ulcer (*Mycobacterium ulcerans*) in an urban Victorian practice: a case series

Word count: 695

Running head: Bairnsdale Ulcer in an urban practice

Authors:

Dr Rebekka Jerjen (corresponding author)

BMSc (Hons), MChD (Dist),

Affiliated with Sinclair Dermatology, East Melbourne, Victoria, Australia

Email: Rebekka.jerjen@gmail.com

ORCHID: 0000-0002-7749-1069

Dr Ameshin Moodley

MBBS, FC Derm

Affiliated with UKZN University of KwaZulu-Natal, Durban, South Africa and Sinclair

Dermatology, East Melbourne, Victoria, Australia

Email: ameshinmoodley@gmail.com

Professor Rodney Sinclair

MBBS, MD, FACD

Affiliated with Sinclair Dermatology, East Melbourne, Victoria, Australia

Email: Rodney.Sinclair@sinclairdermatology.com.au

ORCID: 0000-0001-6751-1428

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/AJD.13361](https://doi.org/10.1111/AJD.13361)

This article is protected by copyright. All rights reserved

No funding was received for this article

No acknowledgments

No conflicts of interest

Patient consented to use of photographs

Key words: *Mycobacterium ulcerans*, Bairnsdale ulcer, Buruli ulcer, Victoria, Australia, Mornington Peninsula, infectious disease, public health

DR. REBEKKA BARBARA JERJEN (Orcid ID : 0000-0002-7749-1069)

PROF. RODNEY SINCLAIR (Orcid ID : 0000-0001-6751-1428)

Article type : Case Letter

The Bairnsdale Ulcer (*Mycobacterium ulcerans*) in an urban Victorian practice: a case series

The Buruli Ulcer, known in Australia as the Bairnsdale or Daintree Ulcer, is a locally necrotising disease of skin and soft tissue caused by *Mycobacterium ulcerans*. It is considered a neglected tropical disease by the World Health Organisation and is endemic to focal parts of coastal Victoria¹. Over the last five years Victoria has seen a steady increase in the number and severity of reported *M. ulcerans* cases^{2,3}. This coincides with a geographic expansion beyond the established endemic area of the Bellarine Peninsula into the Mornington Peninsula, South eastern bayside and Frankston region^{2,3}. Here we present a series of three cases of *M. ulcerans* diagnosed between 2018 and 2019 in a central Melbourne dermatology practice.

Case 1: A 59-year-old female was referred from a central Melbourne hospital with a lower leg ulcer of five weeks duration not responding to oral augmentin, flucloxacillin and topical mupirocin. She had recently travelled to North America and Israel and reported bushwalking at the Mornington Peninsula five months earlier. Wound swab culture and Polymerase Chain Reaction (PCR) testing were negative and no Acid-Fast Bacilli (AFB) were identified. Two punch biopsies also failed to identify a pathogen. Given the history of bushwalking in an endemic area, the lack of response to other antibiotics and the suggestive clinical presentation, we suspected *M. ulcerans* despite negative culture results. She was successfully treated with rifampicin and clarithromycin.

Case 2 (figure 1): An 18 year old female was also referred from a nearby hospital with an ulcer of 6 weeks duration which was persistently erythematous and oedematous despite intravenous and oral antibiotics (ceftriaxone, clindamycin and ciprofloxacin). On examination in our clinic there was mild cellulitis surrounding a one centimetre punched out, sloughing ulcer on the left lateral ankle. Biopsy samples were negative for culture but a PCR and histology confirmed the presence of *M. ulcerans*. She had no history of travel to or contact with an *M. ulcerans* endemic region. She was referred to local infectious diseases services for management.

Case 3: A 70-year-old male with a past history of pyoderma gangrenosum (PG) presented with a new, painful purple papule on his right calf. The lesion was initially treated as PG with prednisone, however, there was rapidly progressive localised oedema, erythema, with only minimal pain. Biopsies were sent for culture and he was commenced on roxithromycin. Histology showed an atypical lymphocytic lobular panniculitis. Culture and PCR confirmed the diagnosis of *M. ulcerans* despite no history of exposure. He was switched to clarithromycin and rifampicin and the wound was surgically debrided. The lesion healed over eight weeks.

Consistent with previous reports on *M. ulcerans*⁴, these three patients presented with symptoms of several weeks duration prior to diagnosis and all had single lesions occurring on the lower limbs. Case 1 had an incubation period of 5 months which correlates with the previously reported mean *M. ulcerans* incubation period of 4 to 5 months⁵. These cases demonstrate the recent geographical expansion of *M. ulcerans* in Victoria as Case 2 and 3 had no identifiable contact with a previously known endemic area. The reason for the expansion and increased disease severity is unclear, possible reasons discussed to date include climate change, evolution of more pathogenic strains and/or population expansion². We speculate that case 2 and 3, who both presented with symptoms in winter, had been exposed to infection during the summer. Increased outdoor activity during warmer weather and subsequent direct exposure to soil (e.g. through gardening) and/or insect vectors (when less clothing is worn) are possible sources of transmission in these cases. Furthermore, a potential zoonotic cycle with possums has been postulated to be involved in the expansion of *M. ulcerans* infections^{2,6}. These mammals are commonly seen in urban Melbourne parks

and suburbs thus could act as a reservoir for the infection which can be transmitted to humans, most likely indirectly through faeces⁶.

The infection is a public health issue and interventions are necessary to prevent transmission, however, this is challenging as the environmental reservoir and mode of transmission remains unknown. Victorian medical practices need to be vigilant when seeing patients with an atypical ulcer or nodule as prompt recognition and treatment will reduce disease severity and complications.

References

1. Maccallum P, Tolhurst JC, Buckle G, Sissons HA. A new mycobacterial infection in man. *J Pathol Bacteriol.* 1948;60(1):93-122. doi:10.1002/path.1700600111
2. Tai AYC, Athan E, Friedman ND, Hughes A, Walton A, O'Brien DP. Increased severity and spread of *Mycobacterium ulcerans*, Southeastern Australia. *Emerg Infect Dis.* 2018;24(1):58-64. doi:10.3201/eid2401.171070
3. Department of Health and Human Services. Cutaneous *Mycobacterium ulcerans* infection. <https://www2.health.vic.gov.au/about/news-and-events/healthalerts/buruli-ulcer-dec2016>. Published 2016. Accessed April 13, 2020.
4. Boyd SC, Athan E, Friedman ND, et al. Epidemiology, clinical features and diagnosis of *Mycobacterium ulcerans* in an Australian population. *Med J Aust.* 2012;196(5):341-344. doi:10.5694/mja12.10087
5. Loftus MJ, Trubiano JA, Tay EL, et al. The incubation period of Buruli ulcer (*Mycobacterium ulcerans* infection) in Victoria, Australia – Remains similar despite changing geographic distribution of disease. *PLoS Negl Trop Dis.* 2018;12(3):1-10. doi:10.1371/journal.pntd.0006323
6. Fyfe JAM, Lavender CJ, Handasyde KA, et al. A major role for mammals in the ecology of *Mycobacterium ulcerans*. *PLoS Negl Trop Dis.* 2010. doi:10.1371/journal.pntd.0000791

Author Manuscript

ajd_13361_f1.png

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Jerjen, R;Moodley, A;Sinclair, R

Title:

The Bairnsdale ulcer (*Mycobacterium ulcerans*) in an urban Victorian practice: A case series

Date:

2020-06-23

Citation:

Jerjen, R., Moodley, A. & Sinclair, R. (2020). The Bairnsdale ulcer (*Mycobacterium ulcerans*) in an urban Victorian practice: A case series. AUSTRALASIAN JOURNAL OF DERMATOLOGY, 61 (4), pp.376-377. <https://doi.org/10.1111/ajd.13361>.

Persistent Link:

<http://hdl.handle.net/11343/278549>