Spatial weighting of land use and temporal weighting of antecedent discharge improves prediction of stream condition.

Christopher J. Walsh • J. Angus Webb

C. J. Walsh • J. A. Webb

Department of Resource Management and Geography

The University of Melbourne

221 Bouverie St

Parkville 3010, Australia

e-mail: cwalsh@unimelb.edu.au

Date of manuscript revision: June 6, 2014

Manuscript word count: 8678

Abstract

Land management to protect streams requires knowing which parts of the landscape most strongly influence stream condition. Understanding how flow through landscapes and along streams affects such land-use impacts requires knowing the period of antecedent discharge that most strongly influences condition. Both considerations require determination of optimal weighting schemes for predictors of stream condition. We calculated forest cover weighted by flow-path distance to 572 urban, peri-urban, and rural sites—in the Melbourne, Australia, region—sampled for macroinvertebrates, and antecedent discharge weighted by time preceding each of 1,723 samples. Using mixed linear models that accounted for spatial dependence, we aimed to determine the weighting curve shape and length that best predicted macroinvertebrate assemblage composition. The best model was a function of mean annual discharge, weighted forest cover, weighted imperviousness, weighted antecedent discharge, and their interactions. Optimal weightings were exponential—half-decay distance 35 m overland (plausible range 26–50 m), and 1.0 km in-stream (0.75–1.3 km) for forest cover—, and linear over ≥ 4 y for antecedent discharge. Model plausibility was more affected by weighting distance than the shape of the weighting function. Regardless of weighting curve shape, riparian forest effects on macroinvertebrate assemblages are strongest within 10^1 – 10^2 m from the stream, and 10^3 m upstream. Although exponential weightings are only marginally more plausible, they are the most realistic representation of physical processes. While our conclusions should not be interpreted as recommendations for buffer widths, they provide valuable insight into the scales of influence in the region and could be used to inform management decisions.

Keywords: riparian forest; impervious; macroinvertebrate assemblages; SIGNAL score; Melbourne; Australia

Introduction

Our ability to use and manage land while protecting streams and other water resources requires knowing which parts of the landscape most strongly influence stream condition. It is widely understood that land use closer to streams is likely to have a greater impact than more distant land (e.g. King et al. 2005). Analogously, the effect of more recent flow conditions has been shown to be greater than conditions of the more distant past (e.g. Bond et al. 2012; Rolls et al. 2012). Thus when modelling the effects of land use and recent flow on stream condition, it is important to quantify land cover and flow variables in ways that capture the spatial and temporal heterogeneity of effects.

Many studies have sought to quantify the effects of land use on stream ecosystems (e.g. Allan 2004; Stephenson and Morin 2009; Peterson et al. 2011; Thomson et al. 2012). A common shortcoming of many such studies has been their quantification of land-use impacts in ways that do not adequately account for spatial arrangement of land cover, and how it is likely to influence the magnitude of effects on stream ecosystems (King et al. 2005). In most large-scale studies of the effects of land cover, land use as a proportion of catchment has been used a candidate predictor, often assessed against a small number of alternative estimates of riparian land cover based on forest cover in stream buffers of one or more widths (from the stream), and lengths (upstream from the site of interest) (e.g. Roth et al. 1996; Strayer et al. 2003; Thomson et al. 2012). In addition to the statistical problems arising from likely collinearity among proportional land cover estimates (Van Sickle and Johnson 2008; an increase in proportional cover of one land cover class necessarily results in decreases in others: Stephenson and Morin 2009), it has been noted that the concept of an abruptly defined riparian buffer beyond which the landscape has no influence on the stream is mechanistically unrealistic (King et al. 2005).

50 A more mechanistically realistic approach is to continuously weight the influence of
different land uses. King *et al.* (2005) and subsequent papers (Van Sickle and Johnson 2008;
Peterson et al. 2011) have demonstrated that weighting of rural land use by its flow-path
distance to- and along- streams using decay functions with a mechanistic underpinning,
increases the predictive power of models of stream condition compared to lumped catchment
55 proportions of land use. Use of near-stream land, and land in drainage lines in which water
accumulates (Peterson et al. 2011), is likely to be an important driver of stream condition in
landscapes in which water drains to streams through natural flow paths. However, human
alteration of drainage lines, which is particularly common in urban landscapes can greatly
increase the spatial extent of catchment land areas that have a strong influence on stream
60 condition. Walsh and Kunapo (2009) demonstrated that proximity of the most downslope
edge of impervious surfaces to urban stormwater drainage lines was a better distance
measure for weighting the influence of such surfaces on stream macroinvertebrate
assemblages than was distance to stream. They therefore argued that impervious runoff
delivered through urban stormwater drainage networks is a likely primary driver of stream
65 degradation, and that the influence of urban impervious surfaces is greatly extended beyond
the riparian zone, potentially far into the catchment. Similar effects of drainage lines on
transport of nutrients have been reported in agricultural landscapes with tile drains (Fraterrigo
and Downing 2008).

 The questions of the optimal weighting schemes for different land uses is related to
70 the question of how best to weight antecedent flow as a predictor of stream condition. The
dynamic nature of stream ecosystems means that antecedent flow conditions are an important
driver of biotic structure and function (Rolls et al. 2012). However, few large-scale studies
have quantified antecedent flow conditions as a predictor of stream condition (Bond et al.
2012).

75 In this paper we aim to identify the most plausible weighting model (shape) and
parameterization (length) for forest cover and antecedent discharge as predictors of
macroinvertebrate assemblage condition in streams of the Melbourne region, Australia. We
also aim to assess if the optimal measure of weighted imperviousness derived by Walsh and
Kunapo (2009) for an eastern portion of our study region remained a superior predictor to
80 total imperviousness for the entire region. Assemblage condition was quantified using
SIGNAL score, a macroinvertebrate biotic index (average score per taxon), which is widely
used in south-eastern Australia. SIGNAL score was developed as an index of water quality
impairment (Chessman 1995), it has been shown to be a strong univariate correlate of
assemblage compositional similarity across gradients of land-use disturbance area in our
85 study region (Walsh 2006). It is therefore a useful measure of the response of
macroinvertebrate assemblages to human disturbance in the Melbourne region.

 We used a large set of macroinvertebrate data across the region surrounding the
metropolis of Melbourne in south-eastern Australia collected over 17 years. We used linear
mixed models and tested optimal weightings across a wide range of potential model
90 structures, including other variables that characterize the physiographic nature of sites to
ensure the robustness of our conclusions. Our primary hypothesis was that the accuracy of
the prediction of SIGNAL score would vary with the distance-weighting functions of forest
cover and impervious area, as well as the temporal weighting function of antecedent
discharge. Our results confirm this, showing that better predictions of stream ecological
95 condition arise from models that take into account the greater influence of more proximate
land use and more recent flow conditions. Incorporation of such information into predictive
models will improve our understanding of land use impacts upon freshwaters.

Methods

Overview

100 We used multi-model inference (Burnham and Anderson 2002) to assess the relative
plausibility of a wide range of models for predicting SIGNAL score across the Greater
Melbourne region. The models used up to 10 predictor variables derived from multiple
sources, which fall naturally into 4 different classes (Table 1). Three variables (process,
number of riffle sample units per sample, number of spring sample units per sample) affect
105 the likelihood of capture of different macroinvertebrate taxa. Four variables (mean annual
discharge, catchment area, elevation, igneous geology) accounted for physiographic variation
across the region likely to affect macroinvertebrate assemblages through effects on water
chemistry and hydrology. Antecedent discharge accounted for temporal hydrologic variation
among samples. We aimed to ensure that influences of human impact were restricted to the
110 land-use variables, which were quantified as imperviousness (as an indicator of urbanization),
and forest cover (as an indicator of land clearance). We sought optimal weightings for land
use variables and antecedent discharge. The other variables were intended as classifiers of
stream or sample type: thus we used modeled measures of discharge, assuming no human
impact. We used mean annual discharge and antecedent discharge as indicators of the
115 general hydrologic state of the streams in the absence of human impacts, and land use
variables as indicators of potential changes to habitat quality and hydrologic patterns.

#Table 1 approximately here#

For forest cover, imperviousness and antecedent discharge, we tested the relative
plausibility of different weighting schemes for these predictor variables in our models. For
120 impervious runoff, we simply compared the plausibility of total imperviousness (TI, a simple
measure of urban density) and attenuated imperviousness (AI, imperviousness weighted by
flow distance to stormwater drains, and thus a more direct indicator of impervious runoff, as

derived by Walsh & Kunapo, 2009). For forest cover, we compared the plausibility of attenuated forest cover calculated using 3 weighting models (exponential decay, linear decay and threshold), across a wide range of decay and threshold distances (to the nearest stream) for each model. For antecedent discharge, we compared its plausibility when calculated for different periods (6 mo to 5 y), either simply summed or weighted so that more recent monthly discharges have greater weight. Full methodological details are provided in Appendices 1 and 2.

Study region

The Melbourne region (Fig. 1) is physiographically diverse, ranging from dry western grassland-dominated basalt plains (mean rainfall 400–500 mm/y) with open *Eucalyptus* woodland along stream valleys, to increasingly wet uplands further east, rising to a mosaic of tall wet-sclerophyll *Eucalyptus* forests and *Nothofagus* rainforests in upland valleys (mean rainfall up to 2000 mm/y). Melbourne (population ~4.5 million) is surrounded by rural lands in which the dominant land use is pasture and non-irrigated cropping. Small areas of the region are used for intensive horticulture, totaling ~4% of the total land area (Melbourne Water et al. 2013). Forest, primarily with *Eucalyptus* spp. as the dominant tree, dominate the uplands of the east and northwest, but elsewhere occur as patches among agricultural and urban land uses (Fig. 1B). The upper Yarra River in the east of the region and some other upland streams of the region are impounded for Melbourne's water supply, but as the river or its tributaries are not used as water supply conduits, seasonal patterns of discharge in downstream waters are largely unchanged by abstraction (Walsh et al. 2007).

#Fig. 1 approximately here#

Data sources

We used five primary data sources:

1. The primary dataset and the dependent variable, SIGNAL score, were derived from the Melbourne Water macroinvertebrate database, which contains data collected from sites across the Greater Melbourne region (Fig. 1A) since 1992 (Melbourne Water,
150 unpublished data).
2. Hydrologic and geological variables were derived from the surface network and catchments data of the Australian hydrological geospatial fabric, a national dataset derived from a 9-second (~220 x 270 m in the study region) digital elevation model, detailing spatial relationships between streams and their catchments (hereafter termed the
155 geofabric: Bureau of Meteorology 2011).
3. Forest cover data were derived from a land-use dataset for the Greater Melbourne region, compiled by Melbourne Water from two sources (Department of Planning and Community Development 2010; Department of Primary Industries 2011).
4. Imperviousness measures and catchment area were derived from 2004 maps of
160 impervious coverage and an associated nested dataset of stream reaches and catchments in the region (Grace Detailed-GIS Services 2012). While Melbourne is growing and expanding, urban infill and expansion primarily over the period of study resulted in only marginal increases in catchment imperviousness in already substantially urban streams, with no expansion into undeveloped catchments that
165 were subsequently sampled.
5. A 10-m-resolution DEM for the Greater Melbourne region (J Kunapo, personal communication) was used to standardize the resolution of the multiple data sources for calculation of predictor variables.

Compilation of the biological dataset

170 We used samples collected before April 2009, the most recent date of hydrologic data
in the geofabric: although parts of the region were affected by wildfires in February 2009,
none of the samples in the data were from fire-affected sites. All samples were collected by
rapid bioassessment methods (RBA: Anon. 1994) either from riffles or pool edges, and either
in autumn (Feb – Jun) or spring (Sep – Dec). 84% of samples were sorted using a standard
175 30-min sort in the field, and 16% were subsampled in the laboratory, and sorted to 10% or
200 individuals, whichever was greater.

Typically two samples were collected at each site on a single day: either one from a
riffle and a second from a pool edge, or two edge samples in the absence of a riffle. To
reduce the effects of sampling error of RBA samples, data from pairs of samples were
180 combined to produce presence-absence data for sample-pairs. 1,382 sample-pairs were
collected on the same date (or for 13 of them, within a month of each other). A further 655
sample-pairs were collected from one site in different seasons of the same year. Sample pairs
are hereafter referred to as samples (and individual samples as sample units).

All data were standardized to family-level identification, except Chironomidae, which
185 were identified to sub-family, and Acarina, Oligochaeta, Polychaeta and Nemertinea, which
were not identified to finer levels. SIGNAL was calculated for each sample using the grades
of EPA Victoria (2003). SIGNAL is the average of ratings (1–10) assigned to
macroinvertebrate families found at a site, with higher ratings indicating more pollution-
sensitive families.

190 While the distribution of sites was well spread across the region, many sites were
close to other sites (Fig. 1). A temporal trend analysis of SIGNAL scores in a subset of our
study sites found no evidence of spatial autocorrelation among them (Webb and King 2009),
but this remains a possibility. To account for any effects of spatial autocorrelation in this

data set, we allocated adjacent sites into groups along the same stream, conservatively ensuring that all members of each group were >5 river-km from sites in any other group (Lloyd et al. 2005). This grouping rule resulted in the exclusion of samples from 136 sites that fell between site groups separated by <10 km. This left 1,723 samples, spread uniformly over the study period (Fig. 2A), from 572 sites in 343 site groups, each containing 1-35 samples to be used in the statistical modeling.

#Fig. 2 approximately here#

Compilation of predictor variables requiring no further calculation

Elevation of each site (m) was calculated from the 10-m DEM. Catchment area (km²) of each site was extracted from the imperviousness dataset. Three variables describing sample characteristics were derived from the macroinvertebrate database:

1. Number of spring sample units per sample (nspring = 0, 1 or 2). Season of collection potentially affects macroinvertebrate assemblage composition as a result of variability in phenology among taxa.
2. Number of riffle samples (nriff = 0, 1, or 2). Macroinvertebrate taxa differ in their tendency to occur in riffle and edge habitats, which potentially has an effect on SIGNAL score.
3. Method by which the samples were sorted (process = field or lab). RBA samples sorted in the field are likely to be biased against some taxa (Humphrey et al. 2000), with potential effects on SIGNAL score.

Two variables were extracted from the geofabric:

1. Mean annual discharge depth in the absence of human impacts (meanQ, mm/y: mean annual accumulated surface water surplus—derived from a simple water balance model using long-term rainfall and potential evapotranspiration data, and strongly correlated with discharge of unregulated rivers of the region (Stein et al. 2002)—

divided by catchment area). MeanQ, as a catchment-standardized measure of annual
stream discharge, distinguishes flow regimes among streams that rise in the drier
western and lowland parts of the region from those that rise in wetter eastern and
upland parts (Fig. 1C).

2. Proportion of catchment area underlain by igneous rock (igneous geology). This
variable is a correlate of electrical conductivity of stream waters across the Melbourne
region (Walsh et al. 2001).

Compilation of variables for optimal weighting searches

a) Antecedent discharge

Antecedent discharge (of different periods) for each sample was calculated from the
geofabric estimates of monthly accumulated surface water surplus in each segment in the
period before each sample. We calculated two weighting schemes for antecedent discharge:

1. unweighted antecedent discharge for x months, calculated as $\frac{\sum_{i=1}^x Q_i}{x Q_{mean.ann}}$, where Q_i is
the discharge in the i th month before the sample date. Dividing the sum of those
discharge values by the mean annual discharge ($Q_{mean.ann}$) adjusted by the number
of months summed expresses the result as a proportion of the long-term mean.
2. linearly weighted antecedent discharge for x months similarly as $\frac{\sum_{i=1}^x Q_i \left(1 + \frac{i}{x}\right)}{x Q_{mean.ann}}$.

b) Forest cover

Our decision to use forest cover as a variable portraying (the converse of) human land
use in the region resulted from preliminary analyses in which we sought to predict SIGNAL
score using a range of agricultural land classes. The patchiness and relative rarity of intensive
agricultural practices in the region resulted in no greater predictive power from multiple
agricultural classes than from a single class of cleared land (or its converse, forest cover).

The land-use data did not permit us to distinguish forest types across the region in our analyses.

To calculate weighted forest cover, we converted forest cover data from the MW land-use dataset and imperviousness data from the impervious dataset to 10-m rasters to match the flow-distance data calculated from the 10-m DEM (see Appendix 1 for more details). We used a distance weighting model after Van Sickle and Johnson (2008), assuming that the influence of forest cover in a grid cell is a non-increasing function of the flow path distance between that cell and the sampling site. Thus the influence of a cell at the sample site is 1, while a more distant cell has a fractional influence, $f_L(d_L, \alpha_L) f_W(d_W, \alpha_W)$, where f_L and f_W are fractions determined by to-stream flow distance d_L and in-stream flow distance d_W (Fig. 3), and the parameters α_L and α_W , which control the degree of attenuation over a set distance.

#Fig. 3 approximately here#

The cumulative distance-weighted forest cover, $F(\alpha_L \alpha_W)$, for the catchment of each site is estimated by dividing the sum of the weighted fractional influence of all forested grid cells ($C_i = 1$ if forested, 0 if not) by the weighted fractional influence of all grid cells.

$$F(\alpha_L \alpha_W) = \frac{\sum_i C_i f_L(d_{Li}, \alpha_L) f_W(d_{Wi}, \alpha_W)}{\sum_i f_L(d_{Li}, \alpha_L) f_W(d_{Wi}, \alpha_W)} \quad \text{eqn. 1}$$

We compared the plausibility of models using distance-weighted forest cover determined by one of three one-parameter weighting functions. These have been considered as explanatory models for distance-attenuated effects in past studies (Van Sickle and Johnson 2008), and span the range of attenuation shape. For example, curves with the most rapid decline near $d = 0$ could portray the mechanisms of pollutant or water uptake or loss along flow paths; whereas those with the least rapid decline near $d = 0$ are similar to those used widely in the assessment of fixed buffer widths.

1) exponential decay, $f(d) = \exp(-\alpha d)$ eqn. 2

2) linear decay to zero, $f(d) = \max(1 - \alpha d, 0)$ eqn. 3

3) threshold, $f(d) = 1$ if $d \leq \alpha$, else $f(d) = 0$ eqn. 4

265 Rather than reporting the values of α_L and α_W , for exponential and linear weighting, we report the half-decay distance (HD): the distance at which a grid cell would have a weighting of 0.5. For exponential decay $HD = -\ln(0.5)/\alpha$, for linear decay, $HD = 0.5\alpha$. For threshold, we report the threshold value, α

c) Imperviousness

270 We compared (unweighted) TI with the most plausible value of (exponentially weighted) AI derived by Walsh and Kunapo (2009) for streams of eastern Melbourne (Fig. 3B). AI is a measure of the influence of impervious surfaces as determined by the anthropogenic stormwater drainage system associated with urban land. The formulation of AI differs from the weighting schemes we apply to forest cover here in two important ways.

275 First the weighting distance used in AI is the overland flow distance from the most downstream point of an impervious surface to the nearest stormwater drain (or stream if the flow path does not cross a drain: Fig. 3A). AI is partly a measure of altered land use, but also a measure of alteration to the hydrologic network. To capture differential effects of altered catchment flow paths among catchments, the denominator of AI is the total catchment area,

280 rather than the weighted fractional influence of all grid cells as in eqn. 1.

Statistical models

Selection of candidate models

We used mixed linear models that accounted for spatial dependence to assess whether models incorporating weighted measures of imperviousness, forest cover and antecedent

285 discharge were more plausible than models with unweighted measures. We included other

predictor variables as described earlier to account for climatic and topographic variability and differences among sample types, independent of the three variables to be weighted. We thus sought to compare the plausibility of a subset of models that combined up to ten predictor variables and their interactions.

290 Candidate models were selected on the following basis. We expected that SIGNAL scores would be predicted well by a combination of forest cover, imperviousness, meanQ (Fig. 1B, C), antecedent discharge (Fig. 2B), and interactions between these variables. We also expected that some variation in SIGNAL would be explained by stream size (indicated by catchment area), elevation (combining the effects of stream size and temperature), igneous
295 geology and the characteristics of the samples (sample type variables in Table 1). Models tested therefore included all four of the main predictor variables and their interactions, at least one of catchment area, elevation and igneous geology, and at least one sample type variable (Table 2).

#Table 2 approximately here#

300 *Comparisons of model plausibility*

 We used the Akaike Information Criterion adjusted for small sample size (AIC_c) to assess the relative plausibility of alternative models of SIGNAL score. We report ΔAIC_c , the difference between a model's AIC_c and that of the overall best model (i.e. ΔAIC_c of the best model = 0). Models with lower AIC_c are more plausible, but models with $\Delta AIC_c \leq 2$ are
305 considered equally plausible as the best model. ΔAIC_c of 4–7 indicates that the model with the lower AIC_c is superior, with $\Delta AIC_c > 10$ indicating that the model with the lower AIC_c is strongly preferred (Burnham and Anderson 2002).

 We took an iterative approach to testing the effect of different weighting schemes for forest cover, imperviousness and antecedent discharge. All models tested included meanQ,
310 one variant of each of imperviousness, forest cover and antecedent discharge, and all

interactions of these four variables (Table 2). All model structures were first tested using unweighted forest cover, total imperviousness, and unweighted 2-y antecedent discharge, as an arbitrary starting model. The most plausible model structure was then used to determine the optimal antecedent discharge measure. This measure was then used in all candidate models and the most plausible model structure was identified again. The most plausible model structure was then used to assess which of unweighted imperviousness (TI) or weighted imperviousness (AI) was the better measure of imperviousness, and that measure was used in all candidate models to identify the most plausible model structure again. This structure was then used to identify the most plausible forest measure, and finally this measure was used in all model structures, to assess the most plausible model structure.

At each iteration, we also assessed models with each of the third- and fourth-order interactions of the four main variables removed: if their removal did not increase AIC_c by >2 , they were removed. The change in weighted variables did not change the interactions removed at this stage in any iteration.

For antecedent discharge, we assessed a) unweighted and b) linearly weighted antecedent discharge for 6, 12, 24, 30, 36, 64, and 72 months. For each weighting function for forest, we calculated $F(\alpha_L \alpha_W)$ for a wide range of values of α_L and α_W . We began with a range of values spaced approximately exponentially (e.g. HD_L (for linear and exponential decay) or α_L (for threshold) = 1, 2, 4, 8, 16, 32, 64, 128, 360, 640 m, and HD_W or α_W = 100, 300, 1000, 3000, 10000, 30000). We then used the region of lowest AIC to test ranges of α at finer scales.

Mixed model structures and assumptions

We transformed variables as necessary to reduce leverage of high values (Table 1). Collinearity among predictor variables was low (variance inflation factors ≤ 3 : Zuur et al.,

2009). The variables meanQ, forest cover, imperviousness, antecedent discharge, elevation, catchment area, igneous geology, process, nspring, and nriff were considered fixed.

The effect of sample groups was incorporated as a random effect. For all fixed effect model structures, we compared a simple linear model with two mixed models each with a different random component: a random intercept determined by site group with constant slope; and a random intercept determined by site group, and slope determined by antecedent discharge (Zuur et al. 2009). The first random structure assumes that SIGNAL score can vary among site groups, but that the modeled relationship with the fixed effects has the same slope. This model accounts for any potential spatial autocorrelation of samples within site groups. The second random structure assumes that the modeled relationship with fixed effects can vary with antecedent discharge among site groups. Models were calculated with the ‘nlme’ package of R (R Core Team 2013), using restricted maximum likelihood. We checked that model assumptions were met following the protocols of Zuur et al. (2009).

Results

Most plausible model structures

Mixed models with a random intercept determined by site group were consistently much more plausible than linear models with the same fixed effect structure but not accounting for dependence among site groups ($\Delta AIC_c > 300$ in all cases). Models with a random intercept determined by site group, and slope determined by antecedent discharge were more plausible again ($\Delta AIC_c > 30$ in all cases). The random intercept and slope structure was thus used in all models for assessment of optimal weighted variables.

The structure of fixed model effects had a relatively small effect on AIC compared to weighting of forest, imperviousness and antecedent discharge (Table 1). The most plausible model structure included elevation, igneous geology, and process in addition to the four main variables. The only 3- or 4-way interaction that improved AIC was the interaction between

meanQ, antecedent discharge and weighted forest cover. This final model structure was substantially more plausible than any other model structure or any model that used unweighted variables, and was used for assessment of optimal weighted variables.

Most plausible weighting schemes for antecedent discharge, imperviousness and forest cover

The most plausible measures of antecedent discharge were linearly weighted, with all antecedent periods ≥ 4 y being equally plausible (Fig. 4). These weighted measures of antecedent discharge were all substantially more plausible than unweighted antecedent discharge of any period. Attenuated (weighted) imperviousness was a much more plausible predictor than total (unweighted) imperviousness in all models tested (Table 1). The most plausible measure of forest cover was exponentially weighted with half-decay distances of 35 m overland and 1.0 km in-stream, with ranges of equal plausibility ($\Delta AIC_c < 2$), of 26–50 m and 0.75–1.3 km, respectively (Fig. 5, 6). Optimal distances for the three decay functions were consistent: optimal half-decay distances were similar for exponential and linear decay, and about half the threshold values. There was strong variation in ΔAIC_c with varying distance parameters (Fig. 5, 6), but less difference between the weighting functions (Fig. 5).

The final model strongly predicted SIGNAL score, with the Pearson correlation coefficient of 0.955 between fitted and observed values.

#Figs 4, 5 and 6 approximately here#

Discussion

Our results suggest that the influence of forests on in-stream macroinvertebrate assemblage composition is greatest along a riparian corridor extending 10^1 – 10^2 m inland and 10^3 m upstream. Similarly, extending the inference of Walsh & Kunapo (2009) to the broader geographic context of this study, the influence of urban impervious runoff is strongest within 10^1 m of a stream, but this influence usually is extended far into upland parts of catchments by extensive stormwater drainage systems (e.g. Fig. 2B). Thus while rural and

urban land uses may have widely different scales of influence, the flow distances over which their impacts can be moderated are comparably short.

Our finding of an optimal antecedent discharge length of ≥ 4 years is likely to be at least partly a function of the timing of our study, over a period of increasing dryness, encompassing a decadal drought (Fig. 2; Bond et al. 2008). This prevents us from

generalizing this finding to periods of increasing wetness. The interaction between meanQ, attenuated forest cover and antecedent discharge (data ranges provided in Table 1) in the best model suggests that the influence of riparian forests differs between streams with different annual discharges, and between periods of different dryness. Such interactions point to new approaches to be explored in future studies, described below.

Optimal weighting of land cover

Exponential attenuation was the most plausible weighting function, and arguably the most mechanistically meaningful of the three functions, particularly in modeling the behavior of contaminant transport (King et al. 2005). However, as found by Van Sickle and Johnson (2008), the optimal parameterizations of the three functions did not differ strongly in their plausibility. In contrast, all three functions varied widely in plausibility with different decay lengths (e.g. half-decay distances $< \sim 540$ m and $> \sim 1750$ m in-stream, and $< \sim 15$ m and $> \sim 200$ m overland were substantially less plausible than the best model). Such a strong influence of parameterization (i.e. weighting distance) suggests that comparing different weighting models (i.e. shape) using a single parameterization for each scheme (e.g. Peterson et al. 2011) is unlikely to provide a reliable discrimination of different mechanisms.

A decline in influence of forest cover to near-zero over 10^2 m (as suggested by a half-decay distance of 26–50 m, Fig. 5) is consistent with the findings of Van Sickle and Johnson (2008), who used similar methods to predict fish assemblage composition. However, they found that the influence of land use on fish assemblages extended for many km upstream,

410 further than we found for macroinvertebrate assemblages. As they posited, the extent of in-stream influence might be shorter for macroinvertebrate assemblages, which are likely to be less mobile and shorter lived than are fish.

The strong influence of riparian forest cover suggested by our analyses concurs not only with recent similar studies (Van Sickle and Johnson 2008; Sheldon et al. 2012), but with
415 the broader field of ecological research identifying the strong influence of riparian zones on the ecological structure and function of streams (Naiman and Décamps 1997; Pusey and Arthington 2003). However, it contrasts with the conclusions of Stephenson and Morin (2009), who reported that catchment forest cover independently explained more variation in a range of structural in-stream ecological measures than did buffered riparian forest cover. Our
420 model produced a much stronger prediction of ecological response than did any of the models of Stephenson and Morin (2009), pointing to the importance of including a range of predictor variables that are likely to be driving variation unexplained by forest cover, as the primary predictor variable of interest. Other studies have reported varied in-stream ecological and water quality responses to riparian vs catchment land uses (e.g. Strayer et al. 2003; Uriarte et
425 al. 2011), but these studies are not easily comparable to ours because they did not clearly distinguish the effects of urban and agricultural land use, nor consider differences in drainage among land uses.

The continued widespread use of lumped measures of land use (e.g. TI), which do not adequately represent the effects of catchment drainage networks, hampers advancement of
430 ecological knowledge and the ability of ecologists to advise managers on the most appropriate actions to reduce land-use impacts on stream ecosystems, particularly urban land use. For instance, arguments about threshold effects of urban land use (Cuffney et al. 2011; King and Baker 2011) are limited by the use of lumped measures that sub-optimally represent the stressor driving the threshold response. Sheldon *et al.* (2012) found that no single scale

of urban land cover was a strongly preferred predictor of in-stream ecological health in south-east Queensland, to which they attributed the likely importance of the broader catchment connectivity of urban catchments with stormwater drainage networks: a mechanism that was not captured by their urban land cover measures. The superiority of AI over TI as a predictor of SIGNAL score that we have demonstrated across the Melbourne region reinforces the importance of drainage networks in driving the urban degradation of stream ecosystems.

Optimal measure of antecedent discharge

The influence of antecedent discharge of ≥ 4 years on SIGNAL score suggests a shift in assemblage composition with increasing dryness over the period of study. The latter decade of the study period was the worst drought since European settlement of the region (Bond et al. 2008). While our large-scale assessment of antecedent discharge cannot provide information on the mechanisms by which drought affects stream biota, our determination of an optimal measure of antecedent discharge points to the temporal scale over which supra-seasonal droughts impact streams in our study region. Weighted antecedent discharge suggests that discharge of most recent months is of greatest influence on macroinvertebrate assemblages. The optimal length of ≥ 4 years suggests that the cumulative effects of a supra-seasonal drought contribute to assemblage composition over at least 4 years.

As our period of study did not encompass a wetter period following the drought, we are unable to infer if the recovery of macroinvertebrate assemblage composition is slower than the rate of change observed during the drying period (Bond et al. 2008). Thus, the broader relevance of a ≥ 4 -year weighted measure of antecedent discharge requires re-assessment using data collected during multiple supra-seasonal periods of drying and wetting.

Further research directions

The possibility that the influence of riparian forests differs between stream types and between periods of differing dryness points to a potential shortcoming of our work: the fixed

460 definition of the stream network, on which to-stream and in-stream distances are based. Our
chosen threshold (catchment area = 1 km²) was based on our observations of the catchment
area of the smallest perennial streams in the eastern part of region. It is likely that regional
gradients of climate and stream discharge mean that the threshold for stream formation varies,
just as it does with antecedent discharge (Baker et al. 2007). Peterson et al. (2011) addressed
465 this shortcoming by adjusting weighting distances by flow accumulation, to allow for less
attenuation of effect as flow paths lengthen. In the earlier development of this study, we
found that similar flow-accumulation weighting approaches did not optimize within a defined
range of decay distances. We hypothesise that weighting narrow (1 cell) drainage lines
increases the sensitivity of results to minor errors in land-use maps.

470 To address the problems of defining the interface between overland and in-stream
flow, we propose two future refinements. Firstly, to account for regional variability in
threshold area, we propose using stream discharge rather than catchment area as the criteria
for determining the threshold area for stream definition. To avoid potential problems with
using flow-accumulation to alter decay distance, we propose using a stream network with a
475 very small threshold discharge, with each reach of the network classified into discharge
ranges. To-stream decay distances can then be altered as a function of reach class.

Our primary objective was to quantify the most plausible average weighting schemes
for a stream condition index, using an analysis across a physiographically diverse region.
Although we have accounted for broad physiographic patterns across the region, and our
480 results accord well with similar research elsewhere, our model was developed only for a
region encompassing several large watersheds. Hence, it remains possible that covariance
between stream types and land use could have biased our results, or obscured variation across
the region in the spatial extent of land use influences. Moreover, dimensions of riparian
forests with greatest influence on stream ecosystems are likely to vary with stream size,

topography, and long-term and antecedent climate (Richardson et al. 2012). Therefore our conclusions should not be interpreted as firm recommendations for a fixed buffer width for all streams across the region; further meta-scale analyses (e.g. Cuevas et al. 2006) would be needed to assess the generality of our results across larger scales, and their ability to be extrapolated to watersheds not included in the analysis.

In addressing our primary purpose, we developed a new approach to determining optimal spatial weights that produced excellent results, and is applicable to any stream network with reasonable land use data, hydrological data, and digital elevation models. The performance of our optimal model in detecting the effects of land use on stream macroinvertebrates argues strongly for the uptake of our approach in other locations. Finally, the contribution of antecedent discharge to model performance reminds us that even with severe land use impacts, stream macroinvertebrate assemblage composition is still largely driven by flow regime.

Acknowledgements

This study was funded by Melbourne Water and CJW by the Melbourne Waterway Research-Practice Partnership. We thank Edward Tsyrlin and Rhys Coleman for commissioning the work, and for their support and patience; Alistair Danger for assistance with compiling land use data; and Joshphar Kunapo for compiling the digital elevation models used in our analyses. The manuscript was improved by suggestions of Nick Bond, Matthew Burns, Sarah Gergel, Michael Sammonds, Edward Tsyrlin and four referees.

References

- Allan JD (2004) Landscapes and riverscapes: the influence of land use on stream ecosystems. *Ann. Rev. Ecol. Evol. Syst.* 35:257–284
- Anon. (1994) National River Processes and Management Program. Monitoring River Health Initiative. River Bioassessment Manual. Version 1.0. Department of the Environment, Sport

510 and Territories; Land and Water Resources Research and Development Corporation;
Commonwealth Environment Protection Authority, Canberra

Baker ME, Weller DE, Jordan TE (2007) Effects of stream map resolution on measures of
riparian buffer distribution and nutrient retention potential. *Landscape Ecol.* 22:973–992

Bond N, Thomson J, Reich P (2012) Macroinvertebrate responses to antecedent flow, long-
515 term flow regime characteristics and landscape context in Victorian rivers. National Water
Commission, Canberra

Bond NR, Lake P, Arthington AH (2008) The impacts of drought on freshwater ecosystems:
an Australian perspective. *Hydrobiologia* 600:3–16

Bureau of Meteorology (2011) Australian hydrological geospatial fabric (geofabric) product
520 guide. Version 2.0 – November 2011. Australian Government, Bureau of Meteorology,
Canberra

Burnham KP, Anderson DR (2002) Model selection and multimodel inference: a practical
information-theoretic approach. Springer, New York

Chessman BC (1995) Rapid assessment of rivers using macroinvertebrates: a procedure
525 based on habitat-specific sampling, family level identification and a biotic index. *Aust. J.*
Ecol. 20:122–129

Cuevas JG, Soto D, Arismendi I, Pino M, Lara A, Oyarzún C (2006) Relating land cover to
stream properties in southern Chilean watersheds: trade-off between geographic scale, sample
size, and explicative power. *Biogeochemistry* 81:313–329

530 Cuffney TF, Qian SS, Brightbill RA, May JT, Waite IR (2011) Response to King and Baker:
limitations on threshold detection and characterization of community thresholds. *Ecol. Appl.*
21:2840–2845

- Department of Planning and Community Development (2010) Planning schemes online: Victoria's planning schemes. The State of Victoria, Melbourne. Available from
535 <http://planningschemes.dpcd.vic.gov.au/>
- Department of Primary Industries (2011) Victorian resources online: land use. State of Victoria, Melbourne. Available from
<http://vro.dpi.vic.gov.au/dpi/vro/vrosite.nsf/pages/landuse-home>
- EPA Victoria (2003) Rapid bioassessment methodology for rivers and streams. Guideline for
540 environmental management, Publication No. 604.1. Environment Protection Authority Victoria, Melbourne
- Fraterrigo JM, Downing JA (2008) The influence of land use on lake nutrients varies with watershed transport capacity. *Ecosystems* 11:1021-1034
- Grace Detailed-GIS Services (2012) Directly connected imperviousness compilation for
545 Melbourne Water selected catchments. Report prepared for Melbourne Water. Melbourne
- Humphrey CL, Storey AW, Thurtell L (2000) AUSRIVAS: operator sample processing errors and temporal variability - implications for model sensitivity. In: Wright J. F., Sutcliffe D. W., Furse M. T. (eds), *Assessing the Biological Quality of Fresh Waters: RIVPACS and Other Techniques*. Freshwater Biological Association, Ambleside, UK, pp. 143–163
- 550 King RS, Baker ME (2011) An alternative view of ecological community thresholds and appropriate analyses for their detection: comment. *Ecol. Appl.* 21:2833–2839
- King RS, Baker ME, Whigham DF et al (2005) Spatial considerations for linking watershed land cover to ecological indicators in streams. *Ecol. Appl.* 15:137–153
- Lloyd NJ, Mac Nally R, Lake PS (2005) Spatial autocorrelation of assemblages of benthic
555 invertebrates and its relationship to environmental factors in two upland rivers in southeastern Australia. *Divers. Distrib.* 11:375–386

Melbourne Water, Department of Environment and Primary Industries, The University of Melbourne (2013) Improving our understanding of water sensitive farm design pollution treatment systems: final position framework–May 2013. RMCG, Camberwell, Victoria

560 Naiman RJ, Décamps H (1997) The ecology of interfaces: riparian zones. *Ann. Rev. Ecol. Syst.* 28:621–658

Peterson EE, Sheldon F, Darnell R, Bunn SE, Harch BD (2011) A comparison of spatially explicit landscape representation methods and their relationship to stream condition. *Freshwater Biol.* 56:590–610

565 Pusey BJ, Arthington AH (2003) Importance of the riparian zone to the conservation and management of freshwater fish: a review. *Mar. Freshwater Res.* 54:1–16

R Core Team (2013) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, <http://www.R-project.org>

Richardson JS, Naiman RJ, Bisson PA (2012) How did fixed-width buffers become standard
570 practice for protecting freshwaters and their riparian areas from forest harvest practices? *Freshwater Sci.* 31:232–238

Rolls RJ, Leigh C, Sheldon F (2012) Mechanistic effects of low-flow hydrology on riverine ecosystems: ecological principles and consequences of alteration. *Freshwater Sci.* 31:1163–1186

575 Roth NE, Allan JD, Erickson DL (1996) Landscape influences on stream biotic integrity assessed at multiple spatial scales. *Landscape Ecol.* 11:141–156

Sheldon F, Peterson EE, Boone E, Sippel S, Bunn SE, Harch BD (2012) Identifying the spatial scale of land use that most strongly influences overall river ecosystem health score. *Ecol. Appl.* 22:2188–2203

- 580 Stein JL, Stein JA, Nix HA (2002) Spatial analysis of anthropogenic river disturbance at regional and continental scales: identifying the wild rivers of Australia. *Landscape Urban Plan.* 60:1–25
- Stephenson JM, Morin A (2009) Covariation of stream community structure and biomass of algae, invertebrates and fish with forest cover at multiple spatial scales. *Freshwater Biol.* 54:2139–2154
- 585 Strayer DL, Beighley RE, Thompson LC et al (2003) Effects of land cover on stream ecosystems: Roles of empirical models and scaling issues. *Ecosystems* 6:407–423
- Thomson JR, Bond NR, Cunningham SC et al (2012) The influences of climatic variation and vegetation on stream biota: lessons from the Big Dry in southeastern Australia. *Global Change Biology* 18:1582–1596
- 590 Uriarte M, Yackulic CB, Lim Y, Arce-Nazario JA (2011) Influence of land use on water quality in a tropical landscape: a multi-scale analysis. *Landscape Ecol.* 26:1151–1164
- Van Sickle J, Johnson CB (2008) Parametric distance weighting of landscape influence on streams. *Landscape Ecol.* 23:427–438
- 595 Walsh CJ (2006) Biological indicators of stream health using macroinvertebrate assemblage composition: a comparison of sensitivity to an urban gradient. *Mar. Freshwater Res.* 57:37–47
- Walsh CJ, Kunapo J (2009) The importance of upland flow paths in determining urban effects on stream ecosystems *J. N. Am. Benthol. Soc.* 28:977–990
- 600 Walsh CJ, Sharpe AK, Breen PF, Sonneman JA (2001) Effects of urbanization on streams of the Melbourne region, Victoria, Australia. I. Benthic macroinvertebrate communities. *Freshwater Biol.* 46:535–551

Walsh CJ, Waller KA, Gehling J, Mac Nally R (2007) Riverine invertebrate assemblages are degraded more by catchment urbanization than by riparian deforestation. *Freshwater Biol.*

605 52:574–587

Webb JA, King EL (2009) A Bayesian hierarchical trend analysis finds strong evidence for large-scale temporal declines in stream ecological condition around Melbourne, Australia.

Ecography 32:215–225

Zuur AF, Ieno EN, Walker N, Saveliev AA, Smith GM (2009) Mixed effects models and

610 extensions in Ecology with R. Springer Science+Business Media

Table 1. Fixed-effect variables, their classes as considered in the text, their codes (used in Table 2) and transformations used in mixed linear models, their source, derivation, and range among the macroinvertebrate samples. Sources: geofabric (Bureau of Meteorology 2011); DCI dataset (Grace Detailed-GIS Services 2012); 10-m DEM (J Kunapo personal communication); MW land-use data (Department of Planning and Community Development 2010; Department of Primary Industries 2011).

Variable	Class	Code	Transformation for analysis	Source	Derivation	Range
Mean annual discharge depth (mm)	Physiographic	meanQ	square-root	geofabric	RUNANNMEAN/CATAREA (variables in geofabric lookup tables "run" and "terrain" respectively)	4–790
Imperviousness (total, attenuated)	Land use	I	log10	DCI dataset	1-3-m DEM and 2004 20-cm resolution imagery	TI: 0–0.56 AI: 0–0.42
Forest cover (many weighted variations)	Land use	F	none	10-m DEM and MW land-use data	Sum of distance-weighted area of gridcells with forest cover divided by sum of distance weighted area of all gridcells	0–1 (under all weighting schemes)
Antecedent discharge (many weighted variations)	Temporal hydrologic	afi	square-root	geofabric	Weighted sum of monthly RUN values for the x- months before each sample/RUNANNMEAN (variables in geofabric look-up table "runmmm_yy")	0–2 (when spanning >72 mo) to 0–2.6 (for unweighted 6-mo) or 0–5.8 (for weighted 6-mo)
Catchment area (km ²)	Physiographic	carea	log10	DCI dataset	1-3-m DEM	0.8–2850

Elevation (m)	Physiographic	elev	fourth-root	DCI dataset	1-3-m DEM	0–900
Igneous geology	Physiographic	catign	none	geofabric	CAT-IGNEOUS (variable in geofabric lookup table "substrate", proportion of catchment)	0–1
Process	Sample type	process	none	sample specification		field-sort or lab-sort
No.riffle sample-units per sample	Sample type	nriff	none	sample specification		0, 1 or 2
No. spring sample-units per sample	Sample type	nspring	none	sample specification		0, 1 or 2

Table 2. Fixed-effect structures of mixed models (with random intercept determined by site group and slope determined by antecedent discharge) and their ΔAIC_c values compared to the best model ($\Delta AIC_c = 0$, in bold). Variable codes and transformations are defined in Table 1. ‘Unweighted’ models used unweighted 2-y afi, TI and total F. ‘Weighted afi; Unweighted I, F’ models used optimal afi (4-y linearly weighted), TI and total F. ‘Weighted afi, I; Unweighted F’ models used optimal afi, AI (optimal fit as determined by Walsh and Kunapo 2009) and total F. ‘Weighted afi, I, F’ models used optimal afi, AI, and optimal F (exponentially attenuated $HDD_L = 50$ m, $HDD_W = 1500$ m). meanQ * I * F * afi denotes each of these four effects and all their interactions. The final model is the best of the full-interaction models with 3- and 4-way interactions removed that did not improve AIC (colons denote simple interactions).

N	Model	Unweighted afi, I F	Weighted afi	Weighted afi, I	Weighted afi, I, F
				Unweighted I, F	Unweighted F
1	smeanannq * ldai9 * Ft * afi + carea + elev + catign + process + nriff + nspring	268.2	206.1	118.4	45.2
2	meanQ * I * F * afi + carea + elev + catign + process + nriff	258.9	197.5	109.7	36.4
3	meanQ * I * F * afi + carea + elev + catign + process + nspring	266.6	201.7	114.5	37.9
4	meanQ * I * F * afi + carea + elev + catign + nriff + nspring	292.4	226.8	137.4	64.8
5	meanQ * I * F * afi + carea + elev + catign + process	257.4	193.1	105.8	29.1
6	meanQ * I * F * afi + carea + elev + catign + nspring	289.4	221.4	132.4	57.1
7	meanQ * I * F * afi + carea + elev + catign + nriff	283.0	218.4	129.0	56.1
8	meanQ * I * F * afi + carea + elev + catign + process	257.4	193.1	105.8	29.1

9	meanQ * I * F * afi + elev + catign + process	258.8	195.4	105.9	21.3
10	meanQ * I * F * afi + carea + catign + process	271.4	209.9	111.9	42.7
11	meanQ * I * F * afi + carea + elev + process	270.0	206.4	127.0	60.8
12	meanQ * I * F * afi + carea + process	282.1	220.7	130.6	69.3
13	meanQ * I * F * afi + elev + process	273.0	210.8	129.3	53.7
14	meanQ * I * F * afi + catign + process	282.3	222.9	119.3	37.2
15	meanQ + I + F + afi + elev + catign + process + meanQ:I + meanQ:F + I:F + meanQ:afi + I:afi + F:afi + meanQ:F:afi	234.2	172.3	87.5	0.0

Fig. 1. A. Sites across the Greater Melbourne region from which macroinvertebrate samples were collected. Adjacent sites with the same colour were grouped as a random factor in the mixed models. Black points are sites that were >5 km from any other site and each one formed a group of its own. The red rectangle in the inset identifies the extent of the map within Australia. B. Distribution of forest cover and impervious surfaces across the region. C. Distribution of mean annual stream discharge depth (meanQ) in streams of the region.

Fig. 2. A. Temporal distribution of macroinvertebrate samples over the study period. B. Variation in mean 48-month antecedent discharge over the study period in 3 representative reaches spanning a range of meanQ values.

Fig. 3. Two alternative weighting schemes for catchment land use. A. The catchment and stream of an example study site, illustrating schematically distance measures used in weighting. Areas of impervious surfaces are weighted as an exponential decay function of d_D , the overland flow distance from the most downstream point of the surface to the nearest stormwater drain (or stream in the absence of drains: after Walsh and Kunapo 2009). For forest cover, each 10 m $\times$ 10 m gridcell was weighted by d_L , the overland flow distance from the centre of the gridcell to the nearest stream, and by d_W , the distance along the stream to the site. The resulting fractional weighting for all points in the catchment using the optimal weighting scheme that was applied to (B) impervious surfaces (half-decay distance along d_D of 9.4 m: because the decay length is so short, the map is dominated by blue gridcells within 10 m of a stormwater drain or stream with high weightings and more distant brown gridcells with a near zero weighting), and (C) forest cover (exponential decay: half-decay distance along d_L of 35 m, and along d_W of 1000 m). Note that both B and C portray

the weighting that would be applied to any gridcell across the catchment and do not show the real forest and impervious cover in this catchment.

Fig. 4. Aikake Information Criterion (expressed as the difference, ΔAIC_c , from the best model) for models with the optimal structure, but with antecedent discharge (as a proportion of mean annual discharge) calculated for different time periods, either unweighted, or linearly weighted. Models with an AIC_c more than 2 larger (i.e. $\Delta AIC_c > 2$: the grey dotted horizontal line) than the best model are less plausible. 48-month (or greater) weighted discharge was the optimal measure of antecedent discharge.

Fig. 5. Differences in Aikake information criteria (ΔAIC_c) for models with weighted forest cover calculated using one of three decay functions (threshold, linear and exponential decay) and different half-decay lengths. The best weighting scheme was exponential decay with a half-decay distances of 35 m overland and 1.0 km in-stream. Horizontal grey dotted lines mark $\Delta AIC_c = 2$, as for Fig. 4. A. Variation in ΔAIC_c with varying in-stream half-decay length when overland half-decay length was set at the optimal value. B. Variation in ΔAIC_c with varying overland half-decay length when in-stream half-decay length was set at the optimal value.

Fig. 6. Contour plot of difference in Aikake Information Criterion (ΔAIC_c) from the optimal model for exponentially weighted forest cover with varying in-stream and overland half-decay distances. The + sign indicates the best combination of weighting distances. The numbers on the contour lines indicate ΔAIC_c from the best model.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Walsh, CJ;Webb, JA

Title:

Spatial weighting of land use and temporal weighting of antecedent discharge improves prediction of stream condition

Date:

2014-08-01

Citation:

Walsh, C. J. & Webb, J. A. (2014). Spatial weighting of land use and temporal weighting of antecedent discharge improves prediction of stream condition. *LANDSCAPE ECOLOGY*, 29 (7), pp.1171-1185. <https://doi.org/10.1007/s10980-014-0050-y>.

Persistent Link:

<http://hdl.handle.net/11343/282694>