Shoaling behaviour enhances risk of predation from multiple predator guilds in a marine fish

John R. Ford¹ & Stephen E. Swearer

Department of Zoology, Melbourne University, Parkville, Victoria 3010, Australia

5 ¹Corresponding author. Email: jford@unimelb.edu.au Phone: +61 3 8344 7986 Fax: +61 3 8344
7909

Author Contributions: JF and SS conceived and designed the experiments. JF performed the experiments and analyzed the data. JF and SS wrote the manuscript.

Abstract

Predicting the consequences of predator biodiversity loss on prey requires an understanding of multiple predator interactions. Predators are often assumed to have independent and additive effects on shared prey survival; however, multiple predator effects can be non-additive if predators foraging together reduce prey survival (risk enhancement) or increase prey survival through interference (risk reduction). In marine communities, juvenile reef fish experience very high mortality from two predator guilds with very different hunting modes and foraging domains - benthic and pelagic predator guilds. The few previous predator manipulation studies have found or assumed that mortality is independent and additive. We tested whether interacting predator guilds result in non-additive prey mortality and whether the detection of such effects change over time as prey are depleted. To do so, we examined the roles of benthic and pelagic predators on survival of a juvenile shoaling zooplanktivorous temperate reef fish, *Trachinops caudimaculatus*, on artificial patch reefs over two months in Port Phillip Bay, Australia. We observed risk enhancement in the first seven days, as shoaling behaviour placed prey between predator foraging domains with no effective refuge. At day 14 we observed additive mortality, and risk enhancement was no longer detectable. By days 28 and 62, pelagic predators were no longer significant sources of mortality and additivity was trivial. We hypothesize that declines in prey density led to reduced shoaling behaviour that brought prey more often into the domain of benthic predators, resulting in limited mortality from pelagic predators. Furthermore, pelagic predators may have spent less time patrolling reefs in response to declines in prey numbers. Our observation of the changing interaction between predators and prey has important implications for assessing the role of predation in regulating populations in complex communities.

Key words: non-additive mortality, predator-prey theory, shoaling, ecosystem stability, multiple predator effects

Introduction

Predation is a ubiquitous force in nature (Sih et al. 1985; Ricklefs 1987), fundamental to both the regulation of prey populations (Chase et al. 2002) and in shaping community structure (Menge and Sutherland 1987; Gurevitch et al. 2000; Schmitz and Suttle 2001; Stachowicz et al. 2007). Predatory effects may be direct and lethal, or indirect through altering prey behaviour (Lima 1998), morphology (Van Buskirk 2001) or resource allocation (Schmitz et al. 1997). Most prey face multiple predators in natural systems (Schmitz 2007), and single species predator-prey responses overlook the potential for predators to interact and cause non-independent responses in prey (Vance Chalcraft et al. 2004). Empirical studies assessing the interactive effects of multiple predators on prey are uncommon (Sih et al. 1998), limiting our understanding of the effects of specific predator declines in systems where predator diversity is high (Schmitz 2007).

Recent experimental manipulations have observed that the effects of multiple predators on prey mortality cannot be predicted by summing the effects of predators in isolation (Vance Chalcraft and Soluk 2005; Vance-Chalcraft and Soluk 2005; Griffen and Byers 2006b; Griffen 2006; Carey and Wahl 2010). Termed a Multiple Predator Effect (MPE, Sih et al. 1998), predators do not act independently when foraging on shared prey and may interact with each other directly, or influence the behaviour of prey (Griffen and Williamson 2008). When predators foraging together are more effective than the sum of their separate effects, this results in *risk enhancement* for prey (Fig. 1a), which occurs when a prey's response to one predator increases its risk to another predator (DeWitt and Langerhans 2003). Where predators foraging together are less effective than the expected addition of their separate effects, their interaction results in *risk reduction* for prey (Fig. 1b). Risk reduction may occur either by interference competition amongst predators (Griffen and Byers 2005) or by direct predation amongst predators (Vance Chalcraft et al. 2007). Additive mortality (Fig. 1c), or the lack of an MPE, can occur in two ways; where predators' interactions with prey are independent and foraging is spatially or temporally discrete (Sokol-Hessner and Schmitz 2002), or where one predator

does not significantly affect prey, and hence there is no difference in prey survival between both predators together and the more effective predator in isolation (Sih et al. 1998), known as a trivial case of additivity.

60 The nature of interactive effects of multiple predators on prey depends on three fundamental traits of the system under study. Firstly, predators may interact in either an antagonistic or facilitative manner, or may not interact at all (Vance-Chalcraft and Soluk 2005a). Secondly, prey can have discrete behavioural responses to different predators, which cannot all be expressed simultaneously when confronted with multiple predators and hence an optimal choice of behaviours must be made
65 (Krupa and Sih 1998; Griffen and Byers 2006a). Thirdly, predator and prey behaviour may respond to changes in density of either group. Increased predator density may increase predator interference and hence prey survival (Griffen and Williamson 2008), whereas increases in prey density may alter prey behaviour (i.e., intraspecific aggression) in ways that could increase the strength of MPEs (Losey and Denno 1998; Vance Chalcraft and Soluk 2005). Consequently, the dynamic changes in density of
70 both predators and prey will likely affect the nature and strength of MPEs through time, although this prediction has yet to be explicitly tested.

In the marine reef environment, shoaling fish face two predator guilds, benthic and pelagic predators, which are fundamentally different in hunting mode and the spatial and temporal scales over which they forage (White et al. 2010). Benthic predators are resident on reefs and other
75 benthic habitats, have foraging areas similar to that of their prey and are primarily ambush predators (Holbrook and Schmitt 2002). Pelagic predators are transient in their interactions with prey, forage over larger spatial scales than their prey and are primarily chase predators (White et al. 2010). Whilst benthic predators are confined to the same habitat as prey and are limited to targeting local prey resources, pelagic predators will make decisions about where they forage according to
80 energetic costs and benefits, which are often influenced by prey density (Connell 2000; Sandin &

Pacala 2005b). Reefs are therefore a useful model system for investigating the effects of two diverse predator guilds with very different foraging relationships with their prey.

Predation of shoaling reef fish by benthic and pelagic predator guilds also presents mechanisms for MPEs. Shoaling provides anti-predator benefits through increased vigilance and reducing per-capita predation risk from pelagic predators (Pitcher and Parrish 1993), and by allowing fish to forage beyond the reach of benthic predators (White et al. 2010). However, when both guilds are present, shoaling fish are 'sandwiched' between two spatially discrete predator guilds and hence have limited predator-free space (Hixon and Carr 1997). This creates the possibility of synergistic predation, a risk enhancement mechanism (Sih et al. 1998). Despite this potential risk enhancement mechanism, the only other study to manipulate predator guilds targeting shoaling prey detected additive mortality (Hixon and Carr 1997).

Declines in prey density can reduce the effectiveness of shoaling (Godin et al. 1988; Morgan 1988; Stier et al. 2012), and either reduce (Connell 2000) or increase (Sandin and Pacala 2005b) targeting by pelagic predators. Therefore the nature of predator prey interactions and potential MPEs could change in experimental situations where prey are depleted over time. Mortality from predation on reef fishes immediately post-settlement (e.g. 55.7% of fish lost in the first 2 days post settlement across 24 studies of coral reef fishes (Almany and Webster 2006)) is a strong regulator of populations (Schmitt and Holbrook 1999a; b; Shima 1999; Doherty et al. 2004; Sandin and Pacala 2005a; Anderson et al. 2007) and thus assessing how changes in prey densities influence MPEs is relevant for understanding how predation might influence prey mortality during this life-history transition.

In this study we examined the interactive effects of benthic and pelagic predator guilds in causing mortality in a shoaling reef fish. Specifically we tested for the presence of MPEs and aimed to identify whether the presence and nature of MPEs changes over time in response to declines in prey

105 densities. We conducted our experiments in the field on artificial patch reefs by manipulating initial
predator and prey abundance. We did not confine predators to reefs, in order to accommodate the
dynamic foraging decisions made by predators in response to changes in abundance of both
predators and prey. Predator abundances continue to decline as a result of human harvest (Parsons
1992; Madin et al. 2010) and habitat loss (Bruno and O'Connor 2005; Dobson et al. 2006). By
110 manipulating entire predator guilds *in situ*, we aimed to better understand the potential impacts of
disturbance to marine predators on wider communities.

Materials and Methods

115 We investigated the effects of benthic and pelagic predator guilds on the mortality of a reef fish
(*Trachinops caudimaculatus*) on artificial patch reefs in Port Phillip Bay, Victoria, Australia. We used
an additive design for the detection of MPEs, by manipulating the initial presence or absence of
benthic predators by physical removals, and excluding pelagic predators by netting reefs, to produce
an initial crossed design of four present-absent predator treatments. Prey survival was monitored
120 over time, along with the abundance of benthic predators, and the nature of predator effects
assessed using multiplicative statistical models.

Study prey species and artificial reefs

The southern hulafish, *Trachinops caudimaculatus*, is an abundant and conspicuous rocky reef fish in
SE Australia, forming very large (>1000 fish) and dense (>100 fish m²) shoals in coastal embayments
125 (Hunt et al. 2011). Pelagic larvae settle to rocky reefs between November and January (late
spring/early summer) and use rocky reef structure as refuge from pelagic predators, diving for cover
when threatened. At dusk, fish retreat to refuge in small cracks and holes where they remain until
dawn (J. Ford, pers. obs). Fish are highly site attached after settlement; in over 400 dives observing
the species, no movement has been recorded across expanses of open water greater than 20m.
130 Previous experiments with calcein tagging also indicated no movement of fish to neighbouring reefs
(M Le Feuvre & J Ford, unpublished data).

The study was conducted near Altona in the north of Port Phillip Bay, Victoria, Australia (37°54'S,
144°49'E). Port Phillip Bay is a sheltered embayment with a water area of 1930km² and connected to
the ocean through a restricted entrance 3km wide. In Nov-Dec 2008 we placed 12 artificial reefs in a
135 linear array at 9-10m depth, each 150m apart, on open sand approximately 2km from natural reef.
Each artificial reef unit was approximately 0.75m³ with a base area of 1.1m² and constructed of ten
modular units stacked two high in an X design (See Electronic supplementary material Fig. S1). Each

module consisted of a plastic milkcrate (Viscount Plastics IH120, 355mm × 355mm × 322mm) lined with 8 bricks (230mm × 110mm × 75mm), filled with cushion foam in the centre and covered by a concrete paver. 24 of the 80 bricks in a reef had eight 20mm diameter holes running the width of the brick to provide a total of 192 small refuge spaces. Each reef unit was covered with a 6mm PVC sheet measuring 1600mm x 1200mm to provide overhangs of 550mm and 150mm on adjacent sides. Larval hulafish settled naturally to artificial reefs during this period and these fish were removed before the experiments commenced. All experiments were conducted after the end of the 2009-2010 summer settlement period and reefs were naturally fouled with encrusting invertebrates and algae.

Predator identification

Predators were first identified and split into benthic and pelagic predator guilds. Taxa identified as benthic predators were fish of the genera *Cristiceps* and *Heteroclinus* (Clinidae), *Lotella rhacina* (Moridae), *Genypterus tigerinus* (Ophidiidae), and invertebrates *Nectocarcinus* spp. (Portunidae), *Haplochlana maculosa* (Octopodidae) and *Sepia apama* (Sepiidae). Species identified as pelagic predators were the fish *Arripis truttacea* (Arripidae), *Trachurus novaezelandiae* (Carangidae), *Chrysophrys auratus* (Sparidae), and the squid *Sepioteuthis australis* (Loliginidae), all of which are commercially and recreationally fished (Department of Primary Industries 2008). In all cases, potential predators were selected either from visual observations of feeding on hulafish (Clinidae, *Nectocarcinus* spp., *S. Apama*, *C. auratus* and *S. australis*), or published literature listing the species as consuming fish (Clinidae, *L. rhancina*, *A. trutacea* and *T. novaezelandiae* (Edgar and Shaw 1995; Hindell 2006)). Whilst we could find no published gut content studies on *H. maculosa*, the shared use of shelter holes with *T. caudimaculatus*, observation of strikes on fish in lab aquaria, and diet studies of other octopuses (Grubert 1999; Smith 2003), suggest that small fish in nocturnal refuges are likely prey.

Benthic predator manipulation

We manipulated the initial presence or absence of benthic predators by clearing all artificial reefs of mobile macrofauna. Divers on SCUBA used a clove oil and ethanol mixture to anaesthetise fish and dip nets to capture them. The modular units and lid were then lifted to the surface by winch onto a boat where they and all remaining cryptic benthic predators removed. Reefs were redeployed and rebuilt at a location 50 m away from the original location but still 150 m from the nearest artificial reef. Predators were returned to six of the reefs in their original abundances (hereafter referred to as B+) and six were left without of benthic predators (B-). Due to the disturbance involved in deconstructing reefs to remove benthic predators, removal occurred only at the beginning of the experiment and benthic predators subsequently colonised all reefs. B+ and B- therefore refer to the initial manipulation, while benthic predator abundance is treated as a continuous variable in all analyses.

Pelagic predator manipulation

Six reefs were covered in 6 mm nylon diamond mesh nets to exclude pelagic predators (referred to as P-), and the remaining were non-netted and access open to pelagic predators (P+). The net stretched over the lid of the reef and approximately 45° downward from the edge of the lid to the substrate, providing approximately 3.8 m³ of enclosed shoaling space. Because we could not prevent the possible colonisation of benthic predators to non-netted reefs that had been cleared of all macrofauna (i.e., the B-P+ treatment) during the course of the experiment, nets were loosely pegged at the base to also allow potential colonisation of benthic predators to netted reefs (i.e., the B-P- treatment).

Cage control experiment

Exclusion devices such as nets have the potential to introduce confounding factors (Doherty and Sale 1985; Steele 1996) by altering prey or predator behaviour. With respect to the current experiment,

the most likely confounding factors were the restriction to the natural shoaling behaviour of prey inside the net, and benthic predators using the nets as structure from which to strike prey. A partial net control, which allowed access by pelagic predators whilst providing net structure, would not provide an accurate estimation of the potential effects of the nets on hulafish survival. While a partial net would allow hulafish to shoal outside the nets, it would still provide structure for benthic predators and could attract more pelagic predators by providing greater coarse structure (see Connell 1997; Hixon and Carr 1997). A partially caged reef control within a larger cage to exclude pelagic predators (see Steele 1996) was not logistically feasible, as tidal currents and wave action in Port Phillip Bay are too strong to permit construction at depth of multiple large net structures able to withstand inclement weather. Therefore, prior to the main experiment, we used a net control to test and correct for the effects of nets on shoaling behaviour and predation. The net controls consisted of larger width mesh net (19mm) made from thin nylon string that allowed prey to shoal away from the net, excluded pelagic predators, and provided little structure for benthic predators.

Net controls ran for 7 days on three B+ and B- reefs, with fish translocated and surveyed as described below. Comparison of survival between the large (M+) and small (M-) mesh B+ reefs provided an indication of whether possible changes in prey shoaling behaviour created greater opportunity for predation by benthic predators in the M- nets. Comparison between the large (M+) and small (M-) mesh B- reefs provided a correction for predation by pelagic predators on prey shoaling outside the nets on the M+ treatment. We note that ideally, such a correction experiment would have run for the full 62 days to incorporate possible changes to net effects through time as prey density declines.

Prey translocation

Juvenile hulafish (<30d post settlement, 20 - 30mm TL) were caught from nearby natural reefs by SCUBA divers by herding fish into a large mouthed (1200 mm square) set net with a 2 m tapered

210 codend. During summer hula fish are distinct size-assortative shoalers, with juveniles shoaling with conspecifics of similar size in large groups (>100 fish). We were careful to target only shoals of the smallest newly settled fish, with otolith analysis of 12 sub-sampled fish identifying them as having settled within 12-30 days prior to capture. Fish were all caught on the same day and all fish were mixed into the same holding container. 50 randomly selected fish were released on reefs from a bag
215 by a SCUBA diver and observed for 2 minutes, noting any sick or injured fish. Fish swam into the reef structure or shoaled under the overhang and in no cases were fish observed swimming away from the reef.

Surveys

Fish survival was visually surveyed by a diver on SCUBA on days 7, 14, 28 and 62. Nets were cleaned
220 of macroalgae each sampling period. Benthic predators were surveyed on days 0, 28 and 62, as it became clear by day 28 that colonisation by benthic predators was occurring. We interpolated day 7 and 14 benthic predator abundances assuming a linear colonisation rate of benthic predators throughout the experiment, which was the best fit to day 0, 28 and 62 data.

Detecting multiple predator effects

225 Studies of MPEs have a number of underlying design concerns and assumptions that strongly affect the interpretation of statistical results and resulting ecological application. In this study we use an additive experiment design in manipulating predator abundance, the multiplicative statistical test for predator interactions, and we assumed a multiplicative or log decline of prey through time.

Experimental design of MPE studies fall into two specific categories; additive and substitutive
230 designs, which differ in the way the predator treatments are combined (Griffen 2006). Additive experiments add together set densities for the two predators, resulting in an overall higher predator density in the combined treatment. A substitutive design holds predator density constant across all treatments, while manipulating species richness. The two approaches answer different questions –

the additive design examines whether non-additive effects occur due to interspecific competition, whilst the substitutive design compares the effects of interspecific competition to that of intraspecific competition (Griffen 2006). We used an additive design, as our primary question concerned the interaction of the two predator guilds in natural abundances and the consequences for prey survival. Also, a substitutive design would require the maintenance of predator densities which was not feasible in our field environment, and impossible for transient pelagic predators.

Tests for MPEs compare the survival of prey in the both-predator treatment to expected values where risk reduction or risk enhancement does not occur. These expected values are derived from the single predator treatments using either an additive or multiplicative model. The additive model is inappropriate for experiments where prey depletion is not prevented (Sih et al. 1998), and hence we chose to use the multiplicative model, which is more appropriate for proportion survival data used in this study, and accounts for background or predator free mortality (Sih et al. 1998).

Multiplicative models are widely used, firstly by Billick and Case (1994) and Wooton (1994), and more recently used by Vonesh and Osenberg (2003), Vance Chalcraft and Soluk (2005a) and Griffen (2006). Non-additive predator effects can be statistically detected through 2-way ANOVA of log transformed survival data, where a significant interaction between predator treatments indicates non-additivity (Billick and Case 1994; Wooton 1994). The log transformation of the data assumes a logarithmic decline in prey density over time (Wooton 1994), which we observed in our data.

Our experimental design differed to the many previous studies of MPEs in that we did not maintain constant densities of either predator group. We manipulated initial benthic predator number, and surveyed as these fluctuated over time. Pelagic predators were either completely excluded or allowed to access the reef, although we could not survey their abundance. Therefore we treated benthic predator abundance as a continuous variable and pelagic presence or absence as a fixed factor. We tested for significant interactions using 1-factor ANCOVAs for each sampling period, with the presence/absence of pelagic predators as the main factor and benthic predator abundance as

the covariate. Per capita survival from day 0 was used instead of survival between surveys to

260 investigate whether the detection of MPEs is related to prey depletion over time and to remove the effects of variable prey depletion across replicate reefs.

Estimating an expected survival relationship where the effects of predators are independent and additive is complicated in this experiment by the variability in benthic predator abundance amongst reefs. All previous studies of additive effects have controlled predator numbers (Vance Chalcraft et
265 al. 2004; Griffen 2006 amongst others), or biomass (Carey and Wahl 2010), and compared expected survival under an additive model to observed survival with both predators present. This approach, however, was not possible in our open plots exposed to natural predator densities.

Instead, where a significant interaction between predator guilds was detected, we produced interaction plots and compared these to predictions of non-additive interactions illustrated in Figure

270 1. From these comparisons we can determine the nature of the non-additivity, either risk enhancement or risk reduction. We also attempted to identify trivial instances of additivity in the system, where one predator guild had a significant effect on survival and the other did not.

Therefore we identified one of four types of predator interactions at each sampling period:

additivity, trivial additivity, risk enhancement or risk reduction.

275

280

Results

Cage control experiment

Survival was lower when prey could not escape the net in the presence of benthic predators (M+B+
285 = 0.64 ± 0.05 SE, M-B+ = 0.55 ± 0.04 SE), although the difference was not significant ($t = 2.15$ $_{df=4}$, $P = 0.1$). Survival was also lower when fish could escape and shoal outside the net in the absence of benthic predators (M-B- = 0.8 ± 0.02 SE, M+B- = 0.73 ± 0.06 SE), but again the difference was not significant ($t = 0.98$ $_{df=4}$, $P = 0.38$). Therefore, the small mesh net used in the main experiment likely only had a small and non-significant effect of reduced survival by limiting shoaling behaviour, but
290 this was countered by eliminating predation on fish shoaling outside the nets. Subtracting the survival difference between the M+B- and the M-B- treatment (i.e. predation on fish shoaling outside nets, 0.07), from the difference between the M+B+ and M-B+ treatments (i.e. effects of restricting shoaling, 0.09), provides an estimate of the cage effect in the presence of benthic predators: a reduction in per capita survival of 0.02, or approximately 3%. Thus, as a conservative
295 measure, we increased survival estimates in all netted (P-) reef treatments by 3% to account for this effect.

Additivity of predator guild mortality

The abundance of benthic predators had a significant negative relationship with prey survival for all sampling periods (Table 1). Pelagic predator presence significantly decreased survival only in the first
300 7 and 14 days of the experiment and was not significant at 28 and 62 days.

A significant interaction between predator treatments was identified in the first 7 days, indicating a non-additive relationship (Table 1). Comparisons of interaction plots (Fig. 2) showed at 7 days the slope of the observed survival in the P+ treatment was significantly less than (i.e. more negative) than the slope of the P- treatments (Fig. 2a), and follows the prediction for risk enhancement (Fig.
305 1a). No significant predator treatment interactions were detected for subsequent sampling dates

and predator effects were considered additive. Due to the lack of a significant effect of pelagic predators at days 28 and 62, observations at these dates are considered trivial instances of additivity.

315 Discussion

This is the first study to observe changes in the detection of MPEs through time as prey are depleted, and is the first to test for and detect non-additivity by predator functional groups with discrete foraging domains. We observed risk enhancement to juveniles of the shoaling reef fish *Trachinops caudimaculatus* by the benthic and pelagic predator guilds in the first seven days of the experiment. Prey mortality was additive at day 14, as both predator guilds significantly affected survival as predicted by the multiplicative model. We observed trivial instances of additivity at days 28 and 62, when only the benthic predator guild significantly affected survival. Changes in the detection of MPEs and the relative effectiveness of predator guilds through time suggests that predator-prey interactions change in response to declining prey abundance and hence mask the original observation of risk enhancement. These results have important implications for how we interpret the results of other studies that manipulated reef fish predators, and provides insight into dynamics of predator-prey relationships in exploited communities.

Risk enhancement and synergistic predation in early post-settlement

Our observation of risk enhancement in the first seven days of the experiment contrasts with the findings of the only other study to manipulate both predator guilds preying on reef fish (Hixon and Carr 1997), and is different to the majority of predator studies using additive designs that have observed risk reduction (Carey and Wahl 2010; Griffen 2006; Griffen and Williamson 2008; Vance-Chalcraft and Soluk 2005a; Vance-Chalcraft and Soluk 2005b). Hixon and Carr (1997) found that predation on *Chromis cyanea* from the benthic and pelagic predator guilds were additive at all sampling periods up to 32 d. They hypothesized that the two predator guilds occupy and feed in discrete spaces on the reef and do not interfere with each other; therefore the authors concluded that the two mortality sources were independent and additive. Risk enhancement, however, is a

result of multiple predator exposure increasing the overall risk to prey, and does not require any
340 direct interactions amongst predators (Schmitz et al. 2007). For example, Losey and Denno (1998)
observed risk enhancement by two spatially discrete predators when foliage-dwelling predators
caused aphids to drop to the ground and were consumed by ground dwelling predators. Therefore,
we hypothesize that where predator foraging domains do not overlap, but provide no predator free
space between domains, risk enhancement can occur.

345 Although Hixon and Carr (1997) describe the predation of the two guilds as additive, the synergistic
predation they describe should result in non-additive risk enhancement. Because the authors
numerically summed the mortality from the single predator treatments to produce an expected
combined predator mortality, and did not use models of predicted multiplicative mortality used in
many studies of non-additivity (e.g. Vonesh and Osenberg 2003; Vance-Chalcraft and Soluk 2005b;
350 Griffen and Williamson 2007; Carey and Wahl 2010), their analysis could have failed to detect a non-
additive effect when one was actually present. Applying the multiplicative model from Wootton
(1994) to the data presented in Hixon and Carr (1997) results in an observed per capita survival after
32 days in the presence of both guilds (0.35) that is less than that predicted by the additive model
(0.41), suggesting prey risk enhancement may be occurring.

355 Differences in the predation rates attributable to the two guilds between the two studies may also
explain the disparity in detection of MPEs. Hixon and Carr (1997) observed a much stronger effect of
pelagic predators on prey, likely due to small shoal sizes (20 *C. cyanea*) reducing the effectiveness of
shoaling behaviour, low prey and benthic predator densities (see Carr and Hixon 1995), and
targeting of small shoals of *C. cyanea* by pelagic predators (see Sandin and Pacala 2005b). Hixon and
360 Carr (1997) may therefore have observed trivial additivity where pelagic predators were much more
effective than benthic predators, similar to that observed at days 28 and 62 of the present study. In
contrast the present study observed strong effects of both predator guilds on prey in the first seven
days when risk enhancement occurred.

The observed risk enhancement is likely driven by two separate mechanisms: shoaling by prey
365 reducing predation risk in the presence of a single predator, and adjacent predator foraging domains
leaving no spatial refuge when both guilds are present. When only benthic predators were present,
prey could shoal away from the reef and avoid predation, and when only pelagic predators were
present prey could shelter close to the reef. For example, Stier et al. (2012) found no evidence for
risk enhancement or risk reduction to shoaling reef fish prey with changes to density of only a single
370 predator type. When both predator guilds were present in our study, such behavioural responses by
prey to the presence of one predator guild may increase the risk of predation by the other predator
guild (Sih et al. 1988). Prey are in effect 'sandwiched' between predators from above and below
(Hixon and Carr 1997), with no effective spatial refuge (Krupa and Sih 1998). Large numbers of prey
all seeking shelter simultaneously leads to a scramble for limited refuges when a pelagic predator
375 attack occurs (Samhouri et al. 2009). Failure to find refuge and having to 'queue' outside of refuges
would likely leave prey at greater risk of predation by both benthic predators waiting in ambush and
the pelagic chase predators. Such situations could feasibly result in synergistic predation and prey
risk enhancement.

380 **Additive predator effects and the decline in pelagic predator effectiveness**

Predator interactions shifted from synergistic predation in the first seven days to an additive
relationship for the remainder of the experiment. At day 14 the effect was genuine additivity under
the multiplicative model, where both predator guilds significantly affected survival. By days 28 and
62, additivity was 'trivial' (Sih et al. 1998) because only the benthic guild significantly affected
385 survival of prey. Benthic predators were in effect the only important predator and the addition of
the pelagic predator guild did not influence prey survival beyond its small (non-significant) effect in
isolation. Such shifts in the detection of MPEs over the course of the experiment confirms the

dynamic nature of predator prey interactions, and highlights important limitations to discrete experiments attempting to observe snapshots in time.

390 We observed two major shifts that require mechanistic investigation; the change from risk enhancement to additive mortality after the first week, and the decline in mortality from pelagic predators over time. Decreasing prey density can result in reduced risk enhancement behaviours by prey (Losey and Denno 1998). Reduced detection of prey is also likely to be an important driver of risk reduction if predators are more likely to detect (Krause et al. 1998) or target (Connell 2000)

395 larger groups of prey. Thus there are two possible explanations for these shifts, changes in prey and predator behaviours as prey density declined.

Prey behaviour can change with conspecific density in response to the effectiveness of social or group strategies (Alexander 1974; Maher and Lott 2000). Shoaling can be more successful at higher densities through attack abatement (Pitcher and Parrish 1993; Ioannou and Krause 2008), predator

400 confusion (Milinski 1979), corporate vigilance (Magurran et al. 1985; Krause and Ruxton 2002) and more efficient food finding abilities (Pitcher et al. 1982). However at low densities or under low predation pressure, shoaling behaviour is less effective and fish feed as individuals, as observed in guppies (Liley and Seghers 1975) and minnows (Magurran and Pitcher 1987). Similar responses have been observed in the wrasse *Thalassoma amblycephalum*, where the proportion of time shoaling

405 increased with shoal size (Stier et al. 2012). At low densities *T. caudimaculatus* shoal closer to the substrate and are more likely to forage individually (Fumei 2011), restricting their foraging domain to areas close to the reef. Therefore the size of prey foraging domains declines with density, and at the smallest group sizes prey no longer overlap with the foraging domain of pelagic predators (Fumei 2011). The cost of not shoaling is increased exposure to, and hence predation by, the benthic guild.

410 This change in foraging behaviour likely contributed to the observation of trivial additivity after four weeks, where only benthic predators significantly affected prey survival.

The reduced exposure of prey to pelagic predation through changes to shoaling may have been compounded by a corresponding reduction in targeting of smaller shoals by pelagic predators. Due to their large foraging domains, which encompass many square kilometres and multiple reef habitats, pelagic predators can actively choose which prey aggregations they target (Overholtzer-McLeod 2004). Pelagic predators tend to congregate on larger prey aggregations (Hixon and Carr 1997, Connell 2000, Stewart and Jones 2001), which could provide a greater return on foraging effort (Ioannou and Krause 2008). Although Sandin and Pacala (2005b) observed a higher per capita incidence of predator visits in smaller shoals, predators had either a positive or independent relationship with shoal size. If pelagic predators reduced their targeting of prey over the course of the experiment in response to declining densities, this also may have contributed to the reduced effect of pelagic predators and the trivial observation of additivity. Consequently, the shift in the type of MPE observed is related to both prey and predator responses to prey depletion over time.

Conclusion

The observation of the changing nature of predator interactions from non-additive to additive effects has important implications for both our understanding of predation and how we interpret the results of predator manipulation studies. Whilst short-term non-additive effects may be strong, changes to prey densities and resulting behaviour may negate the detection of these effects over time. Although the use of an additive experimental design means we cannot answer the question of whether predator guilds are substitutable, we can assess how non-impacted communities are likely to respond to declines in or the loss of either predator guild. In our system, benthic predators likely play an important role in regulating prey populations at low densities, whilst pelagic predators are more effective at higher prey densities. A loss of either key functional group through pelagic fishing or benthic habitat change is likely to significantly alter the dynamics of prey populations. Identifying

non-additive sources of mortality is, therefore, an important step in understanding the potential effects of biodiversity loss on ecosystem function in marine environments.

Acknowledgements

440 We would like to thank Christian Jung, Dean Chamberlain, Matt Le Feuvre, Madhavi Colton, Mal
Lindsay and Jessica Smith for the back-breaking construction of artificial reefs, these again plus Evan
Hallein, Simon Pahor, Dan Corrie and Seann Chia for setups and surveys, and John Ahern for boat
and technical support. We thank Craig Osenberg for his helpful guidance on determining non-
additivity in the system. Operational costs were covered by grants from the Holsworth Wildlife
445 Research Endowment to JRF and the Australia and Pacific Science Foundation to SES. Animal Ethics
approval granted under Melbourne University AEC0810932. The authors declare they have no
conflict of interest.

References

- 450 Alexander RD (1974) The Evolution of Social Behavior. *Annual Review of Ecology and Systematics* 5:325–383
- Almany GR, Webster MS (2006) The predation gauntlet: early post-settlement mortality in reef fishes. *Coral Reefs* 25:19–22. doi:10.1007/s00338-005-0044-y
- 455 Anderson TW, Carr MH, Hixon MA (2007) Patterns and mechanisms of variable settlement and recruitment of a coral reef damselfish, *Chromis cyanea*. *Marine Ecology Progress Series* 350:109–116. doi:10.3354/meps07135
- Billick I, Case TJ (1994) Higher order interactions in ecological communities: what are they and how can they be detected? *Ecology* 75:1529
- 460 Bruno JF, O'Connor MI (2005) Cascading effects of predator diversity and omnivory in a marine food web. *Ecology letters* 8:1048
- Carey MP, Wahl DH (2010) Interactions of multiple predators with different foraging modes in an aquatic food web. *Oecologia* 162:443
- Carr MH, Hixon MA (1995) Predation effects on early post-settlement survivorship of coral reef fishes. *Marine Ecology Progress Series* 124:31–42
- 465 Chase JM, Abrams PA, Grover JP, Diehl S, Chesson P, Holt RD, Richards SA, Nisbet RM, Case TJ (2002) The interaction between predation and competition: a review and synthesis. *Ecology Letters* 5:302–315. doi:Article
- 470 Connell SD (1997) The relationship between large predatory fish and recruitment and mortality of juvenile coral reef-fish on artificial reefs. *Journal of Experimental Marine Biology and Ecology* 209:261–278
- Connell SD (2000) Is there safety-in-numbers for prey? *Oikos* 88:527–532
- Department of Primary Industries (2008) Fishery Status Report 2008. Fisheries Victoria, Melbourne
- 475 DeWitt TJ, Langerhans RB (2003) Multiple prey traits, multiple predators: keys to understanding complex community dynamics. *Journal of Sea Research* 49:143–155. doi:doi: 10.1016/S1385-1101(02)00220-4
- Dobson A, Lodge D, Alder J, Cumming GS, Keymer J, McGlade J, Mooney H, Rusak JA, Sala O, Wolters V, Wall D, Winfree R, Xenopoulos MA (2006) Habitat loss, trophic collapse, and the decline of ecosystem services. *Ecology* 87:1915
- 480 Doherty PJ, Sale PF (1985) Predation on juvenile coral reef fishes: an exclusion experiment. *Coral Reefs* 4:225–234. doi:10.1007/BF00298081

- Doherty PJ, Dufour V, Galzin R, Hixon MA, Meekan MG, Planes S (2004) High Mortality during Settlement Is a Population Bottleneck for a Tropical Surgeonfish. *Ecology* 85:2422–2428
- Edgar GJ, Shaw C (1995) The production and trophic ecology of shallow-water fish assemblages in southern Australia II. Diets of fishes and trophic relationships between fishes and benthos at
485 Western Port, Victoria. *Journal of Experimental Marine Biology and Ecology* 194:83–106.
doi:10.1016/0022-0981(95)00084-4
- Fumei S (2011) Growing together: group living behaviour and the relationship between density and growth in a shoaling fish. Masters of Science Thesis, Melbourne University
- Godin J, Classon L, Abrahams M (1988) Group Vigilance and shoal size in a small characin fish.
490 Behaviour 104:29–40. doi:10.1163/156853988X00584
- Griffen BD (2006) Detecting emergent effects of multiple predator species. *Oecologia* 148:702–709.
doi:10.1007/s00442-006-0414-3
- Griffen BD, Byers JE (2006a) Intraguild predation reduces redundancy of predator species in multiple predator assemblage. *Journal of Animal Ecology* 75:959–966. doi:10.1111/j.1365-
495 2656.2006.01115.x
- Griffen BD, Byers JE (2006b) Partitioning mechanisms of Predator Interference in different Habitats. *Oecologia* 146:608–614. doi:10.1007/s00442-005-0211-4
- Griffen BD, Williamson T (2008) Influence of predator density on nonindependent effects of multiple predator species. *Oecologia* 155:151–159. doi:10.1007/s00442-007-0889-6
- 500 Grubert MA (1999) Diet and feeding strategy of *Octopus maorum* in southeast Tasmania. *Bulletin of marine science* 65:441
- Gurevitch J, Morrison JA, Hedges LV (2000) The interaction between competition and predation: a meta-analysis of field experiments. *American Naturalist*, The 155:435
- Hindell JS (2006) Assessing the trophic link between seagrass habitats and piscivorous fishes. *Marine and Freshwater Research* 57:121. doi:10.1071/MF05082
505
- Hixon MA, Carr MH (1997) Synergistic predation, density dependence, and population regulation in marine fish. *Science* 277:946–949
- Holbrook SJ, Schmitt RJ (2002) Competition for shelter space causes density-dependent predation mortality in damselfishes. *Ecology* 83:2855–2868
- 510 Hunt TL, Ford JR, Swearer SE (2011) Ecological determinants of recruitment to populations of a temperate reef fish, *Trachinops caudimaculatus* (Plesiopidae). *Marine & freshwater research* 62:502–509. doi:10.1071/MF10262
- Ioannou CC, Krause J (2008) Searching for prey: the effects of group size and number. *Animal Behaviour* 75:1383–1388. doi:10.1016/j.anbehav.2007.09.012

- 515 Krause J, Ruxton GD (2002) *Living in groups*. Oxford: Oxford University Press.
- Krause J, Ruxton GD, Rubenstein D (1998) Is there always an influence of shoal size on predator hunting success? *Journal of Fish Biology* 52:494–501. doi:10.1111/j.1095-8649.1998.tb02012.x
- 520 Krupa JJ, Sih A (1998) Fishing spiders, green sunfish, and a stream-dwelling water strider: male-female conflict and prey responses to single versus multiple predator environments. *Oecologia* 117:258
- Liley N, Seghers B (1975) Factors affecting the morphology and behaviour of guppies in Trinidad. *Function and evolution in behaviour* 6: 92–118
- Lima SL (1998) Nonlethal effects in the ecology of predator-prey interactions. *BioScience* 48:25
- 525 Losey JE, Denno RF (1998) Positive predator-predator interactions: Enhance predation rates and synergistic suppression of aphid populations. *Ecology* 79:2143–2152. doi:10.1890/0012-9658(1998)079[2143:PPPIEP]2.0.CO;2
- Madin EMP, Gaines SD, Warner RR (2010) Field evidence for pervasive indirect effects of fishing on prey foraging behavior. *Ecology* 91:3563
- 530 Magurran AE, Pitcher TJ (1987) Provenance, Shoal Size and the Sociobiology of Predator-Evasion Behaviour in Minnow Shoals. *Proceedings of the Royal Society of London Series B Biological Sciences* 229:439 –465. doi:10.1098/rspb.1987.0004
- Magurran AE, Oulton WJ, Pitcher T (1985) Vigilant behaviour and shoal size in minnows. *Journal of Comparative Ethology* 64:167–178
- 535 Maher CR, Lott DF (2000) A Review of Ecological Determinants of Territoriality within Vertebrate Species. *The American Midland Naturalist* 143:1–29. doi:10.1674/0003-0031(2000)143[0001:AROEDO]2.0.CO;2
- Menge BA, Sutherland JP (1987) Community Regulation: Variation in Disturbance, Competition, and Predation in Relation to Environmental Stress and Recruitment. *The American Naturalist* 130:730–757
- 540 Milinski M (1979) Can an experienced predator overcome the confusion of swarming prey more easily? *Animal Behaviour* 27, Part 4:1122–1126. doi:10.1016/0003-3472(79)90060-5
- Morgan MJ (1988) The influence of hunger, shoal size and predator presence on foraging in bluntnose minnows. *Animal Behaviour* 36:1317–1322. doi:10.1016/S0003-3472(88)80200-8
- 545 Overholtzer-McLeod KL (2004) Variance in reef spatial structure masks density dependence in Coral-Reef fish populations on natural versus artificial reefs. *Marine Ecology Progress Series* 276:269–280
- Parsons TR (1992) The removal of marine predators by fisheries and the impact of trophic structure. *Marine pollution bulletin* 25:51

- 550 Pitcher TJ, Parrish JK (1993) Functions of shoaling behaviour in teleosts. In: Pitcher TJ (ed) Behaviour of Teleost Fishes, 2nd edn. Chapman and Hall, London England
- Pitcher TJ, Magurran AE, Winfield IJ (1982) Fish in larger shoals find food faster. *Behavioral Ecology and Sociobiology* 10:149–151. doi:10.1007/BF00300175
- 555 Ricklefs RE (1987) Community diversity: relative roles of local and regional processes. *Science* 235:167
- Samhuri JF, Vance RR, Forrester GE, Steele MA (2009) Musical chairs mortality functions: density-dependent deaths caused by competition for unguarded refuges. *Oecologia* 160:257–265. doi:10.1007/s00442-009-1307-z
- 560 Sandin SA, Pacala SW (2005a) Demographic Theory of Coral Reef Fish Populations with Stochastic Recruitment: Comparing Sources of Population Regulation. *The American Naturalist* 165:107–119
- Sandin SA, Pacala SW (2005b) Fish aggregation results in inversely density-dependent predation on continuous coral reefs. *Ecology* 86:1520–1530
- 565 Schmitt RJ, Holbrook SJ (1999a) Mortality of juvenile damselfish: Implications for assessing processes that determine abundance. *Ecology* 80:35–50
- Schmitt RJ, Holbrook SJ (1999b) Settlement and Recruitment of Three Damselfish Species: Larval Delivery and Competition for Shelter Space. *Oecologia* 118:76–86
- Schmitz OJ (2007) Predator diversity and trophic interactions. *Ecology* 88:2415
- 570 Schmitz OJ, Suttle KB (2001) Effects of top predator species on direct and indirect interactions in a food web. *Ecology* 82:2072–2081
- Schmitz OJ, Beckerman AP, O'Brien KM (1997) Behaviorally mediated trophic cascades: effects of predation risk on food web interactions. *Ecology* 78:1388
- Shima JS (1999) Variability in relative importance of determinants of reef fish recruitment. *Ecology Letters* 2:304–310
- 575 Sih A, Crowley P, McPeck M, Petranks J, Stroheimer K (1985) Predation, competition, and prey communities: a review of field experiments. *Annual review of ecology and systematics* 16:269
- Sih A, Englund G, Wooster D (1998) Emergent impacts of multiple predators on prey. *Trends in Ecology & Evolution* 13:350–355. doi:doi: 10.1016/S0169-5347(98)01437-2
- 580 Smith CD (2003) Diet of *Octopus vulgaris* in False Bay, South Africa. *Marine biology* 143:1127
- Sokol-Hessner L, Schmitz OJ (2002) Aggregate Effects of Multiple Predator Species on a Shared Prey. *Ecology* 83:2367–2372

Stachowicz JJ, Bruno JF, Duffy JE (2007) Understanding the effects of marine biodiversity on communities and ecosystems. *Annual review of ecology, evolution, and systematics* 38:739

585 Steele MA (1996) Effects of predators on reef fishes: separating cage artifacts from effects of predation. *Journal of Experimental Marine Biology and Ecology* 198:249–267. doi:doi:10.1016/0022-0981(96)00011-1

Stewart BD, Jones GP (2001) Associations between the abundance of piscivorous fishes and their prey on coral reefs: implications for prey-fish mortality. *Marine Biology* 138:383–397

590 Stier AC, Geange SW, Bolker BM (2012) Predator density and competition modify the benefits of group formation in a shoaling reef fish. *Oikos*:no–no. doi:10.1111/j.1600-0706.2012.20726.x

Van Buskirk J (2001) Specific induced responses to different predator species in anuran larvae. *Journal of Evolutionary Biology* 14:482–489. doi:10.1046/j.1420-9101.2001.00282.x

595 Vance Chalcraft HD, Soluk DA (2005) Multiple predator effects result in risk reduction for prey across multiple prey densities. *Oecologia* 144:472

Vance Chalcraft HD, Soluk DA, Ozburn N (2004) Is prey predation risk influenced more by increasing predator density or predator species richness in stream enclosures? *Oecologia* 139:117

Vance Chalcraft HD, Rosenheim JA, Vonesh JR, Osenberg CW, Sih A (2007) The influence of intraguild predation on prey suppression and prey release: a meta-analysis. *Ecology* 88:2689

600 Vance-Chalcraft HD, Soluk DA (2005) Estimating the prevalence and strength of non-independent predator effects. *Oecologia* 146:452–460. doi:10.1007/s00442-005-0201-6

Vonesh JR, Osenberg CW (2003) Multi-predator effects across life-history stages: non-additivity of egg- and larval-stage predation in an African treefrog. *Ecology Letters* 6:503–508. doi:10.1046/j.1461-0248.2003.00470.x

605 White JW, Samhouri JF, Stier AC, Wormald CL, Hamilton SL, Sandin SA (2010) Synthesizing mechanisms of density dependence in reef fishes: behavior, habitat configuration, and observational scale. *Ecology* 91:1949–1961

Wootton JT (1994) Putting the Pieces Together: Testing the Independence of Interactions among Organisms. *Ecology* 75:1544–1551

610

Table 1 Results of ANCOVA testing for the effects of pelagic and benthic predator guilds on log transformed per capita survival of juvenile southern hulafish, *Trachinops caudimaculatus*. Pelagic predators were excluded using cages and their presence and absence forms the main term. Benthic predator abundance is the covariate, as colonisation of reefs by benthic predators changed the original experimental densities. Per capita survival from day 0 was measured on 12 reefs on days 7, 14, 28 and 62.

Source	df	MS	F-ratio	P-value
<i>Survival 0-7 days</i>				
Pelagic	1	1.35	28.2	<0.001
Benthic	1	1.59	33.1	<0.001
Pelagic x Benthic	1	0.53	11.1	0.01
Error	8	0.05		
<i>Survival 0-14 days</i>				
Pelagic	1	1.34	14.1	0.005
Benthic	1	2.65	27.8	<0.001
Pelagic x Benthic	1	0.02	0.23	0.65
Error	8	0.10		
<i>Survival 0-28 days</i>				
Pelagic	1	0.95	4.66	0.063
Benthic	1	4.91	24.1	0.001
Pelagic x Benthic	1	0.01	0.028	0.87
Error	8	0.20		
<i>Survival 0-62 days</i>				
Pelagic	1	2.67	3.54	0.09
Benthic	1	7.87	10.46	0.01
Pelagic x Benthic	1	0.64	0.84	0.38
Error	8	0.75		

Fig. 1 Predicted interaction plots of multiple predator effects on prey survival. In all cases the prey survival is on the y-axis on a logarithmic scale, the presence or absence of predator A is on the x-axis and the two lines represents the presence (B+, solid line) or absence (B-, dashed line) of predator B.

a) Prediction of risk enhancement, where prey mortality is greater in the presence of both predators than that predicted by adding their effects in isolation. Here the slope of the relationship where B is present is significantly less than the slope of the relationship when predator B is absent. b) Prediction of risk reduction under a mutual interference model, where the prey survival in the presence of both predators is significantly greater than in the presence of either predator in isolation. c) Prediction of an additive relationship between predators A and B, where the slopes of the lines are statistically similar. We assume significant prey survival effects in the presence of either predator in isolation. If only predator A was significant in isolation, slopes would be similar and significantly different to 0, and y-intercepts similar. If only predator B was significant in isolation, slopes would not be different from 0 and y-intercepts would be significantly different. Both of these scenarios are examples of trivial additivity (*sensu* Sih et al. 1998)

Fig. 2 Interaction plots of the effects of benthic and pelagic predators on the survival of juvenile hulafish on artificial reefs in Port Phillip Bay, Australia over summer-autumn 2010 for: a) days 0-7, b) days 0-14, c) days 0-28 and d) days 0-62. Pelagic predators were present (P+, circles and solid lines) or absent (P-, crosses and dashed lines). P-values correspond to the benthic x pelagic interaction term from each ANCOVA on log transformed per capita survival data (Table 1), with NS representing a non-significant interaction. A significant interaction between predator treatments was detected only in the first 7 days and plots reveal a risk enhancement effect similar to that predicted in Figure 1a, with the slope of the P+ treatment (-0.14 ± 0.02) significantly less than the slope of the P- treatment (-0.04 ± 0.02). Note the log scale used on the y-axis.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Ford, JR;Swearer, SE

Title:

Shoaling behaviour enhances risk of predation from multiple predator guilds in a marine fish

Date:

2013-06-01

Citation:

Ford, J. R. & Swearer, S. E. (2013). Shoaling behaviour enhances risk of predation from multiple predator guilds in a marine fish. *OECOLOGIA*, 172 (2), pp.387-397. <https://doi.org/10.1007/s00442-012-2508-4>.

Persistent Link:

<http://hdl.handle.net/11343/282716>