Abstract:

BACKGROUND: Despite accumulated experience and advancing techniques for laparoscopic hepatectomy, there are still challenging resections that require specific and innovative intraoperative maneuvers ^{1,2,3}. The right posterior sectionectomy presents special concerns about its location, the extensive transection area and difficult access to the pedicle ^{4,5}. Intrahepatic glissonian approach allows safe en masse control of the portal structures without prolonged dissection ². Its association with the half-Pringle maneuver results in less bleeding during parenchymal transection ^{1,6}.

METHODS: 34 years-old woman referred to treat a 8.0 cm hepatocellular adenoma located at segments VI/VII. Patient was placed in a semi-supine position and six ports were located in a distribution that resembles a Makuuchi's incision. The right liver was mobilized and preparation to an anatomic glissonian approach was performed. A vascular clamp was placed to assure that full control of the right posterior pedicle was possible. Then, a vascular stapler replaced it, with division of the right posterior glissonian pedicle. A vascular clamp was inserted from the inferior right flank 5-mm trocar to perform a half-pringle maneuver of the right pedicle in order to minimize blood loss during parenchymal transection. Liver parenchyma was transected with harmonic scalpel and vascular stapler. The right hepatic vein was divided intra-parenchymally with a vascular stapler. Specimen was extracted through a Pfannenstiel incision.

RESULTS: Total surgical time was 210 minutes. Estimated blood loss was 200 ml and no blood transfusion was required. The recovery was uneventful and hospital discharge occurred on the fifth post-operative day. Pathology confirmed the diagnosis of an hepatocellular adenoma.

CONCLUSIONS: Technical issues have initially hindered the development of laparoscopic liver resections ^{7,8,9,10}. Surgeons were concerned about hemostasis, bleeding control, safe and effective parenchymal transection, adequate visualization and the feasibility to work on deeper regions of the liver. During the past decade, a lot of limitations were overcome, but lesions located on the postero-superior liver are still considered tough to beat ^{5,11}. When we look up at large series and extensive reviews ^{12,13,14}, resections located on the posterior segments are still infrequent. Limited access to the portal triad, difficult pedicle control, a large transection area and its anatomically location, hidden from the surgeon's view, attached to the diaphragm and retroperitoneum, makes such resections defying. Our team has made 97 laparoscopic hepatectomies, including 6 lesions resections of the right posterior sector. In our series, half pedicle clamping was used in 12 patients, and we adopt such maneuver as an inflow control when we're operating on peripheric lesions with difficult vascular control (such as enucleations or postero-superiorly located segmentectomies). This technique is safe and useful, as it reduces liver ischemic aggression, a very important issue in diseased livers (e.g steatosis, steato-hepatitis, prolonged chemotherapy, cirrhosis) ^{6,15}. In our series, we applied the glissonian intrahepatic approach in 7 cases (2 left hepatectomies and 5 right hepatectomies).

We understand that laparoscopy applies perfectly to oddly (postero-superior) located tumors and right posterior sectionectomy can be safely accomplished. In fact, we share opinion with other specialized hepatobiliary centers, believing that this may be the preferred approach ¹⁶.

Glissonian approach permits fast and efficient pedicle control, avoiding the risk of falling into the anatomic pitfalls of the extra-hepatic vasculo-biliary elements. Its association to the Half-Pringle maneuver increases the ability to control bleeding during liver transection.

Drs. Paulo Herman, Jaime Arthur Pirola Krüger, Renato Micelli Lupinacci, Fabrício Ferreira Coelho and Marcos Vinícius Perini have no conflicts of interest or financial ties to disclose.

References:

1. Half-Pringle maneuver: a useful tool in laparoscopic liver resection. Herman P, Perini MV, Coelho F, Saad W, D'Albuquerque LA. J Laparoendosc Adv Surg Tech A. 2010 Feb;20(1):35-7.
2. Machado MA, Makdissi FF, Galvão FH, Machado MC. Intrahepatic Glissonian approach for laparoscopic right segmental liver resections. Am J Surg 2008; 196(10):e38-42

3. Yoon YS, Han HS, Choi YS, Jang JY, Suh KS, Kim SW, Lee KU, Park YH. Total laparoscopic right posterior sectionectomy for hepatocellular carcinoma. *J Laparoendosc Adv Surg Tech A*. 2006 Jun;16(3):274-7.
4. Torzilli G, Donadon M, Marconi M, Botea F, Palmisano A, Del Fabbro D, Procopio F, Montorsi M. Systematic extended right posterior sectionectomy: a safe and effective alternative to right hepatectomy. *Ann Surg*. 2008 Apr;247(4):603-11.
5. Buell JS, Cherqui D, Geller DA, O'Rourke N, Iannitti D, Dagher I, Koffron AJ, Thomas M, Gayet B, Han HS, Wakabayashi G, Belli G, Kaneko H, Ker CG, Scatton O, Laurent A, Abdalla EK, Chaudhury P, Dutson E, Gamblin C, D'Angelica M, Nagorney D, Testa G, Labow D, Manas D, Poon RT, Nelson H, Martin R, Clary B, Pinson WC, Martinie J, Vauthey JN, Goldstein R, Roayaie S, Barlet D, Espat J, Abecassis M, Rees M, Fong Y, McMathers KM, Broelsch C, Busuttil R, Belghitti J, Strassberg S, Chari RS. The international position on laparoscopic liver surgery: The Louisville statement, 2008. *Ann Surg*. 2009 Nov;250(5):825-30.
6. Makuuchi M, Mori T, Gunvén P, Yamazaki S, Hasegawa H. Safety of hemihepatic vascular occlusion during resection of the liver. *Surg Gynecol Obstet*. 1987 Feb;164(2):155-8.
7. Cherqui D, Husson E, Hammoud R, Malassagne B, Stephan F, Bensaid S, et al. Laparoscopic liver resections: a feasibility study in 30 patients. *Ann Surg* 2000;232:753-62.
8. Dulucq JL, Wintringer P, Stabilini C, Berticelli J, Mahajna A. Laparoscopic liver resections: a single center experience. *Surg Endosc* 2005;19:886-91.
9. Laurent A, Cherqui D, Lesurtel M, Brunetti F, Tayar C, Fagniez PL. Laparoscopic liver resection for subcapsular hepatocellular carcinoma complicating chronic liver disease. *Arch Surg* 2003;138:763-9; discussion 769.
10. Dagher I, Proske JM, Carloni A, Richa H, Tranchart H, Franco D. Laparoscopic liver resection: results for 70 patients. *Surg Endosc* 2007;21:619-24.
11. Cho JY, Han HS, Yoon YS, Shin SH. Feasibility of laparoscopic liver resection for tumors located in the posterosuperior segments of the liver, with a special reference to overcoming current limitations on tumor location. *Surgery* 2008 Jul;144(1):32-8.
12. Nguyen KT, Gamblin TC, Geller D. World review of laparoscopic liver resection – 2804 patients. *Ann Surg* 2009 Nov;250(5):831-841.
13. Koffron AJ, Auffenberg G, Kung R, Abecassis M. Evaluation of 300 minimally invasive liver resections at a single institution: less is more. *Ann Surg*. 2007;246:385–392; discussion 392–394.
14. Buell JF, Thomas MT, Rudich S, Marvin M, Nagubandi R, Ravindra KV, Brock G, McMasters KM. Experience with more than 500 minimally invasive hepatic procedures. *Ann Surg*. 2008;248:475–486.
15. Chau GY, Lui WY, King KL, Wu CW. Evaluation of effect of hemihepatic vascular occlusion and the Pringle maneuver during hepatic resection for patients with hepatocellular carcinoma and impaired liver function. *World J Surg*. 2005 Nov;29(11):1374-83.

16. Gumbs AA, Gayet B. Video: the lateral laparoscopic approach to lesions in the posterior segments. *J Gastrointest Surg.* 2008 Jul;12(7):1154.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Herman, P;Krueger, J;Lupinacci, R;Coelho, F;Perini, M

Title:

Laparoscopic bisegmentectomy 6 and 7 using a Glissonian approach and a half-Pringle maneuver

Date:

2013-05-01

Citation:

Herman, P., Krueger, J., Lupinacci, R., Coelho, F. & Perini, M. (2013). Laparoscopic bisegmentectomy 6 and 7 using a Glissonian approach and a half-Pringle maneuver. SURGICAL ENDOSCOPY AND OTHER INTERVENTIONAL TECHNIQUES, 27 (5), pp.1840-1841. <https://doi.org/10.1007/s00464-012-2681-x>.

Persistent Link:

<http://hdl.handle.net/11343/282941>