**EXERCISE-INDUCED INHIBITION OF REMODELING IS FOCALLY OFFSET
WITH FATIGUE FRACTURE IN RACEHORSES**

R. Christopher Whitton¹, Michiko Mirams¹, Eleanor J. Mackie¹, Garry A. Anderson¹, and
Ego Seeman²

¹*Faculty of Veterinary Science, University of Melbourne.*

²*Department of Endocrinology and Medicine, Austin Health, University of Melbourne.*

Corresponding Author:

Chris Whitton

Equine Centre

University of Melbourne

250 Princes Hwy

Werribee, Vic, 3030

Ph: +61 3 97312268 Fax: +61 97312246

Email: cwhitton@unimelb.edu.au

The authors have no conflict of interest. This study was supported by the Rural Industries
Research and Development Corporation of the Australian Government and Racing Victoria
Limited.

22 **Mini abstract**

23 Bone remodeling is inhibited by high repetitive loading. However, in subchondral bone of
24 racehorses in training, eroded surface doubled in association with fatigue fracture and there
25 was greater surrounding trabecular bone volume suggesting trabecular modeling unloads the
26 bone focally allowing damage repair by remodeling.

27

Abstract

Remodeling replaces damaged bone with new bone but is suppressed during high magnitude repetitive loading when damage is most likely. However, in cortical bone of racehorses, at sites of fatigue fracture, focal porosity, consistent with remodeling, is observed in proportion to the extent of surrounding callus. Focal areas of porosity are also observed at sites of fatigue damage in subchondral bone. We hypothesized that fatigued subchondral bone, like damaged cortical bone, is remodeled focally in proportion to the modeling of surrounding trabecular bone.

Eroded and mineralizing surfaces and bone area were measured using backscattered scanning electron microscopy of post-mortem specimens of the distal third metacarpal bone in 11 racehorses with condylar fractures (cases) and 8 racehorses in training without fractures (controls).

Cases had a two-fold greater eroded surface per unit area at the fracture site than controls (0.81 ± 0.10 vs. $0.40 \pm 0.12 \text{ mm}^{-1}$, $P = 0.021$) but not at an adjacent site (0.22 ± 0.09 vs. $0.30 \pm 0.11 \text{ mm}^{-1}$, $P = 0.59$). Area fraction of surrounding trabecular bone was higher in cases than controls ($81 \pm 2\%$ vs. $72 \pm 2\%$, $P = 0.0020$) and the eroded surface at the fracture site correlated with the surrounding trabecular area (adjusted $R^2 = 0.63$, $P = 0.0010$).

In conclusion, exercise-induced inhibition of remodeling is offset at sites of fatigue fracture. Modeling of trabecular bone may contribute to unloading these regions allowing repair by remodeling.

50 Key words

51 Electron microscopy, fatigue fracture, modeling, remodeling, subchondral bone

52

53

54 **Introduction**

55 The ability of bone to adapt to prevailing loading circumstances without failing depends on
56 the integrity of the cellular machinery of bone modeling and remodeling [1,2]. In bones
57 exposed to high repetitive loads, modeling increases cortical and trabecular thickness
58 conferring strength, stiffness and resistance to fatigue damage [3-8]. Remodeling removes
59 fatigue damaged bone and replaces it with new bone. Stress fractures are the result of
60 loading of a skeleton that is not adapted to its loading environment. Although most
61 commonly encountered in army recruits and in ballet dancers, stress fractures occur in any
62 circumstance where there is an increase in loading. The effects of a change in loading are
63 most easily studied at the extremes and the horse is a species which habitually subjects its
64 skeleton to the highest loads measured to date [9].

65 Repetitive loading inhibits remodeling allowing the accumulation of fatigue damage [8,10-
66 14]. This is well documented in thoroughbred racehorses in which repetitive loading of
67 training and racing produces fatigue injuries [15]. Unlike the commonly studied rat model
68 of fatigue fracture in which intra-cortical remodelling does not usually occur, bones of the
69 horse undergo intra-cortical remodeling like that observed in human bone [4,16]. Horses
70 have low cortical porosity at sites subjected to high loads, but focal cortical porosity
71 consistent with remodeling is observed at fatigue fracture sites suggesting that the
72 suppressive effect of repetitive loading can be overcome [10,17,18]. Cortical bone fatigue
73 fracture sites are also characterised by surrounding callus formation [17].

74 Subchondral bone is subjected to repeated high magnitude compressive loads generated by
75 wrapping forces from muscles and tendons [19]. In patients with osteoporosis, subchondral
76 bone is susceptible to fatigue damage due to insufficient structure [20]. Subchondral bone
77 fatigue fracture is common in equine athletes which generate extreme loads in joints of the

lower limb [9]. In a previous study we reported that modeling of surrounding trabecular bone is consistent in association with subchondral bone fatigue fracture sites whereas focal areas of porosity at the fracture site are inconsistent [12].

To determine if remodeling occurs in a high load environment, we examined subchondral surfaces from a site predisposed to fatigue fracture and an adjacent less commonly injured site in racehorses in training. We hypothesised that eroded surfaces would be associated with fatigue fracture, consistent with the notion that remodeling inhibition by repeated high loading can be overcome focally, and similar to cortical bone callus formation, this would occur in regions surrounded by greater trabecular bone volume.

Methods

The third metacarpal bone (MC3) in the horse bears all the weight transmitted through the forelimb. The distal end of MC3 is a common site of stress fractures in racehorses; these fractures usually extend proximally from the parasagittal grooves adjacent to the medial or lateral condyles. Left and right third metacarpal bones were obtained postmortem from Thoroughbred racehorses that died or were euthanized from April 2007 to April 2009 and underwent post mortem examination at the University of Melbourne. These horses were a subset of those presented in a previous publication [12]. Different trainers were responsible for the management of the horses in this study. All horses with fractures were euthanased on the day of injury. Horses were assigned to; 1) cases: 11 horses with third metacarpal fractures with a condylar component within the parasagittal groove that occurred spontaneously (i.e. no report of a collision with another horse or object) during a race or a training gallop, and 2) controls: 8 horses in training with no limb fracture at the time of death. Horses with a fracture of a bone other than the MC3 or where the condylar fracture did not involve the parasagittal groove were excluded.

All 11 horses with condylar fractures were male, eight were castrated. Mean age ($\pm$ SD) of cases was 4.2 ± 2.4 years. All 11 had unilateral fractures involving at least one metacarpal condyle. One had biaxial condylar fractures and the remainder were uniaxial. In 5 of the case horses incomplete condylar fractures were identified with high resolution peripheral computed tomography in the contralateral limb, including biaxial incomplete fractures in the horse with contralateral biaxial complete fractures. This is similar to the findings in race horses from the UK where 52% of horses with fractures of the parasagittal groove had grossly visible fissures at the same site in the contralateral limb [21]. Contralateral limbs were therefore not considered suitable controls. Of the uniaxial fractures, seven involved the lateral metacarpal condyle and three the medial metacarpal condyle. Four were simple condylar fractures, one was non-displaced. The remainder were comminuted proximally. Concurrent injuries occurred in three horses; two lateral condylar fractures also had medial sesamoid midbody fractures and one lateral condylar fracture had a lateral sesamoid midbody fracture. For controls, cause of death was cardiac failure during exercise in four horses and euthanasia due to acute gastrointestinal disease ($n=2$), cervical spinal fracture ($n=1$) and tendon laceration ($n=1$). Mean age of controls was 4.4 ± 1.9 years and there were 5 males, 4 of which were castrated and 3 females. Racing and training history were obtained from race records and telephone interviews with the trainers.

Left and right distal forelimbs were removed post-mortem and the metacarpal bones dissected. For fractured MC3s the distal articular surface was reconstructed and held together with cyanoacrylate adhesive applied dorsally. The palmar-distal aspect of the MC3 was removed by cutting the bone at 55° to the frontal plane through the centre of rotation of the condyles and stored in 70% ethanol. Bones were dehydrated, embedded in methyl methacrylate, then cut with a low-speed saw (Isomet saw: Buehler, Lake Bluff, IL, USA) to expose the section of interest. The surface of the block was ground, polished, carbon-coated

and examined with a scanning electron microscope (FEI Quanta FEG [field emission gun] 200, Hillsboro, OR, USA) fitted with an annular solid-state back-scattered electron detector. Consecutive overlapping micrographs of the whole condyle were obtained at a magnification of 70× and of specific regions of interest (ROIs) at 200× and reconstructed using Adobe Photoshop Elements 3.

Backscattered scanning electron microscopy (BSEM) images were analysed using open source 3D imaging software (Image J, NIH, Bethesda, MD). Analysis was performed in the following ROIs; a) the dense subchondral bone of the lateral condyle, b) the lateral condyle trabecular and subchondral bone, c) the medial condyle trabecular and subchondral bone, d) the lateral parasagittal groove, and e) the medial parasagittal groove (Figure 1). Analysis of ROIs b and c were performed at 70x and ROIs a, c and d at 200x magnification. Images were automatically thresholded using the ‘minimum’ algorithm in Image J [22].

All bone surfaces within a region of interest were examined and classified as follows. Eroded surfaces (E.Pm) were defined as those that were sharp, irregular and crossed lamellar lines (surfaces formed by cracking were not included) whereas mineralising surfaces (Md.Pm) were those that did not cross lamellar lines and where there was a layer of darker (low mineral content) bone on the surface (Figure 2). These surfaces were identified and traced using the segmented line tool and the total surface length of each calculated and divided by the area of the region of interest (T.Ar) or expressed as a percentage of total surface (B.Pm). Surfaces per unit area were included because the high volume fraction of equine subchondral bone limited available surfaces. As resorption is the initial step in the remodeling cycle and formation can be due to both modeling and remodeling we used erosion surface as our primary measure of remodeling. Measurements of maximum pore diameter were made using the line tool.

Statistical analysis

For each limb, data from only one condyle for regions of interest b-e were included in analysis. For fractured limbs data from the affected condyle was used unless the fracture was biaxial in which case one was chosen randomly. For controls, condyles were frequency matched, using a procedure of random allocation, so as to provide the same proportion of lateral and medial condyles as in the cases. A general linear model was used to determine the effect of group and other predictors on BSEM variables. For all analyses age, sex, number of career starts and weeks in current training period were tested for significance or confounding. Age was the only additional variable included in a final model, this being for the analysis of erosion surface in the parasagittal groove (E.Pm/T.Ar and E.Pm/B.Pm) where it had a significant effect. All other variables had no effect. The analyses were also performed with the female horses in the control group excluded and all significant results remained significant. Levene's test was used to assess homogeneity of error variances and was not significant. Analysis was performed using IBM SPSS 19.0 for Windows (IBM Corporation, Armonk, NY, USA). Estimated marginal means $\pm$ standard errors are presented. A two-sided P-value <0.05 was considered significant.

Results

Measures of eroded surface in the subchondral bone of the parasagittal groove (fracture site) in cases were greater than controls (ROI d or e) but not in the lateral condyle (ROI a) (Table 1). Resorption spaces associated with these erosion surfaces at the fracture site were often large (100-450 μ m diameter) (Figure 3) whereas resorption spaces in the lateral condylar subchondral bone were small (50-150 μ m diameter). Despite the local loss of bone at the fracture site, bone area fraction in the whole condyle which included the surrounding trabecular and subchondral bone was higher in cases than controls (ROI b or c, Table 1).

Eroded surface at the fracture site was associated with higher surrounding trabecular and subchondral bone area fraction in all horses (adjusted $R^2=0.63$, $P=0.0010$), but the regression coefficient for eroded surface versus bone area fraction was greater in cases than controls ($P=0.029$) (Figure 4). In all horses with high erosion surface, surrounding trabecular and subchondral bone area fraction was also high (Figure 4).

Surface cracks extending through focal areas of highly mineralised calcified cartilage and into the subchondral bone were observed in all horses at the parasagittal groove (Figure 5). At the lateral condyle fewer horses with mineralised cracks were observed: 4 of 11 (36%) cases and 5 of 8 (63%) controls had cracks.

Discussion

Under conditions of repetitive loading we observed approximately double the level of erosion surface at the fatigue fracture site in cases compared with controls. Fractured bones had greater total bone volume and remodeling at the fracture site was associated with a higher surrounding trabecular and subchondral bone area.

High magnitude repetitive loading increases the risk of fatigue damage but reduces remodeling [8,10-14]. However once damage develops in subchondral bone, focal remodeling can occur [23,12,18]. The mechanism for remodeling suppression is uncertain but osteoclast recruitment is inhibited by repeated strains *in vitro* [24].

Remodeling repairs microdamage but may produce a transitory or permanent increase in intracortical porosity and resorption trenches upon trabecular surfaces which concentrate stress predisposing to further microdamage [25]. Focal bone loss as a result of greater erosion surface could contribute to fracture propagation (Figure 3), but net bone volume of the whole condyle is greater due to modeling of surrounding trabecular bone. This

compensatory trabecular modeling can limit the detrimental effect of remodeling at moderate levels of fatigue loading but the interaction is overwhelmed at high rates of damage accumulation where remodeling exacerbates damage [26].

Increased remodeling in response to damage is reported in models in which the response to microdamage is examined after loading has ceased [27-30,16,31]. Therefore the subsequent increased remodeling may be partly a response to unloading following damage rather than damage alone. By contrast, in our study, horses were loading their skeleton continuously up until the time of sample collection. Under similar circumstances others have observed resorption spaces thought to be produced by remodeling activity but only in close association with areas of damage in equine subchondral bone [23,32]. We confirm that these spaces are a result of bone resorption.

Similar to our subchondral bone site, porosity consistent with remodeling activity is observed in cortical bone subject to fatigue fracture but only in association with mature callus [17]. Furthermore in the rat ulna model of fatigue injury the peak of remodeling activity coincides with a mature periosteal woven bone callus that contributes to whole bone strength [29,33]. Callus formation load shields the damaged area [34,35]. We speculate that increased surrounding trabecular and subchondral bone volume may act similarly, facilitating focal remodeling of damaged bone. Modeled equine trabecular bone has a 50% greater modulus than underlying damaged subchondral bone and such differences in modulus directly affect internal stresses resulting in unloading of the damaged area [6,36].

To our knowledge this is the only model of fatigue damage where remodeling activity has been examined under continued repetitive loading. Similar conditions are likely to apply when porotic bone is subject to an increase in loading. Despite the small sample size we were able to reject the null hypothesis and infer that remodeling inhibition associated with

repetitive high loading can be overcome focally, in regions surrounded by greater trabecular bone volume. Horses in this study were sourced from a number of different trainers. Trainers use a variety of training methods and have been reported to be an independent determinant of fracture risk in some but not all studies [37-40]. The cases and controls were also not uniform for age, and sex but we assessed these factors as well as available training data for confounding. Moreover in a separate analysis without female horses, all significant results remained significant. We did not quantify microdamage which may be a more sensitive method of assessing fatigue damage than grouping bones into those with fractures and those without. Based on the presence of mineralised cracks all horses including the controls had some degree of fatigue damage but more at the fracture site [41]. Some bone was lost at fracture sites and although this was limited, it was from the articular surface and therefore eroded and mineralisation surface would be under-estimated in these cases.

In conclusion, despite subchondral remodeling being inhibited by high magnitude repetitive loading, focal remodeling occurs at sites of fatigue failure, perhaps due to modeling in surrounding trabecular and subchondral bone focally unloading the damaged region. Although this model is an extreme example of increased bone loading it demonstrates a mechanism of adaptation that may apply to any change in loading circumstances.

Acknowledgements

Funding was provided by the Rural Industries Research and Development Corporation of the Australian Government and Racing Victoria Limited. We thank Mark Forwood for advice on the manuscript.

References

1. Frost HM (2000) Does bone design intend to minimize fatigue failures? A case for the affirmative. *Journal of bone and mineral metabolism* 18 (5):278-282
2. Burr DB, Robling AG, Turner CH (2002) Effects of biomechanical stress on bones in animals. *Bone* 30 (5):781-786
3. Danova NA, Colopy SA, Radtke CL, Kalscheur VL, Markel MD, Vanderby R, McCabe RP, Escarcega AJ, Muir P (2003) Degradation of bone structural properties by accumulation and coalescence of microcracks. *Bone* 33 (2):197-205
4. Boyde A, Firth EC (2005) Musculoskeletal responses of 2-year-old Thoroughbred horses to early training. 8. Quantitative back-scattered electron scanning electron microscopy and confocal fluorescence microscopy of the epiphysis of the third metacarpal bone. *N Z Vet J* 53 (2):123-132
5. Warden SJ, Hurst JA, Sanders MS, Turner CH, Burr DB, Li J (2005) Bone adaptation to a mechanical loading program significantly increases skeletal fatigue resistance. *J Bone Miner Res* 20 (5):809-816
6. Rubio-Martinez LM, Cruz AM, Gordon K, Hurtig MB (2008) Mechanical properties of subchondral bone in the distal aspect of third metacarpal bones from Thoroughbred racehorses. *Am J Vet Res* 69 (11):1423-1433
7. Rapillard L, Charlebois M, Zysset PK (2006) Compressive fatigue behavior of human vertebral trabecular bone. *J Biomech* 39 (11):2133-2139
8. Jee WS, Li XJ, Schaffler MB (1991) Adaptation of diaphyseal structure with aging and increased mechanical usage in the adult rat: a histomorphometrical and biomechanical study. *Anat Rec* 230 (3):332-338

- 269 9. Harrison SM, Whitton RC, Kawcak CE, Stover SM, Pandy MG (2010) Relationship
270 between muscle forces, joint loading and utilization of elastic strain energy in equine
271 locomotion. *J Exp Biol* 213 (Pt 23):3998-4009
- 272 10. Firth EC, Rogers CW, Doube M, Jopson NB (2005) Musculoskeletal responses of 2-
273 year-old Thoroughbred horses to early training. 6. Bone parameters in the third metacarpal
274 and third metatarsal bones. *N Z Vet J* 53 (2):101-112
- 275 11. Jackson BF, Goodship AE, Eastell R, Price JS (2003) Evaluation of serum
276 concentrations of biochemical markers of bone metabolism and insulin-like growth factor I
277 associated with treadmill exercise in young horses. *Am J Vet Res* 64 (12):1549-1556
- 278 12. Whitton RC, Trope GD, Ghasem-Zadeh A, Anderson GA, Parkin TD, Mackie EJ,
279 Seeman E (2010) Third metacarpal condylar fatigue fractures in equine athletes occur within
280 previously modelled subchondral bone. *Bone* 47 (4):826-831
- 281 13. Frost HM (1987) Bone "mass" and the "mechanostat": A proposal. *Anatomical Record*
282 219:1-9
- 283 14. Jee WS, Li XJ (1990) Adaptation of cancellous bone to overloading in the adult rat: a
284 single photon absorptiometry and histomorphometry study. *Anat Rec* 227 (4):418-426.
285 doi:10.1002/ar.1092270405
- 286 15. Stover SM, Ardans AA, Read DH, Johnson BJ, Barr BC, Daft BM, Kindu H, Anderson
287 ML, Woods LW, Moore J, Stoltz J, Pool RR (1993) Patterns of stress fractures associated
288 with complete bone fractures in racehorses. *Proceedings of the American Association of*
289 *Equine Practitioners* 39:131-132
- 290 16. Kidd LJ, Stephens AS, Kuliwaba JS, Fazzalari NL, Wu AC, Forwood MR (2010)
291 Temporal pattern of gene expression and histology of stress fracture healing. *Bone* 46
292 (2):369-378

- 293 17. Entwistle RC, Sammons SC, Bigley RF, Hazelwood SJ, Fyhrie DP, Gibeling JC, Stover
294 SM (2009) Material properties are related to stress fracture callus and porosity of cortical
295 bone tissue at affected and unaffected sites. *J Orthop Res* 27 (10):1272-1279
- 296 18. Muir P, Peterson AL, Sample SJ, Scollay MC, Markel MD, Kalscheur VL (2008)
297 Exercise-induced metacarpophalangeal joint adaptation in the Thoroughbred racehorse. *J*
298 *Anat* 213 (6):706-717
- 299 19. Shelburne KB, Torry MR, Pandy MG (2006) Contributions of muscles, ligaments, and
300 the ground-reaction force to tibiofemoral joint loading during normal gait. *J Orthop Res* 24
301 (10):1983-1990
- 302 20. Li ZC, Dai LY, Jiang LS, Qiu S (2012) Difference in subchondral cancellous bone
303 between postmenopausal women with hip osteoarthritis and osteoporotic fracture:
304 Implication for fatigue microdamage, bone microarchitecture, and biomechanical properties.
305 *Arthritis Rheum* 64 (12):3955-3962. doi:10.1002/art.34670
- 306 21. Parkin TD, Clegg PD, French NP, Proudman CJ, Riggs CM, Singer ER, Webbon PM,
307 Morgan KL (2006) Catastrophic fracture of the lateral condyle of the third
308 metacarpus/metatarsus in UK racehorses - fracture descriptions and pre-existing pathology.
309 *Vet J* 171 (1):157-165
- 310 22. Prewitt JM, Mendelsohn ML (1966) The analysis of cell images. *Annals of the New*
311 *York Academy of Sciences* 128 (3):1035-1053
- 312 23. Muir P, McCarthy J, Radtke CL, Markel MD, Santschi EM, Scollay MC, Kalscheur VL
313 (2006) Role of endochondral ossification of articular cartilage and functional adaptation of
314 the subchondral plate in the development of fatigue microcracking of joints. *Bone* 38
315 (3):342-349
- 316 24. Rubin J, Fan X, Biskobing DM, Taylor WR, Rubin CT (1999) Osteoclastogenesis is
317 repressed by mechanical strain in an in vitro model. *J Orthop Res* 17 (5):639-645

- 318 25. Hernandez CJ, Gupta A, Keaveny TM (2006) A biomechanical analysis of the effects of
319 resorption cavities on cancellous bone strength. *J Bone Miner Res* 21 (8):1248-1255
- 320 26. van Oers RF, van Rietbergen B, Ito K, Huiskes R, Hilbers PA (2011) Simulations of
321 trabecular remodeling and fatigue: is remodeling helpful or harmful? *Bone* 48 (5):1210-
322 1215. doi:10.1016/j.bone.2011.01.011
- 323 27. Cardoso L, Herman BC, Verborgt O, Laudier D, Majeska RJ, Schaffler MB (2009)
324 Osteocyte apoptosis controls activation of intracortical resorption in response to bone
325 fatigue. *J Bone Miner Res* 24 (4):597-605
- 326 28. Herman BC, Cardoso L, Majeska RJ, Jepsen KJ, Schaffler MB (2010) Activation of
327 bone remodeling after fatigue: differential response to linear microcracks and diffuse
328 damage. *Bone* 47 (4):766-772. doi:10.1016/j.bone.2010.07.006
- 329 29. Hsieh YF, Silva MJ (2002) In vivo fatigue loading of the rat ulna induces both bone
330 formation and resorption and leads to time-related changes in bone mechanical properties
331 and density. *J Orthop Res* 20 (4):764-771
- 332 30. Verborgt O, Gibson GJ, Schaffler MB (2000) Loss of osteocyte integrity in association
333 with microdamage and bone remodeling after fatigue in vivo. *J Bone Miner Res* 15 (1):60-
334 67
- 335 31. Mori S, Burr DB (1993) Increased intracortical remodeling following fatigue damage.
336 *Bone* 14:103-109
- 337 32. Norrdin RW, Stover SM (2006) Subchondral bone failure in overload arthrosis: a
338 scanning electron microscopic study in horses. *J Musculoskelet Neuronal Interact* 6 (3):251-
339 257
- 340 33. Silva MJ, Touhey DC (2007) Bone formation after damaging in vivo fatigue loading
341 results in recovery of whole-bone monotonic strength and increased fatigue life. *J Orthop*
342 *Res* 25 (2):252-261

34. Vetter A, Liu Y, Witt F, Manjubala I, Sander O, Epari DR, Fratzl P, Duda GN, Weinkamer R (2011) The mechanical heterogeneity of the hard callus influences local tissue strains during bone healing: a finite element study based on sheep experiments. *J Biomech* 44 (3):517-523. doi:10.1016/j.jbiomech.2010.09.009
35. Epari DR, Schell H, Bail HJ, Duda GN (2006) Instability prolongs the chondral phase during bone healing in sheep. *Bone* 38 (6):864-870
36. Shirazi R, Shirazi-Adl A (2009) Computational biomechanics of articular cartilage of human knee joint: effect of osteochondral defects. *J Biomech* 42 (15):2458-2465
37. Ely ER, Avella CS, Price JS, Smith RK, Wood JL, Verheyen KL (2009) Descriptive epidemiology of fracture, tendon and suspensory ligament injuries in National Hunt racehorses in training. *Equine Vet J* 41 (4):372-378
38. Verheyen KL, Newton JR, Price JS, Wood JL (2006) A case-control study of factors associated with pelvic and tibial stress fractures in Thoroughbred racehorses in training in the UK. *Prev Vet Med* 74 (1):21-35
39. Verheyen KL, Wood JL (2004) Descriptive epidemiology of fractures occurring in British Thoroughbred racehorses in training. *Equine Vet J* 36 (2):167-173
40. Verheyen K, Price J, Lanyon L, Wood J (2006) Exercise distance and speed affect the risk of fracture in racehorses. *Bone* 39 (6):1322-1330
41. Boyde A (2003) The real response of bone to exercise. *Journal of anatomy* 203 (2):173-189

Table 1. BSEM variables for thoroughbred racehorses that developed metacarpal condylar fractures while in training (cases) and horses in training without fractures (controls). Estimated marginal mean $\pm$ SE are presented.

		Cases (n=11)	Controls (n=8)	P value
Condyle (ROI b or c)	B.Ar/T.Ar %	81 $\pm$ 2	72 $\pm$ 2	0.0020
Lateral condyle	B.Ar/T.Ar %	90 $\pm$ 1	90 $\pm$ 1	0.65
subchondral bone (ROI a)	E.Pm/T.Ar mm ⁻¹	0.22 $\pm$ 0.09	0.30 $\pm$ 0.11	0.59
	E.Pm/B.Pm %	11 $\pm$ 2	9 $\pm$ 3	0.40
	Md.Pm/T.Ar mm ⁻¹	0.24 $\pm$ 0.30	0.25 $\pm$ 0.35	0.76
	Md.Pm/B.Pm %	11 $\pm$ 1	10 $\pm$ 2	0.51
Parasagittal groove (ROI d or e)	B.Ar/T.Ar %	86 $\pm$ 2	84 $\pm$ 2	0.26
	E.Pm/T.Ar mm ⁻¹	0.81 $\pm$ 0.10	0.40 $\pm$ 0.12	0.021
	E.Pm/B.Pm %	17 $\pm$ 2	10 $\pm$ 2	0.020
	Md.Pm/T.Ar mm ⁻¹	1.19 $\pm$ 0.15	0.67 $\pm$ 0.17	0.034
	Md.Pm/B.Pm %	28 $\pm$ 5	19 $\pm$ 5	0.22

Figure 1. Regions of interest in the distal third metacarpal bone imaged with BSEM. a) The dense subchondral bone of the lateral condyle, b) lateral condyle trabecular and subchondral bone, c) medial condyle trabecular and subchondral bone, d) lateral parasagittal groove, e) medial parasagittal groove. Area b) includes a) and d), and area c) includes e).

Figure 2. BSEM image of the subchondral bone of the third metacarpal bone showing eroded surfaces (white arrow heads) and mineralising surfaces (black arrow heads). (Bar = 250µm)

Figure 3. BSEM image of a fracture through the lateral parasagittal groove of an 8-year-old Thoroughbred racehorse (lateral is to the left). Multiple surface cracks are observed adjacent to the fracture and there are numerous resorption surfaces (white arrow heads) in the immediate subchondral bone (lower insert) and deeper subchondral bone (upper insert). (Bar = 2mm)

Figure 4. Association between eroded surface (E.PM/T.AR) within the parasagittal groove of the distal third metacarpal and surrounding bone area fraction (B.AR/T.AR) (ROI b or c) in cases and controls. Cases solid circles, controls open circles.

Figure 5. BSEM images of the parasagittal groove from a control horse (left) and a case horse (right). Surface cracks extending through focal areas of highly mineralised calcified cartilage (white arrowheads) are present in both.

397

398

399

400

401

402

403

404

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Whitton, RC;Mirams, M;Mackie, EJ;Anderson, GA;Seeman, E

Title:

Exercise-induced inhibition of remodelling is focally offset with fatigue fracture in racehorses

Date:

2013-07-01

Citation:

Whitton, R. C., Mirams, M., Mackie, E. J., Anderson, G. A. & Seeman, E. (2013). Exercise-induced inhibition of remodelling is focally offset with fatigue fracture in racehorses. *OSTEOPOROSIS INTERNATIONAL*, 24 (7), pp.2043-2048. <https://doi.org/10.1007/s00198-013-2291-z>.

Persistent Link:

<http://hdl.handle.net/11343/283061>