CURRENT AND FUTURE CLIMATE VARIABILITY ASSOCIATED WITH WINTERTIME PRECIPITATION IN ALPINE AUSTRALIA

Sonya Louise Fiddes¹, Alexandre Bernardes Pezza¹

¹School of Earth Sciences, University of Melbourne,
Parkville, VIC 3010 Australia

Correspondence:

Sonya Fiddes

Address: School of Earth Sciences, University of Melbourne, Parkville, VIC 3010
Australia

Email: sonya.fiddes@unimelb.edu.au

Phone: +61 (4) 3803 8402

Fax: +61 (3) 8344 7761

Keywords: Rainfall, Alpine, Australia, Climate, Variability

Abstract

The Australian Alps, located in the southeast corner of the continent, are home to important ecologies and industries, including water supply, hydroelectricity and ski resorts. Sharp topography and prevailing westerly winds generate a unique microclimate with cool temperatures and abundant precipitation and is crucial to much of greater southeastern Australia's water supply. Here we study the western, high and eastern slopes separately, exploring the global climate drivers associated with wintertime precipitation variability. The results show that while total precipitation is significantly declining on the western and high slopes, the total rain is not significantly changing on the eastern side. These differing trends are thought to be a result of the changing nature of the westerly storm track and the subtropical ridge. Interestingly, the west/high wintertime rainfall decline is seen primarily as a reduction in the intensity of events, as the number of rainfall days per season has remained relatively constant throughout the analysis. The synoptic patterns associated with extreme precipitation are identified and shown to be well correlated with the total seasonal precipitation, suggesting a great importance of the extreme weather signatures in modulating the longer term climate. This correlation is used to calculate a number of climate indices relying on dynamical indicators such as pressure and temperature gradients, helping simulate the rainfall variability within the area. By exploring contrasting CMIP3 models from the CSIRO Representative Future Climate Framework, we estimate using indices of the circulation dynamics that the west/high wintertime rainfall trend will continue to decline whilst rainfall in the eastern region will remain relatively stable. This result adds new light into future precipitation trends for the area, given the intrinsic difficulty of climate models to correctly represent precipitation behavior over the region.

1. Introduction

Past literature clearly shows that the total wintertime precipitation has, over time, declined significantly in the Australian Alps, especially on the western and high slopes (Nicholls 2009; Chubb et al. 2011). Whilst this has been suggested to be due to the changes in the westerly storm track and subtropical ridge directly influencing the rainfall bearing process of the region, little research has been focused on the notable lack of significant decline in the eastern slopes.

In this study, the question as to what may be causing the two differing rainfall trends over the west/high and east slopes is explored, with a focus on connections to climate drivers and atmospheric and oceanic modes. Furthermore, we investigate what the future of precipitation may be in two contrasting climate model projections.

Australia's Great Dividing Range forms an important topographical barrier running down the east coast, causing the region to receive seasonal rainfalls far greater than the surrounding landscapes. With the Australian Alps being key contributors to water industries such as the Snowy-Hydro Electric Scheme, the Murray-Darling Basin and the La Trobe Valley catchments, a better understanding of wintertime rainfall variability and change will enable such industries to manage water more efficiently in the future.

Numerous studies over the last 50 years have looked into the large-scale rainfall variability and trends in Australia and southeastern Victoria (Wright 1989; Pook et al. 2006; Risbey et al. 2009; Nicholls 2009; Risbey et al. 2012), yet significantly less work has been done focusing solely on wintertime rainfall in the Australian Alps. Studies that do have this focus, such as Landvogt et al. (2008), Chubb et al. (2011) and Fiddes et al. (2014) have primarily concentrated on synoptic classification of rainfall bearing systems.

Despite this, Nicholls (2009) notes that a decline in wintertime precipitation in the Snowy Mountains from 1958-2007 of 15% has occurred. When split into rainfall gauges of high, eastern and western locations it was found by Chubb et al. (2011) that in the period of 1990-2009 the western and high regions showed declining trends of 37% and 43% respectively whilst the eastern region was declining (insignificantly) by a rate of 26%. The authors suggest that the differences in rainfall trends and amounts are due to orographic influences upon the exposure of precipitation in each area.

Wright (1989) noted that meteorologists had long been aware that northern Victorian rainfall had a good association with tropical variability, which is quantified by the Southern Oscillation Index (SOI). Australia-wide studies such as Risbey et al. (2009) support this statement and, by using gridded data, further show that on the western side of the Australian Alps a moderate correlation exists between June, July, August (JJA) total rainfall and SOI, however on the eastern side no such correlation exists.

Risbey et al. (2009) also note a similar pattern in relation to the Indian Ocean Dipole (IOD), which is well known to affect the precipitation pattern in southeastern Australia via induced north-west cloud bands (Meyers et al. 2007). Ashok et al. (2003) also found a broad negative correlation between wintertime rainfall and the IOD in southeastern Australia.

In the work done by Hendon et al. (2007), a positive Southern Annular Mode (SAM) was found to be associated with a decline in wintertime rainfall on the western slopes of the Alps. Landvogt et al. (2008) suggested that the SAM did indeed play an important role in northeast Victorian orographic rainfall, however were unable to prove this relationship. Chubb et al. (2011) noted that during a negative SAM phase alpine precipitation amounts were 30% greater than that of a positive phase for the

winter months. Additionally, they noted the importance of SAM and the position of the westerly jet on the trajectories of rain bearing systems and their ability to bring rainfall to the region.

More recent work done in conjunction with this study by Fiddes et al. (2014) focused on the synoptic drivers of extreme wintertime rainfall in the alps. It was shown that over the western and high slopes extreme rain is brought mainly by cutoff lows and embedded lows (westerly processes) whilst the eastern slopes depended more heavily on east coast lows and easterly dips in the Tasman Sea (easterly processes). Full details on the classification of such synoptic events can be found in the Fiddes et al. (2014) paper. It was found that the westerly driven processes had a strong La Nina sea surface temperature (SST) signature, whilst the occurrence of easterly processes is related to a general warming. With this result it was hypothesized that with the occurrence of global warming and increasing SSTs, the subsequent southwards shift of the westerly storm track (Graff and LaCasce 2011) is causing the easterly processes to become more favorable than in previous climates, whilst the occurrence of the westerly processes is declining. This is further supported by studies in which the intensification of the subtropical ridge in the Australian region has been linked to the decrease in wintertime southeast Australian rainfall (Larson and Nicholls 2009, Timbal and Drosowsky 2013). In this study we explore this hypothesis in terms of the seasonal rainfall trends in the Australian Alps, and aim to explain why the western and high slopes are experiencing a decline in rainfall whilst the east region is not.

We then look to the future to see if such trends are likely to continue. Whilst high uncertainties for regional projections remain, the Fourth Intergovernmental Panel on Climate Change Assessment Report (IPCC AR4) suggests that precipitation in

southern Australia's winter and spring is likely to decline in a range of emission scenarios (Christensen et al. 2007). This reduction of rainfall across the southern part of the Australian continent is linked to a polewards shift of the subtropical ridge and westerly storm track. In the winter months the westerly storm track is at its northern most point, hence this southwards displacement is expected to cause a more dramatic decline in future wintertime precipitation.

More recently, Irving et al. (2012), in a comparison report on the Climate Model Intercomparison Project (CMIP) 3 and CMIP5, found that although both sets of models predict declines in winter rainfall of up to 5-25% from the periods 1990-1999 to 2080-2099, uncertainties in the southeast Australian region are high. It is shown for the CMIP3 ensemble that fewer than 67% of the models agree on the magnitude of the changes in southeastern Australia. The more recent CMIP5 ensemble shows only marginally more agreement of models in some areas of the southeastern Australia region. Additionally, the CMIP5 ensemble shows greater declining trends in the area than that of the CMIP3 ensemble (Irving et al. 2012).

In this work rainfall records acquired from the Australian Bureau of Meteorology (BOM) are analysed over the extended southern hemisphere winter: June, July, August and September (JJAS) period, from 1980-2011, exploring the influence of several atmospheric parameters and climate drivers.

The atmospheric parameters that correlate well with rainfall are then combined to create the Optimized Alpine Precipitation Indices, which provide a greater understanding of wintertime rainfall in the alpine region in today's climate. These indices are then applied to future climates as projected by two of the Commonwealth Scientific and Industry Research Organization's (CSIRO) CMIP3 climate models.

2. Data and Methods

2.1 Data

2.1.1 Data Selection Criteria and Quality Control

The alpine area defined in this study is the portion of the Australian Alps constrained by 35-38°S and 145.5-150°E (see Figure 1). All weather stations, obtained from the BOM, must be above 800m in altitude, with a period of focus of 1980-2011. This period was chosen as it has the most consistent amount of observational data and coincides with the beginning of the ERA-Interim reanalysis data. To be included in this study, the data must span 10 years or more, with at least 8 of those years having more than 80% daily records available.

The stations were then split into three geographical groups; the west region, the high altitude region (above 1400m, along the ridgeline) and the east region. Within these groups, the daily average rainfall over the study period was calculated. Each station was subsequently correlated back to its groups average using Pearson's correlation method (Wilks 1995). If the station had a correlation above 0.65 and more than 80% of its data had been deemed reliable by quality control checks by the BOM then the station was used in this study. If the station had not had more than 80% of the data deemed reliable by the BOM then its correlation value had to be above 0.75 for the station to be deemed reliable. After this quality control process was performed, 48 stations were retained in our analysis.

It must be noted in this work that observational data sets are inherently noisy; especially those of the alpine areas where strong winds, freezing conditions and the effects of topography can interfere with recording precipitation. Whilst the authors have attempted to remove some of this noise, observational data must be used with

caution. However, the results of this study are supported by the majority of literature on southeast Australian rainfall, and are believed to be robust.

2.1.2 Reanalysis Data

The European Centre for Medium-range Weather Forecasting's (ECMWF) ERA-Interim data is used at a resolution of $0.75^{\circ} \times 0.75^{\circ}$ to analyse the background state of the climate over events of particular interest; this data was obtained from the ECMWF Data Server (Dee et al. 2011). The National Oceanic and Atmospheric Administration's (NOAA) high-resolution ($0.25^{\circ} \times 0.25^{\circ}$) sea surface temperature (SST) data was used to determine the association of wintertime precipitation with SST anomalies (Reynolds et al. 2007). Anomaly composites from the reanalysis data sets are made using the JJAS average from 1980-2011 (or 1982-2011 for the SST) as the base climates. A one sided student t-test was used as a means of determining the significance of the anomalies calculated in the composites (Wilks 1995).

2.1.3 Climate Drivers

Climate drivers are looked at in conjunction with the above data. The Southern Oscillation Index (SOI) is used in this study as a proxy of the El Niño Southern Oscillation (ENSO), whereby the difference of standardized pressure at Tahiti and Darwin is calculated (Troup 1965). This data set was obtained from the BOM.

The Dipole Mode Index (DMI) has been used in this study as a measure of the IOD; this data was retrieved from the Japan Agency for Marine-Earth Science and Technology database. The DMI is defined as the difference in SST of the western Indian Ocean at $10^{\circ}\text{S} - 10^{\circ}\text{N}$ and $50^{\circ} - 70^{\circ}\text{E}$ from the southeastern Indian Ocean at $10^{\circ}\text{S} - 0^{\circ}$ and $90^{\circ} - 110^{\circ}\text{E}$ (Saji et al. 1999).

Lastly, the SAM index has been derived as the leading mode of empirical orthogonal function analysis of monthly mean 700hPa height anomalies polewards of 20°S (Gong and Wang 1999). This was obtained from the NOAA Climate Prediction Centre for the 1979-2000 period.

2.2 Precipitation composites

Precipitation composites were created from the average climate state experienced during an extreme precipitation event. During such events, the background climate shows clear and defined characteristics that have been found to relate well to seasonal rainfall, providing the basis for analysis. An extreme event is defined as the 95th percentile (or above) for each individual station fitting the criteria discussed above. Once events were identified, within each group (west, high, east) at least 3 stations must agree that an extreme event occurred on that day, and if so that date is recorded for analysis.

For this study, the west and high regions show very similar responses to extreme weather (Fiddes et al. 2014) and are hence averaged as one. The mean sea level pressure (MSLP), 500hPa geopotential height (GH500), temperature at 850hPa (T850), relative humidity at 500hPa (RH500) and SST anomaly composites are analysed for both the west/high and east regions. The 850hPa temperature is used in order to minimise the impacts of complex terrain.

2.3 Future Climate Projections

Having developed an understanding of current seasonal rainfall, we then look into the prediction of future rainfall with a new perspective, by not relying on precipitation projections themselves, which are shown to be generally unreliable, but

instead on projected dynamical changes of atmospheric circulation and temperature. Using this method an alternative view of the future can give a more specific climatic outlook into the precipitation of the Australian Alps.

This is done by using the Representative Climate Futures (RCF) method, developed by CSIRO, to obtain climate models that best represent a range of impact scenarios on rainfall in the Australian Alps (Clarke et al. 2011; Whetton et al. 2012). Whilst reducing the number of models used in the analysis, the RCF method offers a simple selection and a less time consuming process of obtaining and analysing future climate data.

In this work the *warmer, little change in precipitation climate* represented by the Japanese Model for Interdisciplinary Research on Climate (MIROC) Medres and the *hotter, drier climate*, represented by the CSIRO Mk3.5 models were chosen as key future climate scenarios. The emission scenario A1B was chosen, representing a future with rapid economic growth, more efficient technology and a balance across all energy sources (Nakicenovic et al. 2000). These RCFs provide a best and worst case scenario allowing us to infer the potential dynamical changes between forecasts more or less favorable for precipitation in the Australian Alps.

The CSIRO Mk3.5 model has a horizontal grid resolution of $1.875^{\circ} \times 1.875^{\circ}$ with 18 vertical levels (CSIRO 2013). The MIROC Medres, medium resolution model has 1.4° longitude by 0.5° - 1.4° latitude horizontal grid and 43 vertical levels (CSIRO 2013).

It must be noted that whilst the CMIP5 array of models is in use, little difference in current or projected precipitation exists between the CMIP5 and CMIP3 array (Irving et al. 2012). For this reason, this study has used the more easily available CMIP3 dataset.

3. Results

3.1 Precipitation trends and variability

Figure 2 shows the boxplots for winter rainfall for the high, east and west regions from 1980-2011. Immediately, it can be seen that the east region receives far less JJAS rainfall than that of both the high and west region; the average JJAS rainfall for the east group is 213.2mm, while the west group receives 559.2mm and the high receives 647.3mm. This pattern is unsurprising as in southeastern Australia prevailing westerlies are dominant and the complex topography of the Australian Alps provides a strong mechanism for rainfall producing weather. This implies that for the eastern region to receive good rainfall, it must rely on an alternative source of moisture. Additionally, it can be seen that the interquartile range for the east region has a much smaller span than that of the west, whilst the high region's span is the greatest, indicating that the east group experienced much less variability in this period of study compared to that of the west and high groups.

Figure 3 shows the trends of rainfall for the three regions from 1980-2011. It again demonstrates clearly that the east region receives far less rainfall than that of both the west and high regions. Secondly, Figure 3 shows that whilst both the high and west regions are experiencing a declining trend, the east region is not. From the first 10 years of this study (1980-89) to the last ten years (2002-11), in the west region, there has been a 20% decrease in the amount of rain received. Similarly for the high group a decrease of 18% has been experienced. At the same time, the east group has seen an insignificant decline of 6%.

Additionally, Figure 3 indicates that the high and west regions are highly correlated, which is measured at $R = 0.79$. For numerous sections in the remainder of this work, the west and high groups will be combined for simplicity. When correlating the west and high regions to the east region, lower correlations of $R = 0.45$ and $R =$

0.43 are observed respectively. This further supports the idea that the east group's rainfall is being influenced by an alternative source of moisture.

Figure 4 shows the number of days per JJAS season with recorded precipitation greater than 1mm, from 1980-2011. For each of the high, west and east regions there are no trends in the number of days of rain. This result is interesting as the amount of winter rain above is seen to be declining for the west and high regions. This implies that the intensity of rainfall events must be declining due to the simple relationship shown below, where I is intensity, f is frequency and P is the total precipitation over i number of events.

$$\sum_{i=1}^n I_i \times f_i = P_T$$

This is in agreement with other studies of southeast Australia. Risbey et al. (2012) found that in the Mallee region (to the northwest of the Australian Alps), for the period of April-October, the reduction of rainfall from frontal systems was mostly due to a reduction of the amount of rain per event, not a reduction of the number of events occurring. This reduction was associated with a reduced baroclinicity in the region. Simultaneously, Figure 4 shows that the east region receives about 36-40% fewer days of rain than both the west and high groups.

3.2 Climate drivers and alpine precipitation

A statistically significant correlation was found between west total winter rainfall and SOI where $R = 0.46$, as seen by Table 1. Similarly, the high altitude total winter rainfall and total number of rainfall days also correlate well with the SOI where $R = 0.30$ and $R = 0.46$, accounting for 10% and 21% of rainfall variability respectively. It is interesting to note that the east region's rainfall shows a very poor

correlation to SOI. This is supported by Risbey et al. (2009), where the authors also note that ENSO seems to have little effect on rainfall along the coast of NSW.

The correlation between the DMI and the west and high JJAS rainfall is respectively $R = -0.56$ and $R = -0.47$. These correlations are stronger than that of the SOI and are associated with 31% and 22% of rainfall variability. Similarly to that of the SOI, the east group's rainfall showed no relation to the DMI. This again was supported by Risbey et al. (2009) and by the significant correlation found between the DMI and the number of rainfall days in the high region ($R = -0.46$).

Interestingly, significant correlations between the east region's number of days of rain and the SOI and DMI indices are present ($R = 0.50$ and $R = -0.33$ respectively). This implies that whilst the total eastern JJAS rainfall has little correlation with climate drivers, the same cannot be said of rainfall frequency.

Significant correlations between west and high (and east) rainfall and SAM are not found. This result is interesting as previous literature (Landvogt et al. 2008, Chubb et al. 2011) suggests a relationship between winter rainfall and SAM exists. This lack of trend could be due to the relatively short time period analysed.

We will now concentrate on the dynamical influences of the climate via circulation anomalies associated with rain over the west/high and east regions, using the signatures obtained from rainfall extremes as a proxy for the calculation of seasonal anomalies.

3.3 Precipitation Composites

3.3.1 MSLP Anomaly Composites of Extreme Precipitation

Figures 5a and 5b show the MSLP anomaly composite for extreme precipitation. Looking first at the west/high composites (Figure 5a) we can see in the Tasman region a large low-pressure anomaly, whilst over Western Australia there is a

high-pressure anomaly. The high-pressure anomaly magnitude is not as large as that of the low, implying that in this composite the low-pressure anomaly is the more important feature.

For the east group (Figure 5b), the MSLP anomaly is quite different. In this case, there is an anomalous low off the southern part of the Australian east coast and to the south of this low, an intense high anomaly. Both the positive and negative anomalies are characteristic of easterly dips and east coast lows (Holland et al. 1987). It is evident that the synoptic patterns causing the anomaly are significantly different to that of the west/high pattern.

3.3.2 500hPa Geopotential Height Composite of Extreme Precipitation

The GH500 height anomaly composites are displayed in Figure 6. The west/high group (Figure 6a) shows a negative anomaly over the Tasman region, with a weaker positive anomaly to the south of western Australia and an intense positive anomaly to the east of New Zealand. The composite clearly shows a positive-negative alternating pattern, which is indicative of a wave train affecting half of the hemisphere.

Looking at the east region's GH500 anomaly plot (Figure 6b), a negative anomaly is seen in the Tasman Sea with a positive anomaly of similar intensity to the south of Western Australia and to the east of New Zealand. This composite also shows a wave train pattern, however unlike Figure 6a, it looks to be more clearly defined across the entire hemisphere, if not as intense.

Plots of geopotential height from day minus five through to day zero (not shown) show that for the west/high group a clear wave train is apparent leading up to and during the extreme precipitation events. For the east region, a weak wave train exists but is somewhat disorganised and only begins to strengthen a day before the

event. The rapid intensification of atmospheric structure is characteristic of east coast lows (Hopkins and Holland 1997). It is also interesting to observe that in both groups the anomalies are connected to the Antarctic region with negative anomalies in Figure 6a and positive anomalies in Figure 6b.

3.3.3 850hPa Temperature Composites of Extreme Precipitation

The west/high composite in Figure 7a shows that a strong negative T850 anomaly exists over the south and western part of Australia and into the Southern Ocean whilst a positive anomaly runs down the east coast of the continent and across the Tasman Sea. It is also notable that there is a strong temperature gradient running almost east-west over the alpine region, which acts to enhance baroclinicity.

The east region's temperature anomaly composite (Figure 7b) shows a negative anomaly over almost the entire Australian continent, but not extending significantly into oceanic areas. A positive anomaly again runs down the length of the east coast and the Southern Ocean. A similar temperature gradient is apparent in this composite, however of a lesser magnitude than that of the west/high composite. This suggests that although temperature gradients in the atmosphere are still important to eastern region rainfall, they are not as important as that of the west/high regions.

3.3.4 Sea Surface Temperature Composites of Extreme Precipitation

Figure 8 shows the SST anomaly composites associated with extreme rainfall. Whilst the west/high and east plots show similar features including a strong indication of a La Niña and negative IOD pattern, it can be seen that the west/high plot (Figure 8a) shows colder anomalies on the western coast of Australia compared to the east case (Figure 8b). Importantly, however, in both plots warm SST anomalies are in existence running down the majority of the east coast of Australia, indicating good

heat sources available to be transported inland, and possibly adding to the temperature gradients seen in the previous plot. In previous work done by Fiddes et al. (2014) it was shown that when the SST signals are split into westerly rainfall processes and easterly rainfall processes, the westerly processes show an enhanced La Niña/IOD response, whilst the easterly processes show a general hemispheric warming. This work is in support of the findings in this study where the west/high region's rainfall is more significantly related to the IOD and SOI than that of the east region's rainfall.

3.4 The Atmospheric Indices

By studying the above atmospheric variables in relation to extreme rainfall, one thing could clearly be seen; in each of the anomaly composites a significant dipole structure exists across the Australian region. From these composites, atmospheric indices were created using the gradient between the two opposite anomalies. To do this, boxes, 2° - 5° in length and width, were placed around the maxima and minima of the anomaly composites. The location of these boxes can be seen in Table 2. The average anomaly over these boxes is calculated and the difference between the two boxes is determined, creating a gradient index. Using the ERA-Interim data, the gradient between the boxes is calculated for every 6 hourly time step from 1980-2011, in the JJAS season. This then creates the initial index from which seasonal values can be extracted. Correlations between rainfall and climate drivers with the indices were then performed; enabling us to determine the extent that precipitation depends on the four atmospheric indices. Correlations were performed using the Pearson correlation method (Wilks 1995).

The following notation is used for the gradient indices: Δ MSLP, Δ GH and Δ T representing the MSLP, GH500, and T850 parameters. When averaged seasonally, these indices display fairly good relationships with seasonal precipitation parameters

and the background climate drivers, suggesting that the climatic signature behind extreme rainfall events is strong enough to be able to modulate the seasonal signal. This is seen in Table 3 where values in bold are significant to the 90th percentile. It is immediately clear that the MSLP gradient indices perform the best, whilst the GH500 perform the worst. Below the results are discussed in more detail.

3.4.1 Δ MSLP index

Table 3 shows that the correlation between the west/high JJAS rainfall and the Δ MSLP index is strong and highly significant, where $R = 0.52$. This demonstrates that with a more intense gradient in the MSLP there is a greater chance of seasonal rain. To add to this, a high correlation between the number of rainfall days and the MSLP gradient index exists, where $R = 0.74$.

An association between the west/high Δ MSLP index and the DMI is seen in Table 3 where $R = -0.38$. This is supported by the fact that a positive IOD is associated with a decrease in rainfall for the south Australian region (Risbey et al. 2009). Similarly to the IOD, a negative correlation with SAM has been identified. Positive SAM is associated with less rainfall in southeast Australia and literature discussed earlier supports this (Hendon et al. 2007).

Looking now to the east Δ MSLP index it is seen that a weaker relationship exists with rainfall, where $R = 0.34$ and subsequently 12% of the rainfall variability in the east region is associated with variability in the Δ MSLP index. However, a better relationship between the index and the number of rainfall days per JJAS exists, where $R = 0.48$. On the JJAS timescale the east Δ MSLP index shows no significant correlations with any of the climate drivers used in this study.

3.4.2 Δ GH index

The west/high and east Δ GH index, whilst having no significant correlations to any of the observational data or climate drivers, is still considered as an important aspect of rainfall, as it provides some information as to the structure of the upper atmosphere during good rainfall seasons.

3.4.3 Δ T index

The correlations between the west/high Δ T index and the observational data seen in Table 3 show no significant results. However, disregarding significance, each of the correlations with observational data is above ± 0.2 . This indicates that the Δ T index for the west/high group, whilst not as powerful as that of the Δ MSLP index, still has some association with the observations.

Furthermore, Table 3 shows a moderate negative correlation between the SAM index and the west/high Δ T index. In this instance $R = -0.44$, implying that the SAM is associated with almost 20% of the variation of this temperature gradient. A positive SAM is related to drier conditions and drier conditions are associated with higher average temperatures (Hendon et al. 2007). The rainfall data used in the study has been negatively correlated to maximum temperatures (not shown), and hence is consistent with the literature.

Correlating the east Δ T index to the observational data also yields no significant results and all correlations are below that of ± 0.2 , implying that little or no relationship exists. However, similar to the west/high Δ T index, a strong negative correlation between the east index and the SAM ($R = -0.62$) implies that the SAM is connected to about 38% of the variability of the gradient (not the precipitation).

3.5 Future Climates

Having determined that the atmospheric indices above can provide relationships of varying strengths with seasonal rainfall parameters, we now begin to look into what future climate scenarios from the MIROC Medres and CSIRO Mk3.5 models can tell us about the future of precipitation in alpine Australia. Here, we firstly explore what the climate model simulations are telling us, in comparison with what we know about today's climate. In this section the changes from the 1990 30 year average of MSLP, SST, T850, relative humidity at 1000hPa (RH1000) and GH500 are analysed for the 30-year average over the year 2070 and the emission scenario A1B. By looking into where and by how much key regions of the southern hemisphere are changing, we can give more meaning to the optimized alpine indices discussed in the remainder of this paper.

3.5.1 Future MSLP

The general changes for CSIRO's Mk3.5 MSLP future anomalies from the current estimated average, seen in Figure 9a, show a pattern of negative anomalies to the south of about 60°S, over the South American and African continents and over the Atlantic Ocean. In the mid-latitudes, areas of positive anomalies exist, mainly in the southeastern Pacific Ocean and south of Australia and Africa. These patterns have similar characteristics to that of blocking features across the subtropical southern hemisphere and the patterns increase in intensity with time and increasing emissions.

Comparing this to the MIROC Medres model future anomaly composite (Figure 9b) we can see that these blocking-like features are not apparent. Instead, a more annular feature is in existence, where negative anomalies occur to the south of 60°S once again, but a more uniform positive anomaly occurs to the north. The differences between the two model MSLP outputs initially suggest that the dynamical

processes within each model may be causing them to favor particular atmospheric structures, as seen here.

High-pressure anomalies over the south Australian region have been connected to years of poor winter precipitation in today's climate over the alpine region (Budin 1985).

3.5.2 Future SST

The CSIRO SST projection seen in Figure 10a shows a generalized warming of around 2°C. It can be seen that a few key warm pools have developed whereby an El Niño-like response in the Pacific exists and more locally, a warming on the western boundary of the Tasman Sea. The MIROC model, in Figure 10b, shows warming of a generally similar magnitude to the CSIRO model, but develops an even warmer pool in the western Tasman Sea and instead of a strong El Niño response in the Pacific, a more intense warming in the Indian Ocean can be seen. Literature has long related El Niño and positive IOD responses to a lack of rainfall in southeastern Australia (Meyers et al. 2007; Risbey et al 2009), which is also supported in this work for the western and high regions rainfall correlations.

3.5.3 Future T850

The climate projections for the CSIRO and MIROC models for changes in T850 can be seen in Figures 11a and 11b. Whilst generally both plots show a warming that increases with time and emissions, it is noted that the CSIRO model restricts the largest warming to over the Australian continent. On the other hand the MIROC model shows a broader, yet less intense, warming from northwards of about 40°S. Nevertheless, a warmer atmosphere, via the Clausius Clapeyron equation, has the ability to hold more moisture before saturation point is reached, potentially causing an

increase in precipitation. Additionally, in terms of Australia's snowfall, an increase of 1-4°C shown in these plots would be detrimental to the already fragile process.

3.5.4 Future GH500

The changes in the GH500 simulated by the CSIRO model can be seen in Figure 12a. By 2070, an increase of about 40-80m can be seen. Additionally, similar to that of the MSLP projections, wave train features can be seen across the hemisphere. These wave trains are an important aspect of the southern hemisphere's atmosphere and influence many of the synoptic and climatic phenomena (including precipitation) that are characteristic to Australia. Changes in the structure of such wave trains could significantly influence the atmospheric processes in the southern hemisphere.

The MIROC model's geopotential height projections, seen in Figure 12b, is similar to that of the respective MSLP projections, where a much more annular change is noted, with increases of similar magnitudes to that of the CSIRO model seen in the tropics and mid latitudes, whilst around the poles, little change is seen.

3.6 Optimized Alpine Indices

With the above knowledge of how the climate is expected to change, and the atmospheric indices defined and discussed in section 3.4, the logical next step was to attempt to combine these indices into an Optimized Alpine Precipitation Index and project this into the future. By doing this, a new set of combined indices was empirically designed to give the best relationship possible with the total rainfall and number of rainfall days. Using the Δ MSLP, Δ T and Δ GH indices defined above, equations of differing mathematical complexity were tested in process of experimentation in order to optimize the correlation with rainfall observations. Four

equations (Table 4) emerged as having correlations greater than the individual atmospheric indices. Results showed previously that the Δ MSLP index is the most powerful, having the greatest correlations with the most sets of observational data. This is not surprising as MSLP anomalies are an important driver of the atmospheric circulation leading to precipitation variability in southeast Australia. For this reason, the Δ MSLP index field has a deliberately dominant role in each of the indices defined below. The Δ GH and Δ T indices have weaker relationships with the observational data, and hence play a secondary or no role in the equations shown in Table 4.

Subsequently, using a linear regression (shown in Table 5), these new indices were projected into the future.

Figure 13 shows the optimized alpine indices for east and west/high total rainfall and number of days of rainfall for the current climate and for the projected climate of the year 2070 for the A1B emission scenario. The black dots are seasonal average of each year from 1980-2011, the red squares represent today's seasonal average, the green diamonds are CSIRO's projected seasonal average and the blue triangles MIROC's projected seasonal average.

We will firstly concentrate on the east regions optimized future index for the number of rainfall days in the top left of Figure 13. Here we can see that the current average is located roughly in the middle of the plot, whilst CSIRO's projected average is located above and the MIROC's is below. This result implies that under the worst-case scenario of the CSIRO Mk3.5 model, in the eastern region, an increase in the number of rainfall days is predicted; whilst in the best-case scenario of the MIROC Medres model, the number of rainfall days is predicted to decrease. This apparent contradiction reflects the fact that the eastern precipitation regime is largely different to that of the west and high.

This finding supports the hypothesis suggested throughout this work, where a warming of the atmosphere and ocean is leading to a decreased baroclinicity over the Australian region, a polewards shift of the westerly storm track and an increased favorability of easterly propagating rainfall processes such as easterly dips and east coast lows. If we look now to Figure 13 and the number of rainfall days for the west/high region, we can see that the predictions for the number of rainfall days are now what one would intuitively expect. The MIROC model is seen to project fewer days of rainfall than today, whilst the CSIRO model predicts even fewer, both of which continue the current trend of decline discussed in section 3.1.

Moving onto the indices for the total JJAS rainfall, seen in the middle column for Figure 13, and the message is less clear. However, when looking at the raw data (not shown), it can be seen that the same situation is present as that of the number of rainfall days. In the east region the total rainfall of the CSIRO model, in green, whilst less than today's average seen in the red, is marginally greater than that of the MIROC model in the blue. In the west/high region this is once again reversed, whereby the MIROC model predicts more total rainfall than that of the CSIRO model, whilst both are below today's average. It must be noted however, that the projection for the east region's rainfall and number of rainfall day are still of lesser magnitude than that of the west/high region.

4. Discussion and Conclusion

Wintertime precipitation in the Australian Alps shown in this work is seen to have two aspects, dependant on the location relative to the ridgeline. Firstly, the western slopes and the area along the ridgeline are seen to be experiencing a significant decline in total JJAS rainfall in the duration of this study (1980-2011). Whilst on the other side of the Alps, the eastern side displays quite a different picture;

not only is there no significant decline in rainfall but the region also receives far less JJAS rainfall. The fact that these two locations have differing rainfall signatures has been the focus of this study, whereby factors influencing this difference have been sought.

The declining rainfall on the west/high areas of the Alps is in agreement with previous literature such as Nicholls (2009) who calculated a decline in cool season rainfall of 15% in the 50-year period of 1958-2007. In this study the decline over a shorter period was found to be of greater magnitude (19%). This difference may be a result of separating the eastern region from the west and high regions, thereby enhancing the effects of declining rainfall. This decline in rainfall has previously been associated with changes in SAM and its effect on the westerly jet. Interestingly in this study the only relationship found with SAM and rainfall in the west and high regions is with the autumn immediately before the JJAS season, especially for the high region. In this instance a negative SAM in autumn implies more JJAS rainfall, facilitated by an equatorwards shift of the westerly jet (Meneghini et al. 2007).

The SOI additionally shows a good relationship with west and high region rainfall during the JJAS season. This result once again is enhanced in the high altitude region but agrees with the majority of the literature on Australian rainfall that a positive SOI, associated with La Nina events, tends to bring more rainfall to the southeast Australian region. Similarly the IOD, via the DMI, has shown a strong negative relationship in the region. Although the IOD also has some dependence on ENSO, the warming of the eastern tropical Indian Ocean has a strong influence on moisture transport diagonally across the continent through phenomena known as the northwest cloud band. This tropical influence on rainfall in the alpine region has been

commented on several times throughout the literature (Wright 1989, Landvogt et al. 2008) and is in support of such findings.

In the work done by Risbey et al. (2009), using gridded data, a clear boundary can be seen along the ridgeline of the Australian alps in terms of correlations with SOI and DMI, whereby the western side is well correlated and the eastern side is not. Their result is supported by the station data used in this study, as findings have shown that the west and high regions correlated well to several climate drivers, whilst the east side shows no correlations to any climate drivers used, implying that such climate drivers do not influence eastern rainfall as they do the western and high regions.

Projections of future climates hint to a world with warmer sea surface temperatures and a greater prevalence of anomalous high pressure in the Australian region. Such characteristics are indicative of a poleward shift of the westerly storm track and an intensification of the subtropical ridge as found by Graff and LaCasce (2011). Hopkins and Holland (1997) note that easterly dips and east coast low are associated with anomalous (if not blocking) highs as well as warm SST pools in the Tasman region. The projected changes discussed above that enhance these climatic characteristics could lead to a potential increase of easterly events, whilst the westerly events continue to decrease.

The second aim of this work was to explore the relationships of alpine rainfall with atmospheric indices and consequently create an index that can be used as an indicator of good rainfall years. The atmospheric parameters used were the MSLP, GH500 and T850. In agreement with work done by Budin (1985), the Δ MSLP index was found to have a good relationship with not only rainfall but also with the number of days of rainfall (above 1mm) per JJAS season too.

The Δ MSLP, Δ T and in one case the Δ GH indices were combined into indices aimed at optimizing the correlations with rainfall and number of rainfall days. Because the Δ MSLP showed the strongest relationships with the observations in question it became the dominant factor in our equations, whilst Δ T and Δ GH played secondary roles.

The result was 4 indices that were optimally correlated to rainfall and the number of rainfall days per JJAS season, for the east and west/high regions, where increased correlations with observational data can now explain 23%-52% of variability. Whilst these indices were not created for predictive purposes they can provide a basis for understanding the atmospheric conditions needed for a good or bad rainfall season to occur.

These findings are in support of previous results simulating the decline in rainfall across southeastern Australia, which is mainly driven by a decrease in the westerly rainfall processes such as cutoff lows and embedded lows (Christensen et al. 2007; Irving et al. 2012). However, it also supports our hypothesis as to the future of rainfall in the eastern side of the Australian Alps. This result, whilst at first unexpected, is further supported by the findings in this study and Risbey et al. (2009), showing that the eastern region's wintertime rainfall is not linked to any of the climate drivers associated with the western and high regions wintertime rainfall, or in fact southeastern Australia's.

The use of climate models in general has its own inherent uncertainties, causing scientists to err on the side of caution when using such data. Having said this however, at a preliminary stage this work presents some interesting ideas as to what the future may hold for alpine precipitation and is unique not only in its method, but in that it is specifically targeted to the Australian Alps. Whilst it can be said it is in agreement with the majority of literature on the decline on westerly influenced

rainfall, which currently dominates southeastern Australia, the increase of easterly driven processes found here, if proved likely, may become a crucial factor to the future of southeastern Australia's water resources.

Acknowledgments

S. Fiddes and A. Pezza would like to acknowledge and thank Blair Trewin (BOM), Leanne Webb (CSIRO) for access to the rainfall and climate projection data used in this study. A. Pezza would also like to thank the Australian Antarctic Division for funding parts of this work.

References

- Ashok K., Guan Z. & Yamagata T. 2003. Influence of the Indian Ocean Dipole on the Australian winter rainfall. *Geophysical Research Letters* **30**, 1821-1825.
DOI:10.1029/2003GL017926
- Budin G. R. 1985. Interannual variability of Australian snowfall. *Australian Meteorological Magazine* **33**, 145-159.
- Christensen, J.H., Hewitson B., Busuioc A., Chen A., Gao X., Held I., Jones R., Kolli R. K., Kwon W. -T., Laprise R., Magaña Rueda V., Mearns L., Menéndez G. G., Räisänen J., Rinke A., Sarr A. & Whetton P. 2007. Regional Climate Projections. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

- Chubb T. H., Siems S. T. & Manton M. J. 2011. On the Decline of Wintertime Precipitation in the Snowy Mountains of Southeastern Australia. *Journal of Hydrometeorology* 12, 1483-1497. DOI: 10.1175/JHM-D-10-05021.1
- Clarke J.M., P.H. Whetton P. H. & Hennessy K. J. 2011. Providing application-specific climate projections datasets: CSIRO's Climate Futures Framework. In: Proceedings of MODSIM, international congress on modelling and simulation, modelling and simulation society of Australia and New Zealand, 12– 16 December 2011, Perth, Australia
- CSIRO, 2013. *Science*. OzClim, accessed 17th March 2014, <<https://wiki.csiro.au/confluence/display/ozclim/Science#Science-ClimateModels>>
- Dee D. P., Uppala S. M., Simmons A. J. et al. 2011. The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society* **137**, 553–597.
- Fiddes S. L., Pezza A. B. & Barras V. 2014. Synoptic climatology of extreme precipitation in alpine Australia. *International Journal of Climatology*, DOI: 10.1002/joc.3970.
- Graff L. S. & LaCasce J. H. 2011. Changes in the Extratropical Storm Tracks in Response to Changes in SST in an AGCM. *Journal of Climate* **25**, 1854-1870. DOI: 10.1175/JCLI-D-11-00174.1
- Gong D. & Wang S. 1999. Definition of Antarctic oscillation index. *Geophysical Research Letters* **26**: 459–462.
- Hendon H. H., Thompson D. W. J. & Wheeler M. C. 2007. Australian Rainfall and Surface Temperature Variations Associated with the Southern Hemisphere Annular Mode. *Journal of Climate* **20**, 2452-2467.

- Holland G. J., Lynch A. H. & Leslie L. M. 1987. Australian East-Coast Cyclones. Part I: Synoptic Overview and Case Study. *Monthly Weather Review* **115**, 3024-3036.
- Hopkins L. C. & Holland G. J. 1997. Australian Heavy-Rain Days and Associated East Coast Cyclones: 1958–92. *Journal of Climate* **10**, 621-635.
- Irving D. B., Whetton P. & Moise A. F. 2012. Climate projections for Australia: a first glance at CMIP5. *Australian Meteorological and Oceanographic Journal* **62**, 211–225.
- Larsen S.H. and Nicholls N. 2009. Southern Australian rainfall and the subtropical ridge: Variations, interrelationships, and trends. *Geophysical Research Letters*, **36**: L08708. DOI:10.1029/2009GL037786
- Landvogt P. K., Bye J. A. & Lane T. P. 2008. An investigation of recent orographic precipitation events in northeast Victoria. *Australian Meteorological Magazine* **57**, 235-247.
- Meneghini B., Simmonds I. & Smith I. N. 2007. Association between Australian rainfall and the Southern Annular Mode. *International Journal of Climatology* **27**, 109–121. DOI: 10.1002/joc.1370
- Meyers G, McIntosh P, Pigot L, Pook M. 2007. The Years of El Niño, La Niña, and Interactions with the Tropical Indian Ocean. *Journal of Climate* **20**, 2872-2880. DOI: 10.1175/JCLI4152.1
- Nakicenovic N., Davidson O., Davis G and others. 2000. *Special Report on Emissions Scenarios : a special report of Working Group III of the Intergovernmental Panel on Climate Change*. Accessed from <<https://www.ipcc.ch/pdf/special-reports/spm/sres-en.pdf>> 28/8/2013

- Nicholls N. 2009. Local and remote causes of the southern Australian autumn-winter rainfall decline, 1958–2007. *Climate Dynamics* **34**, 835–845. DOI 10.1007/s00382-009-0527-6
- Pook M. J., McIntosh P. C. & Meyers G. A. 2006. The Synoptic Decomposition of Cool-Season Rainfall in the Southeastern Australian Cropping Region. *Journal of Applied Meteorology and Climatology* **45**, 1156-1170.
- Reynolds R. W., Smith T. M., Liu C., Chelton D. B., Casey K. S. & Schlax M. G. 2007. Daily High-Resolution-Blended Analyses for Sea Surface Temperature. *Journal of Climate* **20**, 5473-5496. DOI: 10.1175/2007JCLI1824.1
- Risbey J. S., McIntosh P. C. & Pook M. J. 2012. Synoptic components of rainfall variability and trends in southeast Australia. *International Journal of Climatology* DOI: 10.1002/joc3597
- Risbey J. S., Pook M. J., McIntosh P. C., Wheeler M. C. & Hendon H. H. 2009. On the Remote Drivers of Rainfall Variability in Australia. *Monthly Weather Review* **137**, 3233-3253. DOI: 10.1175/2009MWR2861.1
- Saji N. H., Goswami B. N., Vinayachandran P. N. & Yamagata T. 1999. A dipole mode in the tropical Indian Ocean. *Letters to Nature* **401**, 360-363.
- Timbal B. and Drosowsky W. 2013. The relationship between the decline of Southeastern Australian rainfall and the strengthening of the subtropical ridge. *International Journal of Climatology*, **33**: 1021–1034. DOI: 10.1002/joc.3492
- Troup A. J. 1965. Southern Oscillation. *Quarterly Journal of the Royal Meteorological Society* **91**: 490-506

- Whetton P., Hennessy K., Clarke J., McInnes K. & Kent D. 2012. Use of Representative Climate Futures in impact and adaptation assessment. *Climate Change* **115**, 433-442. DOI 10.1007/s10584-012-0471-z
- Wilks D. S. 1995. *Statistical methods in the Atmospheric Sciences, Volume 59 of International Geophysical Series*. Academic Press: San Diego, London
- Wright W. J. 1989. A synoptic climatological classification of winter precipitation in Victoria. *Australian Meteorological Magazine* **37**, 217-229.

Figure Captions

Fig. 1 Topographical map of the Australian Alps with the locations of weather stations. Rainfall stations from the Bureau of Meteorology (BOM) are shown in blue for the west region, black for high and red for east.

Fig. 2 Box plots of JJAS rainfall for high (left), east (middle) and west (right) groups from 1980-2011. The box depicts the 25th -75th percentiles (the first and third quartiles). The bar in the box shows the median of the data. The whiskers extend to the furthest data point that is not greater than 1.5 times the interquartile range (third minus the first quartile). Any data points outlying the whiskers are drawn as asterisks

Fig. 3 The total rainfall received in the high, east and west groups in the JJAS season from 1980-2011. The high and west group show a significant declining trend to the 95th percentile (bold) where $R = -0.37$ and $R = -0.33$ respectively. The east group shows no significant trend

Fig. 4 The number of days of rainfall received in the high, east and west groups per JJAS season from 1980-2011. There are no significant trends in this plot

Fig. 5 MSLP anomaly composites (hPa) of extreme rainfall for the west/high group (a) where $n = 154$ and the east group (b) where $n = 126$. The overlaid boxes represent the areas averaged to create the atmospheric gradient indices. Significance to the 95th percentile is shown by stippling

Fig. 6 As Figure 5 but for the 500hPa geopotential height anomaly composites (m)

Fig. 7 As Figure 5 but for the 850hPa temperature anomaly composites (°C)

Fig. 8 Sea surface temperature anomaly composites (°C) of extreme rainfall for the west/high group (a) where $n = 141$ and the east group (b) where $n = 126$. Significance to the 95th percentile is shown by stippling

Fig. 9 MSLP anomaly composites (hPa) for the 30 year average over 2070 compared to 30 year average over 1990 for the A1B scenario as projected by the CSIRO Mk3.5 model (a) and the Miroc Medres model (b).

Fig. 10 As Figure 9 but for the SST anomalies in degrees Celsius

Fig. 11 As Figure 9 but for the 850hPa temperature in degrees Celsius

Fig. 12 As Figure 9 but for the 500hPa geopotential height in meters

Fig. 13 Future Optimized Indices of the number of days of rain (left column) and total precipitation (right column) for the east group (top row) and west/high group (bottom row). Black circles indicate the seasonal values from 1980-2011 for the number of days of rain and total rainfall, and 1988-2011 for the snow accumulation. The red square represents the current climates average index value. The green diamond signifies the CSIRO's Mk3.5 models estimated index values, whilst the blue represents that of the MIROC Medres models for the A1B emission scenario for the year 2070

Total JJAS Rainfall 1980-2011

Number of Days of Rainfall per JJAS Season 1990-2011

Mean Sea Level Pressure (hPa) Anomaly Composites

500hPa Geopotential Height (m) Anomaly Composites

850hPa Temperature (c) Anomaly Composites

Sea Surface Temperature (c) Anomaly Composites

A1B 2070
Mean Sea Level Pressure Anomaly Composites

A1B 2070
Sea Surface Temperature Anomaly Composites

WEST/HIGH		
	Box A	Box B
MSLP	25° – 28°S 121° – 124°E	41.5° – 43°S 149° – 151°E
GH500	39° – 43°S 122° – 128°E	43° – 46°S 51° – 155°E

	<i>High Rainfall</i>	<i>West Rainfall</i>	<i>East Rainfall</i>
<i>SOI</i>	0.30	0.46	0.16
<i>SAM</i>	-0.19	-0.12	-0.04
<i>DMI</i>	-0.47*	-0.56*	-0.20
	<i>High Number of Rainfall days</i>	<i>West Number of Rainfall days</i>	<i>East Number of Rainfall days</i>
<i>SOI</i>	0.41*	0.17	0.50*
<i>SAM</i>	-0.09	-0.06	-0.11
<i>DMI</i>	-0.46*	-0.25	-0.33

Table 1 Correlations of precipitation parameters total JJAS rainfall, number of days of rainfall for west, high and east regions with climate drivers. Bold indicates significance to the 90th percentile and an asterisk indicates significance to the 95th percentile

T850	27° - 28.5°S 125.5° - 130°E	30° - 32°S 153° - 155°E
EAST		
	Box A	Box B
MSLP	49° - 51°S 137° - 141°E	32° - 35°S 152° - 154°E
GH500	45° - 47°S 122° - 128°E	46° - 50°S 158° - 162°E
T850	23° - 26°S 131° - 135°E	34° - 37°S 158° - 161°E

Table 2 Locations of boxes used to make the atmospheric index gradients

	West			East		
	$\Delta MSLP$	ΔGH	ΔT	$\Delta MSLP$	ΔGH	ΔT
<i>Rainfall</i>	0.52	0.03	0.20	0.34	0.12	0.07
<i>Number of Rain days</i>	0.74	-0.16	-0.23	0.48	0.00	-0.17
<i>SOI</i>	0.19	0.11	-0.25	0.16	0.03	0.05
<i>SAM</i>	-0.30	0.15	-0.44	0.19	-0.10	-0.62
<i>DMI</i>	-0.38	-0.16	0.22	-0.04	-0.167	-0.01

Table 3 The correlation values of each of the atmospheric indices with rainfall and climate drivers. Values in bold font are significant to the 90th percentile.

	West/High	East
Index for Total Rainfall	$a_r = \Delta MSLP - \log(\Delta T)$	$a_r = \Delta MSLP - \exp(\Delta T)$
Index for Number of days of Rain	$a_n = \Delta MSLP$	$a_n = \Delta T \times \log(\Delta MSLP)$

Table 4 Table of future optimized alpine indices for the west/high region and east region where a_r = Alpine index for rainfall, a_n = Alpine index for number of days of rain and Δ MSLP and Δ T are the atmospheric indices previously created

	West/High	East
Linear regression for Total Rainfall	$r = -13.79 \times a_r + 536.41$	$r = -86.05 \times a_r + 207.94$
Linear regression for Number of days of Rain	$n = 2.57 \times a_n + 60.25$	$n = 1.16 \times a_n + 35.54$

Table 5 Table of the linear regressions used for estimating rainfall parameters with the future optimized alpine indices for the west/high region and east region where r = JJAS rainfall, n = number of days of rainfall, a_r = alpine index for rainfall and a_n = alpine index for number of days of rain

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Fiddes, SL;Pezza, AB

Title:

Current and future climate variability associated with wintertime precipitation in alpine Australia

Date:

2015-05-01

Citation:

Fiddes, S. L. & Pezza, A. B. (2015). Current and future climate variability associated with wintertime precipitation in alpine Australia. CLIMATE DYNAMICS, 44 (9-10), pp.2571-2587. <https://doi.org/10.1007/s00382-014-2177-6>.

Persistent Link:

<http://hdl.handle.net/11343/283176>