Clinical subtypes of core premenstrual disorders: A Delphi survey

Khaled M K Ismail ^{*1,2}, Tracy Nevatte ^{*3} Shaughn O'Brien³, Elena Paschetta², Torbjorn Bäckström⁴, Lorraine Dennerstein⁵, Elias Eriksson⁶, Ellen W. Freeman⁷, Nick Panay⁸, Teri Pearlstein⁹, Andrea Rapkin¹⁰, Meir Steiner¹¹, John Studd⁸ and Inger Sundström-Poromma¹².

* Joint first authors

Corresponding author: Professor Khaled Ismail *MSc, MD, PhD, FRCOG*
Birmingham Women's Hospital NHS Foundation Trust
Edgbaston
Birmingham, B15 2TG
UK

Affiliation:

1. School of Clinical & Experimental Medicine, College of Medical & Dental Sciences, University of Birmingham, UK
2. Birmingham Women's Hospital NHS Foundation Trust, Birmingham, UK
3. Institute for Science and Technology in Medicine, Guy-Hilton Research Centre. Keele University, UK
4. Umeå Neurosteroid Research Center, Department of Clinical Sciences, Umeå University, Sweden
5. Department of Psychiatry, University of Melbourne and National Ageing Research Institute, Australia
6. Department of Neuroscience and Physiology, Sahlgrenska Academy, University of Gothenburg, Sweden
7. Departments of Obstetrics/ Gynaecology and Psychiatry, University of Pennsylvania, USA

8. Department of Obstetrics and Gynaecology, Chelsea and Westminster Hospital, UK
9. Warren Alpert Medical School of Brown University, USA
10. David Geffen School of Medicine at UCLA, USA
11. Departments of Psychiatry & Behavioural Neurosciences and Obstetrics & Gynecology, McMaster University, Canada
12. Department of Women's and Children's Health, Uppsala University, Sweden

Keywords: Premenstrual disorder
Core premenstrual disorder
Premenstrual syndrome
Premenstrual dysphoric disorder
Delphi method

Table 1: Percentage of group voting for each score category and the Mean and median scores for the proposed clinical subtypes in Delphi rounds

(n=12).

Proposed subtypes	First iteration									Second iteration								
	Percentage (%) of group voting for each score category						Total	Mean	Median	Percentage (%) of group voting for each score category						Total	Mean	Median
	1	2	3	4	5	6				1	2	3	4	5	6			
A- Premenstrual Dysphoric Disorder	0	0	0	8.3	8.3	83.3	69	5.75	6	0	0	0	0	8.3	91.7	71	5.92	6
B- Predominantly psychological	0	0	0	8.3	41.7	50	65	5.42	5.5	0	0	0	8.3	25	66.7	67	5.58	6
C- Predominantly physical	0	0	0	8.3	33.3	58.3	66	5.50	6	0	0	0	8.3	25	66.7	67	5.58	6
D- Predominant breast tenderness	33.3	8.3	16.7	8.3	25	8.3	37	3.08	3	25	25	16.7	0	25	8.3	36	3.00	2.5
E-Predominant bloatedness	41.7	8.3	8.3	33.3	8.3	0	31	2.58	2.5	41.7	16.7	33.3	8.3	0	0	25	2.08	2
F- Predominant weight gain	66.7	8.3	25	0	0	0	19	1.58	1	91.7	0	8.3	0	0	0	14	1.17	1
G- Predominant change of appetite	75	8.3	16.7	0	0	0	17	1.42	1	83.3	8.3	8.3	0	0	0	15	1.25	1
H- Predominant swelling	50	16.7	33.3	0	0	0	22	1.83	1.5	66.7	33.3	0	0	0	0	16	1.33	1
I- Predominant irritability	25	8.3	33.3	33.3	0	0	36	3.00	3	33.3	8.3	33.3	16.7	8.3	0	32	2.67	3

J- History of mood disorder	50	8.3	0	8.3	16.7	16.7	34	2.83	1.5		50	16.7	16.7	0	0	16.7	28	2.33	1.5
K- Predominant fatigue	91.7	8.3	0	0	0	0	13	1.08	1		100	0	0	0	0	0	12	1.00	1
L- Predominant depression	50	8.3	16.7	16.7	0	8.3	28	2.33	1.5		50	25	8.3	8.3	0	8.3	25	2.08	1.5

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Ismail, KMK;Nevatte, T;O'Brien, S;Paschetta, E;Backstrom, T;Dennerstein, L;Eriksson, E;Freeman, EW;Panay, N;Pearlstein, T;Rapkin, A;Steiner, M;Studd, J;Sundstrom-Poromaa, I

Title:

Clinical subtypes of core premenstrual disorders: a Delphi survey

Date:

2013-06-01

Citation:

Ismail, K. M. K., Nevatte, T., O'Brien, S., Paschetta, E., Backstrom, T., Dennerstein, L., Eriksson, E., Freeman, E. W., Panay, N., Pearlstein, T., Rapkin, A., Steiner, M., Studd, J. & Sundstrom-Poromaa, I. (2013). Clinical subtypes of core premenstrual disorders: a Delphi survey. ARCHIVES OF WOMENS MENTAL HEALTH, 16 (3), pp.197-201. <https://doi.org/10.1007/s00737-012-0326-7>.

Persistent Link:

<http://hdl.handle.net/11343/283208>