**EVOLVE: The Australian Rheumatology Association's 'top five' list of investigations
and interventions doctors and patients should question**

Kathleen Morrisroe, MBBS (Hons), FRACP

Rheumatologist, Department of Rheumatology St Vincent's Hospital, Melbourne

PhD student, Department of Medicine, University of Melbourne, Melbourne

Ayano Nakayama MBBS, BSc (Med), BA

Fellow in Medicine, Senior Medical Registrar, Rheumatology Department, Canberra

Hospital, PhD student and Clinical Associate Lecturer, Australian National University

Jason Soon BEc (Hons), LLB

Senior Policy Officer, Royal Australasian College of Physicians

PhD candidate, Menzies Centre for Health Policy, Sydney School of Public Health,
University of Sydney

Mark Arnold MBBS FRACP M Bioethics

Rheumatologist and Associate Dean and Head, School of Rural Health, University of
Sydney, Sydney

Les Barnsley B.Med(Hons) Grad.Dip.Epi PhD FRACP FAFRM(RACP)

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/imj.13654](https://doi.org/10.1111/imj.13654)

Staff Specialist, Department of Rheumatology, Concord Repatriation General Hospital,
Concord NSW

Claire Barrett BSc, MBBS, MRCP, FRACP

Rheumatologist, Redcliffe Hospital, Redcliffe, QLD

Prof. Peter M Brooks MBBS, FRACP, PhD

Professorial Fellow, University of Melbourne School of Population and Global Health,
Centre for Health Policy 207 Bouverie St Parkville, VIC, AUS 3010

Stephen Hall MBBS, FRACP

Associate Professor, Department of Medicine, Monash University and Cabrini Health

Patrick Hanrahan BSc, MBBS, FRACP, CCPU (Rheumatology)

Rheumatologist, Clinical Senior Lecturer Medicine, University of Western Australia

Dr Pravin Hissaria MBBS, MD, DM, FRCPA, FRACP

Department of Immunology, SA Pathology, Clinical Immunology Unit, Royal Adelaide
Hospital, Senior Clinical Lecturer, University of Adelaide

Graeme Jones MBBS (Hons), MSc, MD, FRACP, FAFPHM

Rheumatologist and NHMRC Practitioner Fellow, Menzies Institute for Medical Research,
University of Tasmania, private bag 23, Hobart, TAS, Australia

Veera Srividya Katikireddi MBChB (Hons), MRCP (UK), MEpi (Clin), FRACP
Staff Specialist Rheumatologist and Senior Clinical Lecturer, University of Queensland
Department of Medicine, The Prince Charles Hospital, Chermside QLD

Helen Keen MBBS, FRACP, PhD
Senior Lecturer, University of Western Australia, Perth
Consultant Rheumatologist, Fiona Stanley Hospital and Royal Perth Hospital, Perth.

Rodger Laurent MBBS, FRACP, PhD
Senior Staff Specialist, Royal North Shore Hospital, Department of Rheumatology, NSW

Mandana Nikpour MBBS, FRACP, PhD
Rheumatologist, Department of Rheumatology St Vincent's Hospital, Melbourne
Department of Medicine, University of Melbourne, Melbourne

Katherine Poulsen MBBBS, FRACP
Staff Specialist Rheumatologist, Department of Medicine, The Prince Charles Hospital,
Chermside QLD

Philip Robinson MBChB PhD FRACP

University of Queensland, Royal Brisbane Hospital, Bowen Bridge Road, Herston QLD

A/Prof Muriel Soden, MD, FRACP

Director of Rheumatology, The Townsville Hospital, Douglas, Qld 4814 & Academic

Coordinator, MBBS5, James Cook University, Townsville, Qld 4810.

Nigel Wood MBBS, FRACP

Staff Specialist, Rheumatology, University Hospital Geelong

Nicola Cook BM, MRCP, FRACP

Consultant Rheumatologist, Royal Perth Hospital, Perth

Catherine Hill MBBS, MD, MSc, FRACP

Staff Specialist, Rheumatology Unit, The Queen Elizabeth and Royal Adelaide Hospital and

Clinical Professor, University of Adelaide

Rachelle Buchbinder MBBS (Hons), MSc, PhD, FRACP, FAHMS

NHMRC Senior Principal Research Fellow; Director, Monash Department of Clinical

Epidemiology, Cabrini Institute; and Professor, Department of Epidemiology and Preventive

Medicine, School of Public Health and Preventive Medicine, Monash University, Melbourne,

Australia

Corresponding author

Prof Rachelle Buchbinder

Monash Department Clinical Epidemiology, Suite 41 Cabrini Medical Centre

183 Wattletree Rd, Malvern Vic 3144

Email: rachelle.buchbinder@monash.edu

Tel: 03 9509 4445

Acknowledgements

No funding was received for this project. RB is supported by an NHMRC Senior Principal Research Fellowship.

Word count abstract: 249 words

Word count manuscript: 2647 words

ABSTRACT

Background: The EVOLVE (evaluating evidence. enhancing efficiencies) initiative aims to drive safer, higher-quality patient care through identifying and reducing low-value practices.

Aims: To determine the Australian Rheumatology Association's (ARA's) 'top 5' list of low-value practices.

Methods: A working group comprising 19 rheumatologists and three trainees compiled a preliminary list. Items were retained if there was strong evidence of low-value and there was high or increasing clinical use and/or increasing cost. All ARA members (356 rheumatologists and 72 trainees) were invited to indicate their 'top 5' list from a list of 12 items via MonkeySurvey in December 2015 (reminder February 2016) .

Results: 179 rheumatologists (50.3%) and 19 trainees (26.4%) responded. The top 5 list (percent of rheumatologists including item in their top 5 list) was: Do not perform arthroscopy with lavage and/or debridement for symptomatic osteoarthritis of the knee nor partial meniscectomy for a degenerate meniscal tear (73.2%); Do not order ANA testing without symptoms and/or signs suggestive of a systemic rheumatic disease (56.4%); Do not undertake imaging for low back pain for patients without indications of an underlying serious condition (50.8%); Do not use ultrasound guidance to perform injections into the subacromial space as it provides no additional benefit in comparison to landmark-guided injection (50.3%); and Do not order anti dsDNA antibodies in ANA negative patients unless the clinical suspicion of SLE remains high (45.3%).

Conclusions: This list is intended to increase awareness among rheumatologists, other clinicians and patients about commonly used low-value practices that should be questioned.

Key words: EVOLVE, evidence-based practice, implementation, low-value care, rheumatology

Appendix 1 Recommendations included in the survey considered to be low-value care but did not make the ‘top five’ recommendations

Recommendation: Do not repeat dual energy x-ray absorptiometry (DEXA) scans more often than every two years in people with established osteoporosis

Strength of recommendation: C

NHMRC Level of evidence: III-2

Category of evidence: III

Serial bone mineral density (BMD) measurement with DEXA scans is not more predictive of subsequent fracture than the baseline measurement [44]. Due to limitations in the precision of testing, a minimum of two years is needed to reliably measure a change in BMD [45]. As a result of this evidence, many guidelines recommend a minimum of two years before re-testing BMD, and less frequent testing may be appropriate in patients with stable BMD or those at low risk of fracture [45-47].

The MBS already restricts the frequency of subsidised DEXA scans for most indications. Two MBS item codes are already restricted to no more than one service every two years (12306: to confirm low bone mineral density in a person with one or more minimal trauma fractures, or to monitor low bone mineral density proven by bone densitometry at least 12 months previously; and 12315: diagnosis and monitoring of bone loss associated with primary hyperparathyroidism, chronic liver and/or renal disease, proven malabsorptive disorder, rheumatoid arthritis and/or conditions associated with thyroxine excess) and two are restricted to no more than one service per year (12312: diagnosis and monitoring of bone loss associated with prolonged glucocorticoid therapy, conditions associated with excess glucocorticoid secretion, male hypogonadism and/or female hypogonadism lasting more than

Author Manuscript

six months before age 45 years; and 12321: measurement of bone density 12 months following a significant change in therapy for established low bone mineral density or confirmation of presumptive diagnosis of low bone mineral density made on the basis of one or more fractures occurring after minimal trauma). There is one MBS item for people aged 70 and over that currently has no frequency or other restrictions (12323). However in their second report, the Diagnostic Imaging Clinical Committee of the MBS Review Taskforce (released August 2016), have recommended that there be a modification to item 12323 to restrict subsidised repeat testing of BMD for patients with normal or mild osteopenia to once every five years, and repeat testing for patients with moderate to marked osteopenia to once every two years.

Recommendation: Do not use ultrasound guidance to perform injections into the trochanteric bursa as it provides no additional benefit in comparison to landmark-guided injection for lateral hip pain

Strength of recommendation: C

Category of evidence: IV

Lateral hip pain, also referred to as greater trochanteric pain syndrome, gluteal tendinopathy and trochanteric bursitis is a common condition of middle age and older individuals with a female predominance [48]. Local steroid injection into the site of maximal pain or into the trochanteric bursa, while unproven in randomised placebo-controlled trials, is a commonly used treatment [49]. Similar to ultrasound-guided injections for shoulder pain, increased use of ultrasound guidance to inject into the trochanteric bursa has become increasingly popular despite a lack of high quality evidence supporting its superior efficacy over landmark guided injection or injection into the site of maximal tenderness. A randomised controlled trial comparing fluoroscopic-guided injections to landmark-based injections for people with

greater trochanteric pain syndrome did not report any between group differences in outcome [49].

There has been a large increase in subsidised ultrasound-guided injections (Figure 4, main paper) but as discussed above, the exact number and costs of subsidised ultrasound-guided injections for individual conditions is unknown.

Recommendation: Do not undertake shoulder ultrasound for diagnosis of non-specific shoulder pain, which on clinical evaluation is suggestive of rotator cuff pathology and in which surgery is not planned

Strength of recommendation: D

Category of evidence: IV

Asymptomatic individuals commonly have ultrasound evidence of shoulder pathology, and this increases with age [50-53]. For example, the prevalence of rotator cuff tears was observed to be 20.7% in the general population and 36% in patients with shoulder pain in one Japanese study [52]. This means that pathology detected on ultrasound in someone with shoulder pain may not be of clinical relevance. There is no evidence to suggest that ultrasound examination of the shoulder alters primary care management of shoulder pain or improves patient outcomes and may increase the risk of over-diagnosis and/or overtreatment.

There are four item numbers on the MBS for shoulder ultrasound. Two item numbers (55808 and 55809) are used for patients referred by a medical practitioner, while two items (55810 and 55811) are used when they are not. Both 55809 and 55811, introduced in 2011, specify that benefits are not payable when a patient is referred for non-specific shoulder pain alone and to the end of the 2016 financial year there were a total of only 143 Medicare payments

under these item numbers. The numbers of MBS-subsidised shoulder ultrasounds billed under 55808 and 55810 have more than doubled since 2004 from more than 193,000 to more than 466,000 (Appendix Figure 1) and the costs have similarly almost tripled from approximately \$16.4 million to approximately \$47.15 million over that period.

Recommendation: Do not use ultrasound to investigate lateral hip pain thought on clinical assessment to be related to gluteal tendon pathology

Strength of recommendation: D

Category of evidence: IV

There is no evidence that the use of ultrasound in lateral hip pain aids diagnosis, alters management or improves outcomes. While it is expected that ultrasound could lead to unnecessary therapy, there is no evidence to support this.

MBS has two item numbers for hip and groin ultrasound examination (55816 and 55817) and two for buttock and thigh (55824 and 55825). In the 2015/2016 financial year there were 246,973 hip/groin and 35,345 buttock/thigh ultrasounds reimbursed by Medicare at a cost of \$28.55 million. As shown in Appendix Figure 2 these ultrasounds in total have increased at an annual rate of almost 14.5% over the 2004 to 2015 period.

Recommendation: Do not order extractable nuclear antibodies (ENA) testing in patients with negative ANA

Strength of recommendation: D

Category of evidence: IV

International recommendations advise testing for antibodies to ENA only after detecting a positive ANA in patients with symptoms consistent with a rheumatic disease [25]. In patients

who are ANA negative, ENA testing should only be performed in clinical situations where the pre-test probability of a rheumatic condition such as Sjögren's syndrome or inflammatory myopathy is very high. There is no evidence that serial ENA testing is useful for monitoring.

MBS-funded ENA testing has increased at a rate of almost 8.2% p.a. over the 2004 to 2015 period (Appendix Figure 3), while the total MBS benefits paid out for ENA testing has more than doubled in the same period from \$1.4 million in 2004 to \$3.8 million in 2015. As this is not linked to clinical information or ANA testing, it is not known what proportion is low-value care.

Recommendation: Do not order antineutrophil cytoplasmic antibody (ANCA) testing for diagnosis of vasculitis unless one of the consensus guideline indications is present

Strength of recommendation: C

NHMRC Level of evidence: III-3

Category of evidence: III

International consensus recommendations advise testing ANCA when the diagnosis of primary systemic small vessel vasculitides is suspected based on clinical history and examination [54]. A US study retrospectively tested these consensus recommendations and found a 23% reduction in ANCA ordering when following these recommendations without missing one cause of ANCA vasculitis [55]. These findings were subsequently confirmed in a prospective UK study [56]. Serial ANCA testing can be considered for monitoring inflammatory activity in patients with primary systemic small vessel vasculitides [54]. However, ANCA levels do not always fluctuate with disease activity and the relevance of repeat testing should be evaluated on an individual patient basis [57].

Currently MBS-funded ANCA testing is growing at a rate of 11.3% per year (Appendix Figure 4), some of which may be serial ANCA testing in patients with ANCA vasculitis. Noteworthy is that the incidence of ANCA associated vasculitides has remained stable since the early 2000s in epidemiological studies [58]. The 2014-15 cost to the federal government of this MBS funded testing (which excludes public hospital pathology testing), was more than \$4.8 million.

Recommendation: Do not order glucocorticoid injections for non-specific low back pain, facet joint arthritis or spinal canal stenosis.

Strength of recommendation: A

NHMRC Level of evidence: I

Category of evidence: Ia

There is level I evidence that epidural injections for non-specific low back pain and spinal canal stenosis confer limited short term benefit [59-61] or no benefit at all [59, 62] and do not alter long term pain, disability or progress to surgery [59]. There is a risk of complications such as vertebral fractures from spinal epidural injections [63]. There is high quality evidence that intra-articular steroid injections into the facet joints are ineffective [62, 64-66].

The MBS have two item numbers that can be used for glucocorticoid injections into the spine but neither is specific. One item number 39013 can be used for glucocorticoid injection into one or more facet joints or costo-transverse joints or for injection into one or more primary posterior rami of spinal nerves; while 18232 can be used for intrathecal or epidural injections of substances other than anaesthetic, contrast or neurolytic solutions. Use of these two items has increased by 11.7% per annum from 2004 to 2015 (Appendix Figure 5). Total benefits

paid out for these services amounted to almost \$1.9 million in 2004 but grew to almost \$8.8 million in 2015.

References

1. Hillier TA, Stone KL, Bauer DC, Rizzo JH, Pedula KL, Cauley JA, et al. Evaluating the value of repeat bone mineral density measurement and prediction of fractures in older women: the study of osteoporotic fractures. *Arch Intern Med* 2007; 167: 155-160.
2. Screening for osteoporosis: U.S. preventive services task force recommendation statement. *Ann Intern Med* 2011; 154: 356-364.
3. Screening for osteoporosis: Recommendation statement. *Am Fam Phys* 2011; 83: 1197-1200.
4. Osteoporosis: Fragility Fracture Risk: Osteoporosis: Assessing the Risk of Fragility Fracture. National Clinical Guideline 2012 National Institute for Health and Clinical Excellence.
5. Grimaldi A, Mellor R, Hodges P, Bennell K, Wajswelner H, Vicenzino B: Gluteal Tendinopathy: A Review of Mechanisms, Assessment and Management. *Sports Med* 2015; 45: 1107-1119.
6. Cohen SP, Strassels SA, Foster L, Marvel J, Williams K, Crooks M, et al. Comparison of fluoroscopically guided and blind corticosteroid injections for greater trochanteric pain syndrome: multicentre randomised controlled trial. *BMJ* 2009; 338: b1088.
7. Minagawa H, Yamamoto N, Abe H, Fukuda M, Seki N, Kikuchi K, et al. Prevalence of symptomatic and asymptomatic rotator cuff tears in the general population: From mass-screening in one village. *J Orthopaed* 2013; 10: 8-12.
8. Yamaguchi K, Tetro AM, Blam O, Evanoff BA, Teefey SA, Middleton WD. Natural history of asymptomatic rotator cuff tears: a longitudinal analysis of asymptomatic tears detected sonographically. *J Shoulder Elbow Surg* 2001; 10: 199-203.

9. Yamamoto A, Takagishi K, Osawa T, Yanagawa T, Nakajima D, Shitara H, Kobayashi T. Prevalence and risk factors of a rotator cuff tear in the general population. *J Shoulder Elbow Surg* 2010; 19: 116-120.
10. Titchener AG, White JJ, Hinchliffe SR, Tambe AA, Hubbard RB, Clark DI. Comorbidities in rotator cuff disease: a case-control study. *J Shoulder Elbow Surg* 2014; 23: 1282-1288.
11. Agmon-Levin N, Damoiseaux J, Kallenberg C, Sack U, Witte T, Herold M, et al. International recommendations for the assessment of autoantibodies to cellular antigens referred to as anti-nuclear antibodies. *Ann Rheum Dis* 2014; 73: 17-23.
12. Savige J, Gillis D, Benson E, Davies D, Esnault V, Falk RJ, et al. International Consensus Statement on Testing and Reporting of Antineutrophil Cytoplasmic Antibodies (ANCA). *Am J Clin Path* 1999; 111: 507-513.
13. Mandl LA, Solomon DH, Smith EL, Lew RA, Katz JN, Shmerling RH. Using antineutrophil cytoplasmic antibody testing to diagnose vasculitis: can test-ordering guidelines improve diagnostic accuracy? *Arch Intern Med* 2002; 162: 1509-1514.
14. Arnold DF, Timms A, Luqmani R, Misbah SA. Does a gating policy for ANCA overlook patients with ANCA associated vasculitis? An audit of 263 patients. *J Clin Path* 2010; 63: 678-680.
15. Csernok E, Moosig F. Current and emerging techniques for ANCA detection in vasculitis. *Nature Rev Rheumatol* 2014; 10: 494-501.
16. Watts RA, Mahr A, Mohammad AJ, Gatenby P, Basu N, Flores-Suarez LF. Classification, epidemiology and clinical subgrouping of antineutrophil cytoplasmic antibody (ANCA)-associated vasculitis. *Nephrol Dial Transplant* 2015; 30 Suppl 1: i14-22.

17. Friedman JH, Dighe G: Systematic review of caudal epidural injections in the management of chronic back pain. *Rhode Island Med J* 2013; 96: 12-16.
18. Friedly JL, Comstock BA, Turner JA, Heagerty PJ, Deyo RA, Sullivan SD, et al. A randomized trial of epidural glucocorticoid injections for spinal stenosis. *N Engl J Med* 2014; 371: 11-21.
19. Bresnahan BW, Rundell SD, Dagadakis MC, Sullivan SD, Jarvik JG, Nguyen H, Friedly JL. A systematic review to assess comparative effectiveness studies in epidural steroid injections for lumbar spinal stenosis and to estimate reimbursement amounts. *PM R* 2013; 5: 705-714.
20. Radcliff K, Kepler C, Hilibrand A, Rihn J, Zhao W, Lurie J, et al. Epidural steroid injections are associated with less improvement in patients with lumbar spinal stenosis: a subgroup analysis of the Spine Patient Outcomes Research Trial *Spine* 2013; 38: 279-291.
21. Mandel S, Schilling J, Peterson E, Rao DS, Sanders W: A retrospective analysis of vertebral body fractures following epidural steroid injections. *J Bone Joint Surg Am* 2013; 95: 961-964.
22. Slipman CW, Bhat AL, Gilchrist RV, Issac Z, Chou L, Lenrow DA. A critical review of the evidence for the use of zygapophysial injections and radiofrequency denervation in the treatment of low back pain. *Spine J* 2003; 3: 310-316.
23. Carrette S, Marcoux S, Truchon R, Grondin C, Gagnon J, Allard Y, Latulippe M. A controlled trial of corticosteroid injections into facet joints for chronic low back pain. *N Engl J Med* 1991; 325: 1002-1007.

Figure 2: Number of MBS-funded antinuclear antibody tests in Australia, 2004 to 2015

testing*

*MBS Item numbers included in Figure 2

71907: Antinuclear antibodies - detection in serum or other body fluids, including quantitation if required

Figure 3: Number of MBS-funded plain radiographs and CT scans for low back pain and MRI for sciatica and spinal stenosis in Australia, 2004 to 2015*

*MBS Item numbers included in Figure 3

Radiography

58106 and 58111: SPINE, lumbosacral

Computed Tomography (CT)

56223, 56229: COMPUTED TOMOGRAPHY - scan of spine, lumbosacral region, without intravenous contrast medium

56226 and 56232: COMPUTED TOMOGRAPHY - scan of spine, lumbosacral region, with intravenous contrast medium and with any scans of the lumbosacral region of the spine prior to intravenous contrast injection when undertaken

Magnetic Resonance Imaging (MRI)

63176, 63191, 63234, 63262: sciatica

63179, 63192, 63237, 63263: spinal stenosis

Figure 4: MBS-funded ultrasound-guided injections for all musculoskeletal indications in Australia from 2004-2015*

*MBS Item numbers included in Figure 4

55850, 55851: MUSCULOSKELETAL CROSS-SECTIONAL ECHOGRAPHY, in conjunction with a surgical procedure using interventional techniques, inclusive of a diagnostic musculoskeletal ultrasound service, where the referring practitioner has indicated on a referral for a musculoskeletal ultrasound that a ultrasound guided intervention be performed if clinically indicated.

Figure 5: MBS-funded dsDNA testing in Australia from 2004 to 2015*

*MBS Item numbers included in Figure 5

71099: Double-stranded DNA antibodies - quantitation by 1 or more methods other than the Crithidia method

Figure 1: MBS-funded arthroscopic washout, debridement and/or partial meniscectomy*

*MBS Item numbers included in Figure 1

49558: (KNEE, arthroscopic surgery of, involving 1 or more of: debridement, osteoplasty or chondroplasty – not associated with any other arthroscopic procedure of the knee region)

49559: (KNEE, arthroscopic surgery of, involving chondroplasty requiring multiple drilling or carbon fibre (or similar) implant; including any associated debridement or osteoplasty – not associated with any other arthroscopic procedure of the knee region)

49560: (KNEE, arthroscopic surgery of, involving 1 or more of: partial or total meniscectomy, removal of loose body or lateral release – not being a service associated with any other arthroscopic procedure of the knee region)

49561: (KNEE, ARTHROSCOPIC SURGERY OF, involving 1 or more of: partial or total meniscectomy, removal of loose body or lateral release; where the procedure includes associated debridement, osteoplasty or chondroplasty – not associated with any other arthroscopic procedure of the knee region)

49562: (KNEE, ARTHROSCOPIC SURGERY OF, involving 1 or more of: partial or total meniscectomy, removal of loose body or lateral release; where the procedure includes chondroplasty requiring multiple drilling or carbon fibre (or similar) implant and associated debridement or osteoplasty – not associated with any other arthroscopic procedure of the knee region)

Figure 1: Number of MBS-funded shoulder ultrasounds in Australia from 2004 to 2015*

*MBS Item numbers included in Figure 6

55808: SHOULDER OR UPPER ARM, 1 or both sides, ultrasound scan referred by medical practitioner (not a radiologist) for the evaluation of injury to tendon, muscle or muscle/tendon junction, rotator cuff tear/calcification/tendinosis, biceps subluxation, capsulitis and bursitis, evaluation of mass including ganglion, occult fracture and/or acromioclavicular joint pathology

55810: same but not referred by a medical practitioner

NB: Between the 2011/2 and 2015/6 financial year, Medicare paid for 28 ultrasounds under item 55809 and 115 under item 55811. These benefits are not payable when referred for non-specific shoulder pain alone.

Figure 2: Number of MBS-funded hip/groin and buttock/thigh ultrasounds in Australia, 2004-2015*

*MBS Item numbers included in Figure 7

55816 and 55817: HIP OR GROIN, 1 or both sides

55824 and 55825: 55824: BUTTOCK OR THIGH, 1 or both sides

NB: These items are only applicable if requested by medical practitioners other than radiologists.

Figure 3: MBS-funded ENA tests in Australia, 2004-2015

*MBS Item numbers included in Figure 8

71101: Antibodies to one or more extractable nuclear antigens - detection in serum or other body fluids

71103: Characterisation of an antibody detected in a service described in item 71101 (including that service)

Figure 4: MBS funded ANCA tests in Australia, 2004-2015*

*MBS Item numbers included in Figure 9

71153: Investigations in the assessment or diagnosis of systemic inflammatory disease or vasculitis - antineutrophil cytoplasmic antibody immunofluorescence (ANCA test), antineutrophil proteinase 3 antibody (PR-3 ANCA test), antimyeloperoxidase antibody (MPO ANCA test) or antiglomerular basement membrane antibody (GBM test) - detection of 1 antibody

71154: A test described in item 71153, if rendered by a receiving Australian Pathology Practitioner (APP), where no tests in the item have been rendered by the referring APP - 1 test.

71155: Detection of two antibodies described in item 71153

71156: Tests described in item 71153, other than that described in 71154, if rendered by a receiving APP – each test to a maximum of 3 tests

71157: Detection of three antibodies described in item 71153

71159: Detection of four or more antibodies described in item 71153

Figure 5: Number of MBS-funded spinal injections in Australia, 2004-2015

*MBS Item numbers included in Figure 10

39013: INJECTION UNDER IMAGE INTENSIFICATION with 1 or more of contrast media, local anaesthetic or corticosteroid into 1 or more zygo-apophyseal or costo-transverse joints or 1 or more primary posterior rami of spinal nerves (Anaes.)

18232: INTRATHECAL or EPIDURAL INJECTION of substance other than anaesthetic, contrast or neurolytic solutions, not being a service to which another item in this Group applies

Introduction

The cost of health care in Australia is growing faster than population growth. For example there was a near doubling of health expenditure over the decade 2001-02 to 2011-12.¹ This has placed an increased focus on health care quality, affordability and value. The Royal Australasian College of Physicians (RACP) EVOLVE (evaluating evidence. enhancing efficiencies) initiative is a clinician-led partnership between the College and its specialty societies. It aims to drive safer, higher-quality patient care through identifying and reducing low-value medical care, defined as tests, treatments or procedures that are overused, inappropriate or of limited effectiveness and/or potentially harmful.²

Modelled on Choosing Wisely initiatives in the United States and other countries,³ and working in conjunction with Choosing Wisely Australia,⁴ specialist physicians from over 20 medical specialities have completed or are developing their EVOLVE ‘top five’ lists of low-value clinical practices.² The guiding principles of EVOLVE are that the ‘top five’ list should be within or significantly impact the specialists’ domain of practice with the potential to make a real impact in reducing low-value care; the practices should be either growing in use or currently commonly used; and use of the Delphi consensus method,⁵ as the overarching methodology for identifying a ‘top-five’ list.

In this article we present the Australian Rheumatology Association (ARA) ‘top five’ list of the low-value practices.

Materials and Methods

The EVOLVE ARA working group comprised 19 rheumatologists and 3 advanced rheumatology trainees formed after a call for interested ARA members. At a face-to-face

meeting in 2015, the guiding EVOLVE principles were discussed and it was agreed that items should be included if they were either primarily a rheumatologist issue or an issue that rheumatologists should advocate for on behalf of their patients.

A preliminary list of low-value clinical practices was created based upon the working group's clinical experiences, as well as consideration of potentially relevant items identified from lists generated by others.⁶⁻¹⁰ The working group reduced the initial list to twelve items, noting that some items included multiple components. Two items were excluded (Do not prescribe bisphosphonates for patients at low risk of fracture, and Do not perform whole body bone scans for diagnostic screening for peripheral and axial arthritis in the adults), as these were not considered relevant to the Australian context.

Small teams for each topic were formed to review the evidence and determine if the preliminary list of low-value practices met all of the following criteria:

- i) Strong evidence of low-value clinical practice from a literature review; and
- ii) Evidence of high or increasing clinical use and significant and/or increasing cost to the Australian community based upon publicly available Medicare Benefits Schedule (MBS) item usage and cost data relating to each statement from 2004 to 2015.¹¹

Medicare Statistics provides data for MBS item numbers divided by the number of Medicare participants enrolled at the end of each month. For this project, usage data are expressed as number of services per financial year, and costs are expressed as total benefits paid out for these services by financial year. The number of services and costs included in the Medicare Statistics data only relate to services that are performed by a registered provider, qualify for the Medicare benefit, and for which a claim has been processed by Medicare Australia. They

do not include services provided by hospital doctors to public patients in public hospitals or services that qualify for a benefit under the Department of Veterans' Affairs National Treatment Account. Another important caveat of MBS data is that some single items can be used for multiple indications and the specific indication for which that item is used is not collected. For example, while there are MBS item numbers for ultrasound-guided injections, these do not differentiate between ultrasound guided injections for different body parts. In most, but not all instances we excluded item numbers for diagnostic imaging if the site being imaged was not specified.

One item, 'Do not order an HLA-B27 unless spondyloarthritis is suspected based on specific signs or symptoms', was removed from the list after the review revealed it did not fulfil the criteria of high or increasing usage or high cost in Australia. Following review of the evidence, a new item was included: 'Do not order anti-neutrophil cytoplasmic antibodies (ANCA) testing for diagnosis of vasculitis unless one of the consensus guideline indications is present'. We retained two items, 'Do not use ultrasound guidance to perform injections into the sub-acromial space (or trochanteric bursa), as it provides no additional benefit in comparison to landmark-guided injection', because even though it wasn't possible to extract the exact number and cost of these subsidised ultrasound-guided injections, consensus among the working group was that a large and increasing number of ultrasound-guided injections are being performed (inappropriately) into these sites.

The working group refined the 'do-not-do' statements, and wrote brief summaries of the evidence in support of it being a low-value clinical practice using the NHMRC recommendations for summarising the level of evidence, strength of recommendation and quality.¹²

An anonymous survey was created in SurveyMonkey.¹³ All ordinary (356 rheumatologists) and associate (72 rheumatology trainees) ARA members were invited to participate via email on 10 December 2015 with a reminder sent 17 February 2016. The ARA Board approved the survey and ethical approval was not sought.

Respondents were provided with the survey purpose and background information about EVOLVE, presented with the 12 proposed recommendations for not undertaking a particular test, treatment or procedure and a summary of the evidence for each recommendation. They were asked to select the five recommendations for which they considered the evidence to be the strongest. They could also provide comments for any of the statements in free text. Finally they were asked to provide some demographic and clinical practice details: gender, setting in which the majority of hours are worked (public, private, academic, retired, other), fellowship status (fellow for <10 years, 10-20 years, 21-30 years or >30 years), and practice location (urban/metropolitan, large rural centre, small rural centre, remote). For the purposes of our 'top five' list we excluded trainee responses.

Results

Respondents included 179 rheumatologists (50.3% response rate) and 19 trainees (26.4% response rate). The majority of rheumatologists were male (n=115, 64.3%, 4 missing responses) and just over half worked primarily in private practice (n=95, 53.1%, 5 missing responses).

Table 1 presents the proportion of rheumatologists who put each of the 12 statements into their 'top five' list in order of ranking. Endorsement of individual statements ranged from

20.7 to 73.2% of respondents. The highest endorsement was for not performing arthroscopic treatments for knee osteoarthritis and/or degenerative meniscal tears (73.2%), while over half endorsed not performing ANA testing for patients without rheumatic symptoms (56.4%), imaging for low back pain in those without specific indications (50.8%), and ultrasound guidance for shoulder injections (50.3%). Nearly all of the comments indicated that respondents would have liked to endorse more than five statements. Trainee responses were similar with four of the same recommendations chosen in the top 5 although there was even stronger endorsement for not performing arthroscopic treatments for knee osteoarthritis and/or degenerative meniscal tears (84.2%) and not performing ANA testing for patients without rheumatic symptoms (73.7%).

The top five recommendations together with a summary of the evidence that they are a low-value test or treatment and their current use/cost is summarised below. The remaining seven recommendations are described in Appendix 1.

Recommendation one: Do not perform arthroscopy with lavage and/or debridement or partial meniscectomy for patients with symptomatic osteoarthritis of the knee and/or degenerate meniscal tear

Strength of recommendation: A

NHMRC Level of evidence: I

Category of evidence: Ia

There is consistent evidence to indicate that arthroscopic lavage and/or debridement to treat people for symptomatic knee osteoarthritis, and/or partial meniscectomy for patients with a degenerate meniscal tear (with or without underlying osteoarthritis), is no more effective than placebo surgery or non-operative alternatives.¹⁴⁻¹⁹ There appears to be a high rate of

conversion from knee arthroscopy to total knee arthroplasty, which rises with increased age, further suggesting arthroscopic surgery should be avoided in people over the age of 50 years.²⁰⁻²² Additionally, arthroscopy is associated with peri and post-operative risks and considerable cost.^{18, 23}

To determine the trend in performance of knee arthroscopic treatment for knee osteoarthritis over time we considered 5 of 9 MBS codes for knee arthroscopic washout, debridement and/or partial meniscectomy (Figure 1). In total these item numbers, in people with private health insurance, increased in usage from 2004 to 2012 financial years, then appeared to plateau, and reduced by 5.9% between 2012 and 2015. Over the entire period there was an almost 2% p.a. increase. The total benefit paid out for these services was \$17.3 million in 2004 and almost \$27.1 million in 2015, corresponding to an annual growth rate of 4.15%.

Recommendation two: Do not order anti-nuclear antibody (ANA) testing in patients without symptoms and/or signs suggestive of a systemic rheumatic disease

Strength of recommendation: B

Category of evidence: III-2

ANAs are present in healthy individuals and ANA testing is only useful in patients with symptoms and/ or signs of a rheumatic disease where it can aid in the confirmation or exclusion of systemic connective tissues diseases. ANA testing has a very high negative predictive value for excluding connective tissue diseases. However a positive ANA does not have a high positive predictive value for diagnosing these conditions in isolation, and further sub-serology testing is needed to accurately diagnose and classify these conditions.^{24, 25}

Despite guidelines and recommendations not to perform an ANA test in patients without symptoms and/or signs suggestive of a connective tissue disease,²⁶⁻³⁰ there has been a steady increase over the last decade in the number of MBS-funded ANA tests ordered (Figure 2). The total benefits paid out for these services has increased from \$7.76 million in the 2004 financial year to \$10.96 million in the 2015 financial year, corresponding to an annual growth rate of 3.2%.

Recommendation three: Do not undertake imaging for low back pain in patients without indications of a serious underlying condition

Strength of recommendation: A

NHMRC Level of evidence: I

Category of evidence: Ia

Most episodes of low back pain (~90%) do not require imaging. Imaging may identify irrelevant incidental findings and increase the risk of exposure to unnecessary, and sometimes invasive treatment, in addition to increasing costs.³¹⁻³³ For patients with low back pain and no suggestion of serious underlying conditions there are no significant differences in pain or disability outcomes between immediate imaging as compared with usual care without imaging.^{34, 35}

MBS-funded imaging for low back pain has been increasing consistently since 2004 primarily due to increased numbers of CT and MRI scans (Figure 3). The total MBS benefit paid out for MRI imaging has grown from \$14.76 million in 2004 to almost \$27.96 million in 2015, an annual growth rate of almost 6%. The total benefit paid out for the other imaging modalities of CT imaging and radiography has also grown from \$58.4 million in 2004 to \$99.08 million in 2015, an annual growth rate of 4.9%.

Recommendation four: Do not use ultrasound guidance to perform injections into the sub-acromial space, as it provides no additional benefit in comparison to landmark-guided injection

Strength of recommendation: A

NHMRC Level of evidence: I

Category of evidence: Ia

Currently there is no high quality evidence to support the superiority of ultrasound-guided subacromial injections compared with injections guided by landmarks alone. Based upon moderate evidence from five trials, a Cochrane review was unable to find any advantage of ultrasound-guided injection over either landmark-guided or intramuscular injection.³⁶ These results are consistent with a more recent trial.³⁷ In view of the currently available data and the significant added cost, there is little clinical justification in using ultrasound to guide injections for shoulder pain.

The exact number and costs of subsidised ultrasound-guided injections into the subacromial space is unknown as there are two MBS item numbers that include an ultrasound-guided intervention and neither specify a body site. We consider that a substantial number of these procedures are likely to have been performed for shoulder pain. There has been an annual increase of 26.8% in the number of ultrasound-guided injections for the period 2004 to 2015 (Figure 4). In the 2014/2015 financial year the total benefits paid through the MBS for ultrasound-guided injection was almost \$27.5 million.

As a comparison the total benefits paid through the MBS for landmark-guided joint injections (MBS items 50124 and 50125) in the 2008/2009 financial year was \$12.8 million. These

Author Manuscript

were removed from the MBS on the 1st November 2009 due to a Budget decision by the government that these services are minor and routine in nature and can be delivered as part of a standard consultation. While removal of this MBS item may have resulted in a reduction in landmark-guided injection in primary and secondary care, it may have also contributed to the observed increase in more expensive image-guided injections. Several respondents made comments about the lack of reimbursement for injection, subsequent deskilling of GPs, long wait times for public rheumatology clinics, and radiologist-driven referrals as possible reasons for the increase in image-guided injection.

Recommendation five: Do not order anti-double-stranded (ds) DNA antibodies in ANA negative patients unless clinical suspicion of systemic lupus erythematosus (SLE) remains high

Strength of recommendation: B

Category of evidence: III-2

International recommendations advise testing for anti-dsDNA antibodies only after detecting a positive ANA in patients with symptoms consistent with SLE.²⁵ In patients who are ANA negative, anti-dsDNA should only be ordered in clinical situations where the pre-test probability of SLE is very high.³⁰ Where positive, repeating anti-dsDNA antibodies titres is a useful test for monitoring disease activity, especially in lupus nephritis.³⁸

The number of MBS funded anti-dsDNA tests performed over 2004 to 2015 has steadily increased (Figure 5), and the total benefits paid out for these tests more than doubled in the last decade from \$2.1 million dollars in 2004 to \$4.4 million dollars in 2015. This amounts to an average per annum growth of almost 7%. There is no epidemiologic data suggesting that the incidence of SLE is rising. For example over roughly the same time period for which

hospital separations data are available (2004 to 2014), the number of hospital separations with a principal diagnosis of SLE increased by less than 2.8% p.a.³⁹

Discussion and Conclusion

In this paper we report the top five evidence- and consensus-based recommendations for tests and procedures that Australian rheumatologists consider to be low-value care. An additional eight recommendations, while not included in the top five, were also endorsed by a significant number of rheumatologists. The most endorsed recommendation regarding arthroscopy osteoarthritis of the knee and/or degenerate meniscal tear is consistent with the recently launched Australian Clinical Care Standards for Osteoarthritis of the Knee,⁴⁰ as well as new clinical practice guideline published in the BMJ.⁴¹

While we also include similar recommendations regarding ANA, ENA, dsDNA testing and frequency of BMD monitoring to some other countries, other recommendations were not transferrable to the Australian context. For example items such as testing for Lyme disease and prescribing biologic agents prior to methotrexate were not deemed applicable to Australia due to differences in disease prevalence and mandated Medicare restrictions. This highlights the importance of creating recommendations based on local clinical practices.

In order for our 'top five' recommendations to be implemented into daily practice consideration of enablers and barriers will be required. As a first step we intend to disseminate our recommendations widely to clinicians through peer-review publication, news sites, conferences and presentations; and to consumers through the use of social media such as twitter. Additionally, some of our recommendations may be supported by other initiatives that are already taking place such as the MBS review,⁴² and new models of care for back

Author Manuscript

pain.⁴³ While this means that we will not be able to determine the precise impact of the ARA EVOLVE initiative, we plan to monitor the uptake of our recommendations using Medicare statistics data.

Acknowledgments

No funding was received for this study. RB is supported by an NHMRC Senior Principal Research Fellowship.

References

1. Australian Institute of Health and Welfare 2014. Australia's health 2014. Australia's health series number 14. Cat. no. AUS 178. Canberra: AIHW.
2. Soon J, Buchbinder R, Close J, Hill C, Allan S, Turnour C. Identifying low-value care: the Royal Australasian College of Physicians' EVOLVE initiative. *Med J Aust* 2016; 204: 180-181.
3. Choosing Wisely Canada [homepage on the Internet]. Toronto: University of Toronto, Canadian Medical Association, St Michael's [cited 18 Sept 2017]. Available from <http://choosingwiselycanada.org>
4. Choosing Wisely Australia [homepage on the Internet]. Sydney: NPS Medicine Wise [updated 13 Oct 2016; cited 18 Sept 2017]. Available from: <http://choosingwisely.org.au>
5. Linstone AT, Turoff M (eds). The Delphi method: techniques and applications. Michigan: Addison-Wesley Pub Co; 1975.
6. American College of Rheumatology: Five things physicians and patients should question [homepage on the Internet]. Philadelphia: ABIM Foundation [updated 21 Feb 2013; cited 18 Sept 2017]. Available from: <http://www.choosingwisely.org/societies/american-college-of-rheumatology/>
7. Canadian Rheumatology Association: Five things physicians and patients should question [homepage on the Internet]. Toronto: Canadian Rheumatology Association [updated 2 April 2014; cited 18 Sept 2017]. Available from: <http://www.choosingwiselycanada.org/recommendations/rheumatology>
8. Endocrine Society: Five things physicians and patients should question [homepage on the Internet]. Philadelphia: ABIM Foundation [updated 1 Oct 2013; cited 18 Sept 2017]. Available from: <http://www.choosingwisely.org/societies/endocrine-society>

9. American Academy of Orthopaedic Surgeons: Five things physicians and patients should question [homepage on the Internet]. Philadelphia: ABIM Foundation [updated 11 Sept 2013; cited 18 Sept 2017]. Available from: <http://www.choosingwisely.org/societies/american-academy-of-orthopaedic-surgeons>
10. Elshaug AG: Over 150 potentially low-value health care practices: an Australian study. Reply. *Med J Aust* 2013; 198: 597-598.
11. Medical Item Reports [homepage on the Internet]. Canberra: Australian Government Department of Human Services [updated 25 Aug 2017; cited 12th May 2017]. Available from: <http://medicarestatistics.humanservices.gov.au>
12. NHMRC additional levels of evidence and grades for recommendations for developers of guidelines. Canberra: National Health and Medical Research Council [cited 18 Sept 2017] Available at: https://www.nhmrc.gov.au/_files_nhmrc/file/guidelines/developers/nhmrc_levels_grades_evidence_120423.pdf
13. SurveyMonkey Inc. PA, California, USA (<http://www.surveymonkey.com>).
14. Buchbinder R, Richards B, Harris I. Knee osteoarthritis and role for surgical intervention: lessons learned from randomized clinical trials and population-based cohorts. *Curr Opin Rheumatol* 2014; 26: 138-144.
15. Khan M, Evaniew N, Bedi A, Ayeni OR, Bhandari M: Arthroscopic surgery for degenerative tears of the meniscus: a systematic review and meta-analysis. *CMAJ* 2014; 186: 1057-1064.
16. Katz JN, Brophy RH, Chaisson CE, de Chaves L, Cole BJ, Dahm DL, et al. Surgery versus physical therapy for a meniscal tear and osteoarthritis. *N Engl J Med* 2013; 368: 1675-1684.

17. Sihvonen R, Paavola M, Malmivaara A, Itala A, Joukainen A, Nurmi H, et al. Arthroscopic partial meniscectomy versus sham surgery for a degenerative meniscal tear. *N Engl J Med* 2013; 369: 2515-2524.
18. Thorlund JB, Juhl CB, Roos EM, Lohmander LS. Arthroscopic surgery for degenerative knee: systematic review and meta-analysis of benefits and harms. *BMJ* 2015; 350: h2747.
19. Kise NJ, Risberg MA, Stensrud S, Ranstam J, Engebretsen L, Roos EM. Exercise therapy versus arthroscopic partial meniscectomy for degenerative meniscal tear in middle aged patients: randomised controlled trial with two year follow-up. *BMJ* 2016; 354: i3740.
20. Fedorka CJ, Cernyik DL, Tauberg B, Toossi N, Johanson NA. The relationship between knee arthroscopy and arthroplasty in patients under 65 years of age. *J Arthroplasty* 2014; 29: 335-338.
21. Wai EK, Kreder HJ, Williams JJ. Arthroscopic debridement of the knee for osteoarthritis in patients fifty years of age or older: utilization and outcomes in the Province of Ontario. *J Bone Joint Surg Am* 2002; 84-a: 17-22.
22. Hawker G, Guan J, Judge A, Dieppe P. Knee arthroscopy in England and Ontario: patterns of use, changes over time, and relationship to total knee replacement. *J Bone Joint Surg Am* 2008; 90: 2337-2345.
23. Bohensky MA, Ademi Z, deSteiger R, Liew D, Sundararajan V, Bucknill A, et al. Quantifying the excess cost and resource utilisation for patients with complications associated with elective knee arthroscopy: a retrospective cohort study. *The Knee* 2014; 21: 491-496.
24. Solomon DH, Kavanaugh AJ, Schur PH. Evidence-based guidelines for the use of immunologic tests: antinuclear antibody testing. *Arthritis Rheum* 2002; 47: 434-444.

25. Agmon-Levin N, Damoiseaux J, Kallenberg C, Sack U, Witte T, Herold M, et al. International recommendations for the assessment of autoantibodies to cellular antigens referred to as anti-nuclear antibodies. *Ann Rheum Dis* 2014; 73: 17-23.
26. Yazdany J, Schmajuk G, Robbins M, Daikh D, Beall A, Yelin E, et al. Choosing wisely: the American College of Rheumatology's Top 5 list of things physicians and patients should question. *Arthritis Care Res* 2013; 65: 329-339.
27. British Columbia Guidelines: Antinuclear Antibody (ANA) Testing for Connective Tissue Disease [homepage on the Internet] British Columbia: Ministry of Health [updated 1 June 2013; cited 18 Sept 2017]. Available at: <http://www2.gov.bc.ca/gov/content/health/practitioner-professional-resources/bc-guidelines/ana-testing>
28. Tozzoli R, Bizzaro N, Tonutti E, Villalta D, Bassetti D, Manoni F, et al. Guidelines for the laboratory use of autoantibody tests in the diagnosis and monitoring of autoimmune rheumatic diseases. *Am J Clin Path* 2002; 117: 316-324.
29. Qaseem A, Alguire P, Dallas P, Feinberg LE, Fitzgerald FT, Horwitch C, et al. Appropriate use of screening and diagnostic tests to foster high-value, cost-conscious care. *Ann Intern Med* 2012; 156: 147-149.
30. Kavanaugh AF, Solomon DH. Guidelines for immunologic laboratory testing in the rheumatic diseases: anti-DNA antibody tests. *Arthritis Rheum* 2002; 47: 546-555.
31. Suri P, Boyko EJ, Goldberg J, Forsberg CW, Jarvik JG. Longitudinal associations between incident lumbar spine MRI findings and chronic low back pain or radicular symptoms: retrospective analysis of data from the longitudinal assessment of imaging and disability of the back (LAIDBACK). *BMC Musculoskel Dis* 2014; 15: 152.
32. Graves JM, Fulton-Kehoe D, Martin DP, Jarvik JG, Franklin GM. Factors associated with early magnetic resonance imaging utilization for acute occupational low back

- pain: a population-based study from Washington State workers' compensation. *Spine* 2012; 37: 1708-1718.
33. Webster BS, Bauer AZ, Choi Y, Cifuentes M, Pransky GS. Iatrogenic consequences of early magnetic resonance imaging in acute, work-related, disabling low back pain. *Spine* 2013; 38: 1939-1946.
34. Chou R, Fu R, Carrino JA, Deyo RA. Imaging strategies for low-back pain: systematic review and meta-analysis. *Lancet* 2009; 373: 463-472.
35. Jarvik JG, Gold LS, Comstock BA, Heagerty PJ, Rundell SD, Turner JA, et al. Association of early imaging for back pain with clinical outcomes in older adults. *JAMA* 2015; 313: 1143-1153.
36. Bloom JE, Rischin A, Johnston RV, Buchbinder R. Image-guided versus blind glucocorticoid injection for shoulder pain. *Cochrane Database Syst Rev* 2012; 8 :CD009147.
37. Dogu B, Yucel SD, Sag SY, Bankaoglu M, Kuran B. Blind or ultrasound-guided corticosteroid injections and short-term response in subacromial impingement syndrome: a randomized, double-blind, prospective study. *Am J Phys Med Rehab* 2012; 91: 658-665.
38. Linnik MD, Hu JZ, Heilbrunn KR, Strand V, Hurley FL, Joh T. Relationship between anti-double-stranded DNA antibodies and exacerbation of renal disease in patients with systemic lupus erythematosus. *Arthritis Rheum* 2005; 52: 1129-1137.
39. National quality statement: National Hospital Morbidity Database 2013-14 [homepage on the Internet] Canberra: Australian Institute Health and Welfare [updated 3 June 2015; cited 12 May 2017] Available at: <http://meteor.aihw.gov.au/content/index.phtml/itemId/611030>

40. Australian Commission on Safety and Quality in Health Care. Osteoarthritis of the Knee Clinical Care Standard. Sydney: ACSQHC; 2017.
41. Siemieniuk R HI, Thomas A, Poolman R, Brignardello-Petersen, R, van de Velde S, Buchbinder R, et al. Arthroscopic surgery for degenerative knee disease including arthritis and meniscal tears: a clinical practice guideline. *BMJ* 2017; 357: j1982.
42. Medicare Benefits Schedule Review [homepage on the Internet] Canberra: Australian Government Department of Health [updated 2 Aug 2017; cited 12 May 2017]
Available at:
<http://www.health.gov.au/internet/main/publishing.nsf/content/mbsreviewtaskforce>
43. NSW Agency for Clinical Innovation. Management of people with acute low back pain: model of care. Chatswood; NSW Health; 2016.

Table 1. Support for inclusion of each recommendation to be in the top 5 list by all rheumatologists and by gender and public private practice, and by trainees*

Recommendations	Rheumatologists	Male	Female	Public	Private	Trainees
	(N = 179)	(N = 115)	(N = 60)	(N = 78)	(N = 95)	(N = 19)
	%	%	%	%	%	%
Arthroscopy for symptomatic OA or degenerate meniscal tear	73.2	80	63.3	66.7	82.1	84.2
ANA testing without rheumatic symptoms	56.4	56.5	53.3	57.7	52.6	73.7
Imaging for low back pain without red flags	50.8	47.8	60.0	64.1	43.2	42.1
Ultrasound guided shoulder injections	50.3	52.2	46.7	46.2	52.6	36.8
Anti-dsDNA testing in ANA negative patients	45.3	44.4	45.0	43.6	44.2	52.6
DEXA scans more often than 2-yearly	44.1	39.1	55.0	47.4	43.2	31.6
Ultrasound guidance to perform trochanteric bursa injections	39.1	39.1	38.3	32.1	45.3	42.1
Shoulder ultrasound to diagnose non-specific shoulder pain	36.3	40.9	28.3	37.2	34.7	31.6

Ultrasound to investigate lateral hip pain	31.3	32.2	30.0	30.8	32.6	21.1
ENA testing in patients with negative ANA	27.9	28.7	25.0	21.8	31.6	42.1
ANCA testing for diagnosis of vasculitis	24.6	22.6	25.0	24.4	22.1	26.3
Glucocorticoid injections for non-specific low back pain, facet joint arthritis or spinal canal stenosis	20.7	16.5	30.0	28.2	15.8	15.8

*The percentage in each cell relates to the proportion of respondents that rated the item in their 'top five'

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Morrisroe, K;Nakayama, A;Soon, J;Arnold, M;Barnsley, L;Barrett, C;Brooks, PM;Hall, S;Hanrahan, P;Hissaria, P;Jones, G;Katikireddi, VS;Keen, H;Laurent, R;Nikpour, M;Poulsen, K;Robinson, P;Soden, M;Wood, N;Cook, N;Hill, C;Buchbinder, R

Title:

EVOLVE: The Australian Rheumatology Association's 'top five' list of investigations and interventions doctors and patients should question

Date:

2018-02-01

Citation:

Morrisroe, K., Nakayama, A., Soon, J., Arnold, M., Barnsley, L., Barrett, C., Brooks, P. M., Hall, S., Hanrahan, P., Hissaria, P., Jones, G., Katikireddi, V. S., Keen, H., Laurent, R., Nikpour, M., Poulsen, K., Robinson, P., Soden, M., Wood, N. ,... Buchbinder, R. (2018). EVOLVE: The Australian Rheumatology Association's 'top five' list of investigations and interventions doctors and patients should question. INTERNAL MEDICINE JOURNAL, 48 (2), pp.135-143. <https://doi.org/10.1111/imj.13654>.

Persistent Link:

<http://hdl.handle.net/11343/283432>