Article Type: Supplement Article

Oral Cancer

T Wong*

Oral and Maxillofacial Surgeon

D Wiesenfeld*

Oral and Maxillofacial Surgeon

* Head and Neck Tumour Stream

The Royal Melbourne Hospital and

Victorian Comprehensive Cancer Care Centre, Department of Surgery

University of Melbourne

Mr TSC Wong

Oral & Maxillofacial Surgeon

Head & Neck Tumour Stream

The Royal Melbourne Hospital and

Victorian Comprehensive Cancer Care Centre, Department of Surgery

University of Melbourne

Phone: +61 3 9347 3788

Email: tscwong1@gmail.com

Abbreviations and acronyms

AJCC American Joint Commission of Cancer

H & N Head and neck

MDT Multidisciplinary team

SCC Squamous cell carcinoma

CT Computer tomography

MRI Magnetic Resonance imaging

PET Positron emission tomography

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/adj.12594](#)

This article is protected by copyright. All rights reserved

OPG	Orthopantogram
TMN	Tumour (T), nodal (N), metastasis (M)
+	classification
PEG	Percutaneous endoscopic gastrostomy
US	Ultrasound
ENE	Extra-nodal extension

Abstract

The management of oral cancer is a multidisciplinary endeavour, as each patient presents the treating clinicians with a unique set of challenges the management of which impacts on both survival and quality of life. This article focuses on the management of oral cancer. We highlight the epidemiology and risk factors for oral cancer in Australia, the various clinical presentations that occur and the staging of oral cancer. In the vast majority of cases surgery remains the mainstay of treatment. Radiation and medical oncology is usually in an adjuvant context. Dental professionals play a critical role in many stages of management from the initial detection, to optimising pre treatment dental health and managing the short and long term sequelae of treatment. Monitoring for recurrence and the development of second primary tumours is a key role. We also introduce possible future directions of management.

Introduction

Every patient with oral cancer presents the treating clinician with a unique set of challenging, complex and multidisciplinary clinical problems, the solutions to which impact both their survival and quality of life. The management of all oral cavity cancers should occur in a Multidisciplinary Head and Neck Oncology Team. There are many different clinicians that form part of the Head and Neck Multidisciplinary Team (H&N MDT), and

these include (but are not limited to): oral and maxillofacial, ear, nose and throat and plastic & reconstructive surgeons, radiation oncologists, medical oncologists, radiologists, anatomical pathologists, anaesthetists, speech and language therapists, dieticians, head and neck nurses, physiotherapists, oral medicine specialists, prosthodontists, special needs dentists, facial prosthetists and social workers.

The management of cancers of the oral cavity is complex, due to the functional and aesthetic implications of treatment of tumours in this region. Breathing, speech, deglutition, sight, smell, taste, mastication and jaw function, are just several of the critical functions of the head and neck that can be impaired, either temporarily or permanently by the tumour or its treatment. In addition, our facial and dental aesthetics are important in how we are perceived by others; self-esteem and self-confidence may be severely affected by the tumour itself and/or its treatment.

Dentists play a critical role in the management of oral cancer, from the detection of premalignant lesions, early detection of oral cancer, management of the oral cancer patient's dentition both prior to and post definitive treatment, surveillance of recurrent or new primary tumours in conjunction with the treating specialist, and rehabilitation of missing teeth in conjunction with the treating maxillofacial surgeon and prosthodontist.

Definitions

The oral cavity is defined as the anatomical space which lies between an imaginary coronal plane drawn from the junction of the soft and hard palate and the circumvallate papillae of the tongue to the vermillion of the lips. There are seven oral cavity subsites that are used to classify the oral cavity cancer (lip, tongue, floor of mouth, buccal, hard palate, alveolar, retromolar trigone and soft palate). (Fig. 1)

[Figure 1 here]

Epidemiology

In Australia, there are more than 4000 new cases of head and neck cancers (including lip) diagnosed every year.¹ Over 600 of these cancers are oral cavity cancers. Oral cavity cancer is a highly lethal disease with a mortality rate that approaches 50%. The vast majority of oral cavity cancers are squamous cell carcinomas (SCC), other types of oral cavity cancers such as minor salivary gland malignancies, sarcomas, malignant odontogenic tumours, melanoma and lymphoma comprise less than 10% of oral cavity cancers. Dentists and dental specialists are the common referring clinicians for a patient with oral cancer.

Risk Factors

Smoking and excessive alcohol intake (> 5 standard drinks/day) are regarded as the main risk factors for the development of oral SCC in Australia. Smoking confers a 7 x relative risk of the development of oral SCC and alcohol intake of > 50g/day confers a 6 x relative risk of developing oral cancer.³ In subcontinental countries, betel nut chewing is an important risk factor in the development of oral cancer, where oral cancers represent almost 50% of all total cancer diagnoses (compared with <1% in Australia). There is an additional subgroup of non-smoking non-drinking mostly middle age female patients who are also recognised.⁴

Clinical Presentation

The clinical presentation of oral cancer is highly variable, and the presentation of oral cavity cancer is most often related to the primary tumour, with symptoms and signs from cervical or distant metastases much less common. Any oral cavity lesion, which fails to resolve in 2-3 weeks, should raise the suspicion of the treating clinician.

The most common presentations are that of an ulcerated lesion in the oral cavity, patients may also present with mobile teeth, bleeding, pain or numbness in the mouth or face or an ill fitting dental prosthesis.

[Figure 2 here]

White lesions of the oral cavity can represent a variety of diagnoses, including frictional keratosis, oral lichen planus and viral lesions such as warts. However, even 'benign' appearing white lesions can be dysplastic or even frankly malignant on biopsy. (Fig. 2)

[Figure 3 here]

Erythematous lesions must also raise suspicion of an oral cancer. In most studies, the incidence of an oral cancer/carcinoma in-situ on biopsy of erythroplakia approaches 50%². (Fig. 3)

[Figure 4 here]

Lesions that are exophytic, proliferative or papillomatous (wart like) can also be presentations of oral cancer. (Fig. 4)

[Figure 5 here]

Less commonly, oral cavity cancers can present as a cystic lesion around a tooth, which mimics that of an odontogenic cyst. This is the preoperative OPG of a patient with a radiolucent lesion around the impacted 38 extending to the 37 which on biopsy was a SCC. (Fig. 6)

[Figure 6 here]

Alternatively, a non healing extraction socket (> 6 weeks) should also raise suspicion of a possible alveolar carcinoma. (Fig 7)

[Figure 7 here]

Presentations of other oral cavity cancers apart from SCC can include a pigmented lesion (oral melanoma). (Fig. 8)

[Figure 8 here]

If there is any suspicion of a lesion that may represent oral cavity cancer, the lesion site and size should be documented, a clinical photograph taken if possible, and urgent referral made to an Oral & Maxillofacial Surgeon who is involved in a Head and Neck Oncology MDT.

Diagnosis and Staging

The diagnosis of oral cancer is dependent on obtaining a sample of tissue from the lesion, a biopsy. Ideally, the biopsy should be done by an

Oral & Maxillofacial Surgeon as the treating surgeon has the opportunity to complete a full head and neck examination, including exact measurements, palpation of lesion thickness and clinical examination of the cervical nodes +/- clinical photography and in select situations, even obtain some imaging prior to distortion of the lesion through post biopsy changes. This is especially true for early tumours of the oral cavity where the depth of the lesion can be difficult to determine on post-biopsy imaging due to tissue oedema. Broadly speaking, there are 2 types of biopsies that can be employed, incisional and excisional. In almost all situations, an *incisional* biopsy is favoured, at the margin of the lesion with 'normal' tissue, at an adequate depth for the pathologist to assess invasion of the tumour (SCC) through the lamina propria. The diagnosis of other malignancies in the oral cavity (e.g. lymphoma) may require not just histopathologic analysis (biopsy in formalin), but require fresh tissue to be sent for additional tests (e.g. flow cytometry).

A critical component in the diagnosis of oral cancer is the histopathologic analysis by an anatomic pathologist. All maxillofacial surgeons involved with oral cancer will have a close relationship with an anatomical pathologist who has a detailed knowledge of oral pathology, as there can be differences in the interpretation of a biopsy amongst pathologists. Additionally, the treating surgeon will always maintain a low threshold for re-biopsy if the clinical behaviour of the lesion is not in accordance with the 'diagnosis' from the initial biopsy.

Once the tissue diagnosis has been established, the treating surgeon will arrange appropriate radiologic scans to radiologically stage the tumour: i.e. to assess the primary tumour dimensions and invasion of adjacent structures, cervical node involvement, and whether there are distant metastases. The imaging modalities commonly used in oral cancer evaluation are computed tomography (CT), magnetic resonance imaging (MRI), ultrasound (US) and positron emission tomography (PET) to stage

the cancer. An orthopantomogram (OPG) is useful for assessment of the dentition as well as evaluation of mandibular height in the event that part of the mandible will need to be removed due to involvement or close proximity to the cancer.

CT scans of the head/neck/chest are routinely employed in assessment of oral cancer and are excellent in highlight cortical destruction, potential cervical node metastases and pulmonary metastases.

[Figure 9 here]

[Figure 10 here]

[Figure 11 here]

MRI of the neck is excellent at evaluating the soft tissue extent of the tumour, extent of marrow infiltration of the mandible or maxilla and assessment of intra or perineural involvement.

[Figure 12 here]

[Figure 13 here]

Ultrasound, used either intraorally for accessible tumours, or more commonly for assessment of cervical nodes can be combined with a fine-needle aspirate for cytologic assessment of suspicious cervical nodes.

[Figure 14 here]

PET is often combined with an anatomical imaging modality (CT or MR), and is a functional scan, where a radiotracer is administered intravenously to the patient; the tracer is preferentially taken up by cells with a high metabolic rate (a characteristic of many oral cancers). Infection and inflammation however can also provide similar radiologic appearances. It is most often used in advanced (stage 3 or 4 disease, or in salvage/recurrent cases) and the assessment of metastatic disease.

[Figures 15 (a) and (b) here]

Staging

Staging is completed according to the American Joint Commission of Cancer (AJCC) Cancer Staging Manual (version 8 which is described below will be used from 1st January 2018).⁶

Staging of the oral cavity cancer follows the TNM classification: T = tumour, N = nodal status, M = metastases.

Tumour

Tumour stage has traditionally referred to the maximum radial dimension of the tumour, but in the latest classification, the depth of the tumour also affects the T stage.

[Table 1 here]

[Table 2 here].

Accurate tumour staging is vital to determine the type of treatment offered (curative vs palliative), the ‘field’ of treatment (extent of resection for surgical treatment and area of radiation therapy for radiation treatment) as well as providing important information for treating clinicians and patients regarding the prognosis of their disease.

Principles of Management

Every patient with oral cancer should be presented at a Multidisciplinary Tumour Meeting in order that the appropriate, individualised and optimal treatment plan is offered to each and every patient. MDT's have been shown to significantly improve patient outcomes.⁴ The MDT framework allows multiple specialists and clinicians to provide input into each patient's treatment plan, allowing improved decision making, adherence to best clinical practice guidelines and the ability to also limit the impact of treatment on the patients quality of life (speech, swallow, mastication etc).

The format of each multidisciplinary meeting varies slightly between institutions, but the common elements are that the spectrum of clinicians involved in the treatment of the patient examine the patient, discuss and view the radiology and pathologic specimens/results and arrive at an

individualised management plan for the patient. The first treatment decision in each patient's management plan is to determine whether curative or palliative treatment is offered to the patient. In the management of oral cavity cancer, curative treatment is offered if the disease is surgically resectable and confined to the primary site +/- cervical nodes. Potentially curative treatment in oral SCC involves surgery with the possibility of adjuvant therapy (radiation therapy or chemoradiation therapy). Once oral SCC has spread to distant sites (eg lungs), or is surgically unresectable at the primary/cervical nodal sites due to the involvement of vital structures, then palliative treatment is offered.

Many other factors apart from the type, stage and location of the oral cancer influence the proposed management plan, in particular the patient's systemic health and nutritional status and the patient's previous treatment in a recurrent oral cancer or 'in-field' new primary. The patient's treatment commences from the time of diagnosis, with particular attention paid to optimising their systemic health, nutrition, assessment of perioperative risk (often related to cardiovascular and pulmonary function), management of their medications (eg. anticoagulants, diabetes medication etc) and obtaining consent for treatment (discussion about the nature of treatment, benefits and risks of treatment and treatment alternatives). If there is disagreement about the appropriate management plan, then further investigations may be required, or alternatively the management options are discussed with the patient and their family, highlighting the benefits/risks of each particular option and assisting the patient in making their decision. Apart from decision making regarding the treatment of the oral cancer itself, other important treatment decisions surrounding the management of the patient need to be made: type of airway in the peri-operative period (eg. temporary tracheostomy), route of nutrition (eg nasogastric tube vs PEG tube and post-operative treatment setting (ward vs high-dependency unit or intensive care unit)).

Surgery

Surgery remains the primary modality of treatment for oral cancer.

Surgery can broadly be divided into 'resective' and 'reconstructive' components. Resective surgery includes the removal of the primary tumour +/- management of the cervical nodes +/- establishment of a surgical airway (tracheostomy) if required. Reconstructive surgery essentially involves minimising the morbidity of the resection (eg replacement of tissue, minimisation of effects on speech, swallow and mastication).

The goal of the resection surgeon is to remove the oral cancer with a margin of normal tissue around the cancer in all 3 dimensions. Current clinical guidelines,⁵ recommend that a 5mm microscopic margin of normal tissue around the tumour should be the goal of the resective surgeon. To obtain a microscopic margin of >5mm around the tumour, a macroscopic radial margin of 10-15mm around the tumour is marked at the time of surgery, and the deep margin is determined by the preoperative scans and intraoperative palpation. Tumour shrinkage after resection and during pathological preparation is variable depending upon site, and may be as high as 50%.

[Figure 17 here]

The removal of the cervical lymph nodes is the 'neck dissection,' and various types of neck dissections are described depending on the 'levels' of cervical nodes removed and the preservation or sacrifice of certain structures (sternocleidomastoid, internal jugular vein and spinal accessory nerve). The majority of patients with oral cavity SCC will be indicated for a neck dissection, as any oral cavity cancer staged as T2/T3 and T4, and any T1 oral SCC > 3mm thick is recommended for a neck dissection.⁸

[Figure 18 here]

There are several types of neck dissections described, depending on the levels which are dissected and the preservation or sacrifice of certain structures.

[Table 3 here]⁷

[Figure 19 here]

The reconstruction of the surgical defect is then completed and this is well described in the following paper in this supplement.^{9,10}

Perioperative Care

The management of the patient in the immediate post-operative period is often complex and is aimed at optimising nutrition, mobilisation, pulmonary function, flap and wound care as well as airway protection, speech and swallow rehabilitation and prevention of complications (flap failure, venous thromboembolism, pulmonary infection, wound infection etc). Almost all members of the MDT are involved with the post-operative care of the patient, and the quality of this care is critical in achieving successful outcomes for the patient.

Radiation therapy

Adjuvant post-operative radiation therapy is often indicated in oral cavity cancer, however the treatment decision for this largely depends on the final histopathologic result and stage. Of critical importance is the pathologic staging of the tumour, whether there were positive lymph nodes and the status of the surgical margin. Radiation therapy involves the use of ionising radiation to destroy or damage cancer cells. It is a local treatment, and the 'field' and 'dose' of ionising radiation is individually determined for each patient. Radiotherapy in oral cancer can be direct at the primary site and/or cervical nodes and can also be used in the management of pulmonary metastases from oral SCC. There are different ways to administer radiation, but the most common form employed in the management of oral cancer is external beam irradiation where the patient lies in a fixed and repeatable position in a machine that produces high energy x-rays which target the

specific site/s. Radiation therapy can be used both in the curative and palliative setting, with curative radiation doses significantly higher (eg > 55-60Gy) than that which are used in the palliative setting. The dose of radiation is given in fractions, and the treatment period in the curative setting often lasts 6 weeks.

Chemotherapy

Chemotherapy is added to radiation if extra-capsular extension of the nodal disease is identified. Common drug protocols include cisplatin or Epidermal Growth Factor inhibitors such as cetuximab. The development of new agents and protocols is ongoing and requires extensive randomised multi-centre studies to determine efficacy before they are introduced.

[Figure 20 here]

Follow-up

Every patient with oral cancer requires long-term follow up. Clinical and/or radiologic surveillance for new and recurrent cancers is important, however in addition, there is often significant morbidity from treatment that requires further rehabilitation and treatment, including but not limited to speech and swallow rehabilitation, the preservation of the remaining dentition and restoration of missing dentition and the management of xerostomia. The psychologic and social morbidity of the cancer diagnosis and treatment must also not be overlooked and should be addressed.

Summary

Oral cavity cancer is a challenging disease with high mortality rates; dentists and dental specialists play a critical role at all stages in the management of patients. Prevention through education about smoking cessation and safe alcohol consumption is critical, detection and early referral of premalignant lesions and oral cancers and ongoing surveillance, follow up and preservation of oral health are just a few of the many roles of the dental practitioner in the management of oral cancer. Each and every

patient with oral cancer should be managed within a multidisciplinary team specialised in the management of head and neck tumours and early referral for any suspicious lesion should be made to an Oral and Maxillofacial Surgeon or Oral Medicine Specialist.

References

1. Australian Institute of Health and Welfare (AIHW) 2017. Australian Cancer Incidence and Mortality (ACIM) books: Head and neck including lip. Canberra: AIHW. URL '<http://www.aihw.gov.au/acim-books>'. Accessed June 2017.
2. Shafer WG, Waldron CA. Erythroplakia of the oral cavity. *Cancer* 1975;36:1021-8.
3. Blot WJ, McLaughlin JK, Winn DM, Austin DF, Greenberg RS, Preston-Martin S, Bernstein L, Schoenberg JB, Stemhagen A, Fraumeni JF Jr. Smoking and drinking in relation to oral and pharyngeal cancer. *Cancer Res* 1988;48:3282-7.
4. Koo K, Barrowman R, McCullough M, Iseli T, Wiesenfeld D. Non-smoking non-drinking elderly females: a clinically distinct subgroup of oral squamous cell carcinoma patients. *Int J Oral Maxillofac Surg*. 2013;42:929-33.
5. Friedland PL, Bozic B, Dewar J, et al. Impact of multidisciplinary team in the management in head and neck cancer patients. *Br J Cancer* 2011;104:1246–8.
6. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Head and Neck Cancers. V 1.2015. URL: 'http://www.nccn.org/professionals/physician_gls/pdf/head-and-neck.pdf'. Accessed December 2015.

7. American Joint Committee on Cancer Staging Manual, 8th edition. 2016. URL '<https://cancerstaging.org/About/Pages/8th-Edition.aspx>'. Accessed June 2016.
8. Neck Dissection Classification Update. Revisions Proposed by the American Head and Neck Society and the American Academy of Otolaryngology–Head and Neck Surgery Arch Otolaryngol Head Neck Surg. 2002;128:751-758.
9. Spencer K. Implant based rehabilitation options for the atrophic edentulous jaw. Aust Dent J Supplement 2018;S
10. Batstone M. Reconstruction of major defects of the jaws. Aust Dent J Supplement 2018;S.

Tables

Table 1. Oral Cavity

• T0 deleted
• T1: size $\leq 2\text{cm}$ and DOI $\leq 5\text{mm}$
• T2: size $\leq 2\text{cm}$ and DOI 5-10mm or Size 2-4cm and DOI $\leq 10\text{mm}$
• T3: size $> 4\text{cm}$ or $> 10\text{mm}$ DOI
• T4a extrinsic tongue muscle infiltration Now deleted

DOI = depth of invasion

Table 2. Nodal status

N Category	N Criteria
NX	Regional lymph nodes cannot be assessed
N0	No regional lymph node mets
N1	Mets in single ipsilateral node $\leq 3\text{cm}$ in greatest dimension and ENE (-)

N2a	Mets in single ipsilateral or contralateral node >3cm but ≤6cm in greatest dimension and ENE(-)
N2b	Met in multiple ipsilateral nodes, none >6cm in greatest dimension and ENE(-)
N2c	Met in bilateral or contralateral nodes, none >6cm in greatest dimension and ENE(-)
N3a	Met in a node >6cm in greatest dimension and ENE(-)
N3b	Met in single ipsilateral node ENE(+) or mutiple ipsilateral, contralateral, or bilateral nodes, any with ENE(+)

ENE = extranodal extension

Metastases

M0 = no distant metastases

M1 = distant metastases

Table 3. Classification of Neck Dissection^{7,8}

1991 Classification	2001 Classification
1. Radical neck	1. Radical neck

dissection	dissection
2. Modified radical neck dissection	2. Modified radical neck dissection
3. Selective neck dissection	3. Selective neck dissection: Each variation is depicted by "SND" and the use of parentheses to denote the levels or sublevels removed
a. Supraomohyoid	
b. Lateral	
c. Posterolateral	
d. Anterior	
4. Extended neck dissection	4. Extended neck dissection

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Wong, TSC;Wiesenfeld, D

Title:

Oral Cancer

Date:

2018-03-01

Citation:

Wong, T. S. C. & Wiesenfeld, D. (2018). Oral Cancer. AUSTRALIAN DENTAL JOURNAL, 63, pp.S91-S99. <https://doi.org/10.1111/adj.12594>.

Persistent Link:

<http://hdl.handle.net/11343/283825>