PROF. SAMUEL F. BERKOVIC (Orcid ID : 0000-0003-4580-841X)

Article type : Critical Review – Invited Commentary

Can mutation-mediated effects occurring early in development cause long-term seizures susceptibility in genetic generalised epilepsies?

Running title: Early developmental changes in genetic generalised epilepsy

Authors: Christopher Reid¹, Ben Rollo¹, Steven Petrou¹ & Samuel F Berkovic²

¹ The Florey Institute for Neuroscience and Mental Health, The University of Melbourne, Parkville 3052, Victoria, Australia

² Epilepsy Research Centre, Department of Medicine, University of Melbourne, Austin Health, Heidelberg 3084, Victoria, Australia

***Corresponding author:** Samuel F Berkovic

Epilepsy Research Centre, Department of Medicine, University of Melbourne, Austin Health, Heidelberg 3084, Victoria, Australia. Tel: +613 9035 7330

E-mail: s.berkovic@unimelb.edu.au

Running title: Early development in genetic epilepsy

Key words: epilepsy, genetic, GABA_A receptor, MRI, brain structure, development

Number of text pages: 12

Number of words: 3000

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/epi.14077](https://doi.org/10.1111/epi.14077)

This article is protected by copyright. All rights reserved

1 Figure and 0 tables

Abstract (159)

Epilepsy has a strong genetic component, with an ever increasing number of disease-causing genes being discovered. Most epilepsy-causing mutations are germ line and thus present from conception. These mutations are therefore well positioned to have a deleterious impact during early development. Here we review studies that investigate the role of genetic lesions within the early developmental window specifically focusing on genetic generalised epilepsy (GGE). Literature on the potential pathogenic role of sub-mesoscopic structural changes in GGE is also reviewed. Evidence from rodent models of genetic epilepsy support the idea that functional and structural changes can occur in early development leading to altered seizure susceptibility into adulthood. Both animal and human studies suggest that sub-mesoscopic structural changes occur in GGE. The existence of sub-mesoscopic structural changes prior to seizure onset may act as biomarkers of excitability in genetic epilepsies. We also propose that pre-symptomatic treatment may be essential for limiting the long term consequences of disease-causing mutations in genetic epilepsies.

- Genetic contributions are increasingly being recognised in the pathogenesis of epilepsy with mutations mostly present from conception.
- Disease-causing mutations can impact in early development to set seizure susceptibility into adulthood in animal models of disease.
- Sub-mesoscopic structural changes occur in genetic generalized epilepsy and may be biomarkers of excitability.
- Anti-epileptogenic drugs used before overt symptoms are observed may alleviate both seizures and comorbidities.

Epilepsy affects up to 4% of the population at some time in their lives ¹. It is characterized by recurrent paroxysms resulting from hyper-synchronous discharges in the brain. There are major challenges in clinical epilepsy care with at least 30% of patients resistant to current therapies. Even amongst those patients whose seizures are controlled, major issues of drug side-effects and comorbidities often affect quality of life. Although it is difficult to establish an exact proportion, it is becoming increasingly clear that genetic factors play a significant pathogenic role in a majority

of the epilepsies with well over 50 epilepsy genes now identified ²⁻⁵. Knowledge of these causative mutations provides an unprecedented opportunity to investigate the underlying mechanisms of epileptogenesis in the genetic epilepsies. Much effort is placed on understanding the acute impact of genetic variation on neuronal excitability with the view that the protein products of this variation drive cellular dysfunction. However, because of the coordinated control of gene expression, these genetic mutations can exert their effects at different periods throughout life and are therefore positioned to divert the course of brain development and consequently long-term seizure susceptibility (Figure 1). Here we review both clinical and experimental evidence suggesting that genetic mutations can impact early development and discuss this paradigm with specific focus on genetic generalised epilepsy (GGE).

Genetic epilepsies

The new ILAE Classification stipulates that "The concept of a genetic epilepsy is that it directly results from a known or presumed genetic mutation in which seizures are a core symptom of the disorder" ⁶. The archetypical genetic epilepsy is GGE, also known as idiopathic generalized epilepsy (IGE). This form of epilepsy has the highest heritability of the common epilepsies, although the majority of genes that increase the risk for GGE are yet to be identified. GGE is a family of related syndromes that are associated with a characteristic generalized spike-and-wave discharge on electroencephalogram (EEG) and usually beginning in childhood or adolescence ⁷. Common GGE syndromes include childhood absence epilepsy, juvenile absence epilepsy, juvenile myoclonic epilepsy and generalized tonic-clonic seizures alone ⁶. There is also an increasingly recognized genetic component to focal epilepsies ⁸ and the rarer but devastating epileptic encephalopathies have emerged as a group of predominantly *de novo* dominant disorders ³. Although the latter two groups are not the focus of this review it is of interest that genes for the epileptic encephalopathies also raise risk for GGE ² suggesting that biological observations made in GGE may be generally applied.

Functional developmental changes altering long-term seizure susceptibility in models of genetic epilepsy

Rodent models of generalised epilepsy have given insight into the potential role genetic lesions play in early development. Investigations using the Wistar albino

Glaxo rats of Rijswijk (WAG/Rij) model of polygenic GGE revealed that treatment with the antiepileptic drug ethosuximide during the early epileptogenic period has a sustained impact on disease progression⁹. A persistent suppression of seizures was observed even after the therapy was discontinued⁹. This early treatment also blocked changes in the expression of Na_v1.1, Na_v1.6, and HCN1 ion channels that are markers of disease in this rat model⁹. A long term protective effect of early antiepileptic drug treatment on seizure susceptibility was also observed in the Genetic Absence Epilepsy Rats from Strasbourg (GAERS), another inbred model of polygenic GGE¹⁰. Similarly, treatment in early development of a transgenic mouse model, based on a dysfunctional Kv7 channel, with the sodium-potassium-chloride cotransporter 1 (NKCC1) antagonist bumetanide normalised hippocampal structural changes, electrophysiological properties and some behavioural markers of hyperexcitability¹¹.

To test this idea more directly, a conditional mouse model based on the GABA_A $\gamma 2_{R43Q}$ GGE mutation was engineered allowing the investigation of the impact of a single known epilepsy gene in early development^{12; 13}. Although rare, patients harbouring the GABA_A $\gamma 2_{R43Q}$ mutation have epilepsy phenotypes that are common, including childhood absence epilepsy and febrile seizures¹³. Understanding the principal pathogenic mechanisms caused by the GABA_A $\gamma 2_{R43Q}$ are expected to have broader implications, although their relevance for understanding GGE due to other genetic mechanisms is not currently known. Mice in which the mutation had been silenced during early development were less susceptible to a proconvulsant challenge than mice that had expressed the epilepsy-causing mutation from conception¹⁴. This suggests that the GABA_A $\gamma 2_{R43Q}$ mutation is most likely altering neuronal network architecture in the early developmental window to set seizure susceptibility later in adulthood.

Similar findings have been observed in other mouse models of epilepsy. *SLC2A1* encodes for GLUT1 that is responsible for glucose transport across the blood brain barrier with mutations causing deficiency in this protein accounting for up to 1% of GGE¹⁵. GLUT1 is also a cause of early onset CAE¹⁶. A *Slc2a1* haploinsufficient mouse recapitulates many of the features of GLUT1-deficiency including a spike and wave discharge phenotype¹⁷. Recent efforts have demonstrated that viral-mediated gene replacement therapy was able to reverse many of the disease phenotypes in the

Glut1-deficient mouse including epileptiform activity¹⁸. Interestingly, Tang and colleagues show that the effectiveness of gene therapy is largely limited to treatment in a narrow pre-symptomatic postnatal window¹⁸. Although not GGE specific, another example includes a transgenic mouse model of familial nocturnal frontal lobe epilepsy based on a *CHRNA2* mutation that causes a gain-of-function in the encoded $\beta 2$ nicotinic receptor¹⁹. When expressed throughout life mice carrying this mutation display a spontaneous epileptic phenotype with high frequency interictal spikes and seizures¹⁹. In a conditional model of this disease, the silencing of the transgene expression during early development was sufficient to prevent the occurrence of seizures in adulthood¹⁹. Further, seizures in a mouse model of Angelman syndrome (an epileptic encephalopathy) based on the loss of function of the maternal ubiquitin protein ligase E3A (UBE3A) are only suppressed when the protein is reinstated in an early developmental period²⁰. Together, these studies strongly suggest that there are pre-symptomatic changes occurring in neuronal networks that are caused, at least in part, by underlying genetic factors. Altered neuronal network structure during early development is one mechanism which may cause long-term increases in seizure susceptibility and is discussed next.

Structural brain abnormalities in generalized epilepsy

Human studies

There is strong evidence linking structural changes in the brain and seizure genesis in acquired focal epilepsy. Temporal lobe epilepsy, the archetypal acquired epilepsy syndrome, is sometimes associated with structural changes, particularly hippocampal sclerosis²¹. Similarly, for genetic epilepsies such as tuberous sclerosis and familial multiple cavernous malformation syndrome, brain malformations are evident and clearly contribute to hyperexcitability. Surgical removal of the brain malformation in both these genetic diseases can be effective in reducing seizure burden^{22; 23}.

In contrast, GGE mesoscopic scale structural lesions cannot be detected by conventional diagnostic methods such as gross anatomical analysis or magnetic resonance imaging (MRI). Historically this was part of the basis for the designation of these epilepsies as 'idiopathic'. However, studies based on more advanced imaging technologies have begun to resolve subtle differences in brain structure in GGE²⁴⁻²⁷. For example, MRI volumetry and voxel-based morphometry studies have reported

regional changes, particularly in the thalamus, of GGE patients ^{24; 26}. Changes in cortical gray matter volumes have also been reported ^{24; 28}. Whether these more subtle structural changes are a contributing factor to, or a consequence of, seizures is not known. It is also not clear if these structural changes occur in pre-symptomatic patients when they could act as ideal biomarkers of disease progression. Animal models that enable investigation of neuronal morphology changes in the pre-seizure brain will help us to resolve the relationships between these structural changes and disease progression.

Rodent Models

Preclinical MRI studies in rodent models support the emerging view that structural brain changes can occur in GGE. Structural MRI has detected differences in GAERS including increases in amygdala, cortical and ventricular volumes ²⁹. Axonal connectivity changes measured using diffusion tensor imaging suggest that chronic seizures in the WAG/Rij and GAERS rat models of GGE causes reduction in myelin and/or decreased axon fibre density changes in the corpus callosum ³⁰. Functional connectivity studies using MRI have also demonstrated abnormally high degree of cortical-cortical correlations in the WAG/Rij model supporting the idea that neuronal connectivity is altered in these mice ³¹. However, studies reporting structural changes all investigate rodents that are already epileptic making it difficult to assess if these morphological changes are a result of seizure activity or are themselves the cause of seizures. In support of the former, axon fibre density changes are not seen in pre-seizure WAG/Rij rats suggesting that at least some changes occur as a result of the seizures themselves ³⁰. Unfortunately, evidence for structural changes in the absence of overt seizures is limited. In one study, high field MRI in fixed brains of the GABA_A $\gamma 2_{R43Q}$ mouse model of GGE revealed a small increase in dentate granule cell layer volume of the hippocampi of non-seizing mice ³². This marks one of the first indications that morphological changes can occur in GGE due to the direct impact of an epilepsy mutation and not as a consequence of seizure activity. Although these morphological changes are well placed to contribute to enhanced excitability, experiments have not yet been performed which can elucidate their role in pathogenesis. Ongoing imaging studies in other genetic rodent models of GGE are required to assess if similar pre-symptomatic morphological changes are a common hallmark of the disease.

Current imaging techniques are not sensitive to various microscopic changes in neuronal morphology such as changes in neuron shape; and position, arborisation and density of dendritic spines and axonal boutons. Such changes can provide structural substrates for pathological changes in excitability. In the GABA_A $\gamma 2_{R43Q}$ mouse model stereological analysis of seizure free brains showed a 33% increase in dentate granule cell density in the hippocampus³². Further quantitative morphological analysis of the GABA_A $\gamma 2_{R43Q}$ mouse using three-dimensional reconstruction of cortical columns showed that changes in inhibitory neuron numbers and positions presaged the occurrence of seizures³³. Specifically, the densities of GABA-, calretinin-, parvalbumin-, and calbindin-containing inhibitory neurons were increased in the cortex of GABA_A $\gamma 2_{R43Q}$ mice³³. Morphological analysis has also revealed changes in the microarchitecture of GAERS brains. Similar to that observed in the GABA_A $\gamma 2_{R43Q}$ mouse, GABAergic neuron numbers are increased in the lateral geniculate, ventral posteromedial, and ventral posterolateral thalamic nuclei of GAERS model when compared to Wistar controls³⁴. Although this seems paradoxical, others have proposed circuit level increases in GABA_A receptor-mediated transmission can be epileptogenic³⁵. Finally, in the WAG/Rij model, Golgi staining reveals differences in dendritic arborisation in the cortex³⁶. Collectively, this data strongly supports that microarchitecture changes occur in the genetic epilepsies prior to and after the appearance of seizures and that subtle change in brain structure may be part of an underlying epileptogenic and post seizure pathogenic processes. More advanced non-invasive imaging modalities that are capable of detecting these microscopic changes are needed to test these ideas in pre-clinical animal models and humans.

Exploring the impact of a GABA_A epilepsy mutation in early development

To develop new therapeutic approaches targeted within early developmental windows we will need to resolve underlying pathogenic mechanisms. Dysfunctional proteins may exert their impact through a raft of different mechanisms (Figure 1), during different developmental time periods and each mutation may do so in a unique way. However, in the genetic epilepsies it is likely that common mechanistic pathways will emerge that could be used as drug targets to treat patients before seizures occur.

One broad pathogenic mechanism may be the disruption of early developmental oscillatory behaviour that is critical for the refinement of neuronal networks. Discontinuous structure of EEG activity, alternating between periods of oscillatory discharges and network silence, in premature infants are thought to reflect such activity³⁷. Rodents at birth provide ideal models for investigating these early events given they correspond to the second gestational trimester in humans³⁸. Neonatal spindle bursting is an example of an early developmental oscillatory network behaviour in rodents that is similar to that seen in humans³⁷. Giant depolarising potentials (GDPs) are localised synchronised events that can be measured in brain slices cut from rodents early in development allowing more detailed molecular mechanisms to be resolved³⁹. The coordinated activation of a range of ion channels is critical to the generation of GDPs, with excitatory GABA_A receptor transmission a particularly important driver of this activity³⁹. We next discuss structural and functional evidence obtained from the GABA_A $\gamma 2_{R43Q}$ mouse model of GGE to highlight how disruptions in early oscillatory activity could lead to long-term changes in network excitability.

A key difference in the GABAergic system in early development compared to adulthood is how neurons regulate intracellular chloride ions (Cl⁻). At these early developmental time-points neurons primarily express NKCC1. This transporter pumps Cl⁻ into the cell while the K-Cl cotransporter 2 (KCC2) that is expressed in adult neurons efficiently pumps Cl⁻ out⁴⁰. As a result of this expression profile immature neurons have higher internal Cl⁻ concentrations. This high internal concentration shifts the reversal potential of Cl⁻ so that at typical holding potentials Cl⁻ leaves the neuron and the resulting depolarisation takes it closer to the threshold for action potential firing⁴⁰. Synaptic release of GABA acting on GABA_A receptors under these conditions is therefore excitatory. Recurrent spontaneous GDPs are an early developmental neuronal network phenomenon that relies on this excitatory GABA_A receptor-mediated transmission^{41; 42}. These events cause neuronal synchronized firing and set up Ca²⁺ oscillations that help the establishment and refinement of network circuitry⁴³.

Neonatal hippocampal slices from the GABA_A $\gamma 2_{R43Q}$ mouse model of GGE have a lower frequency of GDPs when compared to their wildtype littermates⁴⁴. A similar

reduction in GDP frequency can be achieved pharmacologically using bumetanide which blocks NKCC1 and consequently shifts the reversal potential of Cl^- so that it is no longer excitatory^{45; 46}. Interestingly, wild-type mice treated with bumetanide in the antenatal period have a long-lasting increase in thermogenic seizure susceptibility that is comparable with that observed in the $\text{GABA}_A \gamma 2_{\text{R43Q}}$ mouse⁴⁴. It should be noted that bumetanide given early in development has a protective effect in a genetic mouse model based on dysfunctional Kv7 channels, normalising several cellular and behavioural markers of hyperexcitability¹¹. Related work has also shown that manipulations of GABAergic transmission, using other pharmacological tools, in this early developmental window have a long-lasting impact on neuron circuitry, potentially influencing a range of behaviours including altered seizure susceptibility in adulthood⁴⁷⁻⁴⁹.

Perturbation of the GABAergic system in early development has previously been postulated to influence the migratory behaviour and differentiation of cortical neurons⁵⁰. For instance, GABA-signalling stimulates the migration of embryonic cortical cells with concentration-dependent effects^{47; 51; 52}. A causative link has not yet been established between reductions in GDP frequency and changes in neuron positioning in the $\text{GABA}_A \gamma 2_{\text{R43Q}}$ mouse³³. Further, how changes in neuronal migration, proliferation, differentiation may alter long-term network excitability remains an open question. One speculation is that subtle changes due to the mis-positioning or change in numbers of certain GABAergic neurons may alter the excitatory-inhibitory balance of local circuits leading to a susceptibility to increased excitability. Long-term structural modifications to neuronal morphology may also occur. As an example, reduced dendritic branching has been implicated as part of the pathogenic mechanism of increased neuronal excitability in both acquired and genetic epilepsy^{53; 54}. Although the precise mechanism may remain unclear, the early developmental GABAergic system is well positioned to contribute to morphological changes in the young brain and consequently long-term neuronal hyper-excitability. This system is also likely to be amenable to therapeutic intervention, for example, increasing GDP frequency in $\text{GABA}_A \gamma 2_{\text{R43Q}}$ patients using drugs that increase GABA_A receptor activity may reduce future seizure risk.

Implications for comorbid pathologies in epilepsy

There is a vast literature implicating dysfunction in early development as a predisposing factor for a range of neurological conditions. Interestingly, studies in humans indicate that major depression, bipolar disorders, psychiatric history and cognitive impairment are strongly associated with an increase in the risk of developing epilepsy⁵⁵⁻⁵⁹. These clinical relationships suggest that common network pathologies may underlie both changes in seizure susceptibility and increased risk of psychiatric and cognitive disorders. There is also evidence for this at the neuronal structure level. For example, dynamic changes in dendritic spines are associated with epilepsy; they are also seen in a number of neuropsychiatric and cognitive disorders, including autism spectrum disorders, schizophrenia and Alzheimer's disease^{60; 61}. As a consequence, early therapeutic intervention that normalises network dysfunction may also alleviate comorbid states. One experimental example of this is that the anxiety-like behaviour, that is a comorbid state in GAERS, is reduced in adult rats that had only been treated during the pre-symptomatic period¹⁰.

Precision medicine implications

The understanding of the genetic basis of many of the epilepsies provides an opportunity to generate precision medicine strategies. Part of this strategy will be to understand the early developmental consequences of variants that raise risk for epilepsies and its comorbidities. Here we have highlighted how animal models are beginning to shed light into how they exert their effect on the developing brain. Another particularly promising approach is the use of human stem cell-based neural models. The fundamental mechanisms of neurogenesis can be modelled by the *in vitro* differentiation of neurons from pluripotent cells, such as embryonic stem or patient derived induced pluripotent (iPS) cells. For instance, *in vitro* protocols for neural differentiation enable the faithfully reproduction of the temporal appearance, spatial organisation and gene expression profiles of developing neurons *in utero*⁶²⁻⁶⁵. These methods have been used to model neural microarchitecture changes that accompany epilepsy disease progression, such as impaired neurite outgrowth in iPS-derived neurons from individuals with Ohtahara syndrome, a severe epileptic encephalopathy⁶⁶. Further, iPS cell-derived GABA positive inhibitory neurons carrying a mutation in *SCN1A* have been shown to replicate disease pathology previously identified in animal models including a reduction in voltage dependent Na⁺ currents in inhibitory neurons⁶⁷. Recently, neurodevelopmental and synaptic changes

in a human stem cell-based model of tuberous sclerosis were rescued using pharmacological intervention ⁶⁸. This growing body of research indicates that stem cell-based models will be integral to helping our understanding of early developmental pathogenic changes caused by epilepsy mutations, many of which may be difficult to study in whole organisms, and for which novel therapeutic strategies can be developed.

Taking these ideas to the clinic

The major conclusion of this review is that disease-targeted intervention may need to occur in pre-symptomatic GGE patients to be effective. This will require the identification of patients that will benefit from such early intervention. As our understanding of the genetic architecture improves and pre- and post-natal genetic screening becomes more widely employed it will be possible to identify individuals who have an increased risk of developing epilepsy. However, as incomplete penetrance and variable expressivity is commonly observed it may be difficult to predict accurately the clinical outcome simply from DNA sequence analysis alone ⁶⁹. Additional biomarkers are therefore going to be essential to make informed decisions regarding the suitability of individuals for early therapeutic intervention. Improving human imaging technologies are enabling us to better map the brain from both a structural and functional perspective allowing us to identify morphological and/ or functional markers of excitability in GGE and other genetic epilepsies. Although limited in scope, evidence from animal models suggests that subtle changes in pre-seizure brains can indeed be detected. Finally, it is important to note that we must develop new therapeutic strategies, moving away from the development of classical anti-epileptic drugs that treat the symptoms of epilepsy ('anti-seizure') to those that are anti-epileptogenic and target the underlying disease mechanism to prevent the development of epilepsy and its comorbidities.

Figure 1

Cartoon illustrating the potential that an epilepsy mutation can alter neuronal network structure and function in early development to set seizure susceptibility in adulthood.

Acknowledgments: This work was supported by National Health and Medical Research Council (NHMRC) Program Grant (10915693) to SF, SP and CAR.

Conflict: No authors have any conflicts to declare in relation to this work.

The Authors confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this report is consistent with those guidelines.

References

1. Hesdorffer DC, Logroscino G, Benn EK, et al. Estimating risk for developing epilepsy: a population-based study in Rochester, Minnesota. *Neurology* 2011;76:23-27.
2. Ultra-rare genetic variation in common epilepsies: a case-control sequencing study. *Lancet Neurol* Epi4K consortium, 2017;16:135-143.
3. McTague A, Howell KB, Cross JH, et al. The genetic landscape of the epileptic encephalopathies of infancy and childhood. *Lancet Neurol* 2016;15:304-316.
4. Myers CT, Mefford HC. Advancing epilepsy genetics in the genomic era. *Genome Med* 2015;7:91.
5. Oyrer J, Maljevic S, Scheffer IE, et al. Ion Channels in Genetic Epilepsy: From Genes and Mechanisms to Disease-Targeted Therapies. *Pharmacol Rev* 2018;70:142-173.
6. Scheffer IE, Berkovic S, Capovilla G, et al. ILAE classification of the epilepsies: Position paper of the ILAE Commission for Classification and Terminology. *Epilepsia* 2017;58:512-521.
7. Loiseau J, Loiseau P, Guyot M, et al. Survey of seizure disorders in the French southwest. I. Incidence of epileptic syndromes. *Epilepsia* 1990;31:391-396.
8. Helbig I, Heinzen EL, Mefford HC. Primer Part 1-The building blocks of epilepsy genetics. *Epilepsia* 2016;57:861-868.

9. Blumenfeld H, Klein JP, Schridde U, et al. Early treatment suppresses the development of spike-wave epilepsy in a rat model. *Epilepsia* 2008;49:400-409.
10. Dezsi G, Ozturk E, Stanic D, et al. Ethosuximide reduces epileptogenesis and behavioral comorbidity in the GAERS model of genetic generalized epilepsy. *Epilepsia* 2013;54:635-643.
11. Marguet SL, Le-Schulte VT, Merseburg A, et al. Treatment during a vulnerable developmental period rescues a genetic epilepsy. *Nat Med* 2015;21:1436-1444.
12. Tan HO, Reid CA, Single FN, et al. Reduced cortical inhibition in a mouse model of familial childhood absence epilepsy. *Proc Natl Acad Sci U S A* 2007;104:17536-17541.
13. Wallace RH, Marini C, Petrou S, et al. Mutant GABA(A) receptor gamma2-subunit in childhood absence epilepsy and febrile seizures. *Nat Genet* 2001;28:49-52.
14. Chiu C, Reid CA, Tan HO, et al. Developmental impact of a familial GABAA receptor epilepsy mutation. *Ann Neurol* 2008;64:284-293.
15. Arsov T, Mullen SA, Damiano JA, et al. Early onset absence epilepsy: 1 in 10 cases is caused by GLUT1 deficiency. *Epilepsia* 2012;53:e204-207.
16. Suls A, Mullen SA, Weber YG, et al. Early-onset absence epilepsy caused by mutations in the glucose transporter GLUT1. *Ann Neurol* 2009;66:415-419.
17. Wang D, Pascual JM, Yang H, et al. A mouse model for Glut-1 haploinsufficiency. *Hum Mol Genet* 2006;15:1169-1179.
18. Tang M, Gao G, Rueda CB, et al. Brain microvasculature defects and Glut1 deficiency syndrome averted by early repletion of the glucose transporter-1 protein. *Nat Commun* 2017;8:14152.
19. Manfredi I, Zani AD, Rampoldi L, et al. Expression of mutant beta2 nicotinic receptors during development is crucial for epileptogenesis. *Hum Mol Genet* 2009;18:1075-1088.
20. Silva-Santos S, van Woerden GM, Bruinsma CF, et al. Ube3a reinstatement identifies distinct developmental windows in a murine Angelman syndrome model. *J Clin Invest* 2015;125:2069-2076.

21. Schramm J. Temporal lobe epilepsy surgery and the quest for optimal extent of resection: a review. *Epilepsia* 2008;49:1296-1307.
22. Kannan L, Vogrin S, Bailey C, et al. Centre of epileptogenic tubers generate and propagate seizures in tuberous sclerosis. *Brain* 2016;139:2653-2667.
23. Raychaudhuri R, Batjer HH, Awad IA. Intracranial cavernous angioma: a practical review of clinical and biological aspects. *Surg Neurol* 2005;63:319-328; discussion 328.
24. Ciumas C, Savic I. Structural changes in patients with primary generalized tonic and clonic seizures. *Neurology* 2006;67:683-686.
25. Koepp MJ, Woermann F, Savic I, et al. Juvenile myoclonic epilepsy--neuroimaging findings. *Epilepsy Behav* 2013;28 Suppl 1:S40-44.
26. Pardoe H, Pell GS, Abbott DF, et al. Multi-site voxel-based morphometry: methods and a feasibility demonstration with childhood absence epilepsy. *Neuroimage* 2008;42:611-616.
27. Seneviratne U, Cook M, D'Souza W. Focal abnormalities in idiopathic generalized epilepsy: a critical review of the literature. *Epilepsia* 2014;55:1157-1169.
28. Nuyts S, D'Souza W, Bowden SC, et al. Structural brain abnormalities in genetic generalized epilepsies: A systematic review and meta-analysis. *Epilepsia* 2017;58:2025-2037.
29. Boullieret V, Hogan RE, Velakoulis D, et al. Morphometric abnormalities and hyperanxiety in genetically epileptic rats: a model of psychiatric comorbidity? *Neuroimage* 2009;45:267-274.
30. Chahboune H, Mishra AM, DeSalvo MN, et al. DTI abnormalities in anterior corpus callosum of rats with spike-wave epilepsy. *Neuroimage* 2009;47:459-466.
31. Mishra AM, Bai X, Motelow JE, et al. Increased resting functional connectivity in spike-wave epilepsy in WAG/Rij rats. *Epilepsia* 2013;54:1214-1222.
32. Richards KL, Kurniawan ND, Yang Z, et al. Hippocampal volume and cell density changes in a mouse model of human genetic epilepsy. *Neurology* 2013;80:1240-1246.
33. Wimmer VC, Li MY, Berkovic SF, et al. Cortical microarchitecture changes in genetic epilepsy. *Neurology* 2015;84:1308-1316.

34. Cavdar S, Bay HH, Yildiz SD, et al. Comparison of numbers of interneurons in three thalamic nuclei of normal and epileptic rats. *Neurosci Bull* 2014;30:451-460.
35. Mann EO, Mody I. The multifaceted role of inhibition in epilepsy: seizure-genesis through excessive GABAergic inhibition in autosomal dominant nocturnal frontal lobe epilepsy. *Current Opinion in Neurology* 2008;21:155-160.
36. Karpova AV, Bikbaev AF, Coenen AM, et al. Morphometric Golgi study of cortical locations in WAG/Rij rats: the cortical focus theory. *Neurosci Res* 2005;51:119-128.
37. Lindemann C, Ahlbeck J, Bitzenhofer SH, et al. Spindle Activity Orchestrates Plasticity during Development and Sleep. *Neural Plast* 2016;2016:5787423.
38. Hanganu-Opatz IL. Between molecules and experience: role of early patterns of coordinated activity for the development of cortical maps and sensory abilities. *Brain Res Rev* 2010;64:160-176.
39. Griguoli M, Cherubini E. Early Correlated Network Activity in the Hippocampus: Its Putative Role in Shaping Neuronal Circuits. *Front Cell Neurosci* 2017;11:255.
40. Knoflach F, Hernandez MC, Bertrand D. GABAA receptor-mediated neurotransmission: Not so simple after all. *Biochem Pharmacol* 2016;115:10-17.
41. Ben-Ari Y, Cherubini E, Corradetti R, et al. Giant synaptic potentials in immature rat CA3 hippocampal neurones. *J Physiol* 1989;416:303-325.
42. Leinekugel X, Medina I, Khalilov I, et al. Ca²⁺ oscillations mediated by the synergistic excitatory actions of GABA(A) and NMDA receptors in the neonatal hippocampus. *Neuron* 1997;18:243-255.
43. Zhang LI, Poo MM. Electrical activity and development of neural circuits. *Nat Neurosci* 2001;4 Suppl:1207-1214.
44. Vargas E, Petrou S, Reid CA. Genetic and pharmacological modulation of giant depolarizing potentials in the neonatal hippocampus associates with increased seizure susceptibility. *J Physiol* 2013;591:57-65.

45. Nardou R, Ben-Ari Y, Khalilov I. Bumetanide, an NKCC1 antagonist, does not prevent formation of epileptogenic focus but blocks epileptic focus seizures in immature rat hippocampus. *J Neurophysiol* 2009;101:2878-2888.
46. Tyzio R, Allene C, Nardou R, et al. Depolarizing actions of GABA in immature neurons depend neither on ketone bodies nor on pyruvate. *J Neurosci* 2011;31:34-45.
47. Haas M, Qu Z, Kim TH, et al. Perturbations in cortical development and neuronal network excitability arising from prenatal exposure to benzodiazepines in mice. *Eur J Neurosci* 2013;37:1584-1593.
48. Nicosia A, Giardina L, Di Leo F, et al. Long-lasting behavioral changes induced by pre- or neonatal exposure to diazepam in rats. *Eur J Pharmacol* 2003;469:103-109.
49. Wang DD, Kriegstein AR. Blocking early GABA depolarization with bumetanide results in permanent alterations in cortical circuits and sensorimotor gating deficits. *Cereb Cortex* 2011;21:574-587.
50. Manent JB, Beguin S, Ganay T, et al. Cell-autonomous and cell-to-cell signalling events in normal and altered neuronal migration. *Eur J Neurosci* 2011;34:1595-1608.
51. Behar TN, Li YX, Tran HT, et al. GABA stimulates chemotaxis and chemokinesis of embryonic cortical neurons via calcium-dependent mechanisms. *J Neurosci* 1996;16:1808-1818.
52. Cuzon VC, Yeh PW, Cheng Q, et al. Ambient GABA promotes cortical entry of tangentially migrating cells derived from the medial ganglionic eminence. *Cereb Cortex* 2006;16:1377-1388.
53. Paz JT, Davidson TJ, Frechette ES, et al. Closed-loop optogenetic control of thalamus as a tool for interrupting seizures after cortical injury. *Nat Neurosci* 2013;16:64-70.
54. Reid CA, Leaw B, Richards KL, et al. Reduced dendritic arborization and hyperexcitability of pyramidal neurons in a Scn1b-based model of Dravet syndrome. *Brain* 2014;137:1701-1715.

55. Hesdorffer DC, Hauser WA, Olafsson E, et al. Depression and suicide attempt as risk factors for incident unprovoked seizures. *Ann Neurol* 2006;59:35-41.
56. Hesdorffer DC, Ishihara L, Mynepalli L, et al. Epilepsy, suicidality, and psychiatric disorders: a bidirectional association. *Ann Neurol* 2012;72:184-191.
57. Lin JJ, Mula M, Hermann BP. Uncovering the neurobehavioural comorbidities of epilepsy over the lifespan. *Lancet* 2012;380:1180-1192.
58. Nilsson FM, Kessing LV, Bolwig TG. On the increased risk of developing late-onset epilepsy for patients with major affective disorder. *J Affect Disord* 2003;76:39-48.
59. Taylor J, Kolamunnage-Dona R, Marson AG, et al. Patients with epilepsy: cognitively compromised before the start of antiepileptic drug treatment? *Epilepsia* 2010;51:48-56.
60. Penzes P, Cahill ME, Jones KA, et al. Dendritic spine pathology in neuropsychiatric disorders. *Nat Neurosci* 2011;14:285-293.
61. Wong M, Guo D. Dendritic spine pathology in epilepsy: cause or consequence? *Neuroscience* 2013;251:141-150.
62. Camp JG, Badsha F, Florio M, et al. Human cerebral organoids recapitulate gene expression programs of fetal neocortex development. *Proc Natl Acad Sci U S A* 2015;112:15672-15677.
63. Chambers SM, Fasano CA, Papapetrou EP, et al. Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling. *Nat Biotechnol* 2009;27:275-280.
64. Lancaster MA, Renner M, Martin CA, et al. Cerebral organoids model human brain development and microcephaly. *Nature* 2013;501:373-379.
65. Shi Y, Kirwan P, Livesey FJ. Directed differentiation of human pluripotent stem cells to cerebral cortex neurons and neural networks. *Nat Protoc* 2012;7:1836-1846.
66. Yamashita S, Chiyonobu T, Yoshida M, et al. Mislocalization of syntaxin-1 and impaired neurite growth observed in a human iPSC model for STXBP1-related epileptic encephalopathy. *Epilepsia* 2016;57:e81-86.

67. Sun Y, Pasca SP, Portmann T, et al. A deleterious Nav1.1 mutation selectively impairs telencephalic inhibitory neurons derived from Dravet Syndrome patients. *Elife* 2016;5.
68. Costa V, Aigner S, Vukcevic M, et al. mTORC1 Inhibition Corrects Neurodevelopmental and Synaptic Alterations in a Human Stem Cell Model of Tuberous Sclerosis. *Cell Rep* 2016;15:86-95.
69. Dickinson ME, Flenniken AM, Ji X, et al. High-throughput discovery of novel developmental phenotypes. *Nature* 2016.

epi_14077_f1.tif

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Reid, CA;Rollo, B;Petrou, S;Berkovic, SF

Title:

Can mutation-mediated effects occurring early in development cause long-term seizure susceptibility in genetic generalized epilepsies?

Date:

2018-05-01

Citation:

Reid, C. A., Rollo, B., Petrou, S. & Berkovic, S. F. (2018). Can mutation-mediated effects occurring early in development cause long-term seizure susceptibility in genetic generalized epilepsies?. *EPILEPSIA*, 59 (5), pp.915-922. <https://doi.org/10.1111/epi.14077>.

Persistent Link:

<http://hdl.handle.net/11343/283829>