The Clinical Need for Standardised MDM Assessment Processes

Wing Hing Yau^{1,2}, Bianca Devitt^{1,3}, Soe Yu Aung¹, Phillip Parente^{1,3}, Sharad Sharma², Carmel Pezaro^{1,3}

¹Department of Medical Oncology, Eastern Health, Box Hill, Australia

²Department of Oncology, Launceston General Hospital, Launceston, Australia

³Eastern Health Clinical School, Monash University, Box Hill, Australia

Author Positions:

Wing Hing Yau	Medical Oncology Advanced Trainee, Eastern Health, Box Hill, Victoria, Australia
Bianca Devitt	Medical Oncologist, Eastern Health, Box Hill, Victoria, Australia
Soe Yu Aung ⁱ	Medical Oncology Advanced Trainee, Andrew Love Cancer Centre, Barwon Health, Box Hill, Victoria, Australia
Phillip Parente	Director of Cancer Services, Eastern Health, Box Hill, Victoria, Australia
Sharad Sharma ⁱⁱ	Medical Oncologist, Ballarat Health Services, Ballarat, Victoria, Australia
Carmel Pezaro	Medical Oncologist, Eastern Health, Box Hill, Victoria, Australia

Author contributions:

All authors contributed to the intellectual content and revision of manuscript.

Corresponding author:

Dr Carmel Pezaro

Department of Medical Oncology
Box Hill Hospital, Eastern Health
8 Arnold Street, Box Hill 3128
Victoria, Australia
Telephone: +61 1300 342 255
Email: carmel.pezaro@monash.edu

Acknowledgements: No relevant conflicts of interests to disclose. No financial grants or funding were used for the purpose of this work.

Manuscript word count: 482 words (Maximum 500)

Keywords: Multidisciplinary, cancer, bladder cancer, quality improvement

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/imj.13688](https://doi.org/10.1111/imj.13688)

ⁱ Dr Soe Yu Aung is no longer affiliated with Eastern Health, Australia at the time of submission. He is currently affiliated with Barwon Health, Australia.

ⁱⁱ Dr Sharad Sharma is no longer affiliated with Launceston General Hospital, Australia at the time of submission. He is currently affiliated with Ballarat Health Services, Ballarat, Australia.

We read with interest the article by Johnson et al. regarding the development of a peer-review framework for cancer multidisciplinary meetings (MDMs)¹. MDMs are a critical component of multidisciplinary oncological care, associated with improved patient outcomes and satisfaction, trials recruitment, and interdisciplinary communication²⁻⁵. Whilst national and international guidelines exist relating to the conduct of MDMs^{3,6,7}, processes still vary within and across institutions, differing in MDM frequency, the patients presented, team membership, leadership and communication^{2,4}. There is currently no standardised process within Australia to monitor the quality of MDM across different institutions and regional settings. However, variations between MDMs may affect quality of care.

We recently reviewed neoadjuvant chemotherapy delivery for muscle-invasive bladder carcinoma (MIBC) in a metropolitan and regional setting and found MDM discussion to be a determinant of patients receiving neoadjuvant chemotherapy. Neoadjuvant chemotherapy is standard of care for patients with MIBC⁸. Our retrospective audit analysed 19 metropolitan patients and 23 regional patients who underwent radical cystectomy for MIBC. Regional patients had significantly lower rates of both referral to a pre-operative MDM (84% vs 42%, $p=0.023$) and neoadjuvant chemotherapy (42% vs 13%, $p=0.043$). Nine of the 11 patients (82%) who received neoadjuvant chemotherapy had been discussed at a MDM, demonstrating the association between MDM referral and neoadjuvant treatment. There were no significant differences identified in the patient populations or type of chemotherapy. Optimising the MDM processes could therefore improve delivery of standard of care treatment for these patients.

Whilst it is intuitive to believe that an MDM that meets regularly, with well-considered terms of reference, strong leadership and good team communication will make better decisions, there is minimal literature comparing variations or interventions in MDM processes and their effect on patient outcomes. Lamb et al. demonstrated that use of a checklist during MDM discussion to

ensure adequate information was presented and all team members' views discussed, improved efficiency and treatment decisions. There was no measure as to whether this translated into improved patient outcomes⁹. The MDM peer-review framework described by Johnson et al. is another tool, designed to reduce variation in practice and provide opportunity for quality improvement. It may be limited by the labour-intensive process and the perceived relevance of its recommendations¹⁰.

We strongly support ongoing research into improving the quality and efficiency of the MDM process. We argue that for any MDM intervention, improved patient outcomes should be the primary measure of success. Whilst a peer review process enables assessment of communication and leadership within an MDM team, we wonder whether less costly strategies should be explored first. Our research would suggest that process to ensure that all patients are discussed at an MDM soon after their diagnosis, which are attended (either in person or via teleconference) by all appropriate specialists, would be a welcome first step. It would be interesting to investigate whether a simple checklist used during the MDM to prompt discussion could further improve patient outcomes.

References

1. Johnson CE, Slavova-Azmanova N, Saunders C. Development of a peer-review framework for cancer multidisciplinary meetings. *Intern Med J.* 2017;47(5):529-35.
2. Fleissig A, Jenkins V, Catt S, Fallowfield L. Multidisciplinary teams in cancer care: are they effective in the UK? *Lancet Oncol.* 2006;7(11):935-43.
3. Wright FC, De Vito C, Langer B, Hunter A, Expert Panel on Multidisciplinary Cancer Conference S. Multidisciplinary cancer conferences: a systematic review and development of practice standards. *Eur J Cancer.* 2007;43(6):1002-10.
4. Taylor C, Munro AJ, Glynne-Jones R, Griffith C, Trevatt P, Richards M, et al. Multidisciplinary team working in cancer: what is the evidence? *BMJ.* 2010;340:c951.
5. Kesson EM, Allardice GM, George WD, Burns HJ, Morrison DS. Effects of multidisciplinary team working on breast cancer survival: retrospective, comparative, interventional cohort study of 13 722 women. *BMJ.* 2012;344:e2718.
6. Team NCA, The characteristics of an effective multidisciplinary team, UK National Cancer Intelligence Network 2010.
7. Centre NBC, Multidisciplinary meetings for cancer care: a guide for health service providers, Cancer Australia 2005.

8. Liew MS, Azad A, Tafreshi A, Eapen R, Bolton D, Davis ID, et al. USANZ: Time-trends in use and impact on outcomes of perioperative chemotherapy in patients treated with radical cystectomy for urothelial bladder cancer. *BJU Int.* 2013;112 Suppl 2:74-82.
9. Lamb BW, Green JS, Benn J, Brown KF, Vincent CA, Sevdalis N. Improving decision making in multidisciplinary tumor boards: prospective longitudinal evaluation of a multicomponent intervention for 1,421 patients. *J Am Coll Surg.* 2013;217(3):412-20.
10. Slavova-Azmanova NS, Johnson CE, Platell C, Bydder S, Saunders CM. Peer review of cancer multidisciplinary teams: is it acceptable in Australia? *Med J Aust.* 2015;202(3):144-7.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Yau, WH;Devitt, B;Aung, SY;Parente, P;Sharma, S;Pezaro, C

Title:

Clinical need for standardised multidisciplinary meeting assessment processes.

Date:

2018-02

Citation:

Yau, W. H., Devitt, B., Aung, S. Y., Parente, P., Sharma, S. & Pezaro, C. (2018). Clinical need for standardised multidisciplinary meeting assessment processes.. Intern Med J, 48 (2), pp.230-231. <https://doi.org/10.1111/imj.13688>.

Persistent Link:

<http://hdl.handle.net/11343/283875>