Title

Implications of treatment providers' varying conceptions of the disease model of addiction – A response

Author names and affiliations

Anthony Barnett¹, Wayne Hall^{2,3}, Craig L. Fry⁴, Ella Dilkes-Frayne⁵, Adrian Carter^{1,6}

¹Monash Institute of Cognitive and Clinical Neurosciences and School of Psychological Sciences, Monash University, Melbourne, Australia

²Centre for Youth Substance Abuse Research, The University of Queensland, Brisbane, Australia

³National Addiction Centre, Institute of Psychiatry, Psychology and Neuroscience, Kings College London, London, United Kingdom.

⁴College of Health & Biomedicine, Victoria University, Melbourne, Australia

⁵School of Sociology, Research School of Social Sciences, The Australian National University, Canberra, Australia.

⁶Australia and University of Queensland Centre of Clinical Research, University of Queensland, Brisbane, Australia.

Anthony Barnett B. Psychological Studies (Hons), PhD Candidate, Wayne Hall PhD, Director, Centre for Youth Substance Abuse Research and Professor, National Addiction Centre, Craig L. Fry DrPH, Research Program Leader, Ella Dilkes-Frayne PhD, Research Fellow, Adrian Carter PhD, Senior Research Fellow. Correspondence to Mr Anthony Barnett, Brain and Mental Health Laboratory, Q5 School of Psychological Sciences, Monash University, 770 Blackburn Road, Clayton, Vic. 3800, Australia. Tel: +61 (3) 9905 9322. E-mail: anthony.barnett@monash.edu

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/dar.12844](https://doi.org/10.1111/dar.12844)

People providing treatment for alcohol and other drug (AOD) issues are a diverse group. They come from very different walks of life, with varying educational backgrounds, and bring vastly different experiences to the task of helping people overcome problems with drugs. Unsurprisingly, they also have divergent views about the nature of AOD problems (whether viewing it as a disease, a moral or social problem). As a result, people accessing AOD services may face multiple, even contradictory, explanations of their AOD issues [1].

In their responses to our systematic review of treatment providers' views about the disease model of addiction [1], Savic and Lubman [2] and Storbjörk [3] make compelling cases against attempting to 'standardise' treatments to address contradictory explanations that treatment seekers may face. Savic and Lubman's argument against implementing an 'overarching, universal addiction model' is threefold. Firstly, attempts to translate a standardised treatment model into practice would be challenging. Secondly, a 'one-size fits all' approach would not account for people's complex needs in treatment. Thirdly, implementing an overarching model would paternalistically rely on 'expert' knowledge at the expense of consumer participation and client-centred care. Advocating for the maintenance of multiple treatment models in practice, Storbjörk argues that "different conceptions in the treatment landscape [...] facilitate better matches between service user beliefs and the models and goals available" [3].

However, these arguments against attempts to standardise addiction treatment, run the risk of overlooking the consequences of a system where treatment providers have such varying views about the underlying nature of AOD problems. Savic and Lubman [2] understate the extent to which treatment providers' varying conceptualisations of addiction shape and direct treatment practices. Savic and Lubman argue that clients' needs and beliefs should be the starting point in understanding their concerns and that treatment providers play the role of presenting clients with a range of possible ways of understanding and addressing their problems. While we don't disagree, the findings of our review suggest that treatment providers play a more active, assertive role in directing care in line with their belief systems about addiction and treatment [1].

For example, there is evidence that treatment providers who support the disease model of alcoholism are more likely to insist on abstinence as the only treatment goal [4], less likely to consider controlled drinking [5], more likely to refer to Alcoholics Anonymous [6], and more likely to impose their own treatment goals rather than incorporate the goals of the client [5]. A critical area of future research is the extent to which support for the disease model of addiction influences practice and how discrepant client and treatment provider views of addiction are negotiated in care settings where power and authority is vested in the treatment provider.

Storbjörk [3] discusses the contested medicalization of addiction in the “non-medical stronghold” of the Nordic countries. As she states, in Sweden, key stakeholders within AOD treatment and policy resist abandoning the social tradition in favour of biomedical understandings of addiction problems (e.g. [7]). Storbjörk’s commentary prompts us to consider tensions between biomedical and social frameworks of addiction in the global context. For instance, when examining AOD treatment in Victoria, Australia, we see a highly heterogeneous treatment provision landscape made up of services that differ in treatment philosophies (e.g. pharmacotherapy clinics based on harm reduction approaches; therapeutic communities based on abstinence models). Within such settings there has been tension (e.g. [8]) between a biomedicalised framing of addiction as a ‘chronic, relapsing brain disease’ [9,10], and social models such as the *Social Identity Model of Recovery* [11] that see the solutions to AOD problems residing within individuals’ social networks.

Schmidt [12] also draws attention to conflicts between different models of addiction. She notes that we found that treatment providers appear to endorse disease and other models simultaneously and strategically deploy different models for their presumed therapeutic benefits. Schmidt suggests that moralised formulations of the disease concept were present in Benjamin Rush’s original formulation of alcoholism as a ‘disease of the will’ [13,14] and, because they “have never fully dropped away”, continue to pose problems for clients (e.g. stigma deterring people from accessing treatment). Schmidt sees this as problematic because addiction treatment stakes its claims to legitimacy on evidence-based medical science rather than ideological systems of belief.

In addressing Storbjörk’s [3] and Schmidt’s [12] concerns, we see no reason why different treatments based on social and biomedical models cannot co-exist, particularly given that medical and social factors are so intertwined for people experiencing AOD problems. Treatment clients may benefit from both social interventions (e.g. built on a Social Identity of Model of Recovery approach) and biomedical interventions (e.g. pharmacotherapies). However, matching people to individualised treatments remains a challenge. Addressing this challenge, centralised intake and assessment processes (such as we see in Victoria, Australia) provide one practical way to assess clients’ needs and direct them to appropriate care.

As we elaborated on in our review [1] (p. 21), the ‘hybrid approach’ [15] in which treatment providers endorse varying models of addiction simultaneously, may be useful. In one ‘treatment situation’ the use of a medical model might de-moralise clients, whereas in another, a moral model may encourage clients to take responsibility for their recovery [16,17]. Critically, the decision when to deploy different models and how to frame clients’ AOD use lies within the expertise of treatment providers and their interactions with clients. In view of this, training programmes are vital for treatment providers on the clinical impacts of disease models of addiction.

To conclude, all three commentaries point to a wider debate on how diverse views of addiction cohere and function in treatment services. While it may not be preferable to standardise treatment along the lines of a single approach (e.g. a disease model approach), in order to improve treatment we need to give greater attention to addressing the diversity of treatment providers' views and their impacts on clients seeking treatment and support to address AOD issues.

References

1. Barnett AI, Hall W, Fry CL, Dilkes-Frayne E, Carter A. Drug and alcohol treatment providers' views about the disease model of addiction and its impact on clinical practice: A systematic review. *Drug Alcohol Rev* 2017 [Epub ahead of print].
2. Savic M, Lubman DI. An argument against the implementation of an 'overarching universal addiction model' in alcohol and other drug treatment. *Drug Alcohol Rev* 2018 [Epub ahead of print].
3. Storbjörk J. One model to rule them all? Governing images in the shadow of the disease model of addiction. *Drug Alcohol Rev* 2018 [Epub ahead of print].
4. Hsieh SY, Srebalus DJ. Alcohol treatment issues: Professional differences. *Alcoholism Treatment Quarterly* 1997;15:63-73.
5. Moyers TB, Miller WR. Therapists' conceptualizations of alcoholism: Measurement and implications for treatment decisions. *Psychology of Addictive Behaviors* 1993;7:238.
6. Casswell S, McPherson M. Attitudes of New Zealand general practitioners to alcohol-related problems. *J Stud Alcohol* 1983;44:342-51.
7. Winter K. Coproduction of scientific addiction knowledge in everyday discourse. *Contemp Drug Probl* 2016;43:25-46.
8. Best DW, Lubman DI. The recovery paradigm: A model of hope and change for alcohol and drug addiction. *Aust Fam Physician* 2012;41:593.
9. Leshner AI. Addiction is a brain disease, and it matters. *Science* 1997;278:45-7.
10. Volkow ND, Koob GF, McLellan AT. Neurobiologic advances from the brain disease model of addiction. *N Engl J Med* 2016;374:363-371.
11. Best D, Beckwith M, Haslam C, Alexander Haslam S, Jetten J, Mawson E, Lubman DI. Overcoming alcohol and other drug addiction as a process of social identity transition: the social identity model of recovery (SIMOR). *Addiction Research & Theory* 2016;24:111-23.
12. Schmidt L. A persistent contradiction in treatment providers' views on addiction. *Drug Alcohol Rev* 2018 [Epub ahead of print].
13. Levine HG. The discovery of addiction: Changing conceptions of habitual drunkenness in America. *J Subst Abuse Treat* 1985;2:43-57.
14. Schmidt LA. "A battle not man's but God's": origins of the American temperance crusade in the struggle for religious authority. *J Stud Alcohol* 1995;56:110-21.
15. Karasaki M, Fraser S, Moore D, Dietze P. The place of volition in addiction: Differing approaches and their implications for policy and service provision. *Drug Alcohol Rev* 2013;32:195-204.
16. Brickman P, Rabinowitz VC, Karuza J, Coates D, Cohn E, Kidder L. Models of helping and coping. *American psychologist* 1982;37:368-384.
17. Palm J. The nature of and responsibility for alcohol and drug problems: Views among treatment staff. *Addiction Research & Theory* 2004;12:413-31.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Barnett, A;Hall, W;Fry, CL;Dilkes-Frayne, E;Carter, A

Title:

Implications of treatment providers' varying conceptions of the disease model of addiction: A response

Date:

2018-09-01

Citation:

Barnett, A., Hall, W., Fry, C. L., Dilkes-Frayne, E. & Carter, A. (2018). Implications of treatment providers' varying conceptions of the disease model of addiction: A response. DRUG AND ALCOHOL REVIEW, 37 (6), pp.729-730. <https://doi.org/10.1111/dar.12844>.

Persistent Link:

<http://hdl.handle.net/11343/284202>