DR. BING GAO (Orcid ID : 0000-0003-2069-5279)

PROF. BAOJING GU (Orcid ID : 0000-0003-3986-3519)

PROF. DAVID STEPHEN POWLSON (Orcid ID : 0000-0001-8776-2339)

PROF. PETE SMITH (Orcid ID : 0000-0002-3784-1124)

Article type : Primary Research Articles

Manuscript submitted to *Global Change Biology*

Type of contribution: original research

Chinese cropping systems are a net source of greenhouse gases despite soil carbon sequestration

Running head: Chinese cropping systems are a net C source

Bing Gao^{a,b,c}, Tao Huang^{c,d}, Xiaotang Ju^{c*}, Baojing Gu^{e,f}, Wei Huang^{a,b}, Lilai Xu^{a,b}, Robert M. Rees^g,
David S. Powlson^h, Pete Smithⁱ, Shenghui Cui^{a,b*}

^a Key Lab of Urban Environment and Health, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen 361021, China

^b Xiamen Key Lab of Urban Metabolism, Xiamen 361021, China

^c College of Resources and Environmental Sciences, Key Laboratory of Plant-soil Interactions of MOE, China Agricultural University, Beijing 100193, China

^d College of Geography Science, Nanjing Normal University, Nanjing 210046, China

^e Department of Land Management, Zhejiang University, Hangzhou, 310058, PR China

^f School of Agriculture and Food, The University of Melbourne, Victoria, 3010 Australia

^g SRUC, West Mains Rd. Edinburgh, EH9 3JG, Scotland, UK

^h Department of Sustainable Agriculture Sciences, Rothamsted Research, Harpenden, AL5 2JQ, UK

ⁱ Institute of Biological and Environmental Sciences, University of Aberdeen, Aberdeen AB24 3UU, UK

Bing Gao & Tao Huang contributed equally to this work.

Corresponding author: **Xiaotang Ju and Shenghui Cui**

College of Resources and Environmental Sciences, Key Laboratory of Plant-soil Interactions of MOE,

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/gcb.14425](https://doi.org/10.1111/gcb.14425)

This article is protected by copyright. All rights reserved

China Agricultural University, Beijing 100193, China.

Phone: +86-10-62732006; Fax: +86-10-62731016.

E-mail: juxt@cau.edu.cn

Institute of Urban Environment, Chinese Academy of Sciences, 1799 Jimei Road, Xiamen 361021, China.

Phone: +86-592-6190777; Fax: +86-592-6190977.

E-mail: shcui@iue.ac.cn **Abstract**

Soil carbon sequestration is being considered as a potential pathway to mitigate climate change. Cropland soils could provide a sink for carbon that can be modified by farming practices, however, they can also act as a source of greenhouse gases (GHG), including not only nitrous oxide (N₂O) and methane (CH₄), but also the upstream carbon dioxide (CO₂) emissions associated with agronomic management. These latter emissions are also sometimes termed “hidden” or “embedded” CO₂. In this paper, we estimated the net GHG balance for Chinese cropping systems by considering the balance of soil carbon sequestration, N₂O and CH₄ emissions, and the upstream CO₂ emissions of agronomic management from a life cycle perspective during 2000–2017. Results showed that although soil organic carbon (SOC) increased by 23.2±8.6 Tg C yr⁻¹, the soil N₂O and CH₄ emissions plus upstream CO₂ emissions arising from agronomic management added 269.5±21.1 Tg C-eq yr⁻¹ to the atmosphere. These findings demonstrate that Chinese cropping systems are a net source of GHG emissions, and that total GHG emissions are about 12 times larger than carbon uptake by soil sequestration. There were large variations between different cropping systems in the net GHG balance ranging from 328 to 7567 kg C-eq ha⁻¹ yr⁻¹, but all systems act as a net GHG source to the atmosphere. The main sources of total GHG emissions are nitrogen fertilization (emissions during production and application), power use for irrigation, and soil N₂O and CH₄ emissions. Optimizing agronomic management practices, especially fertilization, irrigation, plastic mulching, and crop residues to reduce total GHG emissions from the whole chain is urgently required in order to develop a low carbon future for Chinese crop production.

Keywords: Agronomic management; Upstream CO₂ emissions; Life cycle analysis; Net greenhouse gas balance; N₂O and CH₄ emission; Soil organic carbon. **Introduction**

Soil is a large reservoir of carbon (C) in terrestrial ecosystems with a pool size of around 1500 Pg C (1 Pg = 10¹⁵ g) (Davidson et al., 1998; Lal, 2004). Cropland soil accounts for 8–10% of this C pool (Eswaran et al., 1993), which plays a significant role in the global C budget (Mahecha et al., 2010). There is an estimated technical potential to sequester 1.6 Pg C equivalents (C-eq) yr⁻¹ into agricultural soils globally (Smith et al., 2007; 2008). Hence, an initiative that aims to increase global agricultural SOC stocks by 0.4% (four per thousand) was launched (<http://4p1000.org>), in order to slow down rising levels of atmospheric CO₂ (Minasny et al., 2017), though doubts have been expressed as to whether this rate of increase is generally achievable (van Groenigen et al., 2017; Poulton et al., 2018).

The SOC stock changes represent the net exchange of CO₂ between soil and atmosphere (Mosier et al., 2006; Robertson and Grace, 2004; Shang et al., 2011). Many agricultural practices, such as optimized

fertilization, reduced tillage, and straw return to the fields have been advocated to mitigate GHG emissions by enhancing removals of CO₂ from the atmosphere (Smith et al., 2008; Snyder et al., 2009). Many field experimental studies have suggested that fertilizer application and straw return can increase soil C and sequester C from the atmosphere (Mosier et al., 2006; Shang et al., 2011). However, these practices may also stimulate nitrous oxide (N₂O) emissions that offset the SOC sequestration benefits (Pathak et al., 2005; Huang et al., 2013). Further, methane (CH₄) emissions can be increased after adding organic materials, especially in rice grown under flooded conditions (Zou et al., 2005; Shang et al., 2011). In addition to SOC sequestration, N₂O and CH₄ emissions from cropland soils are two additional crucial components of the GHG balance because of their high global warming potential (Robertson et al., 2000; Cubasch et al., 2013). The manufacture and transport of fertilizers and pesticides, power use for irrigation and field operations all require fossil fuels; the combustion of which results in GHG emissions (Snyder et al., 2009; Grassini and Cassman, 2012). Fertilization, irrigation, tillage and other management practices in the different cropping systems will affect the upstream C-eq, which is defined as C-eq released from such agricultural inputs (Robertson and Grace, 2004; Lal, 2007; Snyder et al., 2009; Schlesinger, 2010). The climate benefits of SOC sequestration in croplands might be offset by N₂O and CH₄ emissions (Tian et al., 2012; Norse and Ju, 2015), and upstream CO₂ released from the life cycle of agricultural inputs (Lal, 2007; Snyder et al., 2009; Schlesinger, 2010). The C lifecycle approach considers the full C cycle as it includes upstream C-eq release, but often does not consider soil GHG emissions and SOC sequestration (West and Marland, 2002; Robertson and Grace, 2004; Mosier et al., 2006). Elevated SOC storage in croplands will only mitigate climate change if the combined GHG emissions from agronomic practices are lower than the SOC sequestration from a life cycle perspective (Powlson et al., 2011). The climate benefits of a given cropping system should not only focus on the sequestration of SOC, but also on soil GHG emissions and the associated C-eq released from agricultural inputs and management practices (Mosier et al., 2006; Lal, 2007; Schlesinger, 2010). To measure the overall climate effect, the concept of net a GHG balance (kg C-eq ha⁻¹ yr⁻¹) has been proposed, based on the cumulative radiative forcing from all GHGs considered together (Robertson and Grace, 2004; Powlson et al., 2011). Greenhouse gas intensity (GHGI, kg C-eq kg⁻¹ grain) is used to compare the magnitude of GHG emissions to produce the same crop yield (Mosier et al., 2006; Grassini and Cassman, 2012). This concept can assist in solving the global challenges of increasing food production and concomitantly identifying the main targets for mitigation in different cropping systems and regions, which is important when seeking ways to decrease total GHG emissions associated with agricultural production, especially in China.

China covers a broad range of soil-climatic regimes, and corresponding cropping systems. The net GHG balance of different cropping systems will vary with soil-climatic conditions, crops and management practices. Large numbers of studies have investigated soil carbon sequestration, N₂O and CH₄ emissions in different cropping systems in China (Table S1). However, few have reported the net GHG balance associated with all sinks and sources in Chinese cropping systems (Huang et al., 2013; Gao et al., 2015). Furthermore, large uncertainties exist in these previous estimates, e.g. the GHG balance ranges over one

order of magnitude due to uncertainties in upstream CO₂ emissions of agronomic management practices (Zhang and Zhang, 2016). There is an urgent need, therefore, to synthesize literature for calculating the net GHG balance associated with soil N₂O and CH₄ emissions, the upstream CO₂ emissions from agronomic management and the change of SOC storage in Chinese cropping systems.

The aim of this paper is to obtain a national estimate of the net GHG balance in the main Chinese cropping systems. A database was compiled from relevant research published between 2000–2017, totaling 634 results for SOC changes and 233 results for N₂O and CH₄ emissions. Then, a meta-analysis was performed to explore the net GHG balance of each cropping system. We focused on five key questions in our analysis: (i) was there a change in topsoil (0–20 cm) SOC storage? (ii) what were the total GHG emissions associated with soil N₂O and CH₄ emissions and upstream CO₂ emissions from agronomic management? (iii) what were the main sources of total GHG emissions? (iv) what was the net GHG balance? and (v) what are the most effective measures to improve the net GHG balance of Chinese crop production?

Materials and methods

Description of Chinese cropping systems

Rice (*Oryza sativa*), wheat (*Triticum aestivum*), maize (*Zea mays* L.), potato (*Solanum tuberosum*), soybean (*Glycine max*), rapeseed (*Brassica rapa*), cotton (*Gossypium spp*), and vegetables (*Herbas*) are the main crops in China, accounting for 18.2%, 14.5%, 22.9%, 3.3%, 3.9%, 4.5%, 2.3% and 13.2% of the national total crop area sown, respectively (NBSC, 2016). Vegetables mainly included leafy vegetables, cabbages, fruit vegetables, melons, root and stem vegetables, etc. In 2015, China had 12.8 Mha orchard (*Hortus*), including apple, citrus, pear, peach and grape, etc., equivalent to 53.1% of the sown area of wheat (24.1 Mha) (NBSC, 2016). China has multiple crops each year, e.g. winter wheat–summer maize double cropping system, i.e. two harvests in one year, so we provide summary data on a yearly basis. Double cropping is different to annual rotations such as maize–soybean rotations that have a single crop in each year in places such as the US and elsewhere (Mosier et al., 2006). The main principle for classification of cropping systems in this study is based on different agricultural zones and the management practices used by farmers, because soil type, climate and fertilization practices of a given cropping system are similar within an agricultural zone. We collected data for greenhouse vegetables, open-field vegetables, potato and orchard systems across China, because these crops are widely distributed in all agricultural zones and there is insufficient available data on soil GHG emissions and SOC change for these systems within a single agricultural zone. This allows 15 distinct cropping systems to be defined (Table 1). We evaluated the mean GHG emissions and SOC change of a given cropping system in its dominant agricultural zone. The proportions of these cropping systems in relation to total crop sowing area are given in Appendix S1. The spatial patterns of the different cropping systems at a county-level (Fig. 1) were developed from a 30 × 30 m resolution land use map at a county-scale for China in 2010 provided by Wu et al. (2014). The specific combinations and distributions of cropping systems at county-level are shown in Appendix S2.

Data sources

Basal data used in this study, such as crop sowing area and yields, were taken from official statistical yearbooks and bulletins (NBSC, 2016). The data used for calculating the net GHG balance of the cropping systems, including the change of topsoil SOC, soil N₂O and CH₄ emissions, the upstream CO₂ from the manufacture and transportation of the chemical fertilizer (N, P₂O₅ and K₂O), power used for irrigation, fuel combustion in farm operations, application of pesticides, and film used for mulching crops or as a cover for greenhouse or fruits (Robertson et al., 2000; Mosier et al., 2006; Grassini and Cassman, 2012), were collected from the published literature, dissertations or books with data records for the period between 2000–2017.

The SOC data were categorized into three groups: (I) SOC data with monitoring of < 5 years; (II) SOC data with monitoring of ≥ 5 years; (III) data describing the mean changes of SOC for entire Chinese croplands. As most studies reported SOC change over at least a 5 year period in a given cropping system (Mosier et al., 2006; Lemke and Janzen, 2007; Huang et al., 2013); we mainly considered the effect of the duration of the experiment on SOC changes for this group classification. The detailed criteria for collecting SOC data are described in Appendix S3. In total, 647 results were collected, which were divided between each of the cropping systems as follows; WMN (41), WMSW(13), RW (47), RR (32), DR (100), SRNE (28), SRENE(46), SB (21), MNE (63), MNW (34), GV (42), OV (25), PS (30), CS (49), OS (76) and entire Chinese croplands (CC 14). We also obtained information for 582 locations of the collected SOC data under categories I and II (Fig. 2).

The criteria for collecting soil N₂O and CH₄ emissions, fertilizer input, the power used for irrigation, pesticides, fuel, and plastic film are described in Appendix S3. Results from 233 studies reporting N₂O and CH₄ emissions were collected from all cropping systems (Table S1). The detailed criteria for collecting the consumption of power use for irrigation, fuel and plastic film are described in Appendix S4. The adopted carbon emission factors are presented in Table 2.

Calculation of C-eq

An annual topsoil SOC sequestration rate (δSOC , kg C ha⁻¹ yr⁻¹) was estimated on the basis of an increased rate of change of the topsoil SOC density ($d\text{SOC}/dt$, g C kg⁻¹ yr⁻¹) [Eqn (1)] (Robertson et al., 2000; Shang et al., 2011).

$$\delta\text{SOC} = d\text{SOC}/dt \times \gamma \times 20/10 \quad (1)$$

where γ is the bulk density (g cm⁻³) of 0–20 cm depth topsoil, which was directly reported in the soil physical-chemical properties together with SOC concentrations in most of the collected literature. For the few sites with no reported bulk density, we firstly used bulk density of the same cropping system at a nearby site. If this was not available we assumed the mean bulk density values of 1.3 and 1.2 g cm⁻³ reported for upland and rice paddy fields in China, respectively (Pan et al., 2010). Since pedo-transfer functions also introduce uncertainty. The use of mean values for final gap filling was considered adequate

for the objectives of our study, so no pedo-transfer functions were used to estimate missing bulk density values. The values of 20 and 10 in the equation (1) are the topsoil depth and the area conversion coefficient, respectively. Most of the studies in Chinese cropping systems define the 0–20 cm soil depth as the plough layer under long-term conventional tillage practices, and the soil samples were taken to a depth of 20 cm to determine the SOC content, so we adopt a soil depth of 0–20 cm for calculating the SOC stock change.

Values of C-eq from soil N₂O and CH₄ emissions were estimated by multiplying the annual emissions in different cropping systems with the global warming potential (GWP) values over a 100-yr time horizon, which are 34 kg CO₂-eq kg⁻¹ or 9.3 kg C-eq kg⁻¹ for CH₄ and 298 kg CO₂-eq kg⁻¹ or 81.3 kg C-eq kg⁻¹ for N₂O (Cubasch et al., 2013). The C-eq emissions from certain agricultural inputs, i.e. the applied fertilizers, pesticides, plastic film and paper bags (kg ha⁻¹ yr⁻¹), power use for irrigation (kwh ha⁻¹ yr⁻¹), and fossil fuel use in farm operations (L ha⁻¹ yr⁻¹) were estimated by agricultural input, multiplying with the individual carbon intensity (in kg C per unit volume or mass) of manufacture and transportation of synthetic fertilizers, and/or applied for individual agricultural inputs (Table 2).

Calculation of net GHG balance and GHGI

We defined the boundary of the soil-crop system as the carbon gains and emissions per hectare per year between the soil and atmosphere, and calculated the main carbon fluxes from crop sowing to harvest using a life cycle approach (Robertson et al., 2000; Mosier et al., 2006; Smith et al., 2010). The “hidden” or “embedded” CO₂ from upstream production of fertilizers etc. or farmers’ operations are important components to be included in these comparison. The GHG emissions from non-agricultural IPCC sectors (e.g. energy) were included only where they are used specifically for agricultural use.

All the main fates of GHGs including upstream CO₂ emission, soil GHG emissions, CO₂ fixed by crops (photosynthesis) and emitted by crops (respiration) and soil SOC change, have to be considered when assessing a system’s capacity to act as a GHG sink or source. But the actual calculations of net GHG flux depend on the characteristics and nature of different ecosystems, in which only the actual carbon fluxes are included. For example, in forest systems, CO₂ fixed by plant photosynthesis must be included in addition to the change of soil SOC, because the aboveground biomass of forest is cumulative (Tang et al., 2018). However, for cropland ecosystems, the carbon in crops is not included, since these crops are harvested and consumed within a year, so there is no carbon sink; the carbon is simply recycled to the atmosphere within a year, so (very temporary) carbon stocks in crops should not be included in the GHG balance (Smith et al., 2010). The SOC change is the net balance between carbon inputs and outputs of the returned crop residues and soil respiration, and it also represents the net exchange of CO₂ between soil and atmosphere (Mosier et al., 2006; Smith et al., 2010).

Organic fertilizers (manure) accounted for 14.5% of the total N fertilization in China in 2010 (Gu et al., 2015). Our study set the boundary as Chinese cropping systems, which includes SOC stock changes and N₂O and CH₄ emissions induced by manure being applied to cropping systems, but does not include N₂O and CH₄ during storage, treatments (e.g. compost) and transportation of manure, which are regarded as

emissions from animal production (Smith et al., 2010). To evaluate a net GHG balance and yield basis of GHG emissions (GHGI), we used the following equation [Eqn (2) and (3)], which has widely been used in the calculations of net GHG balance and GHGI of different cropping systems (Robertson and Grace, 2004; Mosier et al., 2006; Shang et al., 2011, Grassini and Cassman, 2012; Gao et al., 2015; Zhou et al., 2017). A positive net GHG balance represents a source of C-eq to atmosphere, while a negative value represents a sink of C-eq from the atmosphere (Robertson et al., 2000; Mosier et al., 2006).

$$\text{Net GHG balance (kg C-eq ha}^{-1} \text{ yr}^{-1}) = a \times \text{CE}_{\text{N}_2\text{O}} + b \times \text{CE}_{\text{CH}_4} + c \times \text{CE}_{\text{N}} + d \times \text{CE}_{\text{P}} + e \times \text{CE}_{\text{K}} + f \times \text{CE}_{\text{I}} + g \times \text{CE}_{\text{F}} + h \times \text{CE}_{\text{PE}} + i \times \text{CE}_{\text{PF}} + j \times \text{CE}_{\text{PB}} - \delta \text{SOC} \quad (2)$$

$$\text{GHGI (kg C-eq Mg}^{-1}) = \text{Net GHG balance/Yield} \quad (3)$$

where the different small letters represent the amounts of soil GHG emissions and different agricultural inputs. $\text{CE}_{\text{N}_2\text{O}}$, CE_{CH_4} , CE_{N} , CE_{P} , CE_{K} , CE_{I} , CE_{F} , CE_{PE} , CE_{PF} and CE_{PB} represent the individual C emission equivalents for soil N_2O and CH_4 emissions, inputs of synthetic N, P_2O_5 and K_2O fertilizers, power use for irrigation, fuel, pesticides, plastic films and paper bags used for crop production, respectively (Table 2). 12 and 44 are the molecular weights of C and CO_2 . δSOC is the change of cropland SOC.

Variations at county scale

We converted each source and sink of the net GHG to an annual basis based on the compiled the datasets from relevant publications between 2000 and 2017. We then estimated the spatial patterns of topsoil SOC stock, and GHG emissions (kg C-eq ha^{-1}) from N_2O and CH_4 emissions, N fertilizer input, power use for irrigation and other sources including P_2O_5 and K_2O application, fuel in farm operations, pesticides and plastic film use and total GHG emissions (kg C-eq ha^{-1}) in Chinese cropping systems from the above sources (Fig. S1), and net GHG balance based on the available 30 m $\times$ 30 m land use map and spatial pattern of cropping systems at county-scale for China in 2010, which represents the mean pattern of the study period. The weighted mean SOC stock change and net GHG balance between greenhouse vegetables (342 kg C $\text{ha}^{-1} \text{ yr}^{-1}$, 7567 kg C-eq $\text{ha}^{-1} \text{ yr}^{-1}$) and open-field vegetables (296 kg C $\text{ha}^{-1} \text{ yr}^{-1}$, 4617 kg C-eq $\text{ha}^{-1} \text{ yr}^{-1}$) was calculated based on the average SOC stock change and net GHG balance, and the area of the two types vegetables and total vegetable production, because of lack of the proportions of two types of vegetable production at county scale.

Statistical and uncertainty analysis

The significance of the differences in SOC stock change of different cropping systems were tested with an analysis of variance (ANOVA) using the SPSS16.0 statistical package. Statistical significance was determined at the 95% confidence level at $p < 0.05$. In addition, to minimize the uncertainty of our analysis, we first set up uniform criteria for collecting topsoil SOC change, and other emission sources from different cropping systems. Then the mean and variation of these data were calculated with the 90th

percentile confidence interval. The uncertainty of the net GHG balance was analyzed using the error propagation equation of mathematical statistics (IPCC, 2000) by the uncertainties of the collected data on SOC change, soil GHG emissions and upstream CO₂ emissions from agronomic managements. A detailed description of the error of propagation equation from a mathematical statistical analysis is shown in Appendix S5. The same error propagation equation was used for calculating the uncertainties for total SOC stock change and total GHG emissions.

Results

Change of SOC in Chinese cropping systems

The frequency distribution of the annual SOC change showed an overall increase of the SOC stocks in the topsoil of Chinese cropping systems (Fig. 3) and the SOC changes were significantly ($p < 0.05$) different among the 15 cropping systems (Table S4). Overall, 78.1% of the observations showed an increase in SOC stocks, with decreases in SOC mainly occurring in MNE, OS, CS, SB and PS systems (Table S2). The annual change values of SOC stock were in the range -3,309 to 3,687 kg C ha⁻¹ yr⁻¹, and followed a normal distribution, with the minimum values occurring in the MNE system in the Northeast and maximum values in the DR system in Central China.

All cropping systems except SB showed an increase in SOC stock during the 2000-2017 period. The magnitudes were 170–825 and 50–432 kg C ha⁻¹ yr⁻¹ in groups I and II, respectively (Fig. 4). In group III, the mean change of SOC across all Chinese croplands was 178 ± 27 kg C ha⁻¹ yr⁻¹ with a range of 27–538 kg C ha⁻¹ yr⁻¹. The standard deviations of the observations in group I were much larger than those in groups II and III (Fig. 4), highlighting greater uncertainties of SOC stock changes when the study duration is less than 5 years. Therefore, the changes of SOC from method II were used to estimate the net GHG balance of Chinese cropping systems.

The distribution of SOC stock changes in different cropping systems at county-scale is shown in Fig. 5. The regions with the highest SOC increase have a high proportion of paddy fields (Fig. 1), located in the Middle and Lower Yangtze River, Sichuan Basin, eastern Heilongjiang province, as well as some scattered regions in Southern and Eastern China. The regions with the second-highest SOC increase were dominated by winter wheat, summer maize, cotton, and vegetable production (Fig. 1). These regions are mainly concentrated on the North China Plain, north of Sichuan Basin, southern Shanxi, Shaanxi and Gansu provinces and northwest Xinjiang. The regions with relatively low SOC increases were dominated by single spring maize, potato and orchard planting, located in Northeastern China, southern Inner Mongolia, northern Shanxi and Shaanxi provinces, Sichuan Basin and Southwestern China. A decrease of SOC storage occurred in northeastern Inner Mongolia and western Heilongjiang due to the cultivation of soybean in soils with an initially high SOC stock (Fig. 1).

Net GHG balance, their main sources and GHGI

All cropping systems acted as a net GHG source when considering the combined soil N₂O and CH₄

emissions and upstream CO₂ calculated from the life cycle emissions of agricultural inputs. The net GHG balance was in the range of 328–7567 kg C-eq ha⁻¹ yr⁻¹, with a rank of SB < MNE < PS < WMSW ≈ CS < MNW < OS ≈ WMN < SRNE < SRENE < RR < RW < OV < DR < GV (Table 3). The spatial pattern of the net GHG balance at county-level showed that the regions with the highest net GHG emissions were mainly found in North China due to vegetable planting with very high fertilizer N applications, over-irrigation, and plastic used for greenhouse covering (Fig. 6). The regions with the second-highest net GHG balance were concentrated in Central, Southern and Eastern China, because of high CH₄ emissions from cultivation of rice in those regions. The regions with the lowest net GHG balance were mainly located in Northeastern China, central and southern Inner Mongolia, Gansu and Southwestern China in SB, MNE and PS systems (Fig. 1).

Except for the SB system, manufactured and transported fertilizer N was the largest source of C-eq emissions in upland systems (Fig. 7), accounting for between 29.3–71.8% of the net C-eq emissions. N fertilization induced large emissions of N₂O, accounting for 3.6–37.7% of net C-eq emissions. The total C-eq emissions from power use for irrigation ranged from 129 to 1,178 kg C ha⁻¹ yr⁻¹, accounting for 14.2–34.2% of the net C-eq emissions except for WMSW, SB and MNE systems. Plastic films are mainly used to cover the ground surface to promote crop germination by increasing soil surface temperature when sowing in relatively low temperature periods, and to increase water use efficiency by reducing soil surface evaporation loss, or for use to cover greenhouses. Plastic films are also an important emissions source, accounting for up to 22.7–24.5% of the net C-eq emissions in MNW, GV and PS systems. In paddy fields, CH₄ emissions were the largest contributor to emissions, accounting for 39.4%, 41.9%, 73.7%, 50.8% and 64.0% of the net C-eq emissions in RW, RR, DR, SRNE and SRENE systems, respectively, followed by emissions from irrigation (14.6–40.7%) and fertilizer N (11.5–24.6%). Nitrous oxide was an important source of emissions in RW and RR systems, accounting for 19.7% and 19.1% of the net C-eq emissions, respectively. Methane uptake from WMN, SB, MNE, MNW, GV, OV, PS, CS and OS contributed a small amount of negative emissions (< 1%). Emissions from the application of P₂O₅ and K₂O, fuel in farm operations and pesticides application contributed to 0.8–5.8%, 0.6–4.4%, 0.4–10.3% and 0.7–6.7% of net C-eq emissions, respectively, in all cropping systems except the SB system.

Changes of the SOC sink accounted for only 2.5–28.7% of the total C-eq emissions of the cropland systems (Fig. 7). The decrease in SOC stocks in the northeast SB system acted as a source of CO₂, accounting for 17.0% of the total C-eq emissions in this system.

The magnitude of GHG emissions to produce the same crop yield were in the range of 65–648 kg C-eq Mg⁻¹, with a rank of MNE < SB < WMSW < WMN < MNE < PS < SRNE < RR < RW < OV ≈ CS < SRENE < GV < DR < OS (Table 3).

Soil C sequestration vs total GHG emission

We estimated the total topsoil SOC increase and total GHG emissions of Chinese cropping systems by

multiplying the SOC change and the total C-eq emissions of each cropping system with their sowing area (Table 3). Total topsoil SOC changed at rates of between -0.2 – 4.5 Tg ($1 \text{ Tg} = 10^{12} \text{ g}$) C yr^{-1} , resulting in an accumulation of 23.2 ± 8.6 Tg C yr^{-1} across the whole China, close to the calculated mean increase in the topsoil C stock of China's croplands of 25.5 Tg C yr^{-1} between 1985 and 2006 (Pan et al., 2010). The GHG emissions from different cropping systems was in the range of 1.0 – 35.3 Tg C-eq yr^{-1} , with a rank of $\text{SB} < \text{MNW} < \text{WMSW} < \text{CS} < \text{GV} < \text{PS} < \text{SRNE} < \text{MNE} < \text{SRENE} < \text{RW} < \text{RR} < \text{OS} < \text{DR} \approx \text{WMN} < \text{OV}$, summing to 269.5 ± 21.1 Tg C-eq yr^{-1} , accounting for 13.9–15.2% of total national GHG emissions in 2005–2007 (Yan and Yang, 2010; National Development & Reform Commission of China, 2014). We further calculated the ratio of total GHG emissions from different cropping systems to total soil carbon sequestration. These ratios were 7.4 (WMN), 6.9 (WMSW), 11.9 (RW), 10.3 (RR), 13.4 (DR), 6.9 (SRNE), 9.7 (SRENE), 4.5 (MNE), 6.9 (MNW), 23.1 (GV), 16.5 (OV), 9.2 (PS), 6.0 (CS), and 41.6 (OS), respectively, and about 12 for all Chinese croplands (Table 3).

Discussion

Changes in SOC in Chinese cropping systems

In order to achieve the objectives of the Paris Climate Change Agreement, to keep global temperature increases well below 2°C , it is widely recognized that negative emission technologies will be needed to lower atmospheric concentrations of CO_2 (Rockstorm et al., 2017). Soil carbon sequestration by cropland soils could play a potential role in many regions (Robertson and Grace, 2004; Smith et al., 2008; Powlson et al., 2011; Wollenberg et al., 2016). In this paper, we indeed find that cropland SOC increased in Chinese cropping systems, with a range of 50 to 432 $\text{kg C ha}^{-1} \text{ yr}^{-1}$, close to a suggested global mean rate of 300 to 500 $\text{kg C ha}^{-1} \text{ yr}^{-1}$ (Lal, 2007). The increase of cropland SOC in China in the past forty years can be mainly attributed to increased crop yields resulting from improvements in agronomic management in Chinese (i.e. new crop varieties, fertilizer inputs, improved protection from pests, diseases and weeds, irrigation), and increased yields leading to larger organic matter returns to soil from roots and stubble (Huang and Sun, 2006; Yan et al., 2011; Han et al., 2017; Zhao et al., 2018; Powlson et al., 2018). We found that the average yields per unit area showed a good correlation ($p < 0.05$) with the SOC increase of the different cropping systems (Fig. S2). The increase in SOC has also been attributed to the development of no-tillage and reduced-tillage practices in China (Huang and Sun, 2006; Yan et al., 2011). The initially low average SOC content (11.5 – 12.0 g kg^{-1} for 0 – 20 cm depth) between 1979 and 1982 is another contributory factor for the observed increase in SOC across China (Yan et al., 2011; Yang et al., 2017). Even though the average SOC content in soil had increased to 12.7 – 14.3 g kg^{-1} in 2005–2014, it is still lower than in many European and US cropland soils (Johnston et al., 2009; Yan et al., 2011; Fan et al., 2012; Yang et al., 2017; Zhao et al., 2018). The lower starting point was mainly caused by soil mining, together with low crop residue returns to the soil due to lower yields, in turn caused by low inputs and poor agronomic management before policy changes in 1978 (Huang and Sun, 2006; Fan et al., 2012; Zhao et al., 2018).

The weighted mean rate of increase in SOC was 243 ± 92 $\text{kg C ha}^{-1} \text{ yr}^{-1}$ for the main Chinese cropping

systems (Table 3), higher than the mean annual SOC increase of $178 \pm 27 \text{ kg C ha}^{-1} \text{ yr}^{-1}$, which was 27 to 538 $\text{kg C ha}^{-1} \text{ yr}^{-1}$ derived from reviewing previous studies on total cropland area in China (Table S2). This difference might be explained by the following factors. Firstly, in addition to the main crops (rice, maize, wheat, and vegetables), Chinese croplands also produce other crops such as millet, sorghum, peanut, tobacco, hemp crops, etc. which might lead to smaller SOC increases compared to the OS and PS systems across China, or cause a decrease as in the SB system in Northeastern China, because the new carbon input from crop growth in these systems is lower than in the main cropping systems for production of rice, maize, wheat and vegetables; the quantity of crop residues is a key factor in determining changes in SOC stocks (Yan et al., 2011; Yang et al., 2017). Secondly, long-term field experiments on SOC change are typically established on the main crops, including wheat, rice, maize, and vegetables, which have higher yields compared to those of soybean, millet, sorghum, peanut, tobacco, hemp crops, etc., which may lead to an overestimation of the SOC increase for some relatively low yielding crops. Thirdly, study durations of the SOC data used in this paper are mainly between 5–15 years, shorter than previous long-term studies on total cropland of China such as Yan et al. (2011), in which the study duration was around 30 years. The annual increase rates of SOC in cropland are usually larger in the initial years following a change in management, declining with the duration of study, and reaching an equilibrium after around 20–50 years (Mosier et al., 2006; Lemke and Janzen, 2007; Wang et al., 2017). The SOC change might also be underestimated a little since changes of SOC stock consider only the top 0–20 cm soil layer, and changes of SOC stock below 20 cm might also contribute to soil carbon sequestration (Yan et al., 2011; Zhao et al., 2018), but any such underestimation should not greatly affect our main results and conclusions.

There is still a potential to increase SOC stocks in Chinese croplands. Compared to the high SOC concentrations of $14.5\text{--}23.2 \text{ g kg}^{-1}$ or SOC stocks (0–20 cm) ($40.2\text{--}43.7 \text{ t C ha}^{-1}$) in Europe and the US (Fan et al., 2012; Zhao et al., 2018), over half of the topsoil in China's croplands have SOC concentrations lower than 11.6 g kg^{-1} (Yang et al., 2017), and the estimated 0–20 cm SOC stocks were $26.6\text{--}29.4 \text{ t C ha}^{-1}$ in 1979–1982 and $31.4\text{--}33.5 \text{ t C ha}^{-1}$ in 2007–2011 (Yan et al., 2011; Zhao et al., 2018). Topsoil SOC pools can increase rapidly in the early years after a change of management and then more slowly thereafter, as observed for reduced- and no-tillage (West and Post, 2002; Lal, 2007; Johnston et al., 2009). Conservation tillage practices have only begun recently in China, which means there should be potential to increase SOC in Chinese croplands in the coming years (Yang et al., 2017); however, recent evidence suggests that conservation tillage may largely redistribute carbon within the profile, though net carbon gains are often still observed (Powlson et al., 2014). Changes in SOC stock can be influenced by the availability of nutrients (van Groenigen et al., 2017) and it is possible that more appropriate management of nutrients in Chinese croplands (probably decreased N applications but increased P, K, S or micronutrients) could lead to greater increases in SOC.

Soil N₂O and CH₄ emissions and upstream CO₂ emission

Although topsoil SOC stocks increased in the main cropping systems in China, its net effect on GHG

balance was more than offset by large emissions of soil N₂O and CH₄ and upstream CO₂ emissions from agronomic management (Fig. 6). This emphasizes that climate change mitigation strategies cannot rely only on SOC sequestration in croplands, and more effort is required for reducing total GHG emissions from cropland management (Fig. S1). Indeed, the SOC increase is smaller than the emissions of soil N₂O and CH₄, even without considering upstream CO₂ emissions from cropland management practices.

Emissions from the manufacture and transportation of N fertilizer are the largest contributor to total GHG emissions in China (Fig. 7). Emissions are about 2–31 times greater than that of the major cropping systems in the US, because of the higher N application rate and higher CO₂ emissions associated with the manufacture and transportation of N fertilizer in China, which is 8.3 vs 3.0 kg CO₂ kg⁻¹ N applied in China vs US, respectively (Mosier et al., 2006; Zhang et al., 2013). In China, the cropland SOC sink is mainly caused by N fertilizer applications (Tian et al., 2012), but this also results in large N₂O emissions (Bouwman et al., 2002; Synder et al., 2009; Gao et al., 2015).

Pumping of groundwater for irrigation is one of the most energy consuming on-farm processes and it represents an important source of GHG emissions that has rapidly increased, and which at present is largely unregulated (Wang et al., 2012). The high emissions from irrigation are mainly due to a combination of excessive irrigation, low energy use efficiency for pumping, and high power generation emissions. In China, power demand for pumping per unit of water is 4.3 kwh mm⁻¹ ha⁻¹, which falls into the range of 2.1–6.4 kwh mm⁻¹ ha⁻¹ estimated by Wang et al. (2012), compared to only 0.15 kwh mm⁻¹ ha⁻¹ in the US. Further, CO₂ emissions from electricity generation in China are 1.32 kg CO₂ kwh⁻¹, much higher than the 0.32 kg CO₂ kwh⁻¹ in the US (Mosier et al., 2006; Zhang et al., 2013). This part of the emissions budget will increase with the decline of groundwater table and the increase in air temperature in China if no improvements in water conservation are made (Foster and Garduño, 2004; Liu et al., 2010; Powlson et al., 2018).

Plastic film usage has a large contribution to total GHG emissions in MNW, GV, OV, PS, and CS systems, even exceeding the contribution of N₂O emissions in MNW and PS systems. In the GV system, pesticides contribute to large emissions of about 3.0–33.3 times that of the same source in other cropping systems. More attention needs to be paid to this source, given that the planting area of greenhouse vegetables has been increasing by around 10% per year (Fan et al., 2014).

Except for the OV system, we found that upland soils act as a weak sink for CH₄, but it can be neglected given that it represents <1% of the net GHG balance (Cui et al., 2013; Gao et al., 2015). However, CH₄ emissions are an important contributor to total GHG emissions in Chinese rice-based rotations (Ma et al., 2013; Shang et al., 2011). CH₄ emissions from paddy rice cultivation accounts for 20.3% of total GHG emissions in Chinese agriculture (National Development & Reform Commission of China, 2014).

With population and economic growth, Chinese grain demand is expected to increase by 6.9%, 3.3% and 52.9% for rice, wheat and maize by 2030, respectively, relative to 2012 (Chen et al., 2014). Due to the limitation on arable land expansion in China (Burney et al., 2010; Cui et al., 2013), producing more food might occur at the expense of increasing nutrient inputs if no other improvements in agronomic

management are made. This presents the challenge of producing more grains with fewer inputs, and with reduced environmental and climate impacts.

We showed the weighted mean SOC stock change and net GHG balance for total vegetable production (GV+OV) at county scale, because there is no data on the proportion of the two types of vegetables at this scale. This represents the mean status of SOC change and net GHG emissions of Chinese vegetables production. However, the SOC change and net GHG balance of greenhouse vegetables were about 1.2 and 1.6 times greater than open-field vegetables. Greenhouse vegetables are, mainly distributed around Bohai and Huang-Huai-Hai region, the middle and low reaches of Yangtze river, and Northwest China, and these regions accounted for 60.3%, 19.7% and 7.5% of the total sown area of greenhouse vegetables in 2010, respectively (DAMM, 2017). As a result, this study might have underestimated the SOC change and net GHG emissions of vegetable production in counties with a high proportion of greenhouse vegetables in the three main greenhouse vegetable producing areas mentioned above, and might have overestimated the SOC change and net GHG emissions of vegetable production in counties with low proportions of greenhouse vegetables. The impacts of the production on SOC change and net GHG balance of greenhouse vegetables at a county scale requires further study.

GHGI of different cropping systems

GHGI provides a platform for comparing the overall effects of any given cropping system on GHG emissions per unit of crop yield (Mosier et al., 2006; Grassini and Cassman, 2012). The magnitude of GHG emissions per unit of grain yield ranged from 65 to 648 kg C-eq Mg⁻¹ in different Chinese cropping systems. This is significantly higher than that of 32–61 kg C-eq Mg⁻¹ in conventional irrigated maize systems in Northeastern Colorado and Nebraska of the U.S., which was -35 kg C-eq Mg⁻¹ in conventional no-till corn–soybean rotation in Northeastern Colorado of the U.S (Mosier et al., 2006; Grassini and Cassman, 2012), and 75 kg C-eq Mg⁻¹ in conventional wheat and double-cropped soybean in mid-Atlantic region of the U.S. (Cavigelli et al., 2009). In China, the net GHG balance of rice production systems (DR, SRNE and SRENE) was 2154–5342 kg C-eq ha⁻¹ yr⁻¹, close to that of 1718–5342 kg C-eq ha⁻¹ yr⁻¹ in Japan, USA and Italy. The high net GHG balance of rice production in different countries resulted from high baseline of CH₄ emissions (Hokazono and Hayashi, 2012). However, the GHGI of Chinese rice production (253–535 kg C-eq Mg⁻¹) is lower relative to GHGI for rice in Japan, USA and Italy (398–753 kg C-eq Mg⁻¹), because rice yield reach 7.7–12.9 Mg ha⁻¹ in China (Table 3), but only 4.4–7.3 Mg ha⁻¹ in Japan, USA and Italy (Hokazono and Hayashi, 2012). The high GHGI of upland cropping systems in China was caused by over-use of fertilizer and over-irrigation, but with low yields of maize, wheat and soybean relative to the U.S. (Mosier et al., 2006; Cavigelli et al., 2009; Grassini and Cassman, 2012). Many studies have indicated that China has a large potential to produce more grain with lower GHG emissions by optimizing fertilization and irrigation (Ju et al., 2009; Cui et al., 2013; Chen et al., 2014), and this could close the gap of GHGI for grain production between China and low GHGI countries.

Reducing total cropland GHG emissions

The ratio of total GHG emissions to soil C sequestration was reported in this study, which expresses the ability of soil C sequestration to offset the total GHG emissions of the soil-crop system. This showed that total GHG emissions are about 12 times larger than soil carbon sequestration, indicating that non-CO₂ GHG emissions need to be reduced substantially if Chinese crop production is to become GHG neutral with the help of soil C sequestration. Major increases in yield have been achieved in Chinese cropping systems over the time period for which soil carbon changes were assessed, but the yield increase came at the cost of higher N₂O emissions. By evaluating the overall GHG balance, the environmental cost of these yield increases can be seen, so that improved agronomic management practices can be identified to reduce these impacts. This study shows that the main measures to reduce the net GHG balance of Chinese cropping systems should focus on reducing N₂O emissions, chemical N fertilizer use, GHG emissions from fertilizer manufacture, power use for irrigation, and CH₄ emissions in rice-based cropping systems, while at the same time, increasing soil C sequestration. Moreover, emissions from the use of plastic film and pesticides should also be considered in vegetable cropping systems.

Emissions of N₂O are normally positively correlated with N fertilizer inputs, and with N surpluses (difference between total N input and crop N uptake) in cropping systems (van Groenigen et al., 2010; Grassini and Cassman, 2012; Cui et al., 2013; Gao et al., 2015). Appropriate N fertilization not only reduces N₂O emissions, but also reduces emissions associated with manufacture and distribution of N fertilizer (Mosier et al., 2006; Huang and Tang, 2010; Gao et al., 2015). Chinese cropping systems usually receive excessive amounts of N fertilizer, about two to even tens of times greater than the actual crop demand (Ju et al., 2009; Chen et al., 2014; Nayak et al., 2015). It has been clearly demonstrated that N fertilization rates can be reduced by knowledge-based N management practices without compromising crop yields, and in some cases even causing an increase (Ju et al., 2009; Chen et al., 2014; Xia et al., 2016). In addition to decreasing the quantity of N fertilizer applied, altering the timing of N fertilizer application can increase N use efficiency and decrease N₂O emissions (Ju et al., 2009; Cui et al., 2013; Chen et al., 2014). Practices such as deep placement of N fertilizer could also reduce N₂O emissions by 5–40% (Xia et al., 2016). If a 30% reduction in N fertilization rate was achieved, a potential reduction in GHG emissions would reach 16.4 Tg C-eq from production of paddy rice, wheat, maize and soybean (Cheng et al., 2015). Although knowledge-based practices are available in China, there remain socioeconomic barriers, such as small farm size that need to be addressed to facilitate their widespread implementation (Ju et al., 2016).

The reduction of emissions from power use for irrigation should concentrate on reducing energy consumption, which depends on the efficiency of irrigation and power generation (Mosier et al., 2006; Grassini and Cassman, 2012). Measures for reducing irrigation emissions include testing of soil water content (Meng et al., 2012) and development of fertigation systems (Fan et al., 2014). These knowledge-based irrigation practices could reduce irrigation emissions by 16–43% in Chinese cropping systems (Cabangon et al., 2004; Meng et al., 2012; Fan et al., 2014). Energy use, both for the purposes of irrigation and fertilizer manufacture, is currently associated with high GHG emissions as a consequence of

its dependence on fossil fuel-based energy sources. Therefore, substitution of these energy sources by low emission renewable alternatives could significantly reduce the carbon footprint of food production (Schandl et al., 2016).

Methane emissions were mainly affected by water regimes and organic amendments in paddy fields (Wassmann et al., 2000; Shang et al., 2011; Nayak et al., 2015). Strategies for reducing CH₄ emissions include mid-season drainage and intermittent irrigation (Wassmann et al., 2000; Zou et al., 2005; Ma et al., 2013; Cheng et al., 2014; Nayak et al., 2015). These measures could reduce CH₄ emissions from paddy fields by 36–65% (Zou et al., 2005; Ma et al., 2013). However, a trade-off between CH₄ and N₂O emissions appeared from mid-season drainage and intermittent irrigation (Wassmann et al., 2000; Zou et al., 2005; Ma et al., 2013). The increase of N₂O emissions offset the reduction of CH₄ emissions by 49.2% and 67.6% for plots without and with wheat straw amendment in midseason drainage, respectively. Reductions in CH₄ emissions could be completely offset by increased N₂O emissions when the field was moist but not waterlogged by intermittent irrigation, in comparison with the treatment that was frequently waterlogged within the midseason drainage period (Zou et al., 2005). Therefore, Wassmann et al. (2000) proposed that the changes in water regime are only recommended for rice systems with high baseline emissions of CH₄ from waterlogged and midseason drainage to intermittent irrigation.

There is considerable interest in the possibility of mitigating climate change by sequestering extra C from atmosphere into soil through changes in land management (Smith et al., 2008; Powlson et al., 2011, 2018). Effective measures for increasing SOC stocks mainly include the return of crop residues to soils, the application of biochar, conservation tillage, and mulch plants (Synder et al., 2009; Zhao et al., 2014; Qian et al., 2015; Nayak et al., 2015; Zhou et al., 2017; Powlson et al., 2018). However, the direct return of straw to rice paddy fields is not recommended, because CH₄ emissions are increased by a factor of 1.6–3.7 and the net GHG emissions associated CH₄ and N₂O are greatly increased in rice paddy fields when receiving organic amendments (Zou et al., 2005). Converting straw to biochar then applying it to soils is a possible alternative to soil C sequestration, and could contribute to CH₄ mitigation, and improvements of soil and crop productivity, without increasing N₂O emissions (Zhao et al., 2014). However, there are widely divergent opinions in the scientific community about the practicalities and economics of biochar production and use (Powlson et al., 2018). The global mean rate of SOC sequestration for conversion from conventional tillage to no-till is 100–200 kg ha⁻¹ yr⁻¹ (Lal, 2007). Converting conventional tillage to reduced tillage in rice-based cropping systems in China could sequester 213 kg C ha⁻¹ yr⁻¹ (Nayak et al., 2015) but, again, the small size of most farms in China presents practical and economic barriers to adoption of reduced tillage. Mulching different living plants could significantly increase SOC sequestration in orchards soil (Qian et al., 2015).

Emissions associated with the manufacture of plastic film use in the GV system could be reduced by prolonging its service life and recycling of the abandoned film (Cheng et al., 2011). Further, the high rates of pesticide application in Chinese GV systems could be reduced by controls on the occurrence of disease and insect pests through technologies such as reduced temperature and humidity in greenhouses, and

physical controls (i.e. trapping and insect screens), advanced application equipment and drip irrigation (Wang and Wang, 2016) and integrated pest management (Pretty and Bharucha, 2015).

Despite the soil carbon sink found in Chinese cropland soils, emissions of N_2O and CH_4 and upstream CO_2 -eq emissions associated with agronomic management are about one order of magnitude larger than the soil carbon sink under current farmers' practices. Chinese croplands are therefore a net GHG source. Over-fertilization with N and low energy use efficiency of irrigation and other agronomic management practices are largely responsible for these high GHG emissions. To feed an increasingly wealthy population, Chinese crop production is expected to continue to expand in the future, posing great challenges for reducing GHG emissions. However, there is still much room for improving the net GHG balance of Chinese croplands. Mitigation measures can focus on, but are not limited to, optimizing fertilizer applications, better irrigation practices and conservation tillage.

Acknowledgements

This work was funded by National Basic Research Program of China (2014CB953800), Young Talents Projects of the Institute of Urban Environment, Chinese Academy of Sciences (IUEMS201402), National Natural Science Foundation of China (41471190, 41301237, 71704171), China Postdoctoral Science Foundation (2014T70144) and Discovery Early Career Researcher Award of the Australian Research Council (DE170100423). The work contributes to the UK-China Virtual Joint Centres on Nitrogen "N-Circle" and "CINAg" funded by the Newton Fund *via* UK BBSRC/NERC (grants BB/N013484/1 and BB/N013468/1, respectively).

References

- Bouwman, A. F., Boumans, L. J. M., & Batjes, N. H. (2002). Emissions of N₂O and NO from fertilized fields: summary of available measurement data. *Global Biogeochemical Cycles*, 16, 1080–1088.
- Burney, J. A., Davis, S. J., & Lobell, D. B. (2010). Greenhouse gas mitigation by agricultural intensification. *Proceedings of the National Academy of Science USA*, 107, 12052–12057.
- Cabangon, R. J., Tuong, T. P., Castillo, E. G., Bao, L. X., Lu, G. A., Wang, G. H., Cui, Y. L., Bouman, B. A. M., Li, Y. H., Chen, C. D., & Wang, J. Z. (2004). Effect of irrigation method and N-fertilizer management on rice yield, water productivity and nutrient-use efficiencies in typical lowland rice conditions in China. *Paddy and Water Environment*, 2, 195–206.
- Cavigelli, M. A., Djurickovic, M., Rasmann, C., Spargo, J. T., Mirsky, S. B., & Maul, J. E. (2009). Global warming potential of organic and conventional grain cropping systems in the mid-Atlantic region of the U.S. 2009 Farming Systems Design Proceedings, 23-26 August, Monterey, California, p. 51–52.
- Chen, X. P., Cui, Z. L., Fan, M. S., Vitousek, P., Zhao, M., Ma, W. Q., Wang, Z. L., Zhang, W. J., Yan, X. Y., Yang, J. C., Deng, X. P., Gao, Q., Zhang, Q., Guo, S. W., Ren, J., Li, S. Q., Ye, Y. L., Wang, Z. H., Huang, J. L., Tang, Q. Y., Sun, Y. X., Peng, X. L., Zhang, J. W., He, M. R., Zhu, Y. J., Xue, J. Q., Wang, G. L., Wu, L., An, N., Wu, L. Q., Ma, L., Zhang, W. F., & Zhang, F. S. (2014). Producing more grain with lower environmental costs. *Nature*, 514 (7523), 486–489.
- Cheng, K., Pan, G. X., Smith, P., Luo, T., Li, L., Zheng, J. F., Zhang, X., Han, X., & Yan, M. (2011). Carbon footprint of China's crop production: an estimation using agrostatistics data over 1993–2007. *Agriculture, Ecosystems and Environment*, 142, 231–237.
- Cheng, K., Ogle, S. M., Parton, W. J., & Pan, G. X. (2014). Simulating greenhouse gas mitigation potentials for Chinese Croplands using the DAYCENT ecosystem model. *Global Change Biology*, 20, 948–962.
- Cheng, K., Yan, M., Nayak, D., Smith, P., Zheng, J. F., & Zheng, J. W. (2015). Carbon footprint of crop production in China: an analysis of National Statistics data. *Journal of Agricultural Science*, 153, 422–431.
- Cubasch, U. D., Wuebbles, D., Chen, D. L., Facchini, M. C., Frame, D., Mahowald, N., & Winther, J. G. (Eds.). (2013). *Climate Change 2013: The Physical Science Basis*. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 121–155.
- Cui, Z. L., Yue, S. C., Wang, G. L., Meng, Q. F., Wu, L., Yang, Z. P., Zhang, Q., Li, S. Q., Zhang, F. S., & Chen, X. P. (2013). Closing the yield gap could reduce projected greenhouse gas emissions: a case study of maize production in China. *Global Change Biology*, 19, 2467–2477.
- Davidson, E. A., Belk, E., & Boone, R. D. (1998). Soil water content and temperature as independent or confounded factors controlling soil respiration in temperate mixed hardwood forest. *Global Change Biology*, 4, 217–227.

- DAMM (Department of Agricultural Mechanization, Ministry of Agriculture). (2017). <http://data.sheshiyuanyi.com/AreaData/>.
- Eswaran, H., Van Den Berg, H., & Reich, P. (1993). Organic Carbon in Soils of the World. *Soil Science Society of American Journal*, 57(1), 192–194.
- Fan, M. S., Shen, J. B., Yuan, L. X., Jiang, R. F., Chen, X. P., Davies, W. J., & Zhang, F. S. (2012). Improving crop productivity and resource use efficiency to ensure food security and environmental quality in China. *Journal of Experimental Botany*, 63, 13–24.
- Fan, Z. B., Lin, S., Zhang, X. M., Jiang, Z. M., Yang, K. C., Jian, D. D., Chen, Y. Z., Li, J. L., Chen, Q., & Wang, J. G. (2014). Conventional flooding irrigation causes an overuse of nitrogen fertilizer and low nitrogen use efficiency in intensively used solar greenhouse vegetable production. *Agriculture Water Management*, 144, 11–19.
- Foster, S., & Garduño, H. (2004). China: Towards Sustainable Groundwater Resource Use for Irrigated Agriculture on the North China Plain. Counterpart Organizations: Ministry of Water Resources (MWR), Institute of Water Resources and Hydropower (IWRH) and Guantao County Water Resources Bureau (GCWRB), the World Bank water resources website (www.worldbank.org/gwmate) and the Global Water Partnership website (www.gwpforum.org). The World Bank, Washington D.C., USA, pp. 1–16.
- Gao, B., Ju, X. T., Meng, Q. F., Cui, Z. L., Christie, P., Chen, X. P., & Zhang, F. S. (2015). The impact of alternative cropping systems on global warming potential, grain yield and groundwater use. *Agriculture, Ecosystems and Environment*, 203, 46–54.
- Grassini, P., & Cassman, K. G. (2012). High-yield maize with large net energy yield and small global warming intensity. *Proceedings of the National Academy of Science USA*, 109, 1074–1079.
- Gu, B. J., Ju, X. T., Chang, J., Ge, Y., & Vitousek, P. M. (2015). Integrated reactive nitrogen budgets and future trends in China. *Proceedings of the National Academy of Science USA*, 11(28), 8792.
- Han, D. R., Wiesmeier, M., Conant, R. T., Kuhnel, A., Sun, Z. G., Kogel-Knabner, I., Hou, R. X., Cong, P. F., Liang, R. B., & Zhu, Ouyang. (2017). Large soil organic carbon increase due to improved agronomic management in the North China Plain from 1980s to 2010s. *Global Change Biology*, pp: 1–14.
- Hokazono, S., & Hayashi, K. (2012). Variability in environmental impacts during conversion from conventional to organic farming: A comparison among three rice production systems in Japan. *Journal of Cleaner Production*, 28: 101–112.
- Huang, Y., & Sun, W. J. (2006). Changes in topsoil organic carbon of croplands in mainland China over the last two decades, *Chinese Science Bullion*, 51, 1785–1803.
- Huang, Y., & Tang, Y. H. (2010). An estimate of greenhouse gas (N₂O and CO₂) mitigation potential under various scenarios of nitrogen use efficiency in Chinese croplands. *Global Change Biology*, 16, 2958–2970.
- Huang, T., Gao, B., Christie, P., & Ju, X. T. (2013). Net global warming potential and greenhouse gas intensity in a double cropping cereal rotation as affected by nitrogen and straw management.

645 *Biogeosciences*, 10, 1–15.

646 IPCC. (2000). Good practice guidance and uncertainty management in national greenhouse gas inventories.

647 Published : IGES, Japan.

648 Johnston, A. E., Poulton, P. R., Coleman, K. (2009). Soil organic matter: its importance in sustainable

649 agriculture and carbon dioxide fluxes. *Advances in Agronomy*, 101, 1–57.

650 Ju, X. T., Xing, G. X., Chen, X. P., Zhang, S. L., Zhang, L. J., Liu, X. J., Cui, Z. L., Yin, B., Christie, P.,

651 Zhu, Z. L., & Zhang, F. S. (2009). Reducing environmental risk by improving N management in

652 intensive Chinese agricultural systems. *Proceedings of the National Academy of Science USA*, 106,

653 3041–3046.

654 Ju, X. T., Gu, B. J., Wu, Y. J., & Galloway, J. N. (2016). Reducing China's fertilizer use by increasing

655 farm size. *Global Environmental Change*, 41, 26–32.

656 Lal, L. (2004). Soil carbon sequestration impacts on global climate change and food security. *Science*, 304,

657 1623.

658 Lal, L. (2007). Carbon management in agricultural soils. *Mitigation and Adaptation Strategies for Global*

659 *Change*, 12, 303–322.

660 Lemke, R. L., & Janzen, H. H. (2007). Implications for increasing the soil organic carbon store: calculating

661 the net greenhouse gas balance of no-till farming: in Greenhouse gas sinks. (eds. Reay, D.S., Hewitt,

662 C.N., Smith, K.A. and Garce, J.), Athenaeum Press Ltd, Gateshead, UK.

663 Liu, Y., Wang, E. L., Yang, X. G., & Wang, J. (2010). Contributions of climatic and crop varietal changes

664 to crop production in the North China Plain, since 1980s. *Global Change Biology*, 16, 2287–2299.

665 Ma, Y. C., Kong, X. W., Yang, B., Zhang, X. L., Yan, X. Y., Yang, J. C., & Xiong, Z. Q. (2013). Net global

666 warming potential and greenhouse gas intensity of annual rice–wheat rotations with integrated

667 soil–crop system management. *Agriculture, Ecosystems and Environment*, 164, 209–219.

668 Mahecha, M. D., Reichstein, M., Carvalhais, N., Lasslop, G., Lange, H., Senevirratne, S. I., Vargas, R.,

669 Ammann, C., Arain, M. A., Cescatti, A., Janssens, I. A., Migliavacca, M., Montagnani, L., &

670 Richardson, A. D. (2010). Global convergence in the temperature sensitivity of respiration at ecosystem

671 level. *Science*, 329, 838–840.

672 Meng, Q. F., Sun, Q. P., Chen, X. P., Cui, Z. L., Yue, S. C., Zhang, F. S., & Römheld, V. (2012). Alternative

673 cropping systems for sustainable water and nitrogen use in the North China Plain. *Agriculture,*

674 *Ecosystems and Environment*, 146, 93–102.

675 Minasny, B., Malone, B. P., McBratney, A. B., Angers, D. A., Arrouays, D., Chambers, A., Chaplot, V.,

676 Chen, Z. S., Cheng, K., Das, B. S., Field, D. J., Gimona, A., Hedley, C. B., Hong S. Y., Mandal, B.,

677 Marchant, B. P., Martin, M., McConkey, B. G., Mulder, V. L., O'Rourke, S., Richer-de-Forges, A. C.,

678 Odeh, I., Padarian, J., Paustian, K., Pan, G. X., Poggio, L., Savin, I., Stolbovoy, V., Stockmann, U.,

679 Sulaeman, Y., Tsui, C. C., Vågen, T. G., van Wesemael, B., & Winowiecki, L. (2017). Soil carbon 4 per

680 mille. *Geoderma*, 292, 59–86.

681 MOA (Ministry of Agriculture of China). (2014). China Agriculture Yearbook. China Agriculture Press,

- Beijing, China.
- Mosier, A. R., Halvorson, A. D., Reule, C. A., & Liu, X. J. (2006). Net global warming potential and greenhouse gas intensity in irrigated cropping systems in Northeastern Colorado. *Journal of Environmental Quality*, 35, 1584–1598.
- NBSC (National Bureau of Statistics of China). (2016). China Statistical Yearbook. China Statistics Press, Beijing, China.
- National Development and Reform Commission of China. (2014). The people's republic of China national greenhouse gas inventory of 2005. China Environmental Science Press, Beijing, China.
- Nayak, D., Saetnan, E., Cheng, K., Wang, W., Koslowski, F., Cheng, Y. F., Zhu, W. Y., Wang, J. K., Liu, J. X., Maron, D., Yan, X. Y., Cardenas, L., Newbold, J., Pan, G. X., Lu, Y. L., & Smith, P. (2015). Management opportunities to mitigate greenhouse gas emissions from Chinese agriculture. *Agriculture, Ecosystems and Environment*, 209, 108–124.
- Norse, D., & Ju, X. T. (2015). Environmental costs of China's food security. *Agriculture, Ecosystems and Environment*, 209, 5–14.
- Pan, G. X., Xu, X. W., Smith, P., Pan, W. N., & Lal, R. (2010). An increase in topsoil SOC stock of China's croplands between 1985 and 2006 revealed by soil monitoring. *Agriculture, Ecosystems and Environment*, 136, 133–138.
- Pathak, H., Li, C., & Wassmann, R. (2005). Greenhouse gas emissions from Indian rice fields: calibration and upscaling using the DNDC model. *Biogeosciences*, 2, 113–123.
- Poulton, P., Johnston, J., Macdonald, A., White R., & Powlson, D. S. (2018). Major limitations to achieving “4 per 1000” increases in soil organic carbon stock in temperate regions: Evidence from long-term experiments at Rothamsted Research, United Kingdom. *Global Change Biology*, 00, 1–22. <https://doi.org/10.1111/gcb.14066>.
- Powlson, D. S., Norse, D., & Lu, Y. L. (2018). Agricultural development in China: environmental impacts, sustainability issues and policy implications assessed through China-UK projects under SAIN (UK-China Sustainable Agriculture Innovation Network), 2008–2017, pp, 1–32. SAIN Working Paper no. 1. <http://www.sainonline.org/pages/News/SAIN%20Working%20Paper%20No%201.pdf>.
- Powlson, D. S., Stirling, C. M., Jat, M. L., Gerard, B. G., Palm, C. A., Sanchez, P. A. & Cassman, K. G. (2014). Limited potential of no-till agriculture for climate change mitigation. *Nature Climate Change*, 4, 678–683.
- Powlson, D. S., Whitmore, A. P., & Goulding, W. T. (2011). Soil carbon sequestration to mitigate climate change: a critical re-examination to identify the true and the false. *European Journal of Soil Science*, 62, 42–55.
- Pretty, J., & Bharucha, Z. P. (2015). Integrated pest management for sustainable intensification of agriculture in Asia and Africa. *Insects*, 6, 152–182.
- Qian, X., Gu, J., Pan, H. J., Zhang, K. Y., Sun, W., Wang, X. J., & Gao, H. (2015). Effects of living mulches on the soil nutrient contents, enzyme activities, and bacterial community diversities of apple

- orchard soils. *European Journal of Soil Biology*, 70, 23–30.
- Robertson, G. P., Paul, E. A., & Harwood, R. R. (2000). Greenhouse gases in intensive agriculture: contributions of individual gases to the radiative forcing of the atmosphere. *Science*, 289, 1922–1926.
- Robertson, G. P., & Grace, P. R. (2004). Greenhouse gas fluxes in tropical and temperate agriculture: the need for a full-cost accounting of global warming potentials. *Environment Development Sustainability*, 6, 51–63.
- Rockstrom, J., Gaffney, O., Rogelj, J., Meinshausen, M., Nakicenovic, N., & Schellnhuber, H. J. (2017). A roadmap for rapid decarbonization. *Science*, 355, 1269.
- Schandl, H., Hatfield-Dodds, S., Wiedmann, T., Geschke, A., Cai, Y., West, J., Newth, D., Baynes, T., Lenzen, M., & Owen, A. (2016). Decoupling global environmental pressure and economic growth: scenarios for energy use, materials use and carbon emissions. *Journal of Cleaner Production*, 132, 45–56.
- Schlesinger, W. H. (2010). On fertilizer-induced soil carbon sequestration in China's croplands. *Global Change Biology*, 16, 849–850.
- Shang, Q. Y., Yang, Q. X., Gao, C. M., Wu, P. P., Liu, J. J., Xu, Y. C., Shen, Q. R., Zou, J. W., & Guo, S. W. (2011). Net annual global warming potential and greenhouse gas intensity in Chinese double rice-cropping systems: a 3-year field measurement in long-term fertilizer experiments. *Global Change Biology*, 17, 2196–2210.
- Smith, P., Lanigan, G., Kutsch, W. L., Buchmann, N., Eugster, W., Aubinet, M., Ceschia, E., Béziat, P., Yeluripati, J. B., Osborne, B., Moors, E. J., Brut, A., Wattenbach, M., Saunders, M., & Jones, M. (2010). Measurements necessary for assessing the net ecosystem carbon budget of croplands. *Agriculture, Ecosystems & Environment*, 139, 302–315.
- Smith, P., Martino, D., Cai, Z. C., Gwary, D., Janzen, H., Kumar, P., McCarl, B., Ogle, S., O'Mara, F., Rice, C., Scholes, B., & Sirotenko, O. (2007). Agriculture. In: Metz, B., Davidson, O.R., Bosch, P.R., Dave, R., Meyer, L.A. (Eds.), Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Mitigation of Climate Change 2007. Cambridge University Press, Cambridge/New York.
- Smith, P., Martino, D., Janzen, H., Kumar, P., McCarl, B., Ogle, S., O'Mara, F., Rice, C., Scholes, B., Sirotenko, O., Howden, M., McAllister, T., Pan, G. X., Romanenkov, V., Schneider, U., Towprayoon, S., Wattenbach, M., & Smith, J. (2008). Greenhouse gas mitigation in agriculture. *Philosophical Transactions of the Royal Society B*, 363, 789–813.
- Snyder, C. S., Bruulsema, T. W., Jensen, T. L., & Fixen, P. E. (2009). Review of greenhouse gas emissions from crop production systems and fertilizer management effects. *Agriculture, Ecosystems and Environment*, 133, 247–266.
- Tang, X. L., Zhao, X., Bai, Y. F., Tang, Z. C., Wang, W. T., Zhao, Y. C., Wang, H. W., Xie, Z. Q., Shi, X. Z., Wu, B. F., Wang, G. X., Yan, J. H., Ma, K. P., Du, S., Li, S. G., Han, S. J., Ma, Y. X., Hu, H. F., He, N. P., Yan, Y. H., Han, W. X., Hen, L. L., Yu, G. R., Fang, J. Y., & Zhou, G. Y. 2018. Carbon pools in

- China's terrestrial ecosystems: New estimates based on an intensive field survey. *Proceedings of the National Academy of Science USA*, 115(16): 4021–4026.
- Tian, H. Q., Lu, C. Q., Melillo, J., Ren, W., Huang, Y., Xu, X. F., Liu, M. L., Zhang, C., Chen, G. S., Pan, S. F., Li, J. Y., & Reilly, J. (2012). Food benefit and climate warming potential of nitrogen fertilizer uses in China. *Environment Research Letters*, 7, 044020 (8pp)
- van Groenigen, J. W., Velthof, G. L., Oenema, O., Van Groenigen, K. J., & Van Kessel, C. (2010). Towards an agronomic assessment of N₂O emissions: a case study for arable crops. *European Journal of Soil Science*, 61, 903–913.
- van Groenigen, J. W., van Kessel, C., Hungate, B. A., Oenema, O., Powlson, D. S., & Van Groenigen, K. J. (2017). Sequestering soil organic carbon: a nitrogen dilemma. *Environmental Science & Technology*, 51, 4738–4739.
- Wang, J. X., Rothausen, S. G., Conway, D., Zhang, L. J., Xiong, W., Holman, I. P., & Li, Y. M. (2012). China's water–energy nexus: greenhouse-gas emissions from groundwater use for agriculture. *Environmental Research Letters*, 7, 014035.
- Wang, S. C., Zhao, Y. W., Wang, J. Z., Zhu, P., Cui, X., Han, X. Z., Xu, M. G., & Lu, C. L. (2017). The efficiency of long-term straw return to sequester organic carbon in Northeast China's cropland. *Journal of Integrative Agriculture*, 16(0), 60345–60347.
- Wang, X., & Wang, J. (2016). Facilities vegetables chemical pesticide use decrement measures. *Agricultural Technology & Equipment*, 323, 45–46 (in Chinese with English abstract).
- Wassmann, R., Neue, H. U., Buendia, L. V., Corton, T. M., & Lu, Y. (2000). Characterization of methane emissions from rice fields in Asia. III. Mitigation options and future research needs. *Nutrient Cycling in Agroecosystems*, 58, 23–36.
- West, T. O., & Marland, G. A. (2002). Synthesis of carbon sequestration carbon emissions, and net carbon flux in agriculture: comparing tillage practices in the United States. *Agriculture, Ecosystems and Environment*, 91, 217–232.
- West, T. O., & Post, W. M. (2002). Soil organic carbon sequestration rates by tillage and crop rotation. *Soil Science Society of America Journal*, 66(6), 1930–1946.
- Wollenberg, E., Richards, M., Smith, P., Havlik, P., Obersteiner, M., Tubiello, F. N., Herold, M., Gerber, P., Carter, S., Reisinger, A., Van Vuuren, D., Dickie, A., Neufeldt, H., Sander, B. O., Wassmann, R., Sommer, R., Amonette, J., Falcucci, A., Herrero, M., Opio, C., Roman-Cuesta, R. M., Stehfest, E., Westhoek, H., Ortiz-Monasterio, I., Sapkota, L., Rufino, M. C., Thornton, P. K., Verchot, L., West, P. C., Soussana, J. F., Baedeker, T., Sadler, M., Vermeulen, S., & Campbell, B. M. (2016). Reducing emissions from agriculture to meet the 2 °C target. *Global Change Biology*, 22, 3859–3864.
- Wu, B. F., Yuan, Q. Z., Yan, C. Z., Wang, Z. M., Yu, X. F., Li, A. N., Ma, R. H., Huang, J. L., Chen, J. S., Chang, C., Liu, C. L., Zhang, L., Li, X. S., Zeng, Y., & Bao, A. M. (2014). Land Cover Changes of China from 2000 to 2010. *Quaternary Sciences*, 34(4), 723–731 (in Chinese with English abstract).
- Xia, L. L., Lam, S. K., Chen, D. L., Wang, J. Y., Tang, Q., & Yan, X. Y. (2016). Can knowledge-based N

- management produce more staple grain with lower greenhouse gas emission and reactive nitrogen pollution? A meta-analysis. *Global Change Biology*, 23, 1917–1925.
- Xia, L. L., Lam, S. K., Yan, X. Y., & Chen, D. L. (2017). How does recycling of livestock manure in agro-ecosystems affect crop productivity, reactive nitrogen losses, and soil carbon balance? *Environmental Science & Technology*, 51(13), 7450–7457.
- Yan, X. Y., Cai, Z. C., Wang, S. W., & Smith, P. (2011). Direct measurement of soil organic carbon content change in the croplands of China. *Global Change Biology*, 17, 1487–1496.
- Yan, Y. F., & Yang, L. K. (2010). China's foreign trade and climate change: A case study of CO₂ emissions. *Energy Policy*, 38, 350–356.
- Yan, M., Cheng, K., Yue, Q., Yan, Y., Rees, R. M., & Pan, G. X. (2016). Farm and product carbon footprints of China's fruit production—life cycle inventory of representative orchards of five major fruits. *Environmental Science and Pollution Research*, 23, 4681–4691.
- Yang, F., Xu, Y., Cui, Y., Meng, Y. D., Dong, Y., Li, R., & Ma, Y. B. (2017). Variation of soil organic matter content in croplands of China over the last three decades. *Soils*, 54(5), 1047–1055 (in Chinese with English Abstract).
- Zhang, D., & Zhang, W. F. (2016). Low carbon agriculture and a review of calculation methods for crop production carbon footprint accounting. *Resources Science*, 7, 1395–1405 (in Chinese with English Abstract).
- Zhang, W. F., Dou, Z. X., He, P., Ju, X. T., Powlson, D. S., Chadwick, D., Norse, D., Lu, Y. L., Zhang, Y., Wu, L., Chen, X. P., Cassman, K. G., & Zhang, F. S. (2013). New technologies reduce greenhouse gas emissions from nitrogenous fertilizer in China. *Proceedings of the National Academy of Science USA*, 110, 8375–8380.
- Zhao, X., Wang, X. W., Wang, S. Q., & Xing, G. X. (2014). Successive straw biochar application as a strategy to sequester carbon and improve fertility: A pot experiment with two rice-wheat rotations in paddy soil. *Plant Soil*, 378, 279–294.
- Zhao, Y. C., Xu, S. X., Wang, M. Y., & Shi, X. Z. (2018). Carbon sequestration potential in Chinese cropland soils: review, challenge, and research suggestions. *Bulletin of Chinese Academy of Sciences*, 33(2), 191–197 (in Chinese with English Abstract).
- Zhou, M. H., Zhu, B., Wang, S. J., Zhu, X. Y., Vereecken, H., & Brüggemann, N. (2017). Stimulation of N₂O emission by manure application to agricultural soils may largely offset carbon benefits: a global meta-analysis. *Global Change Biology*, 23(10), 4068–4083.
- Zou, J. W., Huang, Y., Jiang, J. Y., Zheng, X. H., & Sass, R. L. (2005). A 3-year field measurement of methane and nitrous oxide emissions from rice paddies in China: effects of water regime, crop residue, and fertilizer application. *Global Biogeochemical Cycles*, 19, GB2021, doi: 10.1029/200GB002401.

827 **Tables**

828 Table 1 Abbreviations for different Chinese cropping systems in this paper.

Abbreviation	Cropping systems
WMN	winter wheat – summer maize double-cropping in Northern China
WMSW	winter wheat – summer maize double-cropping in Southwestern China
RW	rice – winter wheat double-cropping in Central and Eastern China
RR	rice – rapeseed double-cropping in Central and Southwestern China
DR	double rice per year in Central and Southern China
SRNE	single rice in Northeastern China
SRENE	single rice across China except for Northeastern China
SB	soybeans in Northeastern China
MNE	single spring maize per year in Northeastern China
MNW	single spring maize in Northern and Northwestern China
PS	potato system across China
CS	cotton system in Northern and Northwestern China
GV	greenhouse vegetables across China
OV	open field vegetables across China
OS	orchard system across China

829 Table 2 Emission factors of soil GHG emission and agriculture inputs

Emission source	Abbreviation	Unit	Emission factor (kg C-eq unit ⁻¹)	Literature
N ₂ O	CE _{N2O}	kg	81.3	Cubasch et al., 2013
CH ₄	CE _{CH4}	kg	9.3	Cubasch et al., 2013
Fertilizer	CE _N	kg	2.3	Zhang et al., 2013
	CE _P	kg (P ₂ O ₅)	0.4	Huang et al., 2013
	CE _K	kg (K ₂ O)	0.3	Huang et al., 2013
Power for irrigation	CF _I	kwh	0.4	Zhang et al., 2013
Fuel in farm operations	CF _F	L	0.7	Cheng et al., 2011
Pesticides	CF _{PE}	kg	4.9	West and Marland, 2002
Plastic film	CF _{PF}	kg	5.2	Cheng et al., 2011
Paper bags	CF _{PB}	kg	0.3	Yan et al., 2016

830 Table 3. The total GHG balance and sources of GHGs in different Chinese cropping systems.

Cropping systems	N ₂ O	CH ₄	N	P ₂ O ₅	K ₂ O	Irrigation	Fuel	Pesticide	Plastic film	SOC change	Net GHG balance*	Yield [#]	GHGI [†]	Plant area [‡]	Total SOC change [§]	Total GHGs emission [¶]	Total GHG emission/total SOC change [§]
	kg C-eq ha ⁻¹ yr ⁻¹									kg C ha ⁻¹ yr ⁻¹	kg C-eq ha ⁻¹ yr ⁻¹	t ha ⁻¹	kg C-eq Mg ⁻¹	×10 ⁴ ha	Tg C yr ⁻¹	Tg C-eq yr ⁻¹	
WMN	439±33	-17±3	1148±45	59±6	35±5	606±38	121±12	35±5	-	326±34	2093±294	14.3±0.4	146±28	1371	4.5±2.5	33.2±5.0	7.4±4.3
WMSW	496±122	-34±59	758±32	65±4	32±3	0±0	136±47	20±5	-	223±101	1312±362	9.8±0.8	134±49	205	0.5±0.7	3.1±0.8	6.9±10.1
RW	780±81	1564±23	978±29	49±4	39±4	758±107	121±27	44±10	-	363±60	3970±856	13.3±0.6	299±82	475	1.7±1.6	20.6±4.7	11.9±11.3
RR	712±81	1561±254	758±45	63±5	40±5	824±106	102±39	66±15	-	400±72	3725±776	10.0±0.4	289±71	532	2.1±1.6	22.0±4.9	10.3±7.9
DR	190±33	3935±397	616±36	47±3	52±4	780±56	103±19	51±10	-	432±38	5342±1566	12.9±0.3	535±162	564	2.4±1.9	32.6±10.2	13.4±11.0
SRNE	98±16	1091±228	333±32	18±5	19±3	855±105	67±5	15±5	-	366±74	2154±713	8.5±0.3	253±93	431	1.6±1.3	10.9±4.0	6.9±6.1
SRENE	366±81	2300±262	453±57	43±6	22±6	731±4	74±16	25±5	-	416±45	3596±817	7.7±0.2	466±122	463	1.9±1.2	18.6±4.6	9.7±6.8
SB	86±33	-8±8	68±9	26±4	6±1	0±0	76±11	10±5	-	-56±120	328±371	2.6±0.1	129±246	356	-0.2±1.3	1.0±0.7	/
MNE	211±33	-8±3	457±27	26±4	17±2	30±14	67±13	20±5	2±2	183±47	642±352	9.8±0.5	65±27	1637	3.0±5.2	13.5±3.1	4.5±7.4
MNW	382±154	-25±8	638±59	29±4	9±3	351±103	75±13	25±5	393±108	273±80	1621±824	10.3±0.7	155±70	90	0.2±0.3	1.7±0.7	6.9±9.0
GV	2064±260	-17±8	2214±226	149±24	112±15	1178±140	125±39	370±44	1719±182	342±73	7567±1738	14.5±1.2	523±222	108	0.4±0.4	8.5±2.1	23.1±24.2
OV	1699±236	17±17	1754±161	71±18	47±11	776±147	21±21	104±10	409±32	296±118	4617±1308	11.7±1.7	394±204	719	2.1±2.9	35.3±10.3	16.5±23.5
PS	106±24	-8±2	407±20	48±3	40±3	129±24	24±9	59±10	182±44	110±85	889±391	5.3±0.3	170±77	875	1.0±2.3	8.9±2.4	9.2±22.3
CS	263±60	-15±15	559±68	52±11	9±3	462±66	25±11	89±10	178±42	276±50	1349±472	3.4±0.4	399±250	485	1.4±1.2	7.9±2.3	6.0±5.6
OS	319±114	-23±5	1060±106	118±24	84±17	355±109	55±19	101±39	29±10	50±51	2047±802	3.2±0.3	648±371	1237	0.6±3.3	25.9±9.9	41.6±223.5
Mean/Total										243±92 ^{††}					23.2±8.6	269.5±21.1	11.6±4.4

831 * Net GHG balance calculated by equation 2, values are means and uncertainty ranges.

832 [#] Yields of vegetable and fruit = Fresh yields in the literature × 0.1.

833 [†] Plant areas cited and calculated from China Agriculture Yearbook (MOA, 2014) and Department of Agricultural Machinery Management (DAMM, 2017).

834 [‡] Total SOC change (Tg C yr⁻¹) = SOC change (kg C ha⁻¹ yr⁻¹) × Plant area (× 10⁴ ha)/10⁹.

835 [¶] Total GHGs emissions (Tg C-eq yr⁻¹) = C-eq from N₂O + CH₄ + N + P₂O₅ + K₂O + Irrigation + Fuel + Pesticide + Plastic (kg C-eq ha⁻¹) × Plant area (× 10⁴ ha)/10⁹.

837 [§] Times of total GHG emissions to total SOC change = Total GHG emissions/Total SOC change.

838 ^{††} The weighted mean increase rate of SOC for Chinese main cropping systems.

Figure captions

Fig. 1. Spatial pattern of cropping systems in China. Abbreviations are shown in Table 1, GV+OV represents total vegetable production. These abbreviations are also used in subsequent figures.

Fig. 2. The locations and mean changes of SOC density in Chinese cropping systems from 2000 to 2017.

Fig. 3. The frequency distribution of annual SOC changes in Chinese cropping systems.

Fig. 4. Changes of SOC in Chinese cropping systems. CC represents Chinese total croplands. Different letters in parentheses represent the changes of SOC under methods I, II and III, respectively. Box-and-whisker diagrams show the median, 5th, 25th, 75th and 95th percentiles for relative change in SOC stocks.

Fig. 5. Spatial patterns of the SOC stock change in different cropping systems.

Fig. 6. Spatial pattern of net GHG balance in different cropping systems.

Fig. 7. Sources and allocation of greenhouse gas emissions in different cropping systems in China.

gcb_14425_f2.tif

gcb_14425_f5.tif

gcb_14425_f6.tif

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Gao, B;Huang, T;Ju, X;Gu, B;Huang, W;Xu, L;Rees, RM;Powlson, DS;Smith, P;Cui, S

Title:

Chinese cropping systems are a net source of greenhouse gases despite soil carbon sequestration

Date:

2018-12-01

Citation:

Gao, B., Huang, T., Ju, X., Gu, B., Huang, W., Xu, L., Rees, R. M., Powlson, D. S., Smith, P. & Cui, S. (2018). Chinese cropping systems are a net source of greenhouse gases despite soil carbon sequestration. GLOBAL CHANGE BIOLOGY, 24 (12), pp.5590-5606. <https://doi.org/10.1111/gcb.14425>.

Persistent Link:

<http://hdl.handle.net/11343/284472>