Title Page

Title: Subungual Bowen's disease. Complete remission 14 years post PDT

Short running title: Subungual Bowen's disease

Authors:

1. Dr Louise Photiou

- Sinclair Dermatology, East Melbourne, VIC, Australia
- The Victorian Melanoma Service, The Alfred Hospital, Melbourne, VIC, Australia

2. Prof Rodney Sinclair

- Sinclair Dermatology, East Melbourne, VIC, Australia
- The University of Melbourne, Melbourne, VIC, Australia

None of the authors have any conflict of interest or affiliations to declare.

Corresponding author:

Dr Louise Photiou

48 Perth Street

Prahran, VIC 3181

louise.photiou@gmail.com

0401470403

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as [doi: 10.1111/ajd.12901](https://doi.org/10.1111/ajd.12901)

This article is protected by copyright. All rights reserved

Article type : Correspondence Letter

Letter to the Editor

Subungual Bowen's disease. Complete remission 14 years post PDT.

Dear Sir,

In 2004 we reported the first case of biopsy-proven subungual Bowen's disease, successfully treated with photodynamic therapy (Tan, Sinclair, & Foley, 2004) . We now report long-term follow-up of this gentleman who remains in complete remission 14 years later.

The patient was a police shooting instructor and the Bowen's disease affected his trigger finger. Standard treatment in 2004 was distal amputation of the digit. In view of his occupation he was keen to explore non-surgical alternatives.

After partial nail avulsion and a course of a keratolytic pretreatment (20% salicylic acid in white soft paraffin), the area was photosensitized with 20% 5-amino-levulinic acid (5-ALA) oil in water emulsion applied under occlusion for four hours. Omnilux 630nm was used as the light source, delivering a dose of 63 J/cm². Three treatments were given, two weeks apart and then a fourth treatment was given four weeks later. The patient achieved complete clinical remission and repeat biopsy at 6 months post-treatment confirmed histological remission. He was followed up for 2 years and then discharged from the clinic.

The patient re-presented to our clinic fourteen years later for an unrelated matter. Examination of the trigger finger, (figure 1), showed no evidence of recurrence of the Bowen's disease. He has continued to work in the police force as a shooting instructor.

A second patient with biopsy proven subungual Bowen's disease of the thumb, also treated successfully with PDT, also remains in complete remission 16 years later (figure 2).

PDT, at least in these 2 patients, offers a potentially curative, non-surgical alternative to digital amputation in the management of subungual Bowen's disease.

This article is protected by copyright. All rights reserved

Figure Legend:

Figure 1 - Patient 1 (A) Prior to PDT treatment of the right index finger, (B) Prior to PDT treatment, (C) 6 months post-PDT, (D) 14 years post-PDT

Figure 2 - Patient 2, 16 years post-PDT treatment of the right thumb

Bibliography

ajd_12901_f1.jpg

ajd_12901_f2.jpg

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Photiou, L;Sinclair, R

Title:

Subungual Bowen disease. Complete remission 14 years post-photodynamic therapy (PDT)

Date:

2019-02-01

Citation:

Photiou, L. & Sinclair, R. (2019). Subungual Bowen disease. Complete remission 14 years post-photodynamic therapy (PDT). AUSTRALASIAN JOURNAL OF DERMATOLOGY, 60 (1), pp.74-75. <https://doi.org/10.1111/ajd.12901>.

Persistent Link:

<http://hdl.handle.net/11343/284998>