Systematic review of organisation-wide, trauma-informed care models in out-of-home care (OoHC) settings

Cate Bailey¹ PhD, Anna Klas² PhD, Rachael Cox² PhD, Heidi Bergmeier² PhD, Julie Avery MSocSci 1, Helen Skouteris (corresponding author)¹ PhD

¹ Monash Centre for Health Research and Implementation, School of Public Health and Preventive Medicine, Monash University

² School of Psychology, Deakin University, Geelong, Australia;

Corresponding author: Professor Helen Skouteris, Monash Centre for Health Research and Implementation, School of Public Health and Preventive Medicine, Monash University, Level 1, 43-52 Kanooka Grove Clayton; Locked Bag 29 Clayton Vic 3168

Email: helen.skouteris@monash.edu

Phone: +61 414 409 563

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/hsc.12621](https://doi.org/10.1111/hsc.12621)

DR. CATE BAILEY (Orcid ID : 0000-0001-5030-430X)

Article type : Review Article

Systematic review of organisation-wide, trauma-informed care models in out-of-home care (OoHC) settings

Abstract

Trauma in early childhood has been shown to adversely affect children's social, emotional and physical development. Children living in out-of-home care (OoHC) have better outcomes when care providers are present for children, physically, psychologically and emotionally. Unfortunately, the high turn-over of out-of-home carers, due to vicarious trauma (frequently resulting in burnout and exhaustion) can result in a child's trauma being re-enacted during their placement in OoHC. Organisation-wide (encompassing the whole organisation, from the CEO to all workers including administration staff) therapeutic care models that are trauma-informed and have been developed to respond to the complex issues of abuse and neglect that children enter care having experienced. These models incorporate a range of therapeutic techniques, and provide an overarching approach and common language that is employed across all levels of the organisation. The aim of this study was to investigate the current empirical evidence for organisation-wide trauma-informed therapeutic care models in OoHC. A systematic review searching leading databases was conducted for evidence of organization-wide, trauma-informed, out-of-home care studies, between 2002 and 2017. Seven articles were identified covering three organizational models. Three of the articles assessed the Attachment Regulation and Competency framework (ARC), one study assessed the Children

and Residential Experiences program (CARE), and three studies assessed The Sanctuary Model. Risk of bias was high in 6 of the 7 studies. Only limited information was provided on the effectiveness of the models identified through this systematic review, although the evidence did suggest that trauma-informed care models may have significantly positive outcomes for children in OoHC. Future research should focus on evaluating components of trauma-informed care models and assessing the efficacy of the various organisational care models currently available. (265/300)

Six keywords: OoHC; looked after children; trauma-informed care; organisation-wide; abuse; neglect

What is known about this topic

- Children living in OoHC have frequently experienced complex relational trauma
- When living in OoHC, children have better outcomes when care providers are present for children, physically, psychologically and emotionally.
- Organisation-wide, trauma-informed care models have been proposed to help ameliorate the effects of trauma on children and carers. Although these models are promising, there is a lack of evidence regarding the efficacy of these models.

What this paper adds

- Using implementation science criteria, empirical evidence on organisation-wide trauma informed care models was found to be weak, with high risk of bias.
- The initial evidence presented suggests that the application of trauma-informed care models may have significantly positive outcomes for children in OoHC.
- Future research is urgently required to provide empirical evidence for organisation-wide trauma informed care models.

Previous research has consistently demonstrated that trauma can adversely affect children's social, emotional, physical and neurological development, especially when trauma is experienced during the critical period of early childhood (Siegel, 2007; van der Kolk, 2007). Traumatic life experiences can be detrimental for children's health and social outcomes, behaviour and mental health (Bloom, 2016), and can profoundly impact on the child's well-being across their life-course, (Hummer, Dollard, Robst, & Armstrong, 2010). Trauma occurs when an individual experiences an intense event that harms or threatens harm to their physical or emotional well-being or to someone close to them, e.g., a family member or friend (van der Kolk, 2007). Trauma has commonly been categorised as either a single event trauma (a life threatening event with potential to cause harm or injury) or complex trauma (interpersonal danger, violence or abuse, usually over multiple incidences of extended duration) (van der Kolk, 2003). As early development is embedded in the context of the caregiving relationship, trauma also has a strong influence on the caregiving system, which particularly affects children living outside of the family home (Arvidson et al., 2011).

In 2013-14, around 51,539 Australian children lived in Out of Home Care (OoHC), a rate of 9.8 children per 1,000 (AIHW, 2017). OoHC is a broad term that may refer to a formal or informal arrangement, and long- or short-term care (AIHW, 2017). OoHC is generally defined in Australia and the US as: (1) foster care; (2) relative or kinship care; (3) family group homes; (4) residential care; and (5) independent living (Barth, Greeson, Zlotnik, & Chintapalli, 2011; Department of Families, Housing, Community Services and Indigenous Affairs, 2011). Typically, children and young people are moved into OoHC due to physical, sexual or emotional abuse, neglect, or domestic violence (Department of Families, Housing, Community Services and Indigenous Affairs, 2011). These children are therefore likely to have experienced high rates of complex trauma and adversity (Fraser et al., 2014). Failure to understand and address issues arising for children with a traumatic history can inadvertently aggravate symptoms, and traumatise the child further (Murphy, Moore, Redd, & Malm, 2017). Figure 1, displays the process of a child being placed in OoHC.

[insert Figure 1 here]

Figure 1. Process of child moving into OoHC

Research suggests that survivors of trauma experienced during childhood may be more resilient if they experience positive and considerate care providers (Esaki et al., 2013). However, carers can be at high risk of experiencing vicarious trauma from caring for children in OoHC (Victorian Auditor-General, 2014) (vicarious trauma is defined here as the effect on a direct carer from working with clients' traumatic experiences (AIFS, 2007)). Difficulties arising from caring for traumatised children can create an environment where a child's trauma is reconstructed during OoHC stays, making it harder for caregivers to build meaningful relationships with children (Esaki et al., 2013; Farragher & Yanosy, 2005). Unfortunately, and despite best intentions, there is often a high turnover of out-of-home carers due to burnout and exhaustion, which can directly affect the child's experience of care giving (Middleton & Potter, 2015).

Each person in an organisation, staff member, carer or client, has their own adaptation to the stress, trauma and adversity they have experienced in their lives (Bloom, 2016). Because of the inter-relationship between each level in an organisation, the adaptation to stress that each person brings frequently results in increased levels of stress across an organisation. Under stress, an organisation may respond with a crisis intervention rather than looking more closely at prevention activities. This process is shown in figure 2.

[Insert Figure 2 here]

Figure 2. The impact of trauma on organisational response

Protective services and community agencies worldwide have been challenged in how best to address the needs of children who have been traumatised by abuse, neglect, violence and loss (Macdonald & Millen, 2012). Typically, a therapeutic care approach is recommended, which incorporates multiple types of interventions, often stemming from a variety of therapeutic techniques or frameworks presented and employed in varying ways. Although definitions vary, therapeutic care in Australia is defined as intensive, time-limited care that responds to complex issues of abuse and neglect, as well as separation from family, and is provided to a child or young adult who is in statutory care (McLean, Price-Robertson, & Robinson, 2011).

To better integrate services provided to children in OoHC, it has been proposed that system-wide changes to organisations can be made so that they are trauma-informed at an organisational level. This approach aims to provide an overarching strategy and a common language that is employed across all levels of the organisation, including staff, carers,

children and young people (Wall, Higgins, & Hunter, 2016). Trauma-informed organisations are built on a whole of system recognition of the impact of trauma, and the development of an understanding of trauma in the workforce (including carers) (Murphy et al., 2017). The overall aim of this approach, is to reduce the experience of trauma (including previous, ongoing and vicarious trauma) for staff, carers, and, most importantly, children and young people (Bloom, 2016).

Whilst there are many benefits to employing a trauma-informed approach in OoHC, there had been a lack of definition, and a large degree of flexibility in how trauma-informed approaches are constructed (Wall et al., 2016). In an attempt to define what is meant by trauma-informed care, Hanson and Lang (2016) identified components that target three main domains: 1) 'workforce development', which included workforce awareness, training and understanding of secondary traumatic stress; 2) 'trauma-focused services', encompassing evidence-based practices and standardized screening; and 3) 'organizational environment and practices', including safe physical environments, staff collaboration, defined policies, and leadership, as represented in Figure 3.

[Insert Figure 3 here]

Figure 3. Factors in an organisation-wide trauma-informed model

There has been increased interest in organisation-wide (encompassing the whole organisation, from the CEO to all workers including administration staff) trauma-informed care models over the last decade (Becker-Blease, 2017). Nonetheless, the implementation of these practices is still preliminary (Beyerlein & Bloch, 2014), and there has been a lack of empirical research on these models (Hanson & Lang, 2016). Systems change in organisational culture and communities is complex, requiring effort from multiple stakeholders, and it can be difficult to attribute change from specific interventions within organisations or systems, and how the organisation as a whole may have transformed (Esaki et al., 2013). Continued organisational restructuring takes time, and can result in 'change fatigue' (Murphy et al., 2017). Despite these difficulties, assessing trauma-informed care models is essential, as these models are being widely adopted in OoHC settings (Bloom, 2017).

The overall aim of the current study was to examine the empirical evidence available for organisation-wide, trauma-informed care models. In order to address this aim we systematically evaluated the evidence for organisation-wide, trauma-informed care models in OoHC. Whilst the papers that have been included in this review are important contributions to the literature in describing trauma-informed organisational change initiatives and their outcomes, this review focuses specifically on the strength of the research designs and the empirical evidence that can be derived.

Method

Information sources

Papers were sourced through a search of eight databases: Scopus, Academic Search Complete, CINAHL with full text, Medline complete, Psych Articles, Psych Info, Social Work abstracts and EMBASE for the time-period between April 31st 2002 and April 31st 2017. The search was organised around subheadings that were generated from the research aim by the first author (CB), and reviewed and endorsed by the senior author (HS). Search terms are shown in Box 1. A total of 176 articles were obtained from the database search (after the removal of 222 duplicates). Seven additional articles were obtained from other sources (google scholar, searching reference lists). A total of 183 abstracts were reviewed for suitability by the first author.

INSERT BOX 1 HERE

Eligibility criteria and study selection

Articles were included if they were peer-reviewed, published in English, and referred to children aged from birth to adolescence (0-17 years). The population was children living in OoHC (as defined in the introduction). Studies needed to evaluate system- or organisation-wide programs, and to present empirical evidence. Studies were excluded if they assessed trauma-informed care programs not implemented at the organisational level. Titles and abstracts were checked against the inclusion criteria, and if relevant, the full text version was sourced. The first two authors read and discussed the papers until all the selected papers were mutually agreed upon.

Risk of bias assessment

Papers that met inclusion criteria were assessed for the quality of methodology using the criteria outlined by Thomas et al. (2004). Using this criteria, the assessment of the methodological quality was rated as strong, moderate or weak, for six components: selection bias, design, confounders, blinding, data collection methods, withdrawals, and dropouts. After evaluation on each of the six components, each paper was rated on the combined results. Papers with no weak ratings and four or more strong ratings were considered strong; papers with less than four strong ratings, and one or less weak rating were considered moderate, whilst papers with two or more weak ratings were considered weak. Papers that were rated as either moderate or strong were then assessed on two further components: integrity of the intervention (percentage of study participants receiving the intervention as designed), and utilising an appropriate statistical analysis methodology (including intention to treat analysis).

Data extraction

Thomas and colleagues (2004) recommended only analysing papers rated as either moderate or high in the risk of bias assessment. In the current study, only one study met the risk of bias criteria. However, in order to describe and evaluate the three models covered by the selected papers, information from all papers that met the inclusion criteria have been included in the current review. Data were extracted using the standardized format as per Thomas et al. (2004) for the following variables: funding source; number of participants by group and number of dropouts; descriptions of the target population, intervention and study outcome; plus the length of follow-up. An extra field was added to include methodology in this table. The extracted information is presented in Table 1. Papers were synthesised narratively; meta-analysis was not used as outcome measures in the papers were not comparable.

Results

Of the initial 183 abstracts sourced, evaluation of titles and abstracts revealed 37 papers which may have met the inclusion criteria, of which seven articles were retained after assessing the full text, as shown in Figure 4. Supplementary Table S1 provides details of the excluded studies, with reasons for exclusion. All seven papers included in this review were

from the United States, and all presented empirical data from OoHC populations from one of three organisation-wide trauma-informed models. The three models were: 1) Attachment Regulation and Competency framework (ARC); three papers (Arvidson et al., 2011; Hodgdon, Blaustein, Kinniburgh, Peterson, & Spinazzola, 2016; Hodgdon, Kinniburgh, Gabowitz, Blaustein, & Spinazzola, 2013); 2) the Children and Residential Experiences program (CARE); one paper: (Izzo et al., 2016); and 3) The Sanctuary Model; three papers (Bloom et al., 2003; Kramer, 2016; Rivard, Bloom, McCorkle, & Abramovitz, 2005).

Risk of bias

Only one of the seven papers was rated as moderate on the risk of bias assessment (Izzo et al., 2016), with all other studies rated as weak. Five of the seven papers we assessed had weak study designs (Arvidson et al., 2011; Bloom et al., 2003; Hodgdon et al., 2016, 2013; Kramer, 2016), and only one study accounted for confounding variables (Izzo et al., 2016). No studies mentioned blinding of participants or researchers, and all except one study (Izzo et al., 2016) were rated as weak on withdrawals/dropouts of participants from studies. Data collection methods were rated between weak and moderate (please contact the authors for further information on the risk of bias assessment.).

[Insert Figure 4 here]

Figure 4. PRISMA diagram of the systematic search

Table 1

Data extraction and synthesis

[Insert Table 1 here]

Attachment, Self-Regulation and Competency (ARC) Model. The Attachment, Self-regulation and Competency Model aims to provide a theoretically-driven and flexible framework for organisations that provide services for traumatised children (Hodgdon et al., 2016). Included in the organisation-wide approach are three levels, the individual child, the child's family or caregivers, and the organisational culture. Within this framework there are opportunities to incorporate various interventions for use by practitioners, which may be individual, family, or system based (Kinniburgh, 2005). The ARC model provides guidelines on building therapeutic culture for the child within organisations, so that the child's ability to heal from trauma may be enhanced. The model focusses on: 1) building healthy attachments between children and their caregivers, particularly family members; 2) supporting children to develop skills to manage their emotions and physiological states, and thus increasing the child's self-regulation; 3) building the child's competency, by increasing their capacity and skills; and 4) working with children to integrate experiences of trauma, thereby increasing their self-understanding (Kinniburgh, 2005). These aspects are perceived in an organisational context, as this model highlights the importance of system-wide changes to support effective outcomes for children, as well as recognising that children need individually tailored programs to support their needs.

Three of the seven studies included in the current review evaluated the ARC model (Arvidson et al., 2011; Hodgdon et al., 2016, 2013). In the first of these papers, Arvidson and colleagues used a naturalistic study design to measure preliminary evidence of the effectiveness of the ARC model on children in the Alaskan child protection system. The intervention in this study used the ARC model to help children who had been exposed to trauma to have the opportunity to process their experiences in an appropriate way for their emotional and cognitive development. The ARC framework was applied in three ways: 1) increasing the ability of the caregiver to be attuned with the child so as to build secure attachment, 2) developing the caregiving system to support the child's self-regulation systems, and 3) caregivers supporting the child's development of a positive sense of self and ability to master tasks.

Of the 93 children who received the intervention, only 54% were deemed to have received the ARC model by the date of the article's submission (Arvidson et al., 2011). Of these 50 children, only 26 had completed treatment at discharge due to relocation of the family (26%), drop out, due to family reunion (14%) and loss to follow up (8%). A further five children were excluded due to either lack of data or legal consent, leaving a sample of 21 children. Despite the low sample size, significant improvements in child behaviour were found between baseline and final measurement on the Child Behaviour Checklist (CBLT) scores, whereas children who had not completed the program (transferred, dropped out, lost, other) had no significant reductions in symptoms on their last record for the CBLT. A high proportion of the children who participated fully in the program moved into permanent placements (92%), compared to an average of 40% placements in 'usual practice'. Authors concluded that the ARC model was a promising practice for young children, but required more formal research be conducted. This paper was rated as weak on all but one field (data collection methods was rated as moderate) in the risk of bias assessment.

The second ARC method paper also employed a naturalistic design to provide an empirical basis for the application of the ARC model (Hodgdon et al., 2013). The study measured treatment outcomes for 126 girls (aged 0 to 21 years) in residential care, who had experienced at least one traumatic event. This paper aimed to focus on a description of the initial stages of applying the ARC framework in a residential setting, and presented results from the program evaluation. Significant positive relationships were found between the ARC intervention and lower post-traumatic stress disorder (PTSD) symptomology. Externalising and internalising behaviours were reduced over the course of the intervention, as was the use of restraints. The authors acknowledged that one of the most significant limitations to the study was the study design, which was a naturalistic outcome evaluation, and not an experimental study. A second limitation was intervention fidelity, as the intervention was not fully employed for all clients in the programs. These factors meant that although outcomes were statistically significant, they were only modest from a clinical perspective. The study was rated weak overall in the risk of bias assessment, with two factors (selection bias and data collection methods) rated as moderate, and all other factors rated as weak.

The sample in the final ARC paper was pre- or post-adoptive children who had two or more lifetime traumatic exposures, and their carers (Hodgdon et al., 2016). In this semi-structured naturalistic study, children received 16 weeks of individual and group based ARC treatment, involving 16 individual sessions with the child, plus six group sessions which

included caregiver. Specific guidance on the goals of the sessions, the psychoeducational content, and strategies as defined in the ARC domains, targets and subskills were provided to children. There was a significant reduction in PTSD scores following the intervention: 76% of children were assessed as having clinical PTSD at baseline, compared to 33% at follow up. Significant reductions in children's anxiety, depression dissociation and anger scores were also found, and carers' stress was significantly reduced. Study limitations were listed as lack of control group and low treatment fidelity. Although results could be considered promising, for instance the effect size for the reduction in PTSD symptoms was large (Cohen's $D = 1.88$), authors stated that a more thorough evaluation was required. This study was rated as weak in the risk of bias assessment, with only the factor of selection bias and data collection methods rated as moderate, and all other factors rated as weak.

Children and Residential Experiences (CARE) model. The principal aim of the Children and Residential Experiences (CARE) model was the development of an organisational climate that was therapeutically beneficial; supporting and attending to the needs of each child within the organisation (Holden et al., 2010). This process was termed 'creating a therapeutic milieu', and involved personnel from all levels of the organisation incorporating CARE principles into daily practice. Inherent in the attention on the whole-of-organisation approach is the assumption that a positive organisational climate and positive staff interactions will lead to better services, as well as improved child outcomes and wellbeing (Glisson & Hemmelgarn, 1998). The aims of the model were to direct programming and enhance the dynamics of relationships throughout the organization. The model was based on systemic practices oriented around six core principles, being: 1) relationship based, 2) trauma-informed, 3) developmentally focused, 4) family-involved, 5) competence-centred, and 6) ecologically oriented. In general, agencies that used this model served young people living in residential care who had been referred from child welfare agencies (Izzo et al., 2016).

Only one paper in the current review evaluated the CARE model (Izzo et al., 2016). The aim of this study was to evaluate the implementation of the CARE model in a residential care environment over a three-year period. The authors used a multiple baseline interrupted time series, with five agencies beginning implementation in 2010, and six agencies in 2011. Staff were trained during an initial five-day program and ongoing assistance was provided by the CARE consultants, including observation and feedback, further training, encouraging routines for reflective practice, and addressing barriers at the organisational level. Post

implementation outcomes were compared to the twelve months' time-period prior to implementation. Implementation of the model led to significant reduction in aggression towards staff, property destruction and runaways. Inconclusive results were found for aggression towards peers and self-harm. The evaluation found that a more positive organisational climate predicted less aggression towards peers and less property destruction. This study was the only paper that passed the risk of bias assessment; it was rated as moderate on all fields.

The Sanctuary Model. The Sanctuary Model is an organisation-wide model for changing social service delivery to better respond to the complex needs of children who have experienced trauma (Abramovitz & Bloom, 2003). The model is informed by four knowledge areas: the psychobiology of trauma, actively creating non-violent environments, social learning principles, and understanding complex system change (Bloom et al., 2003). The model aims to implement an organisation-wide approach that involves creating and maintaining an environment that understands how children deal with trauma. A therapeutic community is provided to children that aims to mitigate the adverse effects of trauma. Sanctuary is the only organizational framework that has achieved a Scientific Rating of 3 (Promising Research Practice) by the California Evidence-Based Clearinghouse for Child Welfare (California Evidence-Based Clearinghouse for Child Welfare, 2016).

The theoretical basis of The Sanctuary Model stems from four conceptual frameworks: Trauma Theory, Social Learning Theory, Nonviolence, and Complexity Theory (Abramovitz & Bloom, 2003). Trauma theory is based on several decades of research describing the profound impact of stress on human development. Damage experienced through trauma is thought to be not from the trauma itself, but from the way the individual's mind and body reacts to the experience, combined with how the individual's social group responds. In Social Learning Theory, the active use of the whole environment becomes the grounds for therapeutic change. The incorporation of Nonviolent Practice places attention on safety as an active aspect of organisational life. An understanding of complex adaptive systems, for both individuals as well as organisations, is provided by Complexity Theory.

The model has been designed across four key pillars (Esaki et al., 2013): 1) that trauma can alter brain functioning and behaviour, and can affect whole systems/organisations; 2) the Safety, Emotion, Loss and Future (S.E.L.F) framework, which presents solutions to the complex problems of trauma and stress individually and

organisationally; 3) Sanctuary Tools, which include community meetings, safety plans, red flag meetings, team meetings, psychoeducation and supervision/training; 4) the Seven Sanctuary Commitments of nonviolence, emotional intelligence, social learning, democracy, open communication, social responsibility, and growth and change.

Three papers were found in the search that evaluated the Sanctuary Model (Bloom et al., 2003; Kramer, 2016; Rivard et al., 2005). In the first of these papers, Bloom et al. (2003) evaluated the use of the Sanctuary Model in five diverse organisational settings. Two of these settings were residential treatment programs for children, and a third setting was a group home for disturbed adolescents. In two further settings the sample were adults, and these sections have not been included in the current review. Although information on the study designs, implementation and analyses for the three relevant studies was limited, general outcomes were included. These outcomes included reduction in numbers of seclusions, improved patient satisfaction, improvement in staff interest and pride, the need for leadership to always role model appropriate behaviour, and the need for the organisation to focus on its moral purpose. The quality of methodology for these settings was weak, with only brief descriptions of sample and research methodology included in the paper. The paper was rated as weak overall on the risk of bias assessment.

In 2005, Rivard, Bloom, McCorkle and Abramovitz published a paper that summarised three previous articles on a preliminary evaluation of the implementation and effectiveness of the Sanctuary model for young people in residential treatment (Rivard, 2004; Rivard et al., 2003; 2004). The treatment program was evaluated using qualitative and quantitative methodology (focus groups). Sixteen residential care units were included in the study, with eight of the units receiving the intervention, and eight functioning as controls. Outcomes were measured at baseline, and at three- and six-months follow-up (Rivard et al., 2005). Results of the study included that the environment was significantly improved in the treatment units compared to the controls, and that youth had improved coping skills and sense of control over their lives at the six-month time point (but not at three months). In the risk of bias assessment, this paper was rated as weak overall, with subsections of selection bias and study design designated as moderate.

The most recently published study included in the current review used a qualitative design. Data collection included observation of groups, content analysis of agency documents, and focus groups and interviews with youth residents and staff (Kramer, 2016).

The organisation was a residential unit caring for court-committed adolescents, and the study aimed to investigate the effectiveness of the Sanctuary Model in decreasing symptoms of trauma. The researcher used grounded theory to evaluate interviews and documents, and results suggested that there was evidence that the intervention ameliorated trauma symptoms in the youth. The risk of bias assessment was rated as weak overall, and for all subcategories. While the outcomes from these three studies on the Sanctuary Model indicate the application of this model was promising, further research is required.

Discussion

This review aimed to investigate current evidence for organisation-wide trauma-informed care models in OoHC settings. Trauma-informed care models are currently being incorporated throughout many child protection systems and organisations world-wide, and the evidence points to general improvements from the implementation of these models, and the absence of detrimental outcomes. The current review, however, found that the evidence base for trauma-informed care models was low. Only seven papers met the inclusion criteria for the current review, of empirically measuring trauma-informed care systems in OoHC populations. Of these seven studies, only one was rated as being of moderate quality on the risk of bias assessment, with the other six studies classified as weak.

The second important finding in this systematic review was the difficulty in effectively evaluating outcomes from trauma-informed care models, as shown by the absence of strong study designs utilised in the studies in five of the seven papers. The paper that was assessed as having moderate risk of bias (Izzo et al., 2016), employed a multiple baseline interrupted time-series study design, utilising administrative data. This study design enabled comparisons between pre- and post-implementation, as well as measurements across cohorts, by staggering the introduction of the implementation of the program (five agencies implemented the model in 2010, and six in 2011). Using this method, a comparison could then be made across cohorts so that external environmental factors could be taken into account, whilst also ensuring that the program could be rolled out across all agencies.

Another example of creative methodology was a study by Murphy and colleagues (2017), not included in this review, who described their decision-making process for choosing a longitudinal quasi-experimental study design to evaluate a trauma-informed program (not a system-wide trauma-informed model). The researchers had initially sought to design a randomised controlled trial (RCT); however, after examination of logistic and

contamination concerns, the authors determined that an RCT was neither feasible nor appropriate in this context. Instead, this study used administrative data to compare three years of data collected during implementation to one year of data collected prior to implementation. Further, the dose of trauma-informed care exposure was compared to child outcomes. The study designs utilised in these two papers (Izzo et al., 2016; Murphy et al., 2017) suggest that the use of administrative data is one way that scientifically valid methodology can be used to evaluate system-wide initiatives, and that more standard evaluation strategies, such as randomised-controlled trials, may be neither suitable nor effective in these situations (Dixon et al., 2014).

A further outcome of this systematic review was that despite the slender evidence base evaluating organisation-wide trauma-informed care models, and the difficulties in evaluating organisation-wide processes, outcomes overall provide preliminary support for the efficacy of organisation-wide trauma-informed care models in OoHC populations. This support concords with the extensive anecdotal evidence for trauma-informed care models (such as, Farragher & Yanosy, 2005; Gurwitch et al., 2016; Hanson & Lang, 2014).

The main limitation of the current review was the poor outcome on the risk of bias assessments, suggesting that caution is required when interpreting the effectiveness of these models. The review was limited because of a lack of rigorous empirical evidence but this does not suggest that trauma informed models/frameworks are ineffective. What our findings suggest is that there is promise in adopting these models, but that more rigorous and systematic research is required to build the evidence base to inform implementation for sustainable positive and improved child and family outcomes.

A second limitation to this study was the lack of homogeneity between study outcomes, meaning that it was not possible to conduct a meta-analysis to generate estimates of the effect of organisation-wide trauma-informed care models. These studies have made a large contribution to the literature, and the tensions between the different worlds of research, policy and practice need to be acknowledged. It is important for researchers to bridge the gap to aid evidence-informed and evidence-based practices, and for policy makers and service providers to commit to rigorous evaluation and continual learning. Future research could focus on evaluating the various components of these models, and comparing these components across model types. It is recommended that an implementation science methodology is used to guide future research in this area.

Implementation science is an emerging field which aims to bridge the research-to-practice gap, and improve the quality and effectiveness of services provided. Specifically, implementation science can be defined as “the scientific study of methods to promote the uptake of research findings and other evidence based practices into routine practice” (Bauer, Damschroder, Hagedorn, Smith, & Kilbourne, 2015, p 1). Implementation science can be used to make significant organizational changes whilst attempting to avoid the pitfalls that can be the consequence of managing multiple programs, requirements and priorities (Wilson, Brandes, Ball, & Malm, 2012). When instigating new systems, inadequate planning, insufficient staff and patient input, and a lack of alignment with previous services or priorities can mean that it is difficult to prepare staff to effectively implement an initiative. Implementation science is critical in supporting evidence-based practices in public health (Proctor, 2012).

In conclusion, assessing the efficacy of organisation-wide trauma-informed care models is challenging, and requires creative solutions. The strongest quality criteria have been applied in this study, revealing that the current evidence for trauma-informed care models is limited. Given the amount of resources currently being employed in implementing trauma-informed care models worldwide, more robust evidence is required to show that these types of models are effective and how the models contribute to improved child and organisational outcomes. Despite the limited evidence, the analysis of the papers presented in this review provides promising evidence that the application of trauma-informed care models may have significantly positive outcomes for children in OoHC.

References

- Abramovitz, R., & Bloom, S. L. (2003). Creating Sanctuary in Residential Treatment for Youth: From the “ Well-Ordered Asylum ” To a “ Living-Learning Environment ,” 74(2), 119–135.
- AIHW. (2017). Child protection Australia 2015-2016. Child Welfare Series (Vol. no. 61). <https://doi.org/CWS 41>
- Arvidson, J., Kinniburgh, K., Howard, K., Spinazzola, J., Strothers, H., Evans, M., ... Blaustein, M. E. (2011). Treatment of Complex Trauma in Young Children: Developmental and Cultural Considerations in Application of the ARC Intervention

- Model. *Journal of Child & Adolescent Trauma*, 4(1), 34–51.
<https://doi.org/10.1080/19361521.2011.545046>
- Australian Institute of Family Studies. (2007). *“Feeling heavy”- Vicarious trauma and other issues facing those who work in the sexual assault field*. Melbourne. Retrieved from <https://aifs.gov.au/publications/feeling-heavy/what-vicarious-trauma>
- Barth, R. P., Greeson, J. K. P., Zlotnik, S. R., & Chintapalli, L. K. (2011). Evidence-Based Practice for Youth in Supervised Out-of-Home Care: A Framework for Development, Definition, and Evaluation. *Journal of Evidence-Based Social Work*, 8(5), 501–528.
<https://doi.org/10.1080/15433710903269529>
- Bauer, M. S., Damschroder, L., Hagedorn, H., Smith, J., & Kilbourne, A. M. (2015). An introduction to implementation science for the non-specialist. *BMC Psychology*, 3(1), 32. <https://doi.org/10.1186/s40359-015-0089-9>
- Becker-Blease, K. A. (2017). As the world becomes trauma-informed, work to do. *Journal of Trauma & Dissociation*, 18(2), 131–138.
<https://doi.org/10.1080/15299732.2017.1253401>
- Beyerlein, B. A., & Bloch, E. (2014). Need for Trauma-Informed Care Within the Foster Care System: A Policy Issue. *Child Welfare*, 93(3), 7–21. Retrieved from <http://search.ebscohost.com/login.aspx?direct=true&db=i3h&AN=110870975&site=ehost-live>
- Bloom, S. (2017). The Sanctuary Model.
- Bloom, S., Bennington-Davis, M., Farragher, B., McCorkle, D., Nice-Martini, K., & Wellbank, K. (2003). Multiple opportunities for creating sanctuary. *Psychiatric Quarterly*, 74(2), 173–190. <https://doi.org/10.1023/A:1021359828022>
- Bloom, S. L. (2016). Advancing a national cradle-to-grave-to-cradle public health agenda. *Journal of Trauma & Dissociation*, 17(4), 383–396.
<https://doi.org/10.1080/15299732.2016.1164025>
- California Evidence-Based Clearinghouse for Child Welfare. (2016). California Evidence-Based Clearinghouse for Child Welfare. Retrieved from <http://www.cebc4cw.org/>
- Department of Families, Housing, Community Services and Indigenous Affairs. (2011). An outline of National standards for out-of-home care: A priority project under the

National Framework for Protecting Australia's Children 2009 – 2020. Canberra.

- Dixon, J., Biehal, N., Green, J., Sinclair, I., Kay, C., & Parry, E. (2014). Trials and Tribulations: Challenges and Prospects for Randomised Controlled Trials of Social Work with Children. *British Journal of Social Work*, 44(6), 1563–1581. <https://doi.org/10.1093/bjsw/bct035>
- Esaki, N., Benamati, J., Yanosy, S., Middleton, J. S., Hopson, L. M., Hummer, V. L., & Bloom, S. L. (2013). The Sanctuary Model: Theoretical framework. *Families in Society: The Journal of Contemporary Social Services*, 94(2), 87–95. <https://doi.org/10.1606/1044-3894.4287>
- Farragher, B., & Yanosy, S. (2005). Creating a trauma-sensitive culture in residential treatment. *The International Journal for Therapeutic and Supportive Organizations*, 26(1), 97–113.
- Fraser, J. G., Griffin, J. L., Barto, B. L., Lo, C., Wenz-Gross, M., Spinazzola, J., ... Bartlett, J. D. (2014). Implementation of a workforce initiative to build trauma-informed child welfare practice and services: Findings from the Massachusetts Child Trauma Project. *Children and Youth Services Review*, 44, 233–242. <https://doi.org/10.1016/j.chilyouth.2014.06.016>
- Glisson, C., & Hemmelgarn, A. (1998). The effects of organizational climate and interorganizational coordination on the quality and outcomes of children service systems. *Child Abuse & Neglect*, 22(5), 401–421.
- Gurwitch, R. H., Messer, E. P., Masse, J., Olafson, E., Boat, B. W., & Putnam, F. W. (2016). Child-Adult Relationship Enhancement (CARE): An evidence-informed program for children with a history of trauma and other behavioral challenges. *Child Abuse and Neglect*. <https://doi.org/10.1016/j.chiabu.2015.10.016>
- Hanson, R. F., & Lang, J. (2014). Special focus section: a critical look at trauma informed care (TIC) among agencies and systems serving maltreated youth and their families. *Child Maltreat*, 19, 275. <https://doi.org/10.1177/1077559514557018>
- Hanson, R. F., & Lang, J. (2016). A Critical Look At Trauma-Informed Care Among Agencies and Systems Serving Maltreated Youth and Their Families. *Child Maltreatment*, 21(2), 95–100. <https://doi.org/10.1177/1077559516635274>

- Hodgdon, H. B., Blaustein, M., Kinniburgh, K., Peterson, M. L., & Spinazzola, J. (2016). Application of the ARC Model with Adopted Children : Supporting Resiliency and Family Well Being. *Journal of Child and Adolescent Trauma*, 9(1), 43–53.
<https://doi.org/10.1007/s40653-015-0050-3>
- Hodgdon, H. B., Kinniburgh, K., Gabowitz, D., Blaustein, M. E., & Spinazzola, J. (2013). Development and Implementation of Trauma-Informed Programming in Youth Residential Treatment Centers Using the ARC Framework. *Journal of Family Violence*, 28(7), 679–692. <https://doi.org/10.1007/s10896-013-9531-z>
- Holden, M. J., Izzo, C., Nunno, M., Smith, E. G., Endres, T., Holden, J. C., & Kuhn, F. (2010). Children and Residential Experiences : A Comprehensive for Residential Care. *Child Welfare*, 89(2), 131–150.
- Hummer, V. L., Dollard, N., Robst, J., & Armstrong, M. I. (2010). Innovations in Implementation of Youth Residential Treatment : A Curriculum for Organizational Change. *Child Welfare*, 89(2), 79–96.
- Izzo, C. V., Smith, E. G., Holden, M. J., Norton, C. I., Nunno, M. A., & Sellers, D. E. (2016). Intervening at the Setting Level to Prevent Behavioral Incidents in Residential Child Care: Efficacy of the CARE Program Model. *Prevention Science*, 17(5), 554–564.
<https://doi.org/10.1007/s11121-016-0649-0>
- Kinniburgh, K. (2005). Attachment, Self-Regulation, and Competency: A comprehensive intervention framework for children with complex trauma. *Psychiatric Annals*, 35(5), 424–430.
- Kramer, M. G. (2016). Sanctuary in a residential treatment center: creating a therapeutic community of hope countering violence. *Therapeutic Communities: The International Journal of Therapeutic Communities*, 37(2), 69–83. <https://doi.org/10.1108/TC-01-2015-0005>
- Macdonald, G., & Millen, S. (2012). Therapeutic approaches to social work in residential *child care settings : Literature review*. Belfast. Retrieved from www.scie.org.uk
- McLean, S., Price-Robertson, R., & Robinson, E. (2011). Therapeutic residential care in Australia: Taking stock and looking forward. Melbourne.
- Middleton, J. S., & Potter, C. C. (2015). Relationship Between Vicarious Traumatization and

- Turnover Among Child Welfare Professionals. *Journal of Public Child Welfare*, 9(2), 195–216. <https://doi.org/10.1080/15548732.2015.1021987>
- Murphy, K., Moore, K. A., Redd, Z., & Malm, K. (2017). Trauma-informed child welfare systems and children's well-being: A longitudinal evaluation of KVC's bridging the way home initiative. *Children and Youth Services Review*, 75, 23–34. <https://doi.org/10.1016/j.chidyouth.2017.02.008>
- Proctor, E. (2012). Implementation Science and Child Maltreatment. *Child Maltreatment*, 17(1), 107–112. <https://doi.org/10.1177/1077559512437034>
- Rivard, J. C. (2004). Initial findings of an evaluation of a trauma recovery framework in residential treatment. *Residential Group Care Quarterly*, 5(1), 3–5.
- Rivard, J. C., Bloom, J. C., McCorkle, D., & Abramovitz, R. (2005). Preliminary results of a study examining the implementation and effects of a trauma recovery framework for youths in residential treatment. *The International Journal for Therapeutic and Supportive Organizations*, 26(MARCH), 83–96.
- Rivard, J. C., Bloom, S. L., Abramovitz, R., Pasquale, L. E., Duncan, M., McCorkle, D., & Gelman, A. (2003). Assessing the implementation and effects of a trauma-focused intervention for youths in residential treatment. *Psychiatric Quarterly*, 74(2), 137–154. <https://doi.org/10.1023/A:1021355727114>
- Rivard, J. C., McCorkle, D., Duncan, M. E., Pasquale, L. E., Bloom, S. L., & Abramovitz, R. (2004). Implementing a trauma recovery framework for youths in residential treatment. *Child and Adolescent Social Work Journal*, 21(5), 529–550. <https://doi.org/10.1023/B:CASW.0000043363.14978.e6>
- Siegel, D. (2007). Toward an interpersonal neurobiology of the developing mind: Attachment relationships, “mindsight,” and neural integration. *Infant Mental Health Journal*, 22, 67–94. <https://doi.org/10.1002/1097-0355>
- Thomas, B., Ciliska, D., Dobbins, M., & Micucci, S. (2004). A process for systematically reviewing the literature: providing the research evidence for public health nursing interventions. *Worldviews on Evidence-Based Nursing*, 1(3), 176–184. Retrieved from <http://onlinelibrary.wiley.com/doi/10.1111/j.1524-475X.2004.04006.x/full>
- van der Kolk, B. A. (2003). *Psychological Trauma*. Washington, DC: American Psychiatric

Publishing.

van der Kolk, B. A. (2007). The Developmental Impact of Childhood Trauma. In L. J. Kirmayer, R. Lemelson, & M. Barad (Eds.), *Understanding trauma: Integrating biological, clinical, and cultural perspectives* (pp. 224–241). Cambridge: Cambridge University Press.

Victorian Auditor-General. (2014). *Residential Care Services for Children*. Melbourne. Retrieved from <https://www.audit.vic.gov.au/report/residential-care-services-children>

Wall, L., Higgins, D., & Hunter, C. (2016). *Trauma-informed care in child/family welfare services*. Melbourne. Retrieved from <https://aifs.gov.au/cfca/sites/default/files/publication-documents/cfca37-trauma-informed-practice.pdf>

Wilson, D., Brandes, B., Ball, H., & Malm, K. (2012). Building a Post-Care Service System in Child Welfare: Lessons Learned from the Frontlines of Implementation Science in Catawba County. *Child Trends, Publicatio*(July), 1–8.

Box 1. Search terms

CONCEPT 1 –Trauma

“trauma-informed” OR “trauma informed”

AND CONCEPT 2– Out-of-home care

“foster care” OR “child welfare” OR “out of home care*” OR “looked after child*” OR “residential care*” OR “group home*” OR “kin* care” OR “relative care”

AND CONCEPT 3 – Theory or intervention

“framework*” OR “model” OR “theor*” OR “conceptual framework*” OR “intervention*” OR “program*” OR “strateg*” OR “prevention” OR “treatment” OR “therap*” OR “organi?ation*” OR “organi?ational climate” OR “organi?ational culture” OR “organi?ational social context”

Table 1 Data extraction and synthesis

Author year country	Aim of study and intervention	Number of participants and dropouts	Description of target population	Methodology	Outcomes	Length of follow up	Funding source
Attachment, Self-Regulation and Competency (ARC) Model							
Arvidson et al. 2011 United States	Aim: To provide preliminary evidence of the effectiveness of the ARC model, Intervention: Attachment, Self-Regulation and Competency (ARC) Framework.	93 children treated, but only 26 completed the intervention.	Preschool and school-aged children in the child protective system treated through the Alaska Child Trauma Centre.	Research design: Naturalistic program evaluation. Data analysis: T-tests comparing pre- and post-scores on the Child Behaviour Checklist (CBC) (young child version), including box plot. Percentage comparison of placement after program to children who didn't receive program (transferred, dropped out, lost other)	Mean CBCL t-score at baseline was 62.6 and t-score was 43.6 at discharge. The average drop in CBCL t-scores for children completing treatment was 19 points. 92% children moved to permanent placements compared to usual 40%. ARC found to be a promising practice for young children.	No follow up post the study period.	None stated

				Covariates: none			
Hodgdon, Kinniburgh, Gabowitz, & Blaustein, & Spinazzola 2013 United States	Aim: To provide an empirical basis for the application of the ARC model into clinical and milieu [social environment] programs. Intervention: Attachment, Self-Regulation and Competency (ARC) Framework	126 females from two residential treatment programs.	Participants were a subset of data from residential centres, aged 0 to 21 years, having experienced at least one traumatic event.	Research design: Naturalistic study design using data from the National Child Traumatic Stress Network dataset. Data analysis: Multilevel regression analysis. Covariates: Not stated.	Significant relationships were found between the intervention (ARC) and lowering of Post Traumatic Stress Disorder (PTSD) symptomology, externalizing and internalizing behaviours, and the use of restraints across two programs.	Three follow-up assessments (time frame unclear.).	Partly funded by the Center for Mental Health Services, U.S. Department of Health and Human Services
Hodgdon, Blaustein, Kinniburgh, Peterson, & Spinazzola 2016 United States	Aim: To describe the application of the ARC model with adopted children impacted by complex trauma and their caregivers. Intervention: Attachment, Self-Regulation and Competency (ARC) Framework.	A subset of 481 children and their caretakers.	Pre- or post-adoptive children and carers who had two or more lifetime traumatic exposures, with current Post Traumatic Stress Disorder (PTSD) and	Research design: Semi-structured naturalistic approach, involving 16 weeks of individual and group-based ARC treatment. No control group; low treatment fidelity; raters were not blinded. Data analysis: correlations between	Significant lowering of Child Mental Health Symptoms from pre- to post- treatment, with 76% of children assessed as having PTSD (per the Clinically Administered PTSD [CAPS]) compared to 33.3% at follow up. The effect size for the reduction in PTSD symptoms was large (Cohen's D = 1.88).	12-month follow-up period	Not stated

			functional impairment in two domains.	parent and child outcomes; multilevel regression analysis for all other outcome data. Covariates: not stated.	Significant reductions were found for child anxiety, depression, post-traumatic stress, dissociation and anger. Significant reduction in care-giver stress.		
Children and Residential Experiences program (CARE)							
Izzo et al. 2016 United States	<p>Aim: To evaluate implementation of the CARE model over a three-year implementation period.</p> <p>Intervention: Children and Residential Experiences (CARE), involves systematic practices that are oriented around six principals:</p> <ol style="list-style-type: none"> 1-Relationship based 2-Trauma-informed 3-Developmentally focused 	<p>11 agencies.</p> <p>Incident data per days of care was extracted from administrative sources, with the average number of residents used to estimate per capita rates.</p> <p>Of the 16 agencies that initially agreed to participate in the study, one was no longer eligible due to changes in the</p>	<p>Ten agencies cared for male and female children between the ages of 7 and 18 years.</p> <p>One agency only served males.</p> <p>All agency caregivers lived in the home units for 1-2 week shifts.</p>	<p>Research design: Multiple baseline interrupted time series.</p> <p>Data analysis: A model for each of the five types of behavioural incidents were constructed using a mixed effects negative binomial regression model, to estimate the number of incidents per resident per month. Two slopes (time trends) for the 12 months prior to, and the three years post implementation were modelled.</p> <p>Organisational social</p>	<p>Implementation of the CARE program led to significant declines in the following three types of behavioural incidents:</p> <ul style="list-style-type: none"> - Aggression towards staff - Property destruction - Runaways. <p>Inconclusive results for:</p> <ul style="list-style-type: none"> - Aggression towards peers - Self-harm <p>More positive organisational climate predicted fewer:</p>	<p>The intervention was in place for 48 months, and no information post this time was presented.</p>	<p>The research was partly supported by a multi-year grant from The Duke Endowment, Charlotte, NC, USA</p>

	<p>4-Family-involved</p> <p>5-Competence-centred</p> <p>6-Ecologically oriented.</p> <p>In most agencies, the process of incorporating CARE involves changes in theoretical perspective, in organizational customs and expectations of roles.</p>	<p>resident population type, one closed prior to implementation, one discontinued due to changes in priorities by administration, and two were excluded due to lack of sufficient detail in databases on incident type.</p>		<p>context was determined at the agency level.</p> <p>Covariates: agency, agency's parent organisation, and organizational social context score at initiation of implementation period.</p>	<p>- Aggression towards peers</p> <p>- Property destruction</p> <p>Organisations with poorer conditions at the start had the most consistent implementation process.</p>		
The Sanctuary Model							
<p>Bloom et al. 2003</p> <p>United States</p>	<p>Aim: To describe the implementation of the Sanctuary model in five treatment scenarios.</p> <p>Intervention: The Sanctuary Model.</p>	<p>TP*1- 22 adults**</p> <p>TP2 - 150 children.</p> <p>TP3 - 112 children.</p> <p>TP4 - 10 adolescent children.</p> <p>TP5 - not stated.**</p>	<p>TP*1 - Hospital psychiatric unit.</p> <p>TP2 - Residential and day program for children.</p> <p>TP3 - Residential</p>	<p>Research design:</p> <p>Research design not specified for each TP</p> <p>Data analysis:</p> <p>TP1- Use of seclusion and restraint</p> <p>TP2 - (descriptive only)</p> <p>TP3 - (descriptive only)</p> <p>TP4- Numbers of</p>	<p>TP1 - Over two years, number of seclusions reduced by 87%, and hours spent in seclusion halved. Patient satisfaction surveys improved. Improvements noted in staff interest and pride.</p> <p>TP2 - Discusses process of</p>	Not stated	Not stated

		Drop-outs not stated in TPs	<p>programs for children.</p> <p>TP4 - Small residential facility.</p> <p>TP5 - Long-term residential program for women who are substance abusers and victims of violence.</p>	<p>therapeutic holds and hospitalisations</p> <p>TP5 - (descriptive only)</p> <p>Covariates: None</p>	<p>change at leadership level.</p> <p>Need for leadership to always role model appropriate behaviour.</p> <p>TP3 - Sanctuary helped organisation to focus on their moral purpose.</p> <p>TP4 - Large decreases in numbers of therapeutic holds, and no hospitalisations in second year. Staff demeanour towards children more positive. Difficult to maintain model with high staff turnover.</p> <p>TP5 - Lowered incidence of violence. Changes in how violent incidences were managed.</p>		
Kramer 2016	Aim: To understand the effectiveness of	20 staff participated in focus groups.	Residential staff, and court-	Research design: Qualitative, including	The intervention was found to ameliorate	Not stated	Not stated.

United States	<p>the Sanctuary Model in decreasing trauma symptoms.</p> <p>Intervention: The Sanctuary Model.</p>	<p>Out of 160 residents, 19 consented to participate in focus groups; only 13 participated and 6 withdrew due to concerns with being audio-recorded.</p>	<p>committed male adolescents in a residential treatment program.</p>	<p>observation of groups, content analysis of agency documents, focus groups and interviews with staff.</p> <p>Data analysis: Grounded Theory.</p> <p>Covariates: N/A</p>	<p>symptoms of complex trauma.</p>		
<p>Rivard, Bloom, McCorkle, & Abramovitz 2005</p> <p>United States</p>	<p>Aim: To evaluate the implementation and short-term effects of the Sanctuary Model in residential treatment programs for youth.</p> <p>Intervention: The Sanctuary Model.</p>	<p>Youths (N=158). Staff participated in focus groups (number not stated).</p>	<p>Youths aged 12-20 yrs., mean of 15 yrs. 33% Hispanic; 27% Black; 13% White; 1% other.</p> <p>Staff who worked in the programs and who had agreed to participate in study's focus groups and surveys.</p>	<p>Research design: Comparison Group Design, measured at baseline, 3 and 6 months. Four initial, and four randomly assigned subsequent residential units received the intervention. Eight units provided standard care.</p> <p>Data analysis: Qualitative analysis to assess intervention implementation. COPES scale</p>	<p>Intervention units showed significantly stronger outcomes in the treatment environment, such as support, autonomy, safety, and personal problem orientation, than units without the intervention.</p> <p>Youth gains were made in the areas of coping skills, and sense of control over environment at the six-month time point.</p>	<p>Results were only measured over a short time period of six months, no further follow-up data was presented.</p>	<p>Funded by the National Institute of Mental Health.</p>

				<p>(Community Oriented Programs Environment Scale) used to assess therapeutic environment.</p> <p>Youth outcomes assessed using a series of instruments.</p> <p>Covariates: none</p>			
--	--	--	--	---	--	--	--

* TP= Treatment Place **Only T2, T3 and T4 had suitable populations, and were retained for analysis in this review

hsc_12621_f1.tiff

hsc_12621_f2.tiff

hsc_12621_f3.tiff

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Bailey, C;Klas, A;Cox, R;Bergmeier, H;Avery, J;Skouteris, H

Title:

Systematic review of organisation-wide, trauma-informed care models in out-of-home care (OoHC) settings

Date:

2019-05-01

Citation:

Bailey, C., Klas, A., Cox, R., Bergmeier, H., Avery, J. & Skouteris, H. (2019). Systematic review of organisation-wide, trauma-informed care models in out-of-home care (OoHC) settings. *HEALTH & SOCIAL CARE IN THE COMMUNITY*, 27 (3), pp.E10-E22. <https://doi.org/10.1111/hsc.12621>.

Persistent Link:

<http://hdl.handle.net/11343/285133>