DR. ANSELM WONG (Orcid ID : 0000-0002-6817-7289)

Article type : Original

The NACSTOP trial: A multi-center, cluster-controlled trial of early cessation of acetylcysteine in acetaminophen overdose.

Corresponding author and contact info:

Dr. Anselm Wong MBBS

School of Clinical Sciences, Faculty of Medicine, Nursing and Health Sciences, Monash University

Victorian Poisons Information Centre and Austin Toxicology Service
Austin Hospital

Heidelberg

Victoria 3084

Australia

Tel: +61 3 9496 4509

Email: anselm.wong@austin.org.au

Co-investigators

Dr Richard McNulty PhD

Department of Emergency Medicine, Blacktown and Mount Druitt hospitals
Western Sydney Toxicology Service

NSW

Australia

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/hep.30224](https://doi.org/10.1002/hep.30224)

This article is protected by copyright. All rights reserved

Prof David McD Taylor MD

Emergency Department, Austin Hospital, Heidelberg, Victoria, Australia

Department of Medicine, University of Melbourne, Victoria, Australia

Victoria

Australia

David.Taylor@austin.org.au

Prof Marco L.A. Sivilotti MD MSc

Departments of Emergency and of Biomedical & Molecular Sciences

Queen's University

Kingston, Ontario

Canada

marco.sivilotti@queensu.ca

Dr. Shaun Greene MB ChB

Victorian Poisons Information Centre and Austin Toxicology Service

Austin Hospital

Heidelberg

Victoria

Australia

Shaun.Greene@austin.org.au

A/Prof Naren Gunja MD MS

Western Sydney Toxicology Service

Sydney Medical School

NSW

Australia

naren.gunja@sydney.edu.au

Dr Zeff Koutsogiannis MBBS

Victorian Poisons Information Centre and Austin Toxicology Service

Austin Hospital

Heidelberg

Victoria 3084

Australia

Zeff.Koutsogiannis@austin.org.au

Prof Andis Graudins PhD

Monash Toxicology Service and Monash Emergency Research Collaborative,
Dandenong Hospital

School of Clinical Sciences, Faculty of Medicine, Nursing and Health
Sciences, Monash University

Victoria

Australia

Andis.Graudins@monashhealth.org

Keywords: Paracetamol, N-acetylcysteine, antidotes, hepatotoxicity

List of abbreviations

ALT alanine transaminase

INR International Normalized Ratio

NAC N-acetylcysteine

ED Emergency Department

VAMPIRE Victorian Austin Monash Poisons Information, Research and Education

AST aspartate transaminase

NAARs non-allergic anaphylactic reactions

SAE serious adverse event

APAP acetaminophen

Funding statement

AW is the recipient of an Australian National Health and Medical Research Council (NHMRC) Postgraduate Scholarship ID 1114284. This project was supported by the Morson Taylor Award, Australasian College for Emergency Medicine.

Abstract

Historically, acetylcysteine has been delivered at a fixed dose and duration of 300 mg/kg over 20 hours to nearly every patient deemed to be at any risk for hepatotoxicity following acetaminophen overdose. We investigated a 12-hour treatment regimen for selected low-risk patients. This was a multicenter, open-label, cluster-controlled trial at six metropolitan emergency departments. We enrolled subjects following single or staggered acetaminophen overdose with normal serum alanine transaminase (ALT) and creatinine on presentation and at 12 hours, and acetaminophen $<20\text{mg/L}$ at 12 hours. Patients were allocated to intervention (250 mg/kg over 12-hour) or control (300 mg/kg over 20-hour) regimens by site. Primary outcome: incidence of “hepatic injury” 20 hours post-initiation of acetylcysteine treatment; defined as ALT doubling and peak ALT $> 100\text{ IU/L}$, indicating the need for further antidotal treatment. Secondary outcomes included incidence of hepatotoxicity (ALT $> 1000\text{ IU/L}$), peak INR and adverse drug reactions. Of 449 acetaminophen overdoses receiving acetylcysteine, 100 were recruited to the study. Time to acetylcysteine (median 7 hours [IQR 6,12] vs 7 hours [6,10]) and initial acetaminophen (124 mg/L [58,171] vs 146 mg/L [66,204]) were similar between intervention and control groups. There was no difference in ALT (18 IU/L [13,22] vs 16 IU/L [13,21]) or INR (1.2 vs 1.2) 20 hours after starting acetylcysteine between groups. No patients developed hepatic injury or hepatotoxicity in either group (OR 1.0 [95% CI 0.02,50]). No patients re-presented with liver injury, none died and 96/96 were well at 14-day telephone follow-up. **Conclusion:** Discontinuing acetylcysteine based on laboratory testing after 12 hours of treatment is feasible and likely safe in selected patients at very low risk of liver injury from acetaminophen overdose.

Trial registration: Clinical Trial Registration

ACTRN12615000938505 www.anzctr.org.au Registered 8/9/2015

Acetaminophen is one of the most common medications taken in overdose around the world (1). Acetylcysteine (*N*-acetylcysteine or NAC) is the antidote used to treat patients at risk of developing hepatotoxicity following acetaminophen overdose (2). The decision to treat cases of single acute acetaminophen overdose with acetylcysteine is based upon a timed acetaminophen concentration plotted on a treatment nomogram (3). The standard acetylcysteine treatment regimen, used for several decades, calls for 300mg/kg infused over a fixed interval of 20 to 21 hours (i.e. 150 mg/kg of acetylcysteine intravenously (IV) over 15-60 minutes, followed by 50 mg/kg acetylcysteine over 4 hours and 100 mg/kg acetylcysteine over 16 hours (4)). To simplify acetylcysteine dosing and reduce adverse reactions during the rapid initial load (5-7), we recently instituted a two-bag acetylcysteine regimen consisting of 200 mg/kg over 4 hours followed by a further 100 mg/kg over 16 hours (8, 9). While the total dose and duration is similar to the standard acetylcysteine regimen,(4) the incidence of adverse reactions is substantially lower (Wong et al.(8), McNulty et al.(9), Schmidt (10)). However, the fixed duration of 20 hours, which typically necessitates hospital admission, likely represents overtreatment for patients at very low risk, and ignores an opportunity for further risk stratification based on retesting midway through treatment (11).

The administration of acetylcysteine within eight hours of an acute single acetaminophen overdose prevents hepatotoxicity in nearly all patients (4, 12-14). More recent work has shown that patients with normal liver function tests when treatment is started are unlikely to develop hepatotoxicity (15-17).

Moreover, very few patients in whom acetylcysteine is discontinued prematurely develop hepatotoxicity, provided treatment is started within eight hours of overdose and at least 200mg/kg are administered (17). The initial three-bag, 20-hour intravenous acetylcysteine regimen was first proposed in the 1970's. The 20-hour duration was arbitrarily selected and based upon a crude estimate of acetaminophen elimination (i.e. 5 half-lives). This treatment recommendation was made at a time when the serum acetaminophen concentration was not readily measurable in most hospitals (18). Serial acetaminophen and liver function assays are now easily performed, and can provide reassurance that hepatic function is preserved. Indeed, an acetaminophen concentration that falls by at least 90% over 12 hours (i.e. greater than three half-lives) is the first readily available test of liver function following overdose, and provides a mathematically simple benchmark for early risk stratification(17). Acetylcysteine elimination is slower than acetaminophen in low-risk patients, and antidote remains in the circulation for a few hours after cessation of the infusion (19). A previous randomized trial of 300 mg/kg acetylcysteine administered over 12 hours demonstrated a decrease in adverse reactions to acetylcysteine when compared to the original 20-hour three-bag regimen, with no signal of diminished efficacy (6).

Our aim was to test the feasibility and safety of early discontinuation of the two-bag regimen compared to the full 20-hour course in selected low-risk adult patients based on serial laboratory testing. These low-risk features included normal serum ALT and creatinine both at presentation and after 12 hours of treatment, as well as an acetaminophen concentration that fell into the therapeutic range or lower by 12 hours.

Methods and analysis

Study setting

This was a multi-center, cluster-controlled open-label trial of patients with acute single or staggered acetaminophen overdose who presented to the emergency departments (EDs) of six Australian metropolitan hospitals from February 2016 to February 2018. There were three toxicology units, part of the VAMPIRE (Victorian Austin Monash Poisons Information, Research and Education) network and Western Sydney Toxicology Service, involved in managing patients at each hospital.

Study eligibility and recruitment

Study investigators and treating clinicians aware of the study approached eligible patients during the first 12 hours of acetylcysteine treatment, since the initial 12-hour infusion was identical for all subjects and standardized between sites. All subjects provided written informed consent to participate in the study.

The inclusion criteria were:

- age 16 years or older, AND
- acute single or staggered (i.e. over > 1 hour) acetaminophen ingestion, requiring acetylcysteine as per the Australian and New Zealand acetaminophen guidelines (i.e. above the 150 mg/L at 4 hour treatment nomogram line for single ingestions) (3), AND
- serum alanine transaminase (ALT) less than the local laboratory upper limit of normal (i.e. 40 IU/L) on presentation and after 12 hours of acetylcysteine infusion, AND
- serum acetaminophen concentration < 20 mg/L (132 µmol/L; i.e. the upper therapeutic concentration) after 12 hours of acetylcysteine infusion; AND
- serum creatinine within normal laboratory limits on presentation and after 12 hours of acetylcysteine infusion.

The exclusion criteria were known pregnancy, acetaminophen modified-release ingestion, other supratherapeutic ingestions (i.e. non-intentional ingestions of > 10 grams or > 200 mg/kg over 24 hours, > 6 grams/day over 48 hrs). Modified-release ingestions were excluded because of their erratic absorption, less predictable pharmacokinetics and that peak acetaminophen concentrations may be delayed in overdose(20). Supratherapeutic ingestions were excluded because the current Australian and New Zealand acetaminophen management guideline allows for acetylcysteine discontinuation based on laboratory testing eight hours after initiation of acetylcysteine. Those requiring treatment with acetylcysteine and excluded from the study received management as per the guideline (3).

Study Procedures

Six hospitals were assigned to the two study arms in a 1:1 fashion (i.e. three intervention and three control sites). The toxicology inpatient units were physically based at the intervention sites, hence available to oversee more closely the changed acetylcysteine regimen and to prevent dosing errors including protocol violations, but investigators also consented and consulted

in person on subjects at the control sites. All eligible patients were started on the two-bag infusion acetylcysteine regimen (200 mg/kg over 4 hours followed by 6.25 mg/kg/hr prepared in 1000 mL of glucose 5% to provide 100mg/kg over the next 16 hours]). After 12 hours of treatment, that is after 8 hours of the second bag and 250 mg/kg of acetylcysteine, the serum acetaminophen concentration, liver and renal function tests were repeated. If the serum ALT was < 40 IU/L, serum creatinine was normal, and the acetaminophen was < 20 mg/L (132 micromol/L), acetylcysteine was discontinued at intervention sites and 1 L Ringer's lactate was infused instead over the remaining 8 hours. At control sites, the remaining 50 mg/kg acetylcysteine was continued in all cases to provide 300mg/kg total over 20 hours (Figure 1). If the 12-hour laboratory tests did not meet eligibility criteria (i.e. were abnormal), the patient was excluded from the study and the remaining dose of acetylcysteine was continued for the full 300mg/kg regardless of site.

By study protocol, all patients had serum acetaminophen concentration, electrolytes, creatinine, aspartate transaminase (AST) and ALT measured on presentation and again after 12 and 20 hours of acetylcysteine infusion.

Study Follow-up

A structured telephone interview performed by either of two study investigators (AW/RM) after at least 14 days was performed to identify any unscheduled hospital visits for symptoms suggestive of liver injury including abdominal pain and jaundice, for another overdose, or for admission to hospital for any reason, including surgical procedures, laboratory testing, or referral to the regional liver transplant unit. Electronic medical records from each of the six study hospitals were also reviewed for this information at 14 days post enrollment. We queried the Australian and New Zealand liver transplant registries to identify whether any subject had been listed as a potential recipient within 14 days of enrollment. We also searched the study hospital records prior to study completion to establish vital status for the few patients who could not be contacted by phone.

Outcome Measures

The primary outcome was the incidence of “hepatic injury” at 20 hours post-initiation of acetylcysteine treatment, defined as ALT doubling and peak ALT > 100IU/L (17). Secondary outcomes included the incidence of hepatotoxicity (peak ALT > 1000 IU/L), change in ALT, peak International Normalized Ratio (INR both absolute, and at a cutpoint of 1.3), and adverse reactions to acetylcysteine (grouped as gastrointestinal symptoms and non-allergic anaphylactic (NAARs or “anaphylactoid”) reactions). NAARs were subclassified into cutaneous (flushing, rash, urticaria, wheals, itch), and more severe reactions namely respiratory (bronchospasm, wheeze, dyspnea, shortness of breath), angioedema or cardiovascular instability (hypotension).

Data collection/Management/Monitoring

Data were entered on study-specific case report forms by study investigators. Data included demographic details, reported dose, formulation and time of overdose, results of blood tests as per protocol (acetaminophen, ALT, AST, INR, creatinine), amount of acetylcysteine received, duration of acetylcysteine treatment, adverse reactions to acetylcysteine, length of stay and discharge disposition.

Clinical data were retrieved from hospital electronic medical records (Cerner Millennium®, Cerner, Kansas City, MO, USA) and Symphony®, Version 2.29, Ascribe, United Kingdom) and transcribed manually into a secure password-protected research dataset residing on secure hospital computer servers. All blood specimens were collected and analyzed at the participating hospital laboratories as per standard laboratory practice.

The study was approved by each institution’s review board, with a requirement to immediately review and report any serious adverse events (SAEs) including subject death deemed to be secondary to the altered acetylcysteine regimen or morbidity from dosing errors.

This study was registered with the Australian and New Zealand Clinical Trials Registry (ACTRN12615000938505).

Statistical analysis

A sample size of 100 patients was selected based on expected recruiting efficiency and on the feasibility of the trial given the number of sites participating. The first fifty patients to meet the inclusion criteria in each study arm were included. A low primary outcome event rate in such a sample would also provide some confidence with which to pursue a larger clinical trial, while the secondary outcomes could still have sufficient power to identify an important but subclinical safety signal.

Descriptive data are reported as medians and IQR, as appropriate. Odds ratios (95%CI) were used for binary outcomes, the Mann Whitney U test to compare non-parametric data, and the Wilcoxon matched-pairs signed rank test to compare ALT pairs taken at presentation and after 20 hours post initiation of acetylcysteine. For subjects with an acute ingestion at a known time, we calculated the ratio between each subject's acetaminophen concentration at presentation ($APAP_s$) and the threshold concentration for treatment based on the time interval post ingestion using the Rumack-Matthew nomogram ($APAP_t$) as customary in risk stratification (21). This is to allow comparison of acetaminophen concentrations in relation to time post ingestion. We also calculated the acetaminophen X ALT multiplication product (22) at presentation and after 12 hours of acetylcysteine, and the ψ parameter (23) at the start of acetylcysteine, for further risk stratification. GraphPad Prism (V7, GraphPad Software, California, USA) was used to perform the statistical analysis.

Results

During the study period, 1411 patients presented with acetaminophen overdose at the six study sites, of whom 449 received acetylcysteine (Figure 2). One hundred subjects were enrolled into the trial (50 intervention and 50

control group) and all were intentional overdose. The majority (77%) were female, aged 23 years (median, [IQR 18-32]). The two groups were well balanced with regards to demographics including age, sex, ethnicity, as well as ingested acetaminophen dose, initial acetaminophen concentration (Figure 3), initial serum ALT and time to acetylcysteine treatment (Table 1). The $APAP_s:APAP_t$ ratio was also similar between groups (1.25 [1,1.6] vs 1.3 [1.1,1.7], $p=0.38$); almost all (94%) had a ratio between 1 and 2. All study subjects were admitted to a short stay observation unit for acetylcysteine treatment, and adherence to protocol was excellent.

No subjects met the primary outcome of hepatic injury (OR 1.0 [95% CI 0.02,50]). There was no significant difference in 20-hour ALT between intervention and control groups (median 18 IU/L [13,22] vs 16 [13,21], $p=0.51$) (Figure 4, nor in change from baseline (3 IU/L [0,5.5] vs 2 IU/L [0, 4.25], $p=0.48$). No subjects demonstrated a doubling of ALT regardless of peak. The 20-hour creatinine was also similar between groups (61 μ mol/L [52,70] vs 61 μ mol/L [51,68], $p=0.44$). In addition, no subjects developed hepatotoxicity, died or received a liver transplant.

There was no difference in 20-hour INR between groups (1.2 [1.1-1.3] in both groups, $p = 0.98$). However, there were more subjects with an INR >1.3 in the control group ($n=8$, 16%) versus the intervention group ($n=3$, 6%; OR 3.0 [95%CI 0.74,12]) at 20 hours. Median acetylcysteine duration was 13 hours (IQR 13-13.5) in the intervention group and 20 hours (20-20) in those receiving the full course. No subjects required an extension in acetylcysteine treatment. Those who did not require an inpatient psychiatric admission were discharged with community follow up.

With regards to adverse reactions to the acetylcysteine infusion, only one subject in the control group developed a NAAR, namely dyspnea two hours after commencing the infusion. This resolved spontaneously within 15 minutes and acetylcysteine was ceased temporarily. No subjects in either group developed systemic or cutaneous reactions. The most common adverse

reactions were gastrointestinal (Table 2). There was no significant difference in the incidence of gastrointestinal symptoms between groups (28% vs 24%, OR 1.23 [95%CI 0.50, 3.01], $p=0.64$). Five (10%) subjects in the intervention group and four (8%) in the control group had gastrointestinal symptoms prior to starting acetylcysteine. There were no serious adverse events noted during the trial.

At the 14-day followup, 96 (96%) subjects were contacted by telephone, of whom only one reported returning to hospital with abdominal pain on day two following the overdose. Repeated liver function tests and abdominal ultrasound were both normal at that time, and the patient was discharged. No other subjects reported any persisting gastrointestinal symptoms on phone follow up. Of the four who could not be contacted, all were known to be alive based on subsequent and unrelated presentations to hospital.

Discussion

For many years, treatment of acetaminophen poisoning with acetylcysteine has taken a “one size fits all” approach(11). Improvements in our understanding of risk stratification following acetaminophen overdose (14-16, 24) and wide availability of rapid and accurate laboratory testing present an opportunity for more individualized dosing of antidotal therapy. It has long been recognized that the treatment threshold for acetylcysteine is very conservative (11, 13, 18). Moreover, many patients present and are treated within hours of an acute overdose, and demonstrate persistently normal liver function including acetaminophen elimination (14, 16). We believe such patients to be an ideal target for abbreviated therapy, and set out to systematically describe their outcomes in a carefully controlled prospective and comparative trial.

In this trial, subjects who had persistently normal serum ALT at the start and after 12 hours of treatment, and in whom the serum acetaminophen fell at least into the therapeutic range within 12 hours of treatment, had identical and entirely benign clinical and biochemical outcomes whether the abbreviated or the complete two-bag acetylcysteine regimen was administered. Moreover, there was no indication of any subclinical biochemical differences in any of the secondary outcomes, including any doubling of the ALT regardless of peak, or slowed elimination of acetaminophen. The only exception was a marginally higher proportion of subjects with an INR >1.3 at 20 hours in the control group, almost certainly due to the well-known effect of acetylcysteine on prothrombin time (25).

Also, with a median time of 7 hours to initiation of acetylcysteine, all patients had ALTs performed greater than 24 hours post ingestion which were normal. A normal ALT on presentation has a high negative predictive value (100%) of individuals developing any serious liver injury (ALT >1000 IU/L) in those receiving acetylcysteine (26).

Of note, we identified eligible patients for abbreviated treatment after 12 hours of antidotal therapy. We required that they have, by definition, a 12-hour APAP x ALT product below 800 mg/L x IU/L, (15, 24) a value substantially lower than is seen in patients who go on to develop hepatotoxicity or hepatic failure (20, 22, 24, 27). Other risk stratification measures were also reassuring in these patients: the ψ parameter at the start of treatment was generally ≤ 2 mmol/L x hour (23), , and the serum acetaminophen concentration almost always had fallen by at least 90% in 12 hours.

Other lines of reasoning provide further reassurance of the safety of abbreviated antidotal therapy. Population-pharmacokinetic modelling of the two-bag acetylcysteine regimen suggests that when acetylcysteine is discontinued at 12 hours, it remains detectable in the bloodstream at appreciable concentrations for up to eight hours (28). This may provide some degree of ongoing liver protection once acetylcysteine is ceased, although neither the minimum nor optimum therapeutic serum concentration of acetylcysteine is known.

In addition, preliminary work in a subset of NACSTOP trial subjects has shown similar peak microRNA-122 expression between groups, a novel measure of hepatic injury following acetaminophen overdose (29). MicroRNA-122 can detect subclinical liver injury prior to rises in ALT (30).

The current acetaminophen treatment nomogram in Australia uses a 25% lower acetaminophen concentration threshold for instituting acetylcysteine treatment (150 mg/L or 1000 micromol/L at 4 hours) when compared to the original acetaminophen treatment nomogram used in the U.K. until recently (200 mg/L or 1300 micromol/L at 4 hours) (18) and in Australia prior to 2008 (31). Patients in this so-called “possible hepatotoxicity” zone (i.e. in the absence of treatment) represent a majority of acetaminophen overdoses receiving treatment, and are clearly ideal candidates for abbreviated therapy. However, one-fifth of subjects in our trial had an initial acetaminophen concentration >200 mg/L yet still met inclusion criteria after 12 hours of

acetylcysteine by virtue of rapid acetaminophen metabolism and complete absorption, with a fall in acetaminophen of at least 90%.

In addition, almost half the subjects in our study who received the abbreviated acetylcysteine regimen had treatment commenced more than eight hours post-overdose. Therefore, using this low-risk feature as an inclusion criterion would substantially reduce the number eligible for abbreviated treatment, and demonstrates the advantage of using serial testing instead. We did not observe any biochemical evidence of liver injury even in these subjects in whom the ALT was normal at presentation and after 12 hours of treatment. Another important benefit of serial testing is reduced reliance on knowing with accuracy the time of ingestion, or insisting that the ingestion be taken at a single point in time.

Efforts to abbreviate acetylcysteine protocols have made slow gains in the past, but may now be gathering momentum. Previous modestly-sized studies were part of the evidence base used to abbreviate oral acetylcysteine in the United States when the 72-hour oral course was routine.(32, 33) There have now been at least three studies of abbreviated IV acetylcysteine regimens(6, 17, 34). A randomized clinical trial (6) of a 300mg/kg over 12-hour Scottish Newcastle Acetylcysteine Protocol resulted in a decreased incidence of adverse reactions to antidotal treatment, but was not powered for clinical efficacy. Lucyk et al. reported that hepatic injury or hepatotoxicity were uncommon when the 20-hour intravenous acetylcysteine protocol was discontinued early, perhaps because of adverse effects, especially if the aminotransferases were normal and acetaminophen falling (17). A retrospective review of supratherapeutic acetaminophen ingestions receiving an eight-hour acetylcysteine regimen reported liver injury in 0 out of 46 patients if ALT was < 50 IU/L on presentation (34).

In practice, just under half of patients requiring acetylcysteine treatment would be eligible for the NACSTOP protocol (Figure 2) and more in countries with lower treatment thresholds. Benefits of a decreased duration of medical treatment might include an earlier opportunity for mental health assessment

and interventions, fewer hospitalizations, and perhaps lower incidence of acetylcysteine dosing errors. These advantages may be especially important in jurisdictions which apply an even lower treatment threshold of 100 mg/L at four hours,(35) or which treat empirically (36), or which suffer increasingly crowded emergency departments due to hospital access block (37, 38).

This study also confirmed the very low rate of non-IgE mediated “anaphylactoid” reactions by modifying the load to 200 mg/kg acetylcysteine infusion over four hours. Other studies now support this administration regimen and have also demonstrated lower adverse event rates to acetylcysteine (8-10). The incidence of gastrointestinal adverse events remains high and was similar in both study groups, and included a substantial number with symptoms prior to receiving acetylcysteine. Gastrointestinal adverse rates have been up to 40% of subjects in previous studies, and can often be attributed to the ingestion itself.

Limitations

There are several limitations of this study. We cannot discount the potential for selection bias in this open-label study, yet refusals and missed enrollments were similar in frequency between arms. The study was not randomized at the individual subject level, and a cluster design for the study was chosen for both simplicity and to minimize protocol violations and crossover. While the study was open-label, the primary endpoint was based on biochemical test results and hence unlikely to be unduly biased. The sample size in this study was small, but appropriate for an initial safety and feasibility study. We believe that future studies can now test our abbreviated acetylcysteine regimen using comparable inclusion criteria.

Conclusion

An abbreviated 12-hour acetylcysteine regimen is feasible and likely safe in selected patients at low risk of liver injury from acetaminophen overdose based on laboratory testing after 12 hours of treatment.

Competing interests

The authors declare no competing interest.

Acknowledgements

We thank our colleagues in the treating hospitals who assisted with recruitment.

Funding statement

AW is the recipient of an Australian National Health and Medical Research Council (NHMRC) Postgraduate Scholarship ID 1114284. This project was supported by the Morson Taylor Award, Australasian College for Emergency Medicine.

References

1. Mowry JB, Spyker DA, Cantilena LR, Jr., McMillan N, Ford M. 2013 Annual Report of the American Association of Poison Control Centers' National Poison Data System (NPDS): 31st Annual Report. Clin Toxicol (Phila). 2014;52(10):1032-283.
2. Prescott LF. Paracetamol poisoning. Prevention of liver damage. Med Chir Dig. 1979;8(5):391-3.
3. Chiew A, Fountain JS, Graudins A, Isbister GK, Reith D, Buckley NA. Summary statement: new guidelines for the management of paracetamol poisoning in Australia and New Zealand. Medical Journal of Australia. 2015;203(5):215-8.
4. Prescott LF, Illingworth RN, Critchley JA, Stewart MJ, Adam RD, Proudfoot AT. Intravenous N-acetylcysteine: the treatment of choice for paracetamol poisoning. Br Med J. 1979;2(6198):1097-100.
5. Sandilands EA, Bateman DN. Adverse reactions associated with acetylcysteine. Clin Toxicol (Phila). 2009;47(2):81-8.
6. Bateman DN, Dear JW, Thanacoody HK, Thomas SH, Eddleston M, Sandilands EA, et al. Reduction of adverse effects from intravenous acetylcysteine treatment for paracetamol poisoning: a randomised controlled trial. Lancet. 2014;383(9918):697-704.
7. Yarema M, Chopra P, Sivilotti MLA, Johnson D, Nettel-Aguirre A, Bailey B, et al. Anaphylactoid Reactions to Intravenous N-Acetylcysteine during Treatment for Acetaminophen Poisoning. J Med Toxicol. 2018;14(2):120-7.
8. Wong A, Graudins A. Simplification of the standard three-bag intravenous acetylcysteine regimen for paracetamol poisoning results in a lower incidence of adverse drug reactions. Clin Toxicol (Phila). 2016;54(2):115-9.
9. McNulty R, Lim JME, Chandru P, Gunja N. Fewer adverse effects with a modified two-bag acetylcysteine protocol in paracetamol overdose. Clin Toxicol (Phila). 2017:1-4.
10. Schmidt LE, Rasmussen DN, Petersen TS, Macias-Perez IM, Pavliv L, Kaelin B, et al. Fewer adverse effects associated with a modified two-bag intravenous acetylcysteine protocol compared to traditional three-bag regimen in paracetamol overdose. Clin Toxicol (Phila). 2018:1-7.

11. Dart RC, Rumack BH. Patient-tailored acetylcysteine administration. *Ann Emerg Med.* 2007;50(3):280-1.
12. Doyon S, Klein-Schwartz W. Hepatotoxicity despite early administration of intravenous N-acetylcysteine for acute acetaminophen overdose. *Acad Emerg Med.* 2009;16(1):34-9.
13. Buckley N, Eddleston M. Paracetamol (acetaminophen) poisoning. *BMJ Clin Evid.* 2007;2007.
14. Sivilotti ML, Yarema MC, Juurlink DN, Good AM, Johnson DW. A risk quantification instrument for acute acetaminophen overdose patients treated with N-acetylcysteine. *Ann Emerg Med.* 2005;46(3):272-4.
15. Wong A, Sivilotti ML, Dargan PI, Wood DM, Greene SL. External validation of the paracetamol-aminotransferase multiplication product to predict hepatotoxicity from paracetamol overdose. *Clin Toxicol (Phila).* 2015;53(8):807-14.
16. Green TJ, Sivilotti ML, Langmann C, Yarema M, Juurlink D, Burns MJ, et al. When do the aminotransferases rise after acute acetaminophen overdose? *Clin Toxicol (Phila).* 2010;48(8):787-92.
17. Lucyk SN, Yarema MC, Sivilotti ML, Johnson DW, Nettel-Aguirre A, Victorino C, et al. Outcomes of Patients With Premature Discontinuation of the 21-h Intravenous N-Acetylcysteine Protocol After Acute Acetaminophen Overdose. *J Emerg Med.* 2016;50(4):629-37.
18. Rumack BH. Acetaminophen hepatotoxicity: the first 35 years. *J Toxicol Clin Toxicol.* 2002;40(1):3-20.
19. Sansone RA, Sansone LA. Getting a Knack for NAC: N-Acetyl-Cysteine. *Innov Clin Neurosci.* 2011;8(1):10-4.
20. Wong A, Sivilotti MLA, Graudins A. Accuracy of the paracetamol-aminotransferase multiplication product to predict hepatotoxicity in modified-release paracetamol overdose. *Clin Toxicol (Phila).* 2017;55(5):346-51.
21. Marks DJB, Dargan PI, Archer JRH, Davies CL, Dines AM, Wood DM, et al. Outcomes from massive paracetamol overdose: a retrospective observational study. *Br J Clin Pharmacol.* 2017;83(6):1263-72.

22. Sivilotti MLA, Green TJ, Langmann C, Yarema M, Juurlink D, Johnson D. Multiplying the serum aminotransferase by the acetaminophen concentration to predict toxicity following overdose. *Clinical Toxicology*. 2010;48(8):793-9.
23. Sivilotti ML, Yarema MC, Juurlink DN, Good AM, Johnson DW. A risk quantification instrument for acute acetaminophen overdose patients treated with N-acetylcysteine. *Ann Emerg Med*. 2005;46(3):263-71.
24. Wong A, Graudins A. Risk prediction of hepatotoxicity in paracetamol poisoning. *Clin Toxicol (Phila)*. 2017;55(8):879-92.
25. Jang DH, Weaver MD, Pizon AF. In vitro study of N-acetylcysteine on coagulation factors in plasma samples from healthy subjects. *J Med Toxicol*. 2013;9(1):49-53.
26. Al-Hourani K, Mansi R, Pettie J, Dow M, Bateman DN, Dear JW. The predictive value of hospital admission serum alanine transaminase activity in patients treated for paracetamol overdose. *QJM*. 2013;106(6):541-6.
27. Wong A, Sivilotti MLA, Gunja N, McNulty R, Graudins A. Accuracy of the paracetamol-aminotransferase product to predict hepatotoxicity in paracetamol overdose treated with a 2-bag acetylcysteine regimen. *Clin Toxicol (Phila)*. 2018;56(3):182-8.
28. Wong A, Landersdorfer C, Graudins A. Pharmacokinetic modelling of modified acetylcysteine infusion regimens used in the treatment of paracetamol poisoning. *Eur J Clin Pharmacol*. 2017;73(9):1103-10.
29. Wong A, Nejad C, Gantier M, Choy KW, Doery J, Graudins A. The potential utility of microRNA for comparing efficacy of acetylcysteine regimens in paracetamol overdose. Abstracts (no.207) from the 38th International Congress of the European Association of Poisons Centres and Clinical Toxicologists (EAPCCT) 22-25 May 2018, Bucharest, Romania. *Clin Tox*. 2018;56(6):453-608.
30. Dear JW, Clarke JI, Francis B, Allen L, Wraight J, Shen J, et al. Risk stratification after paracetamol overdose using mechanistic biomarkers: results from two prospective cohort studies. *Lancet Gastroenterol Hepatol*. 2018;3(2):104-13.
31. Daly FF, Fountain JS, Murray L, Graudins A, Buckley NA, Panel of A, et al. Guidelines for the management of paracetamol poisoning in Australia and New Zealand--explanation and elaboration. A consensus statement from clinical

toxicologists consulting to the Australasian poisons information centres. Medical Journal of Australia. 2008;188(5):296-301.

32. Betten DP, Cantrell FL, Thomas SC, Williams SR, Clark RF. A prospective evaluation of shortened course oral N-acetylcysteine for the treatment of acute acetaminophen poisoning. Ann Emerg Med. 2007;50(3):272-9.
33. Woo OF, Mueller PD, Olson KR, Anderson IB, Kim SY. Shorter duration of oral N-Acetylcysteine therapy for acute acetaminophen overdose. Ann Emerg Med. 2000;35(4):363-8.
34. Wong A, Gunja N, McNulty R, Graudins A. Analysis of an 8-hour acetylcysteine infusion protocol for repeated supratherapeutic ingestion (RSTI) of paracetamol. Clin Toxicol (Phila). 2018;56(3):199-203.
35. McQuade DJ, Dargan PI, Keep J, Wood DM. Paracetamol toxicity: What would be the implications of a change in UK treatment guidelines? Eur J Clin Pharmacol. 2012;68(11):1541-7.
36. Schmidt LE. Identification of patients at risk of anaphylactoid reactions to N-acetylcysteine in the treatment of paracetamol overdose. Clin Toxicol (Phila). 2013;51(6):467-72.
37. Sprivulis PC, Da Silva JA, Jacobs IG, Frazer AR, Jelinek GA. The association between hospital overcrowding and mortality among patients admitted via Western Australian emergency departments. Med J Aust. 2006;184(5):208-12.
38. McCarthy ML. Overcrowding in emergency departments and adverse outcomes. BMJ. 2011;342:d2830.

Figure 1: Treatment groups in NACSTOP trial. ALT = alanine transaminase, AST = aspartate transaminase.

Figure 2. Trial consort diagram

Figure 3: Acetaminophen concentrations in intervention group and control group post initiation of acetylcysteine. N.B. Time zero is approximate time of acetylcysteine initiation (within 2 hours of presentation blood tests). Median acetaminophen concentrations at 20 hours post initiation of acetylcysteine are below limit of detection (<1 mg/L).

Table 1: Demographics and clinical features from NACSTOP enrollments. IQR= Interquartile range, ALT= alanine aminotransferase, AST = aspartate transaminase, INR= International Normalized Ratio

Figure 4: Alanine transaminase i) (ALT) and ii) aspartate transaminase (AST) concentrations in trial patients post initiation of acetylcysteine. N.B. Time zero is approximate time of acetylcysteine initiation (within 2 hours of presentation blood tests).

Table 2: Gastrointestinal symptoms in the intervention and control groups during hospital admission.

Author Manuscript

Table 1: Demographics and clinical features from NACSTOP enrollments. IQR= Interquartile range, ALT= alanine aminotransferase, AST = aspartate transaminase, INR= International Normalized Ratio

	Intervention group (n=50)	Control group (n=50)	P value
Participants enrolled at each site, (n)	Austin (24) Dandenong (18) Westmead (8)	Monash Medical (23) Casey (19) Blacktown (8)	
Median age (IQR), years	23 (19,38)	22 (18,32)	NS
Female	37 (74%)	40 (80%)	NS
Time from ingestion to treatment, <8hr	27 (54%)	30 (60%)	NS
Alcohol co-ingested	3 (6%)	3 (6%)	NS
Other drugs co-ingested	21 (42%)	13 (26%)	
Opiates	8	3	
Antihistamines	2	1	
Median presentation ALT (IQR), IU/L	19 (14,25)	18 (13,22)	NS
Median presentation AST (IQR), IU/L	21 (16,28)	21 (16,28)	NS
Median 20 hour post acetylcysteine ALT (IQR), IU/L	18 (13,22)	16 (13,21)	NS
Median 20 hour post acetylcysteine AST (IQR), IU/L	18 (14,20)	17 (14,21)	NS
Median 20 hour post acetylcysteine INR, (IQR)	1.2 (1.1,1.3)	1.2 (1.1,1.3)	NS
Ingestion Type			
Single	49 (98%)	46 (92%)	
Staggered	1 (2%)	4 (8%)	

Median dose ingested acetaminophen (IQR), mg/kg	250 (180,385)	269 (198,346)	NS
Median presentation acetaminophen concentration (IQR), mg/L	124 (58,171)	146 (66,204)	NS
Presentation acetaminophen concentration above 200 mg/L (%)	9 (18%)	11 (22%)	NS
Acetaminophen concentration fall >90% from presentation to 12 hour post acetylcysteine initiation tests, N (%)	48 (96%)	47 (94%)	NS
Median acetaminophen concentration 12 hours post initiation of acetylcysteine (IQR), mg/L	2 (<1,7)	<1 (<1, 11)	NS
Median acetaminophen concentration 20 hours post initiation of acetylcysteine (IQR), mg/L	< Lower limit of detection	< Lower limit of detection	NS
Median acetylcysteine duration (IQR), hours	13 (13,13.5)	20 (20,20)	<0.001
Median time to acetylcysteine from overdose (IQR), hours	7 (6,12)	7 (6,10)	NS
Median total	250 (250,250)	300 (300,300)	<0.001

acetylcysteine dose during admission (IQR), mg/kg			
Median acetaminophen x ALT (IQR), IU X mg/L			NS
-Presentation	2595 (1323,3680)	2614(4120,10164)	
-After 12 hours acetylcysteine	109 (46,160)	145 (98,230)	
Median ψ parameter (IQR), mmol/L x hour	0.499 (0.001,1.328)	0.465 (0.001,1.124)	NS
Median Length of stay for medical treatment (ie. excluding psychiatric admission) (IQR), days	1 (1,1)	1 (1,1)	NS
Discharge disposition post acetylcysteine			
-Home, n	45 (90%)	37 (74%)	
-Admitted psychiatric ward, n	5 (10%)	13 (26%)	

NS not significant ($P>0.05$)

Table 2: Gastrointestinal symptoms in the intervention and control groups during hospital admission.

Gastrointestinal symptoms	Intervention group (n=14 or 28%)	Control group (n=12 or 24%)
Nausea	7 (14%)	3 (6%)

Vomiting	5 (10%)	7 (14%)
Both nausea and vomiting	2 (4%)	2 (4%)

Author Manuscript

Single or staggered acetaminophen overdose requiring treatment

Presentation and 12 hours post acetylcysteine initiation bloods – Acetaminophen level, ALT, AST, Creatinine.

Meets inclusion criteria: Requiring acetylcysteine, ALT <40 IU/L at presentation and at 12 hours of acetylcysteine, normal creatinine and acetaminophen concentration < 20 mg/L (132 micromol/L) at 12 hours of acetylcysteine.

Depending on hospital assignment patients were enrolled into either:

Control group

Acetylcysteine regimen:

- 200mg/kg over 4 hours, then
- 100mg/kg over 16 hours IV

Intervention group

Acetylcysteine regimen:

- 200mg/kg over 4 hours, then
- 50mg/kg over 8 hours IV, followed by
- 1 L Ringer's lactate IV over 8 hours

Both groups: repeat ALT, AST, Acetaminophen concentration, INR, Creatinine at 20 hours post infusions

- If ALT <40 IU/L and Creatinine normal, no further treatment
- If abnormal, then continue or reinstitute acetylcysteine (100mg/kg over 16 hours)

Those excluded at 12 hours of acetylcysteine infusion had repeat ALT/AST/INR/Creatinine/Acetaminophen taken at end of 20 hour infusion and twice daily if requiring further treatment.

hep_30224_f1.tiff

hep_30224_f2.tiff

hep_30224_f3.tiff

hep_30224_f4.tiff

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Wong, A;McNulty, R;Taylor, D;Sivilotti, M;Greene, S;Gunja, N;Koutsogiannis, Z;Graudins, A

Title:

The NACSTOP Trial: A Multicenter, Cluster-Controlled Trial of Early Cessation of Acetylcysteine in Acetaminophen Overdose

Date:

2019-02-01

Citation:

Wong, A., McNulty, R., Taylor, D., Sivilotti, M., Greene, S., Gunja, N., Koutsogiannis, Z. & Graudins, A. (2019). The NACSTOP Trial: A Multicenter, Cluster-Controlled Trial of Early Cessation of Acetylcysteine in Acetaminophen Overdose. HEPATOLOGY, 69 (2), pp.774-784. <https://doi.org/10.1002/hep.30224>.

Persistent Link:

<http://hdl.handle.net/11343/285302>