Lavoie Suzie (Orcid ID: 0000-0001-8372-2399)

Lavoie-EEG in Major Psychiatric Disorders

Title: Staging model in psychiatry: review of the evolution of electroencephalography abnormalities in major psychiatric disorders

Authors: Suzie Lavoie^{1,2}, Andrea R Polari^{1,3}, Sherilyn Goldstone^{1,2, 1}, Barnaby Nelson^{1,2}, Patrick McGorry^{1,2}

1 Orygen, The National Centre of Excellence in Youth Mental Health, Parkville, Australia

2 Centre for Youth Mental Health, The University of Melbourne, Parkville, Australia

3 Orygen Youth Health, Melbourne Health, Parkville, Australia

Corresponding author:

Suzie Lavoie
Orygen
35 Poplar road
Parkville Victoria 3052
Australia
Email: suzie.lavoie@orygen.org.au
Phone: 1300 679 436

Running title: EEG in major psychiatric disorders

¹ Author's present address: NHMRC Clinical Trials Centre, University of Sydney, Sydney, Australia

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/eip.12792](https://doi.org/10.1111/eip.12792)

Abstract

Aim: Clinical staging in psychiatry aims to classify patients according to the severity of their symptoms, from stage 0 (increased risk, asymptomatic) to stage 4 (severe illness), enabling adapted treatment at each stage of the illness. The staging model would gain specificity if one or more quantifiable biological markers could be identified. Several biomarkers reflecting possible causal mechanisms and/or consequences of the pathophysiology are candidates for integration into the clinical staging model of psychiatric illnesses. **Methods:** This review covers the evolution (from stage 0 to stage 4) of the most important brain functioning impairments as measured with electroencephalography (EEG), in psychosis spectrum and in severe mood disorders. **Results:** The present review of the literature demonstrates that it is currently not possible to draw any conclusion with regard to the state or trait character of any of the EEG impairments in both major depressive disorder and bipolar disorder. As for schizophrenia, the most promising markers of the stage of the illness are the pitch mismatch negativity as well as the p300 event-related potentials, as these components seem to deteriorate with increasing severity of the illness. **Conclusions:** Given the complexity of major psychiatric disorders, and that not a single impairment can be observed in all patients, future research should most likely consider combinations of markers in the quest for a better identification of the stages of the psychiatric illnesses.

Key Words: Bipolar Disorder, First episode, High risk, Major Depressive Disorder, Psychosis,

Introduction

It is widely believed that intervening early in major psychiatric disorders can lead to significantly better clinical and functional outcome for those affected (Bertelsen et al., 2008; Hegelstad et al., 2012; McGorry, Nelson, Goldstone, & Yung, 2010; Norman et al., 2011). Clinical staging in psychiatry aims to classify patients according to their severity of the symptoms, enabling adapted treatment at every stage of the illness. Most particularly, the staging model suggests that more benign treatment can be used in the case of help-seeking individuals presenting with non-specific symptoms, typically observed in the early phases of psychiatric disorders. The clinical staging model for psychotic and severe mood disorders classifies patients according to the severity of their clinical phenotype, from stage 0 (increased risk, asymptomatic) to stage 4 (severe illness; McGorry et al., 2010). However, the use of the staging model in general medicine (e.g., cancer, arthritis) not only relies on the clinical presentation of the patient, but also on measurable biomarkers of their illness. Several biomarkers reflecting causal mechanisms and/or consequences of the pathophysiology are candidates for integration into the clinical staging model of psychiatric illnesses. This review will cover the evolution (from stage 0 to stage 4) of the most important brain functioning impairments, as measured with electroencephalography (EEG) in psychosis spectrum and in severe mood disorders.

EEG measures voltage fluctuations resulting from ionic current flows within the neurons of the brain. Electrodes are used to record the summation of electrical dipoles

along the scalp. EEG has been used to measure the activity of the resting brain during sleep and wakefulness, as well as the brain's response to stimuli. At rest, scalp EEG activity shows oscillations at a variety of frequencies. Several of these oscillations have characteristic frequency ranges and spatial distributions, and are associated with different states of brain functioning. Following the presentation of stimuli, averaging the EEG activity time-locked to the onset of the presentation leads to a stereotyped electrophysiological response called an event-related potential (ERP). ERP waveforms consist of a series of positive and negative voltage deflections, and the amplitude of these waves depends on the strength of the underlying generators. Abnormalities in both resting state EEG and ERPs have been observed in schizophrenia, major depressive disorders (MDD) and bipolar disorder (BD). The aim of this literature review is to provide an overview of the value of various EEG components impairment as markers of major psychiatric illness stages, with the view to help inform treatment.

Methods

Pubmed database searches were conducted using the following criteria: resting EEG [or] gamma-phase synchrony [or] P50 [or] mismatch negativity [or] P300 [and] early stage [or] high risk [or] first episode [and] psychosis [or] schizophrenia [or] bipolar disorder [or] major depressive disorder. If critical literature reviews were available for stages 3 and 4 (recurrence or relapse of symptoms and severe, persistent or unremitting illness) of any of the illnesses, they are presented in the present work. For the early stages, i.e. high risk and first episode, original researches are presented.

The research was conducted in accordance with the Helsinki Declaration.

Resting EEG

For the measurement of resting brain activity, participants are instructed to sit quietly and comfortably for a few minutes with their eyes open, then a few minutes with their eyes closed. The raw EEG data recorded is subsequently Fourier-decomposed into a voltage (power) by frequency spectral graphing. Each defined frequency band, i.e. delta (<4 Hz), theta (4–7 Hz), alpha (8–15 Hz), beta (16–31 Hz) and gamma (>32 Hz) is associated with a variety of physiological functions, and the power calculated for each of these frequency bands is compared between studied groups in research settings.

No single pattern of quantitative EEG abnormalities has been identified in schizophrenia. The most consistent observation obtained from spectral analysis in both medicated and unmedicated patients suffering from schizophrenia is the increased activity in the slow wave bands (delta and theta), mainly in the frontal area (See meta-analyses: Boutros et al., 2008; Galderisi, Mucci, Volpe, & Boutros, 2009), consistent with the decrease in the frontal brain activity observed in schizophrenia. Activity in the delta and theta bands is associated with slow wave sleep (stages 3 and 4) and is potentially pathological in the waking state. It may therefore indicate some brain pathology in schizophrenia (Y. H. Chen et al., 2015). Chronicity has a significant effect on both the delta and theta bands, with the effect size of the difference between patients and controls being much higher in chronic than in first-episode psychosis (FEP) patients and intermediate in mixed samples (Galderisi et al., 2009). EEG spectral power analyses performed in individuals at ultra-high risk (UHR) of psychosis showed no

difference when compared to a control population and did not predict transition to psychosis (S. Lavoie et al., 2012; Ranlund et al., 2014; Zimmermann et al., 2010). The increase in slow wave activity has been associated with more severe negative symptoms in schizophrenia (Gattaz, Mayer, Ziegler, Platz, & Gasser, 1992; Gerez & Tello, 1995; Gross, Joutsiniemi, Rimon, & Appelberg, 2006; Sponheim, Clementz, Iacono, & Beiser, 2000; Venables, Bernat, & Sponheim, 2009) in FEP (Gschwandtner, Zimmermann, Pflueger, Riecher-Rossler, & Fuhr, 2009) and in UHR (S. Lavoie et al., 2012; Zimmermann et al., 2010), making this observation a fairly robust marker of the illness. It is however, not clear what this association means in terms of the staging model, as negative symptoms exist across all stages (Carrion, Correll, Auther, & Cornblatt, 2017). The worsening of negative symptoms with duration of illness and as secondary effect of treatment is likely to make the association more robust in later stages.

Studies have reported large decrease in alpha activity in drug-free BD euthymic patients (Basar et al., 2012; Kano, Nakamura, Matsuoka, Iida, & Nakajima, 1992; Ozerdem, Guntekin, Tunca, & Basar, 2008) compared to healthy controls, whereas in schizophrenia, spontaneous alpha activity is only slightly reduced (Basar et al., 2016). In BP patients with acute hypomania or depression, alpha activity was rather increased (Moeini, Khaleghi, & Mohammadi, 2015) and the authors concluded that the increased alpha power, which corresponds to a decrease in the thalamic metabolism leading to diminished attention, is consistent with BD presentation. Another group demonstrated increased alpha activity in BD patients, however the current state of the patient at the

time of EEG recording was not specified (Narayanan et al., 2014). In this study, all BD patients presented with psychotic features, and their EEG profile (increased delta, theta and alpha) matched that of the schizophrenia group. It could therefore be hypothesized that resting EEG activity is a reflection of the psychotic symptoms rather than a specific marker of psychiatric illness. Indeed, in a heterogeneous BD group with only nine out of 30 patients showing psychotic symptoms, no resting EEG impairment was observed (Venables et al., 2009). In first-episode BP patients with psychotic features, increase low-frequency and alpha power was observed (Clementz, Sponheim, Iacono, & Beiser, 1994), while only the increase in alpha band was reported in first-episode BP patients in acute hypomania or depressive state (Moeini et al., 2015). However, in light of the results available, it could be that the increased slow activity is specific to psychotic symptoms rather than schizophrenia itself.

In patients with MDD, higher alpha power with the eyes closed and greater alpha suppression with the eyes open were observed when compared to healthy controls (for reviews see Olbrich & Arns, 2013; Pollock & Schneider, 1990). An increase in alpha power is thought to reflect a relative decrease in the proportion of local cortical neurons engaged in a particular task performance. Impaired alpha activity is considered an endophenotype for MDD, mediating the pathway between the brain-derived neurotrophic factor BDNF Val66Met polymorphism and depressed mood (Gatt et al., 2008; Zoon et al., 2013). However, the most reproducible resting EEG observation in MDD is the increased frontal alpha asymmetry (See review by Thibodeau, Jorgensen, & Kim, 2006). Indeed, this meta-analytic review demonstrates that studies on resting EEG

in depression show relative right-sided resting frontal alpha asymmetry among adults with MDD, but also in infants of afflicted mothers. While there are conflicting findings, it appears that alpha asymmetry may be a marker for developing a depressive disorder. Indeed, alpha asymmetry indicative of relatively less right than left parieto-temporal activity has been observed in second-generation (Bruder et al., 2005) and third-generation (Bruder, Tenke, Warner, & Weissman, 2007) offspring of parents and/or grand-parents suffering from depression. These results, including data collected in children without a lifetime history of depression, support the hypothesis that right parieto-temporal hypoactivation is an endophenotypic marker of vulnerability to the disease.

Gamma phase synchrony

The synchronous activity of neurons, mediated by oscillations in the gamma-band (30–100 Hz) of the EEG, has been proposed to play an important role in the linking of neurons into cell assemblies that code information in the brain. This synchronisation (phase-locking) has been proposed as a mechanism for the integration, or binding, of activity in distributed neural networks (Peter J. Uhlhaas & Singer, 2006). Research using EEG in both animals and humans have shown that gamma-band synchronisation contributes to variety of cognitive functions including attention and memory (For a review, see Rieder, Rahm, Williams, & Kaiser, 2011). Reduced gamma activity and a lack of lateralisation in response to various stimuli have been repeatedly observed in patients suffering from schizophrenia (For reviews see Lee, Williams, Breakspear, & Gordon, 2003; P. J. Uhlhaas & Singer, 2010). Decreased gamma phase synchrony

relative to a pre-stimulus baseline has also been demonstrated in FEP (Slewa-Younan et al., 2004; Spencer, Salisbury, Shenton, & McCarley, 2008; Symond, Harris, Gordon, & Williams, 2005). In another study with FEP patients, it was shown that the absolute magnitude of gamma synchrony is enhanced, markedly in the left centro-temporal region, when the reference to this baseline period is removed (Flynn et al., 2008). The authors suggested that diminished neural synchronisation to stimuli, regardless of their task relevance, is occurring in an environment of generally excessive synchrony. Tada et al. (2014) investigated the time-course of the auditory gamma-band response in a FEP and in a UHR group and they showed a reduction of early-latency (0–100ms) gamma-band in the FES group and a reduction of late-latency (300–500ms) gamma-band in both UHR and FEP groups. Therefore, the early gamma-band response may be reduced after the onset of psychosis in accordance with the staging model, whereas the late gamma-band response may be impaired before the onset of psychosis and considered an endophenotype. However, these results are not supported by the other gamma-band investigations conducted in UHR to date (Leicht et al., 2016; Perez et al., 2013). In those studies, impairment of early-latency gamma-band was observed in UHR individuals. There is therefore no clear story in respect to staging in schizophrenia and the gamma-band synchrony.

Although the study of gamma-band oscillations in BD has received less attention than in schizophrenia, the same reduced gamma power in response to auditory stimuli has been observed repeatedly (For a review, see Onitsuka, Oribe, & Kanba, 2013).

Therefore, reduced gamma-band activity may not represent an efficient diagnostic tool,

although the mechanisms underpinning this impairment in schizophrenia involves the cortical inhibition by GABA inter-neurons, while this is not the case in BD (Farzan et al., 2010).

In contrast with schizophrenia and BD, MDD individuals displayed an increased gamma-band power sustained during eight seconds following presentation of negative versus neutral words, compared to control participants and to individuals with schizophrenia (Siegle, Condray, Thase, Keshavan, & Steinhauer, 2010). Similarly, in individuals with a first-episode of depression, the gamma-band showed increased cortical activity both at rest and during the performance of an arithmetic counting task and a spatial imagination task (Strelets, Novototsky-Vlasov, & Golikova, 2002).

Event-related potentials

P50

During an auditory dual-click paradigm, a positive polarity on the EEG trace occurs approximately 50ms post-stimulus; this is the so-called P50. The relative decrease of the P50 wave to the second click compared with the first is thought to reflect a sensory gating mechanism, aiming to protect against information overload. The abnormal auditory gating observed in schizophrenia, i.e., a decreased attenuation in the amplitude of the second P50, is a robust candidate endophenotype of the illness (See meta-analyses: Elvira Bramon, Rabe-Hesketh, Sham, Murray, & Frangou, 2004; Patterson et al., 2008). Indeed, a reduced P50 suppression has been linked to several gene loci and was found in healthy relatives of schizophrenia patients (for a review see Turetsky et al.,

2007). A central hypothesis proposed to account for perception and attention deficits in schizophrenia is that these patients cannot inhibit, or “gate,” irrelevant sensory input, leading to an overload of information reaching the brain. Impaired sensory gating has been demonstrated in FEP patients (Devrim-Ucok, Keskin-Ergen, & Ucok, 2008b; Hong et al., 2009), but not by all (de Wilde, Bour, Dingemans, Koelman, & Linszen, 2007; Morales-Munoz et al., 2016). In UHR, the results are also inconsistent with some studies showing impaired P50 (A. Brockhaus-Dumke et al., 2008; Myles-Worsley, Ord, Blalles, Ngiralmu, & Freedman, 2004), while others did not (Cadenhead, Light, Shafer, & Braff, 2005; Hsieh et al., 2012; van Tricht et al., 2015). While P50 reflects pre-attentional gating mechanisms, recent studies have extended the evaluation of sensory gating to later stages, namely N1 and P2, for which deficits have also been observed in UHR, FEP and schizophrenia (A. Brockhaus-Dumke et al., 2008; Morales-Munoz et al., 2016). However, these component, similar to the P50, are rather viewed as endophenotypes of the illness (Hsieh et al., 2012; Morales-Munoz et al., 2016; van Tricht et al., 2015) and are therefore not good candidate as contributors to the staging model in schizophrenia. As a consequence, they are not reviewed any further here.

A recent meta-analysis has shown that sensory gating impairment, as indexed by the decrease in P50 suppression, is present in BP patients and it is likely to worsen with a history of psychosis (Cheng, Chan, Liu, & Hsu, 2016). BP is associated with the decrease in P50 suppression with a high genetic component (Hall, Rijdsdijk, Kalidindi, et al., 2007; Sanchez-Morla et al., 2008) and therefore, P50 suppression impairment is a good candidate for an endophenotype of this mental disease. To our knowledge, only

two studies have looked at the P50 in MDD and they have both demonstrated sensory gating impairment (Kuang, Tian, Yue, & Li, 2016; Y. Wang et al., 2009). Interestingly, impaired P50 sensory gating measured in 40 months old infants was predictive of attention problems and anxious/depressed symptoms three years later (Hutchison et al., 2017), indicative of the neurophysiological dysfunction being present long before the behaviour or symptoms manifestation.

Mismatch negativity

Mismatch negativity (MMN) is a change in the activity of the brain induced by the occurrence of novel stimuli, leading to a switch of attention in the subject (For a review see Naatanen, Kujala, & Winkler, 2011). MMN is thought to reflect the functioning of NMDA receptors (Javitt et al., 1995; Daniel Umbricht, Koller, Vollenweider, & Schmid, 2002). A meta-analysis of 32 studies reports that a decrease in the amplitude of MMN has been consistently replicated in schizophrenia (D. Umbricht & Krljes, 2005). This MMN deficit is a remarkably robust finding, and is one of the most replicable findings in schizophrenia (For a review, see Naatanen, Shiga, Asano, & Yabe, 2015). Impaired MMN has been associated with poor social cognition (Wynn, Sugar, Horan, Kern, & Green, 2010) and poor global functioning (Kawakubo & Kasai, 2006; Light & Braff, 2005a, 2005b; Rasser et al., 2011). Auditory memory encoding deficits in schizophrenia patients could lead to a reduced capacity for short-term storage of verbal information (Kiang et al., 2007). This impairment could in turn interfere not only with abstract thinking, but also with everyday tasks. During a classic Auditory Oddball Paradigm (AOP), MMN is induced by the occurrence of a deviant sound in an otherwise

contiguous stream of events. The most widely used deviant stimuli in schizophrenia research are sounds differing in their duration or pitch frequency.

Frequency MMN (fMMN) illustrates the theoretical concept of the staging model.

Indeed, the Umbricht and Krljes (2005) meta-analysis showed that the effect sizes of fMMN were significantly correlated with the duration of illness, indicating that the fMMN amplitude attenuation could reflect disease progression. While this impairment is well established in chronic schizophrenia and it may be present in FEP (Bodatsch et al., 2011; Oades et al., 2006), most FEP studies show that the fMMN is intact in these patients (Devrim-Ucok, Keskin-Ergen, & Ucok, 2008a; Magno et al., 2008; Mondragon-Maya et al., 2013; D. F. Salisbury, Kuroki, Kasai, Shenton, & McCarley, 2007; Dean F. Salisbury, Shenton, Griggs, Bonner-Jackson, & McCarley, 2002; Todd et al., 2007; D. S. Umbricht, Bates, Lieberman, Kane, & Javitt, 2006; Valkonen-Korhonen et al., 2003). However, a significant decrease in the fMMN amplitude was observed in the same participants in the post-acute phase (Devrim-Ucok et al., 2008a) or one year after their first psychotic episode (D. F. Salisbury et al., 2007). Worsening of the MMN generation has been correlated with the grey matter volume reduction in the primary auditory cortex, a correlation that was also observed in chronic schizophrenia (Rasser et al., 2011). Very few studies have measured fMMN in young individuals presenting with an at-risk mental state, although the amplitude was shown to be intact compared to that seen in controls in three studies (Bodatsch et al., 2011; Anke Brockhaus-Dumke et al., 2005; Mondragon-Maya et al., 2013), while it was decreased in another study (Perez et al., 2014). Finally, the generation of fMMN was shown to be intact in first-degree

relatives (Magno et al., 2008) and in twins of affected patients (Ahveninen et al., 2006), making it a poor contender as a biomarker for the illness. Altogether, it appears that fMMN impairment worsens with deterioration of illness, and this hypothesis has been recently tested using longitudinal study designs. One study supported the hypothesis showing a significant decline in fMMN following transition (S. Lavoie et al., 2018), while the other did not (Atkinson et al., 2017).

As for the amplitude of the MMN in response to a change in the duration of the sound (dMMN), it is equally attenuated in schizophrenia and FEP patients (Atkinson, Michie, & Schall, 2012; Bodatsch et al., 2011; Hermens et al., 2010; Kaur et al., 2011; Oades et al., 2006; Solis-Vivanco et al., 2014), although this was not reported in all studies (Magno et al., 2008; D. S. Umbricht et al., 2006). Almost all studies conducted in UHR individuals showed a decrease in the dMMN amplitude in these individuals compared to controls (Atkinson et al., 2012; Bodatsch et al., 2011; Higuchi et al., 2014; Jahshan, Cadenhead, et al., 2012; Perez et al., 2014; Shaikh et al., 2012; Shin et al., 2009; Solis-Vivanco et al., 2014), apart from two studies (Higuchi et al., 2013; Mondragon-Maya et al., 2013). Most interestingly, studies where UHR individuals who transitioned to psychosis were compared to UHR individuals who did not transition have revealed that the decrease in dMMN amplitude was significant only in those who transitioned (Atkinson et al., 2012; Bodatsch et al., 2011; Higuchi et al., 2014; Higuchi et al., 2013; Perez et al., 2014; Shaikh et al., 2012), and therefore, this impairment could be used as a predictor of the illness.

dMMN deficit seems to be related to the genetic disposition of the patient. Indeed, a modest but significant genetic correlation between the amplitude of dMMN and schizophrenia has been observed, suggesting that it is a potentially valid endophenotype for this disease (Hall, Rijdsdijk, Picchioni, et al., 2007). On the other hand, it was shown that fMMN can be improved by the administration of N-acetylcysteine in schizophrenia patients (Suzie Lavoie et al., 2007), reinforcing its status as a state rather than a trait marker.

MMN has not been extensively studied in both BD and MDD, and most of the studies on this topic have been conducted in stages 3 or 4 of the illnesses. A meta-analysis including seven studies looking at MMN in BD compared to controls showed that BD patients display a significantly reduced frontal MMN of moderate effect size (Chitty, Lagopoulos, Lee, Hickie, & Hermens, 2013). However, close examination of the results obtained from each study included in the meta-analysis reveals highly inconsistent results. Indeed, only three studies showed a significant decrease in MMN amplitude in BD patients compared to controls (Andersson, Barder, Hellvin, Løvdahl, & Malt, 2008; Jahshan, Wynn, et al., 2012; Kaur et al., 2012), while it was not significantly decreased in the four other studies (Catts et al., 1995; Hall et al., 2009; D. F. Salisbury et al., 2007; Daniel Umbricht et al., 2003). A recent comprehensive review of the literature on MMN in BD, also demonstrates that MMN appears to be perturbed in BD but to a lesser degree than that observed in schizophrenia (Hermens, Chitty, & Kaur, 2017). The heterogeneity, not only of the BD population, but also of the methodology followed by the various studies, can possibly explain these discrepancies. For example, while a

great majority of the studies present results from a mix of bipolar I and bipolar II, regardless of the phase they are in, i.e., manic, depressive or euthymic, other studies only state that they have recruited patients with BP. To our knowledge, the MMN impairment in BD has not been rigorously studied in the early stages of the illness.

Similar to BD patients, no real consensus can be reached regarding MMN impairment in MDD patients. In schizophrenia research, the paradigm used to elicit the MMN and the methods used to measure the induced brain activity have become relatively standardised over the years. However, in MDD populations various methodologies have been used, such as visual paradigms instead of auditory paradigms, magnetoencephalography (MEG) instead of EEG, midline electrodes instead of temporal or fronto-central ones, etc. It appears that duration MMN is decreased in MDD when compared to controls (J. Chen et al., 2014; Naismith et al., 2012; Qiao et al., 2013; Takei et al., 2009), while the amplitude of the fMMN remains the same (Daniel Umbricht et al., 2003) or even increases in some MDD patients (He et al., 2010; Kahkonen et al., 2007). At first sight, nothing striking can differentiate the participants in the various studies and it seems like the MNN amplitude is not related to depressive symptoms (He et al., 2010; Takei et al., 2009).

Until recent years, the MMN impairment was thought to be specific to schizophrenia. However, studies conducted in BD and MDD over the last decade or so have been contradicting this hypothesis, therefore threatening the use of the MMN impairment as a diagnostic tool for schizophrenia.

P300

The P300 (P3) waveform has been conceptualised as an index of endogenous cognitive processes appearing on EEG traces as a large positive component peaking around 300ms post-stimulus. The preferred paradigm to elicit the P3 is an active button-press oddball paradigm. A meta-analysis including 46 studies (Elvira Bramon et al., 2004) and another including 104 studies (Jeon & Polich, 2003) showed large pooled effect sizes of 0.85 and 0.89 for P3 amplitude differences between schizophrenia patients and controls at midline electrodes, and the P3 latency was also significantly delayed. Abnormalities in P3 may reflect a failure to allocate attention resources to a stimulus and/or deficits in information processing and cognitive updating. P3 impairments are also consistently observed in neuroleptic-naïve FEP patients (Brown, Gonsalvez, Harris, Williams, & Gordon, 2002; de Wilde et al., 2008; Demiralp et al., 2002; Hermens et al., 2010; Hirayasu et al., 1998; Kaur et al., 2011; McCarley et al., 2002; Ozgurdal et al., 2008; Renoult et al., 2007; van der Stelt, Lieberman, & Belger, 2005; J. Wang et al., 2009; Xiong et al., 2010) and in individuals at clinical high risk for psychosis (E. Bramon et al., 2008; Frommann et al., 2008; Nieman et al., 2014; Ozgurdal et al., 2008; van der Stelt et al., 2005). Interestingly, it was shown that P300 peak amplitude significantly predicted later improvement in negative and general symptoms in high risk individuals who did not transition to psychosis within two years of EEG recording (Kim, Lee, Lee, Kim, & Kwon, 2015). A meta-analysis including 11 studies on P3 amplitudes in unaffected relatives of schizophrenia patients showed that these healthy relatives have reduced P3 amplitudes and increased latency compared to controls (E. Bramon et al., 2005). Its high heritability

and stability makes it a candidate of choice in the research of an endophenotype (For a review see Turetsky et al., 2007). P3 amplitude is indeed a strong contender as a biomarker for psychosis as demonstrated by a study that used Cox regression analysis to estimate the best predictors for transition to psychosis (Nieman et al., 2014). In their Cox model, poor social-personal adjustment and reduced P3 parietal amplitude at midline site predicted transition to a first psychotic episode with a sensitivity of 88.9% and a specificity 82.5%.

In light of these results, it appears that the P3 impairment, although present before the onset of psychosis, worsens with the establishment of the illness, showing better established localisation of the damaged source and slower processes. A study showing a decrease in P3 amplitude in at-risk, FEP and schizophrenia patients compared to controls, revealed a progressive course from prodromal to chronic schizophrenia in terms of P3 amplitude impairment (Ozgurdal et al., 2008). A relationship with illness duration has been established for the P3 latency increase (Mathalon, Ford, Rosenbloom, & Pfefferbaum, 2000; O'Donnell et al., 1995). However, it is important to note that impaired P3 is not specific to schizophrenia, being impaired in other mental or brain disorders, but with less consistency (For a review see Duncan et al., 2009).

P3 is delayed and its amplitude is reduced in BD (Hall et al., 2009; O'Donnell, Vohs, Hetrick, Carroll, & Shekhar, 2004; D. F. Salisbury, Shenton, & McCarley, 1999; Schulze et al., 2008). Unaffected relatives of BD patients also present with the same P3 impairments (Hall et al., 2009; Pierson, Jouvent, Quintin, Perez-Diaz, & Leboyer, 2000;

Schulze et al., 2008). Similar to P50 suppression, impairments in the generation of the P3 should be considered as a potential biological marker of BD.

Findings of deviant P3 are less consistent in depression than in schizophrenia and BD, and they may be related to patient subtypes or to the severity of the disease (For a review see Duncan et al., 2009). For example, the increase in P3 latency seems to be associated with the presence of melancholic symptoms (Bruder et al., 1991; Kemp et al., 2010; Schlegel, Nieber, Herrmann, & Bakauski, 1991) or with cognitive impairments (Vandoolaeghe, van Hunsel, Nuyten, & Maes, 1998). Melancholia was not only associated with an increased P300 latency, but also with an exaggerated P200 (Kemp et al., 2010). Vandoolaeghe et al. (1998), who observed the same profile of impairments, associated the latter to non-responders to antidepressant therapy. Another proposition is that P3 is normal in depressed individuals, although in families with multiple members with schizophrenia, relatives who suffer from MDD may show the same P3 abnormalities as do their relatives with schizophrenia (Blackwood, St Clair, Muir, & Duffy, 1991). However, when the amplitude of P3 was measured in the offspring of parents suffering from depression, a decrease was observed when compared to the P3 amplitude of control subjects without a family history of psychiatric disorders (Zhang, Hauser, Conty, Emrich, & Dietrich, 2007). These inconsistent results regarding ERPs in major depression do not allow us to draw any conclusion. A given impairment might be genetically linked to the disease and/or related to a given symptom or abnormality observed in major depression.

Conclusion

This review attempted to present some the EEG impairment observed in some major psychiatric illnesses in relation to the stage of the illness. Our first observation is that although these impairments have been extensively studied in all stages of schizophrenia, the results remain highly inconsistent in chronic BD and MDD, and almost nonexistent for the early stages of these disorders. The clinical stages of MDD and BD have been defined (Berk et al., 2007; Hetrick et al., 2008), and EEG studies in the early stages of these illnesses are needed in order to obtain a better understanding of the course of the observed impairments. It is currently not possible to draw any conclusion with regard to the state or trait character of any of the EEG impairments reviewed here in both MDD and BD. In schizophrenia, the most promising markers of the stage of the illness are pitch MMN as well as p300, as these components seem to deteriorate with the increasing severity of illness. However, in the current state of the literature, it is not possible to conclude that any brain dysfunction as indexed by impaired EEG components can be used as a marker of stage for any of the major psychiatric illness stage.

Longitudinal studies will be necessary to follow the course of the observed impairments in at-risk individual who, after experiencing a first-episode of psychosis, are later diagnosed with schizophrenia. Most studies are designed with follow-up periods too short to fully appreciate the changes in the brain functioning as indexed by EEG components. Furthermore, many studies follow up at-risk individuals who later transition to a stage 2 disorder, but rare are the studies that follow-up the patients who are later

diagnosed with a stage 3 or 4 disorder. Differential ascertainment across stages of illness in cross-sectional studies can lead to spurious conclusions about staging.

Finally, and most importantly, if these markers are going to be used as diagnostic tools in the establishment of the stage of a psychotic disorder, a quantifying parameter will need be identified as well as normal and pathological ranges.

The main limitation of the current review of the literature is that it does not include all the measurable EEG components available in the literature. Indeed, time-frequency analyses of the ERPs, visual or somato-sensorial ERPs, parameters such as latency, or later components indexing information processing, have not been included in this review. Source localisation, global field power and principal component analyses have also been ignored. It is therefore possible that another EEG marker may be more efficient at identifying stages of illness.

Given the complexity of major psychiatric disorders, and that not a single impairment can be observed in all patients, future research should most likely consider combinations of markers in the quest for a better identification of the stages of the psychiatric illnesses.

Conflict of interest statement

The authors have no conflict of interest to disclose.

References

- Ahveninen, J., Jaaskelainen, I. P., Osipova, D., Huttunen, M. O., Ilmoniemi, R. J., Kaprio, J., . . . Cannon, T. D. (2006). Inherited auditory-cortical dysfunction in twin pairs discordant for schizophrenia. *Biological psychiatry*, 60(6), 612-620. doi:10.1016/j.biopsych.2006.04.015
- Andersson, S., Barder, H. E., Hellvin, T., Løv Dahl, H., & Malt, U. F. (2008). Neuropsychological and electrophysiological indices of neurocognitive dysfunction in bipolar II disorder. *Bipolar Disorder*, 10, 888-899.
- Atkinson, R. J., Fulham, W. R., Michie, P. T., Ward, P. B., Todd, J., Stain, H., . . . Schall, U. (2017). Electrophysiological, cognitive and clinical profiles of at-risk mental state: The longitudinal Minds in Transition (MinT) study. *PLoS One*, 12(2), e0171657. doi:10.1371/journal.pone.0171657
- Atkinson, R. J., Michie, P. T., & Schall, U. (2012). Duration mismatch negativity and P3a in first-episode psychosis and individuals at ultra-high risk of psychosis. *Biological psychiatry*, 71(2), 98-104. doi:10.1016/j.biopsych.2011.08.023
- Basar, E., Guntekin, B., Atagun, I., Turp Golbasi, B., Tulay, E., & Ozerdem, A. (2012). Brain's alpha activity is highly reduced in euthymic bipolar disorder patients. *Cogn Neurodyn*, 6(1), 11-20. doi:10.1007/s11571-011-9172-y
- Basar, E., Schmiedt-Fehr, C., Mathes, B., Femir, B., Emek-Savas, D. D., Tulay, E., . . . Basar-Eroglu, C. (2016). What does the broken brain say to the neuroscientist? Oscillations and connectivity in schizophrenia, Alzheimer's disease, and bipolar disorder. *Int J Psychophysiol*, 103, 135-148. doi:10.1016/j.ijpsycho.2015.02.004
- Berk, M., Conus, P., Lucas, N., Hallam, K., Malhi, G. S., Dodd, S., . . . McGorry, P. (2007). Setting the stage: from prodrome to treatment resistance in bipolar disorder. *Bipolar Disord*, 9(7), 671-678. doi:10.1111/j.1399-5618.2007.00484.x
- Bertelsen, M., Jeppesen, P., Petersen, L., Thorup, A., Ohlenschlaeger, J., le Quach, P., . . . Nordentoft, M. (2008). Five-year follow-up of a randomized multicenter trial of intensive early intervention vs standard treatment for patients with a first episode of psychotic illness: the OPUS trial. *Arch Gen Psychiatry*, 65(7), 762-771. doi:10.1001/archpsyc.65.7.762
- Blackwood, D. H., St Clair, D. M., Muir, W. J., & Duffy, J. C. (1991). Auditory P300 and eye tracking dysfunction in schizophrenic pedigrees. *Arch Gen Psychiatry*, 48(10), 899-909.
- Bodatsch, M., Ruhrmann, S., Wagner, M., Muller, R., Schultze-Lutter, F., Frommann, I., . . . Brockhaus-Dumke, A. (2011). Prediction of psychosis by mismatch negativity. *Biol Psychiatry*, 69(10), 959-966. doi:S0006-3223(10)01103-0 [pii]
- 10.1016/j.biopsych.2010.09.057
- Boutros, N. N., Arfken, C., Galderisi, S., Warrick, J., Pratt, G., & Iacono, W. (2008). The status of spectral EEG abnormality as a diagnostic test for schizophrenia. *Schizophr Res*, 99(1-3), 225-237. doi:S0920-9964(07)00532-4 [pii]
- 10.1016/j.schres.2007.11.020
- Bramon, E., McDonald, C., Croft, R. J., Landau, S., Filbey, F., Gruzeliier, J. H., . . . Murray, R. M. (2005). Is the P300 wave an endophenotype for schizophrenia? A meta-analysis and a family study. *Neuroimage*, 27(4), 960-968. doi:S1053-8119(05)00326-5 [pii]
- 10.1016/j.neuroimage.2005.05.022
- Bramon, E., Rabe-Hesketh, S., Sham, P., Murray, R. M., & Frangou, S. (2004). Meta-analysis of the P300 and P50 waveforms in schizophrenia. *Schizophrenia Research*, 70(2-3), 315-329.
- Bramon, E., Shaikh, M., Broome, M., Lappin, J., Berge, D., Day, F., . . . McGuire, P. (2008). Abnormal P300 in people with high risk of developing psychosis. *Neuroimage*, 41(2), 553-560. doi:10.1016/j.neuroimage.2007.12.038

- Brockhaus-Dumke, A., Schultze-Lutter, F., Mueller, R., Tendolkar, I., Bechdorf, A., Pukrop, R., . . . Ruhrmann, S. (2008). Sensory gating in schizophrenia: P50 and N100 gating in antipsychotic-free subjects at risk, first-episode, and chronic patients. *Biol Psychiatry*, 64(5), 376-384. doi:S0006-3223(08)00167-4 [pii]
10.1016/j.biopsych.2008.02.006
- Brockhaus-Dumke, A., Tendolkar, I., Pukrop, R., Schultze-Lutter, F., Klosterkötter, J., & Ruhrmann, S. (2005). Impaired mismatch negativity generation in prodromal subjects and patients with schizophrenia. *Schizophrenia Research*, 73(2-3), 297-310.
- Brown, K. J., Gonsalvez, C. J., Harris, A. W. F., Williams, L. M., & Gordon, E. (2002). Target and non-target ERP disturbances in first episode vs. chronic schizophrenia. *Clinical Neurophysiology*, 113(11), 1754-1763.
- Bruder, G. E., Tenke, C. E., Warner, V., Nomura, Y., Grillon, C., Hille, J., . . . Weissman, M. M. (2005). Electroencephalographic measures of regional hemispheric activity in offspring at risk for depressive disorders. *Biol Psychiatry*, 57(4), 328-335. doi:S0006-3223(04)01181-3 [pii]
10.1016/j.biopsych.2004.11.015
- Bruder, G. E., Tenke, C. E., Warner, V., & Weissman, M. M. (2007). Grandchildren at high and low risk for depression differ in EEG measures of regional brain asymmetry. *Biol Psychiatry*, 62(11), 1317-1323. doi:S0006-3223(06)01557-5 [pii]
10.1016/j.biopsych.2006.12.006
- Bruder, G. E., Towey, J. P., Stewart, J. W., Friedman, D., Tenke, C., & Quitkin, F. M. (1991). Event-related potentials in depression: influence of task, stimulus hemifield and clinical features on P3 latency. *Biol Psychiatry*, 30(3), 233-246. doi:0006-3223(91)90108-X [pii]
- Cadenhead, K. S., Light, G. A., Shafer, K. M., & Braff, D. L. (2005). P50 suppression in individuals at risk for schizophrenia: the convergence of clinical, familial, and vulnerability marker risk assessment. *Biol Psychiatry*, 57(12), 1504-1509. doi:10.1016/j.biopsych.2005.03.003
- Carrion, R. E., Correll, C. U., Auther, A. M., & Cornblatt, B. A. (2017). A Severity-Based Clinical Staging Model for the Psychosis Prodrome: Longitudinal Findings From the New York Recognition and Prevention Program. *Schizophr Bull*, 43(1), 64-74. doi:10.1093/schbul/sbw155
- Catts, S. V., Shelley, A. M., Ward, P. B., Liebert, B., McConaghy, N., Andrews, S., & Michie, P. T. (1995). Brain potential evidence for an auditory sensory memory deficit in schizophrenia. *American Journal of Psychiatry*, 152(2), 213-219.
- Chen, J., Zhang, Y., Wei, D., Wu, X., Fu, Q., Xu, F., . . . Zhang, Z. (2014). Neurophysiological handover from MMN to P3a in first-episode and recurrent major depression. *J Affect Disord*, 174C, 173-179. doi:10.1016/j.jad.2014.11.049
- Chen, Y. H., Stone-Howell, B., Edgar, J. C., Huang, M., Wootton, C., Hunter, M. A., . . . Canive, J. M. (2015). Frontal slow-wave activity as a predictor of negative symptoms, cognition and functional capacity in schizophrenia. *Br J Psychiatry*. doi:10.1192/bjp.bp.114.156075
- Cheng, C. H., Chan, P. Y., Liu, C. Y., & Hsu, S. C. (2016). Auditory sensory gating in patients with bipolar disorders: A meta-analysis. *J Affect Disord*, 203, 199-203. doi:10.1016/j.jad.2016.06.010
- Chitty, K. M., Lagopoulos, J., Lee, R. S., Hickie, I. B., & Hermens, D. F. (2013). A systematic review and meta-analysis of proton magnetic resonance spectroscopy and mismatch negativity in bipolar disorder. *Eur Neuropsychopharmacol*, 23(11), 1348-1363. doi:10.1016/j.euroneuro.2013.07.007
- Clementz, B. A., Sponheim, S. R., Iacono, W. G., & Beiser, M. (1994). Resting EEG in first-episode schizophrenia patients, bipolar psychosis patients, and their first-degree relatives. *Psychophysiology*, 31(5), 486-494.
- de Wilde, O. M., Bour, L. J., Dingemans, P. M., Koelman, J. H., Boeree, T., & Linszen, D. H. (2008). P300 deficits are present in young first-episode patients with schizophrenia and not in their healthy young siblings. *Clin Neurophysiol*, 119(12), 2721-2726. doi:S1388-2457(08)00973-5 [pii]
10.1016/j.clinph.2008.08.024

- de Wilde, O. M., Bour, L. J., Dingemans, P. M., Koelman, J. H., & Linszen, D. H. (2007). Failure to find P50 suppression deficits in young first-episode patients with schizophrenia and clinically unaffected siblings. *Schizophr Bull*, 33(6), 1319-1323. doi:sbm001 [pii]
- 10.1093/schbul/sbm001
- Demiralp, T., Ucok, A., Devrim, M., Isoglu-Alkac, U., Tecer, A., & Polich, J. (2002). N2 and P3 components of event-related potential in first-episode schizophrenic patients: scalp topography, medication, and latency effects. *Psychiatry Res*, 111(2-3), 167-179. doi:S0165178102001336 [pii]
- Devrim-Ucok, M., Keskin-Ergen, H. Y., & Ucok, A. (2008a). Mismatch negativity at acute and post-acute phases of first-episode schizophrenia. *Eur Arch Psychiatry Clin Neurosci*, 258(3), 179-185. doi:10.1007/s00406-007-0772-9
- Devrim-Ucok, M., Keskin-Ergen, H. Y., & Ucok, A. (2008b). P50 gating at acute and post-acute phases of first-episode schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry*, 32(8), 1952-1956. doi:S0278-5846(08)00296-0 [pii]
- 10.1016/j.pnpbp.2008.09.018
- Duncan, C. C., Barry, R. J., Connolly, J. F., Fischer, C., Michie, P. T., Naatanen, R., . . . Van Petten, C. (2009). Event-related potentials in clinical research: guidelines for eliciting, recording, and quantifying mismatch negativity, P300, and N400. *Clin Neurophysiol*, 120(11), 1883-1908. doi:S1388-2457(09)00518-5 [pii]
- 10.1016/j.clinph.2009.07.045
- Farzan, F., Barr, M. S., Levinson, A. J., Chen, R., Wong, W., Fitzgerald, P. B., & Daskalakis, Z. J. (2010). Evidence for gamma inhibition deficits in the dorsolateral prefrontal cortex of patients with schizophrenia. *Brain*, 133(Pt 5), 1505-1514. doi:10.1093/brain/awq046
- Flynn, G., Alexander, D., Harris, A., Whitford, T., Wong, W., Galletly, C., . . . Williams, L. M. (2008). Increased absolute magnitude of gamma synchrony in first-episode psychosis. *Schizophr Res*, 105(1-3), 262-271. doi:S0920-9964(08)00267-3 [pii]
- 10.1016/j.schres.2008.05.029
- Frommann, I., Brinkmeyer, J., Ruhrmann, S., Hack, E., Brockhaus-Dumke, A., Bechdorf, A., . . . Wagner, M. (2008). Auditory P300 in individuals clinically at risk for psychosis. *Int J Psychophysiol*, 70(3), 192-205. doi:S0167-8760(08)00751-4 [pii]
- 10.1016/j.ijpsycho.2008.07.003
- Galderisi, S., Mucci, A., Volpe, U., & Boutros, N. (2009). Evidence-based medicine and electrophysiology in schizophrenia. *Clin EEG Neurosci*, 40(2), 62-77.
- Gatt, J. M., Kuan, S. A., Dobson-Stone, C., Paul, R. H., Joffe, R. T., Kemp, A. H., . . . Williams, L. M. (2008). Association between BDNF Val66Met polymorphism and trait depression is mediated via resting EEG alpha band activity. *Biol Psychol*, 79(2), 275-284. doi:10.1016/j.biopsycho.2008.07.004
- Gattaz, W. F., Mayer, S., Ziegler, P., Platz, M., & Gasser, T. (1992). Hypofrontality on topographic EEG in schizophrenia. Correlations with neuropsychological and psychopathological parameters. *Eur Arch Psychiatry Clin Neurosci*, 241(6), 328-332.
- Gerez, M., & Tello, A. (1995). Selected quantitative EEG (QEEG) and event-related potential (ERP) variables as discriminators for positive and negative schizophrenia. *Biol Psychiatry*, 38(1), 34-49. doi:10.1016/0006-3223(94)00205-H
- Gross, A., Joutsiniemi, S. L., Rimon, R., & Appelberg, B. (2006). Correlation of symptom clusters of schizophrenia with absolute powers of main frequency bands in quantitative EEG. *Behav Brain Funct*, 2, 23. doi:10.1186/1744-9081-2-23
- Gschwandtner, U., Zimmermann, R., Pflueger, M. O., Riecher-Rossler, A., & Fuhr, P. (2009). Negative symptoms in neuroleptic-naïve patients with first-episode psychosis correlate with QEEG parameters. *Schizophr Res*, 115(2-3), 231-236. doi:S0920-9964(09)00284-9 [pii]
- 10.1016/j.schres.2009.06.013

- Hall, M. H., Rijdsdijk, F., Kalidindi, S., Schulze, K., Kravariti, E., Kane, F., . . . Murray, R. M. (2007). Genetic overlap between bipolar illness and event-related potentials. *Psychol Med*, 37(5), 667-678. doi:S003329170600972X [pii]
- 10.1017/S003329170600972X
- Hall, M. H., Rijdsdijk, F., Picchioni, M., Schulze, K., Ettinger, U., Touloupoulou, T., . . . Sham, P. (2007). Substantial shared genetic influences on schizophrenia and event-related potentials. *Am J Psychiatry*, 164(5), 804-812. doi:164/5/804 [pii]
- 10.1176/appi.ajp.164.5.804
- Hall, M. H., Schulze, K., Rijdsdijk, F., Kalidindi, S., McDonald, C., Bramon, E., . . . Sham, P. (2009). Are auditory P300 and duration MMN heritable and putative endophenotypes of psychotic bipolar disorder? A Maudsley Bipolar Twin and Family Study. *Psychol Med*, 39(8), 1277-1287. doi:S0033291709005261 [pii]
- 10.1017/S0033291709005261
- He, W., Chai, H., Zheng, L., Yu, W., Chen, W., Li, J., & Wang, W. (2010). Mismatch negativity in treatment-resistant depression and borderline personality disorder. *Progress in neuro-psychopharmacology & biological psychiatry*, 34(2), 366-371. doi:10.1016/j.pnpbp.2009.12.021
- Hegelstad, W. T., Larsen, T. K., Auestad, B., Evensen, J., Haahr, U., Joa, I., . . . McGlashan, T. (2012). Long-term follow-up of the TIPS early detection in psychosis study: effects on 10-year outcome. *Am J Psychiatry*, 169(4), 374-380. doi:10.1176/appi.ajp.2011.11030459
- Hermens, D. F., Chitty, K. M., & Kaur, M. (2017). Mismatch negativity in bipolar disorder: A neurophysiological biomarker of intermediate effect? *Schizophr Res*. doi:10.1016/j.schres.2017.04.026
- Hermens, D. F., Ward, P. B., Hodge, M. A., Kaur, M., Naismith, S. L., & Hickie, I. B. (2010). Impaired MMN/P3a complex in first-episode psychosis: cognitive and psychosocial associations. *Prog Neuropsychopharmacol Biol Psychiatry*, 34(6), 822-829. doi:S0278-5846(10)00112-0 [pii]
- 10.1016/j.pnpbp.2010.03.019
- Hetrick, S. E., Parker, A. G., Hickie, I. B., Purcell, R., Yung, A. R., & McGorry, P. D. (2008). Early identification and intervention in depressive disorders: towards a clinical staging model. *Psychother Psychosom*, 77(5), 263-270. doi:000140085 [pii]
- 10.1159/000140085
- Higuchi, Y., Seo, T., Miyanishi, T., Kawasaki, Y., Suzuki, M., & Sumiyoshi, T. (2014). Mismatch negativity and p3a/reorienting complex in subjects with schizophrenia or at-risk mental state. *Front Behav Neurosci*, 8, 172. doi:10.3389/fnbeh.2014.00172
- Higuchi, Y., Sumiyoshi, T., Seo, T., Miyanishi, T., Kawasaki, Y., & Suzuki, M. (2013). Mismatch negativity and cognitive performance for the prediction of psychosis in subjects with at-risk mental state. *PLoS One*, 8(1), e54080. doi:10.1371/journal.pone.0054080
- Hirayasu, Y., Asato, N., Ohta, H., Hokama, H., Arakaki, H., & Ogura, C. (1998). Abnormalities of auditory event-related potentials in schizophrenia prior to treatment. *Biological Psychiatry*, 43, 244-253.
- Hong, X., Chan, R. C., Zhuang, X., Jiang, T., Wan, X., Wang, J., . . . Weng, B. (2009). Neuroleptic effects on P50 sensory gating in patients with first-episode never-medicated schizophrenia. *Schizophr Res*, 108(1-3), 151-157. doi:S0920-9964(08)00519-7 [pii]
- 10.1016/j.schres.2008.11.016
- Hsieh, M. H., Shan, J. C., Huang, W. L., Cheng, W. C., Chiu, M. J., Jaw, F. S., . . . Liu, C. C. (2012). Auditory event-related potential of subjects with suspected pre-psychotic state and first-episode psychosis. *Schizophr Res*, 140(1-3), 243-249. doi:10.1016/j.schres.2012.06.021
- Hutchison, A. K., Hunter, S. K., Wagner, B. D., Calvin, E. A., Zerbe, G. O., & Ross, R. G. (2017). Diminished Infant P50 Sensory Gating Predicts Increased 40-Month-Old Attention, Anxiety/Depression, and Externalizing Symptoms. *J Atten Disord*, 21(3), 209-218. doi:10.1177/1087054713488824

- Jahshan, C., Cadenhead, K. S., Rissling, A. J., Kirihaara, K., Braff, D. L., & Light, G. A. (2012). Automatic sensory information processing abnormalities across the illness course of schizophrenia. *Psychological Medicine*, 42(1), 85-97. doi:10.1017/S0033291711001061
- Jahshan, C., Wynn, J. K., Mathis, K. I., Altshuler, L. L., Glahn, D. C., & Green, M. F. (2012). Cross-diagnostic comparison of duration mismatch negativity and P3a in bipolar disorder and schizophrenia. *Bipolar Disorders*, 14(3), 239-248. doi:10.1111/j.1399-5618.2012.01008.x
- Javitt, D. C., Schroeder, C. E., Steinschneider, M., Arezzo, J. C., Ritter, W., & Vaughan, H. G., Jr. (1995). Cognitive event-related potentials in human and non-human primates: implications for the PCP/NMDA model of schizophrenia. *Electroencephalogr Clin Neurophysiol Suppl*, 44, 161-175.
- Jeon, Y. W., & Polich, J. (2003). Meta-analysis of P300 and schizophrenia: patients, paradigms, and practical implications. *Psychophysiology*, 40(5), 684-701.
- Kahkonen, S., Yamashita, H., Rytala, H., Suominen, K., Ahveninen, J., & Isometsa, E. (2007). Dysfunction in early auditory processing in major depressive disorder revealed by combined MEG and EEG. *J Psychiatry Neurosci*, 32(5), 316-322.
- Kano, K., Nakamura, M., Matsuoka, T., Iida, H., & Nakajima, T. (1992). The topographical features of EEGs in patients with affective disorders. *Electroencephalography and clinical neurophysiology*, 83(2), 124-129.
- Kaur, M., Battisti, R. A., Lagopoulos, J., Ward, P. B., Hickie, I. B., & Hermens, D. F. (2012). Neurophysiological biomarkers support bipolar-spectrum disorders within psychosis cluster. *J Psychiatry Neurosci*, 37(5), 313-321. doi:10.1503/jpn.110081
- Kaur, M., Battisti, R. A., Ward, P. B., Ahmed, A., Hickie, I. B., & Hermens, D. F. (2011). MMN/P3a deficits in first episode psychosis: comparing schizophrenia-spectrum and affective-spectrum subgroups. *Schizophrenia Research*, 130(1-3), 203-209. doi:10.1016/j.schres.2011.03.025
- Kawakubo, Y., & Kasai, K. (2006). Support for an association between mismatch negativity and social functioning in schizophrenia. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 30(7), 1367-1368.
- Kemp, A. H., Pe Benito, L., Quintana, D. S., Clark, C. R., McFarlane, A., Mayur, P., . . . Williams, L. M. (2010). Impact of depression heterogeneity on attention: an auditory oddball event related potential study. *J Affect Disord*, 123(1-3), 202-207. doi:S0165-0327(09)00371-1 [pii]
- 10.1016/j.jad.2009.08.010
- Kiang, M., Light, G. A., Prugh, J., Coulson, S., Braff, D. L., & Kutas, M. (2007). Cognitive, neurophysiological, and functional correlates of proverb interpretation abnormalities in schizophrenia. *J Int Neuropsychol Soc*, 13(4), 653-663. doi:S1355617707070816 [pii]
- 10.1017/S1355617707070816
- Kim, M., Lee, T. Y., Lee, S., Kim, S. N., & Kwon, J. S. (2015). Auditory P300 as a predictor of short-term prognosis in subjects at clinical high risk for psychosis. *Schizophr Res*, 165(2-3), 138-144. doi:10.1016/j.schres.2015.04.033
- Kuang, W., Tian, L., Yue, L., & Li, J. (2016). Effects of escitalopram with a Chinese traditional compound Jiuweizhenxin-keli on mismatch negativity and P50 in patients with major depressive disorders. *Neuropsychiatr Dis Treat*, 12, 1935-1941. doi:10.2147/NDT.S104020
- Lavoie, S., Jack, B. N., Griffiths, O., Ando, A., Amminger, P., Couroupis, A., . . . Whitford, T. J. (2018). Impaired mismatch negativity to frequency deviants in individuals at ultra-high risk for psychosis, and preliminary evidence for further impairment with transition to psychosis. *Schizophr Res*, 191, 95-100. doi:10.1016/j.schres.2017.11.005
- Lavoie, S., Murray, M. M., Deppen, P., Knyazeva, M. G., Berk, M., Boulat, O., . . . Do, K. Q. (2007). Glutathione precursor, N-Acetyl-Cysteine, improves mismatch negativity in schizophrenia patients. *Neuropsychopharmacology*, 33, 2187-2199.
- Lavoie, S., Schafer, M. R., Whitford, T. J., Benninger, F., Feucht, M., Klier, C. M., . . . Amminger, G. P. (2012). Frontal delta power associated with negative symptoms in ultra-high risk individuals who transitioned to psychosis. *Schizophrenia Research*, 138, 206-211. doi:10.1016/j.schres.2012.03.033

- Lee, K. H., Williams, L. M., Breakspear, M., & Gordon, E. (2003). Synchronous gamma activity: a review and contribution to an integrative neuroscience model of schizophrenia. *Brain Res Brain Res Rev*, 41(1), 57-78. doi:S0165017302002205 [pii]
- Leicht, G., Vauth, S., Polomac, N., Andreou, C., Rauh, J., Musmann, M., . . . Mulert, C. (2016). EEG-Informed fMRI Reveals a Disturbed Gamma-Band-Specific Network in Subjects at High Risk for Psychosis. *Schizophr Bull*, 42(1), 239-249. doi:10.1093/schbul/sbv092
- Light, G. A., & Braff, D. L. (2005a). Mismatch Negativity Deficits Are Associated With Poor Functioning in Schizophrenia Patients. *Archives of General Psychiatry*, 62(2), 127-136.
- Light, G. A., & Braff, D. L. (2005b). Stability of Mismatch Negativity Deficits and Their Relationship to Functional Impairments in Chronic Schizophrenia. *American Journal of Psychiatry*, 162(9), 1741-1743.
- Magno, E., Yeap, S., Thakore, J. H., Garavan, H., De Sanctis, P., & Foxe, J. J. (2008). Are auditory-evoked frequency and duration mismatch negativity deficits endophenotypic for schizophrenia? High-density electrical mapping in clinically unaffected first-degree relatives and first-episode and chronic schizophrenia. *Biol Psychiatry*, 64(5), 385-391. doi:S0006-3223(08)00363-6 [pii]
- 10.1016/j.biopsych.2008.03.019
- Mathalon, D. H., Ford, J. M., Rosenbloom, M., & Pfefferbaum, A. (2000). P300 reduction and prolongation with illness duration in schizophrenia. *Biological Psychiatry*, 47, 413-427.
- McCarley, R. W., Salisbury, D. F., Hirayasu, Y., Yurgelun-Todd, D. A., Tohen, M., Zarate, C., . . . Shenton, M. E. (2002). Association between smaller left posterior superior temporal gyrus volume on magnetic resonance imaging and smaller left temporal P300 amplitude in first-episode schizophrenia. *Arch Gen Psychiatry*, 59(4), 321-331. doi:yao20418 [pii]
- McGorry, P. D., Nelson, B., Goldstone, S., & Yung, A. R. (2010). Clinical staging: a heuristic and practical strategy for new research and better health and social outcomes for psychotic and related mood disorders. *Can J Psychiatry*, 55(8), 486-497.
- Moeini, M., Khaleghi, A., & Mohammadi, M. R. (2015). Characteristics of Alpha Band Frequency in Adolescents with Bipolar II Disorder: A Resting-State QEEG Study. *Iran J Psychiatry*, 10(1), 8-12.
- Mondragon-Maya, A., Solis-Vivanco, R., Leon-Ortiz, P., Rodriguez-Agudelo, Y., Yanez-Tellez, G., Bernal-Hernandez, J., . . . de la Fuente-Sandoval, C. (2013). Reduced P3a amplitudes in antipsychotic naive first-episode psychosis patients and individuals at clinical high-risk for psychosis. *Journal of psychiatric research*, 47(6), 755-761. doi:10.1016/j.jpsychires.2012.12.017
- Morales-Munoz, I., Jurado-Barba, R., Fernandez-Guinea, S., Rodriguez-Jimenez, R., Jimenez-Arriero, M. A., Criado, J. R., & Rubio, G. (2016). Sensory Gating Deficits in First-Episode Psychosis: Evidence From Neurophysiology, Psychophysiology, and Neuropsychology. *The Journal of nervous and mental disease*, 204(12), 877-884. doi:10.1097/NMD.0000000000000572
- Myles-Worsley, M., Ord, L., Blaises, F., Ngiralmu, H., & Freedman, R. (2004). P50 sensory gating in adolescents from a pacific island isolate with elevated risk for schizophrenia. *Biol Psychiatry*, 55(7), 663-667. doi:10.1016/j.biopsych.2003.12.006
- Naatanen, R., Kujala, T., & Winkler, I. (2011). Auditory processing that leads to conscious perception: a unique window to central auditory processing opened by the mismatch negativity and related responses. *Psychophysiology*, 48(1), 4-22. doi:10.1111/j.1469-8986.2010.01114.x
- Naatanen, R., Shiga, T., Asano, S., & Yabe, H. (2015). Mismatch negativity (MMN) deficiency: A break-through biomarker in predicting psychosis onset. *Int J Psychophysiol*. doi:10.1016/j.ijpsycho.2014.12.012
- Naismith, S. L., Mowszowski, L., Ward, P. B., Diamond, K., Paradise, M., Kaur, M., . . . Hermens, D. F. (2012). Reduced temporal mismatch negativity in late-life depression: an event-related potential index of cognitive deficit and functional disability? *J Affect Disord*, 138(1-2), 71-78. doi:10.1016/j.jad.2011.12.028
- Narayanan, B., O'Neil, K., Berwise, C., Stevens, M. C., Calhoun, V. D., Clementz, B. A., . . . Pearlson, G. D. (2014). Resting state electroencephalogram oscillatory abnormalities in schizophrenia and psychotic bipolar patients and their relatives from the bipolar and schizophrenia network on intermediate phenotypes study. *Biol Psychiatry*, 76(6), 456-465. doi:10.1016/j.biopsych.2013.12.008

- Nieman, D. H., Ruhrmann, S., Dragt, S., Soen, F., van Tricht, M. J., Koelman, J. H., . . . de Haan, L. (2014). Psychosis prediction: stratification of risk estimation with information-processing and premorbid functioning variables. *Schizophr Bull*, 40(6), 1482-1490. doi:10.1093/schbul/sbt145
- Norman, R. M., Manchanda, R., Malla, A. K., Windell, D., Harricharan, R., & Northcott, S. (2011). Symptom and functional outcomes for a 5 year early intervention program for psychoses. *Schizophr Res*, 129(2-3), 111-115. doi:10.1016/j.schres.2011.04.006
- O'Donnell, B. F., Faux, S. F., McCarley, R. W., Kimble, M. O., Salisbury, D. F., Nestor, P. G., . . . Shenton, M. E. (1995). Increased rate of P300 latency prolongation with age in schizophrenia. Electrophysiological evidence for a neurodegenerative process. *Arch Gen Psychiatry*, 52(7), 544-549.
- O'Donnell, B. F., Vohs, J. L., Hetrick, W. P., Carroll, C. A., & Shekhar, A. (2004). Auditory event-related potential abnormalities in bipolar disorder and schizophrenia. *Int J Psychophysiol*, 53(1), 45-55. doi:10.1016/j.ijpsycho.2004.02.001
- S0167876004000169 [pii]
- Oades, R. D., Wild-Wall, N., Juran, S. A., Sachsse, J., Oknina, L. B., & Ropcke, B. (2006). Auditory change detection in schizophrenia: sources of activity, related neuropsychological function and symptoms in patients with a first episode in adolescence, and patients 14 years after an adolescent illness-onset. *BMC Psychiatry*, 6, 7. doi:10.1186/1471-244X-6-7
- Olbrich, S., & Arns, M. (2013). EEG biomarkers in major depressive disorder: discriminative power and prediction of treatment response. *Int Rev Psychiatry*, 25(5), 604-618. doi:10.3109/09540261.2013.816269
- Onitsuka, T., Oribe, N., & Kanba, S. (2013). Neurophysiological findings in patients with bipolar disorder. *Suppl Clin Neurophysiol*, 62, 197-206.
- Ozerdem, A., Guntekin, B., Tunca, Z., & Basar, E. (2008). Brain oscillatory responses in patients with bipolar disorder manic episode before and after valproate treatment. *Brain Res*, 1235, 98-108. doi:10.1016/j.brainres.2008.06.101
- Ozgurdal, S., Gudlowski, Y., Witthaus, H., Kawohl, W., Uhl, I., Hauser, M., . . . Juckel, G. (2008). Reduction of auditory event-related P300 amplitude in subjects with at-risk mental state for schizophrenia. *Schizophr Res*, 105(1-3), 272-278. doi:S0920-9964(08)00240-5 [pii]
- 10.1016/j.schres.2008.05.017
- Patterson, J. V., Hetrick, W. P., Boutros, N. N., Jin, Y., Sandman, C., Stern, H., . . . Bunney, W. E., Jr. (2008). P50 sensory gating ratios in schizophrenics and controls: a review and data analysis. *Psychiatry Res*, 158(2), 226-247. doi:S0165-1781(07)00032-7 [pii]
- 10.1016/j.psychres.2007.02.009
- Perez, V. B., Roach, B. J., Woods, S. W., Srihari, V. H., McGlashan, T. H., Ford, J. M., & Mathalon, D. H. (2013). Early auditory gamma-band responses in patients at clinical high risk for schizophrenia. *Suppl Clin Neurophysiol*, 62, 147-162.
- Perez, V. B., Woods, S. W., Roach, B. J., Ford, J. M., McGlashan, T. H., Srihari, V. H., & Mathalon, D. H. (2014). Automatic auditory processing deficits in schizophrenia and clinical high-risk patients: forecasting psychosis risk with mismatch negativity. *Biol Psychiatry*, 75(6), 459-469. doi:10.1016/j.biopsycho.2013.07.038
- Pierson, A., Jouvent, R., Quintin, P., Perez-Diaz, F., & Leboyer, M. (2000). Information processing deficits in relatives of manic depressive patients. *Psychol Med*, 30(3), 545-555.
- Pollock, V. E., & Schneider, L. S. (1990). Quantitative, waking EEG research on depression. *Biol Psychiatry*, 27(7), 757-780. doi:0006-3223(90)90591-O [pii]
- Qiao, Z., Yu, Y., Wang, L., Yang, X., Qiu, X., Zhang, C., . . . Yang, Y. (2013). Impaired pre-attentive change detection in major depressive disorder patients revealed by auditory mismatch negativity. *Psychiatry Res*, 211(1), 78-84. doi:10.1016/j.psychresns.2012.07.006

- Ranlund, S., Nottage, J., Shaikh, M., Dutt, A., Constante, M., Walshe, M., . . . Bramon, E. (2014). Resting EEG in psychosis and at-risk populations--a possible endophenotype? *Schizophr Res*, 153(1-3), 96-102. doi:10.1016/j.schres.2013.12.017
- Rasser, P. E., Schall, U., Todd, J., Michie, P. T., Ward, P. B., Johnston, P., . . . Thompson, P. M. (2011). Gray matter deficits, mismatch negativity, and outcomes in schizophrenia. *Schizophr Bull*, 37(1), 131-140. doi:sbp060 [pii]
- 10.1093/schbul/sbp060
- Renoult, L., Prevost, M., Brodeur, M., Lionnet, C., Joober, R., Malla, A., & Debruille, J. B. (2007). P300 asymmetry and positive symptom severity: a study in the early stage of a first episode of psychosis. *Schizophr Res*, 93(1-3), 366-373. doi:S0920-9964(07)00158-2 [pii]
- 10.1016/j.schres.2007.03.024
- Rieder, M. K., Rahm, B., Williams, J. D., & Kaiser, J. (2011). Human gamma-band activity and behavior. *Int J Psychophysiol*, 79(1), 39-48. doi:10.1016/j.ijpsycho.2010.08.010
- Salisbury, D. F., Kuroki, N., Kasai, K., Shenton, M. E., & McCarley, R. W. (2007). Progressive and interrelated functional and structural evidence of post-onset brain reduction in schizophrenia. *Arch.Gen.Psychiatry*, 64(5), 521-529.
- Salisbury, D. F., Shenton, M. E., Griggs, C. B., Bonner-Jackson, A., & McCarley, R. W. (2002). Mismatch Negativity in Chronic Schizophrenia and First-Episode Schizophrenia. *Archives of General Psychiatry*, 59(8), 686-694.
- Salisbury, D. F., Shenton, M. E., & McCarley, R. W. (1999). P300 topography differs in schizophrenia and manic psychosis. *Biol Psychiatry*, 45(1), 98-106. doi:S0006-3223(98)00208-X [pii]
- Sanchez-Morla, E. M., Garcia-Jimenez, M. A., Barabash, A., Martinez-Vizcaino, V., Mena, J., Cabranes-Diaz, J. A., . . . Santos, J. L. (2008). P50 sensory gating deficit is a common marker of vulnerability to bipolar disorder and schizophrenia. *Acta Psychiatr Scand*, 117(4), 313-318. doi:ACP1141 [pii]
- 10.1111/j.1600-0447.2007.01141.x
- Schlegel, S., Nieber, D., Herrmann, C., & Bakauski, E. (1991). Latencies of the P300 component of the auditory event-related potential in depression are related to the Bech-Rafaelsen Melancholia Scale but not to the Hamilton Rating Scale for Depression. *Acta Psychiatr Scand*, 83(6), 438-440.
- Schulze, K. K., Hall, M. H., McDonald, C., Marshall, N., Walshe, M., Murray, R. M., & Bramon, E. (2008). Auditory P300 in patients with bipolar disorder and their unaffected relatives. *Bipolar Disord*, 10(3), 377-386. doi:BDI527 [pii]
- 10.1111/j.1399-5618.2007.00527.x
- Shaikh, M., Valmaggia, L., Broome, M. R., Dutt, A., Lappin, J., Day, F., . . . Bramon, E. (2012). Reduced mismatch negativity predates the onset of psychosis. *Schizophrenia Research*, 134(1), 42-48. doi:10.1016/j.schres.2011.09.022
- Shin, K. S., Kim, J. S., Kang, D. H., Koh, Y., Choi, J. S., O'Donnell, B. F., . . . Kwon, J. S. (2009). Pre-attentive auditory processing in ultra-high-risk for schizophrenia with magnetoencephalography. *Biol Psychiatry*, 65(12), 1071-1078. doi:S0006-3223(08)01630-2 [pii]
- 10.1016/j.biopsycho.2008.12.024
- Siegle, G. J., Condray, R., Thase, M. E., Keshavan, M., & Steinhauer, S. R. (2010). Sustained gamma-band EEG following negative words in depression and schizophrenia. *Int J Psychophysiol*, 75(2), 107-118. doi:10.1016/j.ijpsycho.2008.04.008
- Slewa-Younan, S., Gordon, E., Harris, A. W., Haig, A. R., Brown, K. J., Flor-Henry, P., & Williams, L. M. (2004). Sex differences in functional connectivity in first-episode and chronic schizophrenia patients. *Am J Psychiatry*, 161(9), 1595-1602. doi:10.1176/appi.ajp.161.9.1595
- 161/9/1595 [pii]
- Solis-Vivanco, R., Mondragon-Maya, A., Leon-Ortiz, P., Rodriguez-Agudelo, Y., Cadenhead, K. S., & de la Fuente-Sandoval, C. (2014). Mismatch Negativity reduction in the left cortical regions in first-episode psychosis

- and in individuals at ultra high-risk for psychosis. *Schizophr Res*, 158(1-3), 58-63. doi:10.1016/j.schres.2014.07.009
- Spencer, K. M., Salisbury, D. F., Shenton, M. E., & McCarley, R. W. (2008). Gamma-band auditory steady-state responses are impaired in first episode psychosis. *Biol Psychiatry*, 64(5), 369-375. doi:S0006-3223(08)00268-0 [pii]
- 10.1016/j.biopsych.2008.02.021
- Sponheim, S. R., Clementz, B. A., Iacono, W. G., & Beiser, M. (2000). Clinical and biological concomitants of resting state EEG power abnormalities in schizophrenia. *Biol Psychiatry*, 48(11), 1088-1097. doi:S0006-3223(00)00907-0 [pii]
- Strelets, V. B., Novototsky-Vlasov, V. y., & Golikova, J. V. (2002). Cortical connectivity in high frequency beta-rhythm in schizophrenics with positive and negative symptoms. *International Journal of Psychophysiology*, 44, 101-115.
- Symond, M. P., Harris, A. W., Gordon, E., & Williams, L. M. (2005). "Gamma synchrony" in first-episode schizophrenia: a disorder of temporal connectivity? *Am J Psychiatry*, 162(3), 459-465. doi:162/3/459 [pii]
- 10.1176/appi.ajp.162.3.459
- Tada, M., Nagai, T., Kirihaara, K., Koike, S., Suga, M., Araki, T., . . . Kasai, K. (2014). Differential Alterations of Auditory Gamma Oscillatory Responses Between Pre-onset High-risk Individuals and First-episode Schizophrenia. *Cereb Cortex*. doi:10.1093/cercor/bhu278
- Takei, Y., Kumano, S., Hattori, S., Uehara, T., Kawakubo, Y., Kasai, K., . . . Mikuni, M. (2009). Preattentive dysfunction in major depression: a magnetoencephalography study using auditory mismatch negativity. *Psychophysiology*, 46(1), 52-61. doi:PSYP748 [pii]
- 10.1111/j.1469-8986.2008.00748.x
- Thibodeau, R., Jorgensen, R. S., & Kim, S. (2006). Depression, anxiety, and resting frontal EEG asymmetry: a meta-analytic review. *J Abnorm Psychol*, 115(4), 715-729. doi:10.1037/0021-843X.115.4.715
- Todd, J., Michie, P. T., Schall, U., Karayanidis, F., Yabe, H., & Naatanen, R. (2007). Deviant Matters: Duration, Frequency, and Intensity Deviants Reveal Different Patterns of Mismatch Negativity Reduction in Early and Late Schizophrenia. *Biol.Psychiatry*.
- Turetsky, B. I., Calkins, M. E., Light, G. A., Olincy, A., Radant, A. D., & Swerdlow, N. R. (2007). Neurophysiological endophenotypes of schizophrenia: the viability of selected candidate measures. *Schizophr Bull*, 33(1), 69-94. doi:sbl060 [pii]
- 10.1093/schbul/sbl060
- Uhlhaas, P. J., & Singer, W. (2006). Neural Synchrony in Brain Disorders: Relevance for Cognitive Dysfunctions and Pathophysiology. *Neuron*, 52(1), 155-168.
- Uhlhaas, P. J., & Singer, W. (2010). Abnormal neural oscillations and synchrony in schizophrenia. *Nat Rev Neurosci*, 11(2), 100-113. doi:nrn2774 [pii]
- 10.1038/nrn2774
- Umbricht, D., Koller, R., Schmid, L., Skrabo, A., Grubel, C., Huber, T., & Stassen, H. (2003). How specific are deficits in mismatch negativity generation to schizophrenia? *Biological Psychiatry*, 53(12), 1120-1131.
- Umbricht, D., Koller, R., Vollenweider, F. X., & Schmid, L. (2002). Mismatch negativity predicts psychotic experiences induced by nmda receptor antagonist in healthy volunteers. *Biological Psychiatry*, 51(5), 400-406.
- Umbricht, D., & Krljes, S. (2005). Mismatch negativity in schizophrenia: a meta-analysis. *Schizophr.Res.*, 76(1), 1-23.
- Umbricht, D. S., Bates, J. A., Lieberman, J. A., Kane, J. M., & Javitt, D. C. (2006). Electrophysiological indices of automatic and controlled auditory information processing in first-episode, recent-onset and chronic schizophrenia. *Biol Psychiatry*, 59(8), 762-772. doi:S0006-3223(05)01242-4 [pii]
- 10.1016/j.biopsych.2005.08.030

- Valkonen-Korhonen, M., Purhonen, M., Tarkka, I. M., Sipila, P., Partanen, J., Karhu, J., & Lehtonen, J. (2003). Altered auditory processing in acutely psychotic never-medicated first-episode patients. *Cognitive Brain Research*, 17(3), 747-758.
- van der Stelt, O., Lieberman, J. A., & Belger, A. (2005). Auditory P300 in high-risk, recent-onset and chronic schizophrenia. *Schizophr Res*, 77(2-3), 309-320. doi:S0920-9964(05)00173-8 [pii]
10.1016/j.schres.2005.04.024
- van Tricht, M. J., Nieman, D. H., Koelman, J. T., Mensink, A. J., Bour, L. J., van der Meer, J. N., . . . de Haan, L. (2015). Sensory gating in subjects at ultra high risk for developing a psychosis before and after a first psychotic episode. *World J Biol Psychiatry*, 16(1), 12-21. doi:10.3109/15622975.2012.680911
- Vandoolaeghe, E., van Hunsel, F., Nuyten, D., & Maes, M. (1998). Auditory event related potentials in major depression: prolonged P300 latency and increased P200 amplitude. *J Affect Disord*, 48(2-3), 105-113. doi:S0165-0327(97)00165-1 [pii]
- Venables, N. C., Bernat, E. M., & Sponheim, S. R. (2009). Genetic and disorder-specific aspects of resting state EEG abnormalities in schizophrenia. *Schizophr Bull*, 35(4), 826-839. doi:sbn021 [pii]
10.1093/schbul/sbn021
- Wang, J., Tang, Y., Li, C., Mecklinger, A., Xiao, Z., Zhang, M., . . . Li, H. (2009). Decreased P300 current source density in drug-naïve first episode schizophrenics revealed by high density recording. *Int J Psychophysiol*. doi:S0167-8760(09)00295-5 [pii]
10.1016/j.ijpsycho.2009.12.005
- Wang, Y., Fang, Y. R., Chen, X. S., Chen, J., Wu, Z. G., Yuan, C. M., . . . Cao, L. (2009). A follow-up study on features of sensory gating P50 in treatment-resistant depression patients. *Chin Med J (Engl)*, 122(24), 2956-2960.
- Wynn, J. K., Sugar, C., Horan, W. P., Kern, R., & Green, M. F. (2010). Mismatch negativity, social cognition, and functioning in schizophrenia patients. *Biol Psychiatry*, 67(10), 940-947. doi:S0006-3223(09)01417-6 [pii]
10.1016/j.biopsych.2009.11.024
- Xiong, P., Zeng, Y., Zhu, Z., Tan, D., Xu, F., Lu, J., . . . Ma, M. (2010). Reduced NGF serum levels and abnormal P300 event-related potential in first episode schizophrenia. *Schizophr Res*. doi:S0920-9964(10)01158-8 [pii]
10.1016/j.schres.2010.02.1063
- Zhang, Y., Hauser, U., Conty, C., Emrich, H. M., & Dietrich, D. E. (2007). Familial risk for depression and p3b component as a possible neurocognitive vulnerability marker. *Neuropsychobiology*, 55(1), 14-20. doi:000103571 [pii]
10.1159/000103571
- Zimmermann, R., Gschwandtner, U., Wilhelm, F. H., Pflueger, M. O., Riecher-Rossler, A., & Fuhr, P. (2010). EEG spectral power and negative symptoms in at-risk individuals predict transition to psychosis. *Schizophr Res*, 123(2-3), 208-216. doi:S0920-9964(10)01502-1 [pii]
10.1016/j.schres.2010.08.031
- Zoon, H. F., Veth, C. P., Arns, M., Drinkenburg, W. H., Talloen, W., Peeters, P. J., & Kenemans, J. L. (2013). EEG alpha power as an intermediate measure between brain-derived neurotrophic factor Val66Met and depression severity in patients with major depressive disorder. *J Clin Neurophysiol*, 30(3), 261-267. doi:10.1097/WNP.0b013e3182933d6e

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Lavoie, S;Polari, AR;Goldstone, S;Nelson, B;McGorry, PD

Title:

Staging model in psychiatry: Review of the evolution of electroencephalography abnormalities in major psychiatric disorders

Date:

2019-12-01

Citation:

Lavoie, S., Polari, A. R., Goldstone, S., Nelson, B. & McGorry, P. D. (2019). Staging model in psychiatry: Review of the evolution of electroencephalography abnormalities in major psychiatric disorders. *EARLY INTERVENTION IN PSYCHIATRY*, 13 (6), pp.1319-1328. <https://doi.org/10.1111/eip.12792>.

Persistent Link:

<http://hdl.handle.net/11343/285371>