Robertson Oliver (Orcid ID: 0000-0002-4101-5343)

Robertson et al, 2018

Title: Putative neuroprotective pharmacotherapies to target the staged progression of mental illness

Authors:

Oliver D'Arcy Robertson^{1,2}, Nieves Gómez Coronado³, Rickinder Sethi⁴, Michael Berk^{1,2,5,6,7}, Seetal Dodd^{1,2,5,6}

Author affiliations:

¹ Deakin University, Geelong, IMPACT Strategic Research Centre, Victoria, Australia

² University Hospital Geelong, Barwon Health, Geelong, Victoria, Australia

³ Unidad de Gestión Clínica Salud Mental, Hospital Universitario Virgen del Rocio, Sevilla, Spain

⁴ Department of Psychiatry, Western University, London, Ontario, Canada

⁵ Department of Psychiatry, The University of Melbourne, Parkville, Victoria, Australia

⁶ Orygen, the National Centre of Excellence in Youth Mental Health, Parkville, Victoria, Australia

⁷ Florey Institute of Neuroscience and Mental Health, Parkville, Victoria, Australia

Correspondence to: Oliver D'Arcy Robertson.

Email: Oliver.Robertson@BarwonHealth.org.au Address: Barwon Health, PO Box 281, Geelong, Victoria, 3220

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/eip.12775](https://doi.org/10.1111/eip.12775)

Keywords:

Neuroprotection, neuroprogression, bipolar, schizophrenia, neurostructural integrity, oxidative stress, tryptophan, TRYCATs, inflammation, cytokines, depression, mood, antipsychotics, antidepressants, stabilizers, lithium, clozapine, olanzapine, aripiprazole, paliperidone, risperidone, quetiapine, fluoxetine, escitalopram, sertraline, nutritional, supplement, minocycline, aspirin, cyclooxygenase-2 inhibitors, statins, ketamine, alpha-2-delta, pregabalin, gabapentin, erythropoietin, oestrogen, leptin, N-acetylcysteine, curcumin, melatonin, ebselen

Title: 11 words

Abstract: 218 words

Body: 7,708

Abstract

Objective: Neuropsychiatric disorders including depression, bipolar and schizophrenia frequently exhibit a neuroprogressive course from prodrome to chronicity. There are a range of agents exhibiting capacity to attenuate biological mechanisms associated with neuroprogression. This review will update the evidence for putative neuroprotective agents including clinical efficacy, mechanisms of action and limitations in current assessment tools, and identify novel agents with neuroprotective potential.

Method: Data for this review were sourced from online databases PUBMED, Embase and Web of Science. Only data published since 2012 was included in this review, no data was excluded based on language or publication origin.

Results: Each of the agents reviewed inhibit one or multiple pathways of neuroprogression including: inflammatory gene expression and cytokine release, oxidative and nitrosative stress, mitochondrial dysfunction, neurotrophin dysregulation and apoptotic signalling. Some demonstrate clinical efficacy in preventing neural damage or loss, relapse or cognitive/functional decline. Agents include: the psychotropic medications lithium, atypical-antipsychotics and antidepressants; other pharmacological agents such as minocycline, aspirin, cyclooxygenase-2 inhibitors, statins, ketamine and alpha-2-delta ligands; and others such as erythropoietin, oestrogen, leptin, N-acetylcysteine, curcumin, melatonin and ebselen.

Conclusions: Signals of evidence of clinical neuroprotection is evident for a number of candidate agents. Adjunctive use of multiple agents may present a viable avenue to clinical realisation of neuroprotection. Definitive prospective studies of neuroprotection with multimodal assessment tools are required.

Abbreviations

5HT - Serotonin

AA – Arachidonic acid

Akt – Protein Kinase B

AMPK - Adenosine monophosphate activated protein kinase

ADNP - Activity dependent neuroprotective protein

BAD – Bcl-2-associated death promoter

Bcl-2 – B-cell lymphoma 2

BDNF – Brain Derived Neurotrophic Factor

BrdU - 5-bromo-2'-deoxyuridine

cAMP - Cyclic adenosine monophosphate

CA – Cornu Ammonis

CaMKII - Ca²⁺ Calmodulin dependent protein kinase II

COX – Cyclooxygenase

DHA - Docosahexaenoic acid

EPA - Eicosapentaenoic acid

ERK1/2 – Extracellular Signal Regulated Kinases

GSK-3 β - Glycogen synthase kinase 3 beta

GPR30 - G-protein couple receptor 30

IL – Interleukin

IMPase - Inositol monophosphatase

INF α - Interferon alpha

MRI – Magnetic Resonance Imaging

MT₁/MT₂ – Melatonin receptors 1 and 2

mTOR – Mechanistic target of rapamycin

NMDA - N-methyl-D-aspartate

NRF2 - Nuclear transcription factor (erythroid-derived 2)-like 2

NAC – N-acetylcysteine

PET – Positron emission tomography

PGE₂ – Prostaglandin E₂

PI3K - Phosphoinositide 3 kinase

PKA – Protein kinase A

ROS – Reactive oxygen species

SIR – Social isolation rearing

Th1/2 – T-helper cell 1 and 2

TNF – Tumour Necrosis Factor

TRYCAT – Tryptophan catabolite

VBM – Voxel based morphometry

Introduction

Neuropsychiatric disorders including depression, bipolar and schizophrenia manifest a trajectory and course that frequently but not invariably encompasses a prodromal phase, first episode, recurrence and potential chronicity. Beginning even earlier than the prodrome, a cascade of disease processes may begin and lead ultimately to neurostructural and consequent functional changes. This may leave sufferers more vulnerable to relapse, less responsive to treatment and burdened with cognitive impairment and functional decline.

Neuroprogression is a theoretical construct which defines the underpinning elements of this process, characterised by cumulative insults to the brain over a lifetime of neuropsychiatric disease. Neurostructural abnormalities may represent the apex of neuroprogression with multiple convergent pathways contributing to this including: immune challenges, risk genes, inflammation, oxidative stress and mitochondrial dysfunction.

Neurostructural abnormalities

The most tangible evidence for neuroprogressive changes are the alterations of brain structure associated with neuropsychiatric disease. The advent of magnetic resonance imaging (MRI) derived voxel-based morphometry (VBM) has allowed comprehensive analysis of this phenomenon. In first-episode schizophrenia, cortical volume loss is most predictably seen in the anterior cingulate cortex (Brugger & Howes, 2017), while in treatment-resistant schizophrenia the left middle frontal, right precentral and right middle temporal gyri lose cortical volume (Mouchlianitis, McCutcheon, & Howes, 2016). Mood

disorders are associated with complex patterns of grey matter loss. A recent meta-analysis of depression (n=4101) and bipolar disorder (n=2407) showed both conditions are associated with grey matter loss in dorsomedial and ventromedial prefrontal cortex, including the anterior cingulate cortex and bilateral insula (Wise et al., 2017). However, depression was more strongly associated with grey matter loss in the right dorsolateral prefrontal cortex and left hippocampus, as well as, cerebellar, temporal and parietal regions.

White matter volume and integrity, assessed with MRI based diffusion tensor imaging, is also compromised in diverse neuropsychiatric disorders. Regarding overall white matter volume in schizophrenia, findings are inconsistent with studies showing both loss and gain of volume; however, disrupted fibre tract integrity appears to be a consistent feature (Mouchlianitis et al., 2016). In depression and bipolar disorder, white matter integrity is consistently impaired in the genu of the corpus callosum and posterior cingulum fibres (Dong et al., 2017).

The gross neurostructural changes mentioned above may represent the apex of neuroprogression resulting from an imbalance between neurogenesis and neurodegeneration – an imbalance to which dysregulation of neurotrophins may directly contribute. Serum levels of brain derived neurotrophic factor (BDNF), the most commonly assayed neurotrophin, are reduced in patients suffering from bipolar disorder (Munkholm, Vinberg, & Kessing, 2016). Furthermore, a small post-mortem analysis of brain tissue from elderly patients with major depression suggests BDNF and neurotrophin 3 (NT3) are reduced compared to healthy controls (Sheldrick, Camara, Ilieva, Riederer, & Michel, 2017). In schizophrenia, peripheral BDNF levels are lower compared to controls but BDNF levels do not correlate with disease severity (Fernandes et al., 2015). Animal models of neuropsychiatric disease demonstrate aberrant neurotrophin levels, which have been associated with local upregulation of pro-apoptotic pathways such as the B-cell

lymphoma 2 (Bcl2)/Bcl2-associated X protein (Bax)/Caspase 3 pathway; representing an environment favouring neurodegeneration and associated neurostructural abnormalities (Nowakowska, Kus, Ratajczak, Cichocki, & Wozniak, 2014; G. Reus et al., 2012). Directly influencing neurostructural integrity, these pathways are likely triggered by unchecked inflammation as well as redox stress upstream.

Immune challenges, risk genes and inflammation

In the case of schizophrenia, immune challenges in utero may be the first hit in the life course of inflammatory reactions (Davis, Moylan, Harvey, Maes, & Berk, 2014). First hits include seasonality of birth, obstetric complications, maternal infections or metabolic disturbances, which could influence gene expression leading to inflammation and abnormal development. When combined with further hits like trauma and unfavourable lifestyle factors, an increased inflammatory set point may be propagated into later life. This hypothesis was supported in a multicentre epigenome wide analysis of schizophrenic patients versus controls (n=1714) (Hannon et al., 2016). This study identified clusters of DNA methylation, indicating altered gene expression, in regions coding for immunity and neuroplasticity/neurodevelopment.

Meta-analysis of 68-studies provides further evidence for the role of inflammation in mental illnesses (Goldsmith, Rapaport, & Miller, 2016). They demonstrated peripheral inflammatory cytokines interleukin-6 (IL-6) and tumour necrosis factor alpha (TNF α), as well as, soluble cytokine receptor (sIL-2R) and cytokine antagonist (IL-1RA) are significantly increased during acute exacerbations of adult schizophrenia, bipolar disorder and depression; IL-6 is increased in chronically ill patients of all three

conditions; and IL- β and sIL-2R seem elevated only in patients with chronic schizophrenia and euthymic bipolar disorder.

This milieu of circulating inflammatory cytokines may activate and then be propagated by cellular immunity. A small human blood sample (n=60) identified low T-helper cell 1 (Th1)/Th2 ratio and preferential differentiation of macrophages to the pro-inflammatory M1 phenotype in bipolar disorder but not schizophrenia (Brambilla *et al.*, 2014). Others have suggested schizophrenia may be related to irregular Th1/Th2 signalling, though further work is required to confirm this (Davis *et al.*, 2014). Positron emission tomography (PET) imaging can be used to assess activation of microglia, the macrophages of the brain, as a measure of cellular immune activation. In schizophrenia, studies are conflicted with some showing microglial over-activation in the patients with established disease and those at ultra-high risk of psychosis (Selvaraj *et al.*, 2017); however, others suggest microglia are normal or even under active at similar stages of disease (Di Biase *et al.*, 2017; Holmes *et al.*, 2016). Microglia were shown to be over-active in major depression (Holmes *et al.*, 2017). This was associated with cortical volume loss suggesting a relationship between chronic inflammatory states and neuroprogression.

Adding to this positive feedback loop of inflammatory consequences is abnormal tryptophan metabolism. Tryptophan is the essential pre-cursor for serotonin. Intermediates in the tryptophan pathway known as tryptophan catabolites (TRYCATs) are either pro-inflammatory and neurotoxic or anti-inflammatory and neuroprotective (Morris, Carvalho, Anderson, Galecki, & Maes, 2016). Pro-inflammatory and neurotoxic TRYCATs include quinolinic acid (QA) and 3OH-kynurenine (3HK) which are over-produced in an inflammatory milieu such as is seen in depression and schizophrenia, correlating with worsening of negative symptoms including cognition (Kanchanatawan *et al.*, 2017).

Oxidative stress and mitochondrial dysfunction

The activation of immune-inflammatory pathways is tightly linked to oxidative and nitrosative stress (O&NS) and lowered antioxidant defences. Circulating cytokines, macrophage activity and TRYCATs contribute to reactive oxygen species (ROS) and reactive nitrogen species (RNS) production culminating in cellular damage and possibly neuroprogression (Moylan et al., 2014). Meta-analysis of medication naïve depressed patients showed increased oxidative stress with raised malondialdehyde and reduced antioxidant uric acid levels (Jimenez-Fernandez et al., 2015). In bipolar disorder, there is similarly increased lipid peroxidation, DNA/RNA damage and nitric oxide levels (N. C. Brown, Andreazza, & Young, 2014). In schizophrenia, antioxidant defences are weakened, as indicated by decreased super oxide dismutase (Flatow, Buckley, & Miller, 2013). Finally, reduced peripheral glutathione levels, a master anti-oxidant, are associated with symptom severity in bipolar disorder and schizophrenia (Nucifora et al., 2017). Heavily reliant on natural antioxidant defences, mitochondria can become dysfunctional in disease states producing excess ROS and contributing to the inflammatory cycle (Morris & Maes, 2014). Mitochondrial dysfunction has been implicated in all three neuropsychiatric conditions (Hjelm et al., 2015; Kasahara & Kato, 2017).

In summary, the above-mentioned processes outline the key elements of neuroprogression. Attenuation of these processes is the aim of neuroprotection via agents that target the operative pathway elements. This paper aims to identify agents with neuroprotective potential in neuropsychiatric disorders and review evidence for these agents including clinical efficacy, mechanisms of action and limitations in our current assessment tools.

Method

Data for this review were sourced from online databases PUBMED, Embase and Web of Science. A Boolean search was carried out in October 2017 with no language or publication origin restriction using search terms consistent with those provided in supplementary material. Only data published since 2012 was included in this review.

1. Psychotropic Agents

1.1 Lithium

Lithium has received more clinical assessment of its neuroprotective capacity than any other agent, most notably in bipolar disorder. Cross-sectional studies of bipolar patients have associated lithium treatment with: 1) attenuation of cortical grey matter loss and 2) maintained white matter integrity (**Table 1: Psychotropic Agents**) (Giakoumatos et al., 2015; Gildengers et al., 2015; Hajek et al., 2014; Hajek et al., 2012; Hartberg et al., 2015; Pfennig et al., 2014; Poletti, Locatelli, Radaelli, Colombo, & Benedetti, 2014; Simonetti et al., 2016; van Erp et al., 2012; Zung et al., 2016). Though few have correlated lithium's structural benefits with a similar improvement in brain function.

MRI studies implicate the hippocampus, among other regions, as an area susceptible to cortical atrophy in bipolar with particular relevance to cognition and mood dysregulation (Giakoumatos et al., 2015; Hajek et al., 2014; Simonetti et al., 2016; van Erp et al., 2012; Zung et al., 2016). Three of these studies identify

hippocampal subfields cornu ammonis (CA)1, CA2/3, CA4/dentate gyrus, presubiculum and subiculum as being preferentially affected (Giakoumatos et al., 2015; Simonetti et al., 2016; van Erp et al., 2012). In these studies, lithium treatment was associated with ameliorated bipolar disorder associated cortical atrophy, restoring the volume of the hippocampus and other affected regions to the level of healthy controls. Furthermore, chronicity of lithium treatment is positively associated with the maintenance of cortical thickness (Giakoumatos et al., 2015; Hajek et al., 2014; Hartberg et al., 2015; Poletti et al., 2014; Simonetti et al., 2016; van Erp et al., 2012; Zung et al., 2016). This occurs without global increase in cortical thickness, suggesting lithium's neuroprotective qualities may be targeted at disease affected areas only. This is important as the reliability of MRI in the setting of lithium therapy has been questioned. A study of healthy human subjects (n=12) suggested lithium treatment alters grey matter T₁ relaxation time to produce a false positive reading of global cortical hypertrophy when assessed via VBM (Cousins, Aribisala, Ferrier, & Blamire, 2013). As lithium appears to prevent atrophy with specificity to bipolar disorder affected regions, the probability of this being artefact, which would occur globally, is low.

In addition to cortical grey matter atrophy, poor white matter integrity may be a feature of neuroprogression in bipolar disorder (Berk et al., 2017; Gildengers et al., 2015). MRI based comparison of elderly patients with bipolar disorder (n=58) and age matched controls (n=21) demonstrated that bipolar disorder patients had poorer white matter integrity (M. F. Kraus et al., 2007). Lithium treated bipolar patients (n=33) had better white matter integrity than those not on lithium (n=27). Additionally, duration of lithium therapy had a positive association with white matter integrity. In the only available prospective clinical trial of lithium with neuroprotection as a primary outcome, lithium was shown to preserve white matter volume in the left internal capsule following a first episode of mania (Berk et al., 2017). First episode mania patients were treated with either lithium (n=20) or quetiapine (n=19) monotherapy for a period of twelve-months and brain structure was assessed by MRI at baseline, three-

months and twelve-months. Quetiapine, unlike lithium, did not impact structural brain change in this time-frame (Berk et al., 2017). In this same cohort of patients, early lithium treatment was associated with preserved verbal fluency, a marker of preserved cognition (Daglas et al., 2016). However, this connection between improved brain structure and function has proven elusive to other studies.

Despite lithium showing more capacity for maintaining neurostructural integrity than other agents in bipolar, there is only sparse evidence connecting this with superior functional outcomes. Increased grey matter volume was not associated with fewer symptoms of mania or depression at the time of assessment (Hartberg et al., 2015), reduced rates of relapse when looked at retrospectively (Hajek et al., 2014) nor improved cognition compared to bipolar disorder patients on other agents (Giakoumatos et al., 2015). Improved white matter integrity seen with lithium therapy was not associated with improved cognition in elderly bipolar disorder patients (Gildengers et al., 2015). In one of the few studies assessing lithium's effect on cognition as a primary outcome, Pfennig et al. (2014) compared euthymic bipolar disorder patients managed on either lithium (n=58) or other agents (n=31) and healthy controls (n=53). The single significant difference in cognition between bipolar disorder patients and healthy controls was slower early visual processing, a measure of information processing speed. Lithium correlated with significantly slower early visual processing than other treatments. Whether this represents an adverse effect of lithium, or poorer attenuation of functional decline is unclear. Furthermore, as the subjects were middle aged (mean age=47 years), it may be that bipolar disorder associated cognitive deficit has not yet presented itself, therefore, confounding interpretation of this work.

Mechanistically, lithium may reduce the oxidative stress burden seen in bipolar disorder (De Sousa, Zanetti, Talib, Gattaz, & Machado-Vieira, 2013; de Sousa et al., 2014). Furthermore, rat studies suggest lithium abrogates inflammatory signalling pathways including those associated with toll-like receptor 4

(TLR4) (Khan et al., 2017). Lithium inhibition of TLR4 may reduce phosphorylation of nuclear factor- κ B (NF κ B) to suppress inflammatory gene expression, quieten microglial activity and inhibit neuronal apoptosis indicated by decreased caspase-3. Lithium may also have a neurotrophic effect by increasing hippocampal BDNF in the setting of a neurotoxic insult (Motaghinejad, Seyedjavadein, Motevalian, & Asadi, 2016) - a finding replicated in rat hippocampal neurons *in vitro* (Dwivedi & Zhang, 2014).

1.2 Second-generation antipsychotics and neurostructural integrity

A meta-analysis of 317 studies and over 9000 patients demonstrated a 2.6% reduction in total brain volume in patients with schizophrenia versus healthy controls (Haijma et al., 2013). Total grey matter loss was most pronounced in those on high dose second-generation antipsychotics at the time of scanning. In isolation, this finding would suggest a correlation between second-generation antipsychotic treatment and brain atrophy. However, antipsychotic therapy is standard across the disorder and may be a marker rather than a mediator; grey matter loss was also associated with duration of illness. Antipsychotic naïve patients had a substantially lower duration of illness than their treated counterparts. This hinders the interpretability of the effect of antipsychotic use on brain volumes. It cannot be definitively ascertained whether the atrophy observed represents: 1) natural disease progression with no effect from antipsychotics, 2) slowed progression due to neuroprotection from antipsychotics, or 3) accelerated progression due to neurotoxicity from antipsychotics. A more recent meta-analysis suggests that antipsychotic exposure is associated with greater volume loss independent of duration of illness (Fusar-Poli et al., 2013). However, this group did not differentiate between early and second-generation antipsychotics. A third meta-analysis found that early antipsychotics accelerate loss of grey matter while

second-generation antipsychotics attenuate this loss in a dose dependent manner (Vita, De Peri, Deste, Barlati, & Sacchetti, 2015).

In summary, without well-powered prospective clinical studies with appropriate control groups, it is difficult to ascertain with relative certainty, whether antipsychotics potentiate or attenuate cortical volume loss.

1.2.1 Clozapine

Robust human data relating clozapine with neuroprotection is limited. In a two-study series, it was suggested clozapine does not slow grey matter atrophy or loss of white matter integrity in treatment resistant schizophrenic patients; despite universal clinical improvement (Ahmed et al., 2015; Forde et al., 2012). However, these studies compared treatment resistant schizophrenic patients on clozapine (n=33 and n=21 respectively) to age matched healthy controls (n=31 and n=21 respectively) without a third group of antipsychotic naïve patients or patients on other therapies; making interpretation problematic.

Clozapine is thought to induce neurogenesis; proliferation of 5-bromo-2'-deoxyuridine (BrdU) positive cells (labelled putative neurons) in the hippocampus of healthy mice treated with clozapine is indicative of this effect (Chikama et al., 2017). This neurotrophic action is likely occurring through increased expression of BDNF in the hippocampus of clozapine treated mice (Gumuslu et al., 2015; H.-W. Kim, Cheon, Modi, Rapoport, & Rao, 2012). This is a poignant finding as serum levels of BDNF are decreased in antipsychotic naïve schizophrenic patients and increased in clozapine responders (Krivoy et al., 2017; Toll & Mane, 2015). Importantly, clozapine induced neuroproliferation is organised and functional. In a chronic mild stress model of depression in mice, clozapine restored neuronal architecture in the dentate

gyrus with dendritic length equalling that of healthy controls (Morais et al., 2017). Furthermore, neuron cell survival was significantly increased with clozapine.

The anti-inflammatory properties of clozapine highlight its potential for neuroprotection. The brains of clozapine treated rats exhibited preferential throughput of the anti-inflammatory docosahexaenoic acid (DHA) cascade over the pro-inflammatory arachidonic acid (AA) cascade (H.-W. Kim et al., 2012). As evidenced by increased DHA protein levels, decreased cyclooxygenase (COX) activity and lower concentrations of prostaglandin E₂ (PGE₂), a pro-inflammatory AA metabolite. Additionally, Clozapine both alone and in combination with N-acetylcysteine (NAC) normalises tryptophan metabolism, diverting away from production of neurotoxic TRYCAT QA in social isolation reared (SIR) rats – a rat model of schizophrenia (Moller, Du Preez, Emsley, & Harvey, 2012; Moller et al., 2013; Moller, Du Preez, Viljoen, Berk, & Harvey, 2014).

1.2.2 Aripiprazole

There is a preliminary body of evidence for aripiprazole in neuroprotection. Aripiprazole has been shown to raise plasma BDNF in patients (n=50) following first episode psychosis (Yoshimura et al., 2012). Chronic aripiprazole increases hippocampal levels of BDNF and ameliorates cognitive deficit in prenatally stressed rats (Nowakowska et al., 2014). A cross-sectional study of schizophrenic patients (n=101) compared the cognitive impact of olanzapine, risperidone and aripiprazole (Hori et al., 2012). Only aripiprazole was found to have a neutral impact on cognition, while other agents were associated with cognitive decline. Using N-methyl-D-aspartate (NMDA) receptor blockade in rats as a model of schizophrenia, Deiana et al. (2015) showed aripiprazole is superior to olanzapine and risperidone when rescuing cognition.

Collectively these studies do not address mechanistic pathways in detail. Considering aripiprazole's differing receptor affinity to most second-generation antipsychotics which includes partial agonism of dopamine (D2) and serotonin (5HT_{1A}) receptors, studies investigating the mechanisms by which these differences may influence the neuroprotective potential of aripiprazole would be valuable.

1.2.3 Olanzapine

Olanzapine did not increase plasma BDNF in a 6-week prospective study of schizophrenic patients (n=50) (Kudlek Mikulic et al., 2017). A study of ethnically homogenous schizophrenic patients correlated olanzapine treatment with shortened leukocyte telomere length and advanced metabolic age (Monroy-Jaramillo et al., 2017). Furthermore, olanzapine therapy in rats is associated with raised circulating inflammatory cytokines. Elevated TNF α , IL1 β and IL6 were found in the hypothalamus (a structure lacking a blood brain barrier) and adipose tissue of rats treated with olanzapine (Q. Zhang, He, Deng, Wang, & Huang, 2014). Whether this peripheral promotion of inflammation from olanzapine leads to central neurotoxicity is unclear, though the animal studies described below would suggest not.

Olanzapine up-regulates cellular survival pathways in the pre-frontal cortex, hippocampus and striatum of rats (G. Reus et al., 2012). Specifically, olanzapine had a dose dependent preference for increasing expression of BDNF and Bcl-2 over Bcl-2 associated death promoter (BAD) expression, or, neurogenic and anti-apoptotic over pro-apoptotic pathways. These effects were enhanced when combined with fluoxetine. Like aripiprazole, olanzapine improves cognition and increases hippocampal BDNF in prenatally stressed rats (Nowakowska et al., 2014). In light of these neuroprotective qualities of

olanzapine, strategies to mitigate olanzapine's peripheral toxicity would themselves be of neuroprotective benefit.

1.2.4 Risperidone and paliperidone

A study of antipsychotic naïve first episode psychosis patients (n=51) found 12-weeks of risperidone correlates with reduced oxidative stress in the periphery, irrespective of clinical response (Noto et al., 2015). Unlike olanzapine, evidence suggests risperidone's peripheral anti-inflammatory actions may correlate directly with central effects. Risperidone upregulates Bcl-2 expression and BDNF in the brain of SIR rats (Yang, Yang, Wan, Huang, & Liu, 2017). Risperidone prevents the neuropathological changes seen in the offspring born to dams subjected to maternal immune activation, which is a well validated preclinical model of schizophrenia (Piontkewitz et al., 2013). These neuropathological changes include impaired neurogenesis, disturbed micro-vascularisation and loss of parvalbumin expressing hippocampal interneurons (associated with pathogenesis of schizophrenia). Paliperidone has also prevented changes induced by maternal immune activation though in mice (MacDowell et al., 2017). In this model, paliperidone had a net antioxidant effect through at least two mechanisms: 1) upregulation of nuclear transcription factor (erythroid-derived 2)-like 2 (NRF2), which drives antioxidant gene expression and 2) polarization of microglia to their antioxidant phenotype M2. Paliperidone decreases activity of antioxidant enzymes adenosine deaminase, xanthine oxidase and catalase in rat brain tissue indicating reduced burden of oxidative stress (Demirci et al., 2015).

1.2.5 Quetiapine

Efficacious in mood disorders and psychosis, quetiapine has a mixed body of evidence relating to neuroprotection. In a prospective clinical trial of quetiapine versus lithium following first episode of mania, quetiapine was inferior to lithium in preserving white matter volume (Berk et al., 2017). However, quetiapine was efficacious against cuprizone induced demyelination and schizophrenia like behaviour in mice (Wang et al., 2016; Y. Zhang et al., 2012). Quetiapine rescued demyelination with associated stimulation of notch signalling, a regulator of oligodendrocyte maturation and myelination. Protection of myelin may also occur through inhibition of microglia via normalising intracellular Ca^{2+} homeostasis (H. Wang et al., 2015).

Quetiapine may also be an effective antioxidant. Across two studies in mice, acute and chronic quetiapine increased mitochondrial respiratory chain activity and reduced markers of oxidative stress in the pre-frontal cortex, nucleus accumbens, amygdala and hippocampus (Ignácio et al., 2017; Ignacio et al., 2015).

→ Table 1: Psychotropic Agents

1.3 Antidepressants

The roles of specific antidepressants - selective serotonin re-uptake inhibitors (SSRIs), selective serotonin noradrenaline re-uptake inhibitors (SNRIs), tricyclic antidepressants (TCAs) and others - in neuroprotection is a discussion that warrants its own review. Therefore, the central theories of antidepressant driven neuroprotection, as well as, emerging evidence will be discussed here.

A recent systematic review and meta-analysis of depressed patients (n=2384 antidepressant free patients and n=1249 antidepressant treated patients) versus healthy controls (n=2982) suggests antidepressant

treatment rectifies deficits in peripheral BDNF (Molendijk et al., 2014); possibly as a secondary result of central neurogenesis (Sachs & Caron, 2014). Antidepressants have been shown to alter mitochondrial function, increase mitochondrial biogenesis and enhance capacity for dealing with oxidative stress in animal models of depression (Demirdağ, Nazıroğlu, & Övey, 2017; Glombik et al., 2017; Villa et al., 2017). A meta-analysis (n=1137 total subjects) shows antidepressant therapy correlates with a net decrease in peripheral inflammatory cytokines (Wiedłocha et al., 2017). Specifically, antidepressant therapy decreased IL-4, IL-6, IL-10 and IL- β . Though evidence of this correlating with central anti-inflammatory effects is scarce. Duloxetine has been shown to suppress pro-inflammatory cytokines in the hippocampus of adolescent rats (X. Zhang et al., 2016). Sertraline prevents induction of IL- β and TNF- α by seizure activity in the rat hippocampus (Sitges, Gómez, & Aldana, 2014). Antidepressants may augment the TRKAT pathway to favour neuroprotective conditions. A study of patients with depression exhibits increased production of neurotoxic TRKATs and subsequent reduction with escitalopram treatment (Halaris et al., 2012). These pathways represent multiple convergent mechanisms thought to be neuroprotective, though emerging evidence suggests the actions of antidepressants may be context dependent and therefore more difficult to elucidate than previously thought.

Antidepressants, in particular SSRIs, may increase neural plasticity and therefore induce undirected susceptibility to change. Meaning the SSRI treated brain is hyper-responsive to environmental influence. In an environment that promotes recovery, SSRI treatment may prime the brain to adopt the positive neurostructural changes leading ultimately to an improved mental state. However, in an environment promoting mental illness, SSRIs may propagate further maladaptive changes leading to a worsened mental state. Across two studies, previously stressed mice (24-days of stress) treated with fluoxetine in an enriched environment (21-days of treatment) exhibit lower levels of anhedonia, dramatically increased hippocampal and hypothalamic BDNF and lower plasma corticosterone levels to vehicle treated mice

(Alboni et al., 2017; Branchi et al., 2013). However, fluoxetine treatment in a stressed environment exacerbates the depressed phenotype, reduces hippocampal and hypothalamic BDNF, slows hippocampal proliferation and increases plasma corticosterone. In a third study of similar design, the effects of fluoxetine on inflammatory cytokines and microglial function were also dependent on environmental conditions (Alboni et al., 2016). Paradoxically in the enriched environment, fluoxetine increases hippocampal expression of the pro-inflammatory cytokine IL- β and activates microglia. While in the stressed environment, fluoxetine decreases hippocampal inflammatory cytokines IL- β and IL- γ and downregulates microglia activity. The anti-inflammatory properties SSRI's during stress have been shown by other studies (Cheng et al., 2016; Wilson et al., 2014). However, to associate effective fluoxetine treatment with inflammation in the hippocampus is somewhat at odds with the field. Authors postulate effective neuroplastic change requires some degree of inflammatory signalling to occur, though replication of these findings by other groups is required to prove this hypothesis. As SSRI's may share convergent pathways with other classes of agents, these data highlighting the influence of the environment on neuroprotective versus neurotoxic outcomes should be considered in future work.

2. Other pharmaceutical agents

2.1 Minocycline

Minocycline is a tetracycline antibiotic with psychotropic effects being applied to a range of neuropsychiatric conditions. Used as an adjunct to standard therapies, minocycline improved negative symptoms in early schizophrenia (n=46) (Chaudhry et al., 2012). Extrapolating this to include measures of neuroprotection, Chaves et al. (2015) led a small double-blind randomised placebo-controlled trial of

add-on minocycline for early stage schizophrenia. Minocycline patients (n=16) exhibited improved symptom control, preserved grey matter volume and increased cerebral perfusion at twelve months. MRI data showed preserved grey matter in midposterior cingulate cortex and in the precentral gyrus compared to placebo. Additionally, single photon emission computed tomography displayed increased uptake of contrast in frontotemporal regions; indicative of enhanced cerebral perfusion. These are promising results despite the small size of the study.

In depression, a double blind randomised clinical placebo-controlled trial (n=71) found adjuvant minocycline led to improved clinical global impression, quality of life and social and occupational function (Dean et al., 2017). However, no significant difference in the Montgomery–Asberg Depression Rating Scale was observed between the two groups. A smaller double blind randomised clinical placebo-controlled trial of adjuvant minocycline in depression (n=41) found clinical improvement as per Hamilton Depression Rating Scale in addition to other measures (Husain et al., 2017). Lastly, a 6-week open label study of adjuvant minocycline in combination with SSRI's to treat psychotic depression (n=25) demonstrated minocycline's safety in this setting (Miyaoaka et al., 2012). However, lack of control group and open-label design limit interpretation of this particular work.

As the clinical benefits of minocycline continue to be realised, understanding of its mechanism deepens also. Minocycline may be a selective Cytochrome P450 (CYP) inhibitor altering the pharmacokinetics of antipsychotics such as clozapine to raise their circulating levels and enhancing overall effect (Wehring et al., 2017). Nevertheless, minocycline appears to have direct actions of its own.

In maternal immune challenged to mice, Mattei et al. (2014) reported elevated microglial IL- β and TNF α in the hippocampus, indicative of chronic microglial activation. Furthermore, the pro-proliferative TNF α

Receptor Type 2 (TNFR2) was downregulated in hippocampal neural progenitor cells. Minocycline treatment attenuated these changes (**Table 2: Other Pharmaceutical Agents**). In a follow-up study, Mattei et al. (2017) applied RNA-sequencing to hippocampal microglia isolated from these mice. This showed a significantly altered transcriptome, which was almost entirely normalised by minocycline. This suggests minocycline could be inhibiting activated microglia, a finding that has been replicated in multiple other pre-clinical studies (Kreisel et al., 2014; Tanra, 2013; Zhu et al., 2014).

It is unclear exactly how minocycline inhibits microglial activity. It may occur by reducing oxidative stress and inflammatory cytokines which is perpetuated by aberrant microglial hyperactivity. Minocycline normalises glutathione levels and thiobarbituric acid reactive substances (TBARS) in disease models (markers of oxidative stress and lipid peroxidation respectively) (Monte et al., 2013; G. Z. Reus et al., 2015). In a mouse model of post-traumatic stress disorder, minocycline attenuate a rise in IL-1, IL-6 and TNF α in the hippocampus and frontal cortex (Levkovitz, Fenchel, Kaplan, Zohar, & Cohen, 2015).

2.2 Statins

Like minocycline, statins continue to be explored in various clinical settings as adjunct therapies. Meta-analysis suggests add-on statin therapy improves depressive symptoms with good tolerability (Salagre, Fernandes, Dodd, Brownstein, & Berk, 2016); however, larger prospective trials are required to confirm this. In patients after an acute coronary event, statins were comparable to escitalopram in reduction of depression, with the combination numerically superior to monotherapy with either (S. W. Kim et al., 2015). Simvastatin was shown to be more effective than atorvastatin at reducing depressive symptoms following coronary artery bypass grafting (CABG) surgery in a small cohort of patients (n=46) (Abbasi et al., 2015). Neither pravastatin nor lovastatin add-on therapies were shown to lessen symptom severity

in schizophrenia, though pravastatin reduced peripheral inflammatory markers (Ghanizadeh, Rezaee, Dehbozorgi, Berk, & Akhondzadeh, 2014; Vincenzi et al., 2014).

Co-administration of rosuvastatin with paroxetine and citalopram influences glutathione homeostasis and may enhance capacity for coping with oxidative stress in rats (Herbet, Izdebska, Pi tkowska-Chmiel, Poleszak, & Jagiełło-Wójtowicz, 2016). Simvastatin was used to reverse the loss of pyramidal neurons in the hippocampus of high fat diet fed mice (Can, Ulupinar, Ozkay, Yegin, & Ozturk, 2012). This morphological benefit was accompanied by alleviation of depression-like symptoms. Simvastatin shows comparable antidepressant effect to imipramine in a chronic mild stress model of depression in rats (Lin, Chang, & Lin, 2014). The antidepressant effects of statins may involve intracellular pro-survival signalling through the phosphoinositide 3 kinase (PI3K)/protein kinase B (Akt)/glycogen synthase kinase 3 beta (GSK-3 β)/mechanistic target of rapamycin (mTOR) cascade. This pathway, an indicator of anabolic states, is upregulated in hippocampal neurones following an antidepressant dose of atorvastatin in mice (Ludka et al., 2016). This could be driven, at least partly, by increased mature BDNF in this brain region (Ludka et al., 2017).

2.3 Aspirin

As an irreversible inhibitor of COX1 and COX2 signalling, aspirin stimulates endogenous anti-inflammatory signals leading to reduced levels of inflammatory biomarkers and less oxidative stress (Berk et al., 2013). Aspirin does not cross the blood brain barrier but theoretically should exert neuroprotective effects through reducing overall inflammation. For example, aspirin decreases peripheral IL-6 and TNF α while also exerting antidepressant effects; a result comparable to fluoxetine and

imipramine (Guan, Shao, Xie, Chen, & Wang, 2014). These peripheral anti-inflammatory actions correlate with a central reduction in inflammatory cytokines (Li et al., 2017). This may in turn suppress microglia and protect white matter from degeneration. Co-treatment with Aspirin and DHA suppressed activation of microglia *in vitro* and increased the microglial concentration of glutathione; therefore, increasing the anti-oxidant capacity of microglia cells (Pettit, Varsanyi, Tadros, & Vassiliou, 2013). Two studies using a combination of an *in vivo* model of white matter lesions and *in vitro* oligodendrocyte precursor cells demonstrated aspirin can protect white matter from insult (Chen et al., 2014; Huang et al., 2016). Aspirin appears to promote oligodendrocyte differentiation to repair white matter lesions possibly via inhibition of Wnt/ β -Catenin signalling, a regulator of cell fate.

→ Table 2: Other Pharmaceutical Agents

2.4 Cyclooxygenase-2 Inhibitors

The serum expression of COX-2 is elevated in depression (Galecki et al., 2012). A 2014 meta-analysis suggests celecoxib, a selective COX-2 inhibitor, is effective add-on therapy in depression (Kohler et al., 2014). In the years since, celecoxib has shown promise as adjunctive therapy in young manic patients (Arabzadeh et al., 2015; Mousavi et al., 2017). Adjunctive celecoxib may also improve symptoms in schizophrenia according to a recent meta-analysis (W. Zheng et al., 2017).

Highlighting the role of COX-2 inhibition in neuroprotection, Luo, Kuang, Li, Ran, and Yang (2017) inhibited COX-2 signalling via two methods in chronically stressed rats: 1) meloxicam, a COX-2 selective non-steroidal anti-inflammatory and 2) via RNA-interference. COX-2 inhibition reversed the

pro-inflammatory and anti-neurotrophic changes induced by chronic stress. This included reducing hippocampal prostaglandins and increasing hippocampal signalling through cyclic adenosine monophosphate (cAMP), protein kinase A (PKA), cAMP response binding element (CREB) and BDNF. Another study showed COX-2 inhibition with celecoxib reduces hippocampal concentrations of inflammatory markers TNF α and nuclear factor-B (NF- κ B), as well as, the apoptotic marker caspase-3 in mice (Elnahas, Abou Zeid, Kawy, Hendawy, & Baher, 2016).

Collectively these human and animal data imply COX-2 inhibition may play a role in targeting neuropsychiatric symptoms with the added benefit of protecting brain structures from inflammatory insult.

2.5 Ketamine

It is suggested that altered glutamatergic signalling is a component of the pathophysiology of schizophrenia (Howes, McCutcheon, & Stone, 2015). Ketamine, a selective NMDA antagonist, is therefore used as a pre-clinical model of the disease (Monte et al., 2013). High-dose ketamine, such as is used for anaesthesia, is possibly neurotoxic to the developing brain (X. Zheng, Zhou, & Xia, 2015). Furthermore, chronic ketamine abuse is associated with an irreversible decline in brain function (Morgan et al., 2014). However, low-dose acutely administered ketamine is an emerging therapy in treatment resistant depression and suicidality (C. Kraus et al., 2017). These effects have led to interest in ketamine as a possible neuroprotective agent.

In rat models of depression, low-dose ketamine acutely influences anti-inflammatory pathways thought to be neuroprotective. Ketamine reduces hippocampal and serum levels of pro-inflammatory cytokines

IL1 β , IL6 and TNF α while concurrently reducing depressive symptoms (N. Wang et al., 2015; Xie et al., 2017). This culminates in an overall reduction in ROS (Demirdas, Naziroglu, & Ovey, 2017). Additionally, ketamine influences neurotrophic pathways in the acute phase. Ketamine increases hippocampal levels of BDNF, likely via activation of adenosine monophosphate activated protein kinase (AMPK), to alleviate depression in rats (Xu et al., 2013). This rise in BDNF has been linked to maintenance of normal neuron morphology in the hippocampus of chronically stressed rats (W. X. Liu et al., 2016). Inhibition of mTOR, an intracellular mediator of BDNF and driver of synaptogenesis, significantly reduces ketamine's antidepressant effect (G. Z. Reus et al., 2016). In another pathway, ketamine increased the hippocampal level of the novel neurotrophic transcription factor activity-dependent neuroprotective protein (ADNP). This occurred simultaneously with decreased apoptosis marker, caspase-3 (B. P. Brown et al., 2015).

Collectively these data indicate a shift towards neurogenesis acutely following low-dose ketamine treatment but data on chronic outcomes is lacking. Until the lasting effects of low-dose ketamine have been properly assessed, ketamine's neuroprotective potential remains largely theoretical.

2.6 Alpha 2 Delta ($\alpha 2\delta$) Ligands

The $\alpha 2\delta$ ligands pregabalin and gabapentin are used as treatments for neuropathic pain and have recently become known for their abuse potential (Chiappini & Schifano, 2016; Peckham, Fairman, & Sclar, 2017). Regarding neuroprotection their efficacy remains largely unknown but a small body of pre-clinical evidence indicates activation of neurogenesis as one possible avenue. Valente et al. (2012) demonstrated the neurogenic capacity of the $\alpha 2\delta$ ligands with both pregabalin and gabapentin driving the maturation of

hippocampal neuron progenitor cells *in vitro*. Interaction with $\alpha 2\delta$ subunit 2 voltage gated calcium channels and upregulation of NF- $\kappa\beta$ signalling was suggested as a possible mechanism; NF- $\kappa\beta$ signalling is a downstream regulator of neuron differentiation. They went on to show that pregabalin therapy ameliorated the depressed phenotype of chronically restrained mice with associated hippocampal neurogenesis *in vivo* (Valente et al., 2012). Gabapentin's positive effect on neuron differentiation has been replicated in cortical stem cells harvested from rats (Paknejad, Kebriaeezadeh, Ghahremani, Gharghabi, & Ostad, 2015). Moreover, chronic administration of gabapentin prevents behavioural abnormalities correlated with social isolation rearing in mice via an unknown mechanism (Amiri et al., 2017). Cross-sectional analysis of people receiving $\alpha 2\delta$ ligands with neuroprotective end-points would be an expedited way of assessing their potential.

3. Nutritional agents

3.1 Melatonin

Aberrant melatonin levels in the serum and cerebrospinal fluid have been observed in mood disorder patients (Bumb et al., 2016). In severe depression, serum melatonin levels are decreased in proportion to serum BDNF and are significantly lower than healthy controls (Ogłodek, Just, Szromek, & Araszkiewicz, 2016). Adjunctive melatonin has already been shown to improve the metabolic side-effects of antipsychotics (Modabbernia et al., 2014; Romo-Nava et al., 2014).

In the setting of chronic ketamine administration in mice, melatonin increases hippocampal BDNF leading to phosphorylation of Akt and extracellular signal regulated kinases 1 and 2 (ERK1/2) (pro-

survival pathways) (**Table 3: Nutritional, Hormonal and Other Agents**) (Choudhury, Singh, Palit, Shukla, & Ganguly, 2016). These actions were found to be melatonin receptor 1 and 2 (MT₁/MT₂) dependent through blockade with luzindole, an MT₁/MT₂ receptor antagonist. However, a study on hippocampal slice cultures found melatonin induced dendritogenesis through Ca²⁺ calmodulin-dependent protein kinase II (CaMKII) activation independent of MT₁/MT₂; indicating melatonin may influence neuronal structures through other mechanisms (Dominguez-Alonso, Valdes-Tovar, Solis-Chagoyan, & Benitez-King, 2015). Melatonin has demonstrated capacity to preserve hippocampal monoamine handling in disease states, in particular noradrenaline, though whether this is neuroprotective is unclear (Stefanovic et al., 2016). Finally, melatonin may also influence redox equilibrium. Either through the oxygen free radical scavenging properties of its metabolites, or, regulation of antioxidant gene transcription as was seen in a rat model of mania (Galano, Tan, & Reiter, 2013; Souza et al., 2014).

3.2 Oestrogen

The sex hormone oestrogen has a developing evidence base for neuroprotection. Using bone mineral density as a marker of cumulative oestrogen exposure, a study of female psychosis patients (n=14) found a positive correlation between oestrogen exposure and cortical thickness (van der Leeuw et al., 2013). A meta-analysis showed delayed age of menopause and longer reproductive period, also representative of cumulative oestrogen exposure, lowers risk of developing depression (Georgakis et al., 2016). Demonstrating that exogenous oestrogen can also be beneficial, ovariectomized rats were treated with escitalopram and adjunctive oestrogen. Depression-like symptoms and hippocampal architecture were both improved compared to escitalopram alone (Ibrahim, Safar, Khattab, & Agha, 2016). Mouse derived cortical cell cultures suggest oestrogen's neuroprotective effects are dependent on G-protein coupled receptor 30 (GPR30) and ERK1/2 signalling (S. B. Liu et al., 2012).

3.3 Curcumin

There is preliminary evidence supporting curcumin as a therapy for depression; however, pre-clinical data implies neuroprotective potential (Lopresti, Maes, Maker, Hood, & Drummond, 2014). Curcumin has demonstrated efficacy in ameliorating depressive-like symptoms and cognitive deficit in mice submitted to CUMS. Additionally, curcumin pre-treatment raises hippocampal BDNF with upregulation of the pro-survival ERK pathway (D. X. Liu et al., 2014; L. Zhang et al., 2014). Curcumin drives production of DHA in rats leading to increased levels in the brain; therefore, presenting another potential avenue of neuroprotection (Wu et al., 2015). To date, curcumin's poor bioavailability has proven a limitation to human applications; however, ongoing efforts such as nanoformulation are working to address this (Naksuriya, Okonogi, Schiffelers, & Hennink, 2014).

→ Table 3: Nutritional, Hormonal and Other Agents

3.4 Erythropoietin

Erythropoietin (EPO) can be used as an adjunctive treatment in neuropsychiatric disease. In bipolar disorder (n=43) and major depression (n=36), add-on EPO leads to improved objective cognition, which lasts six weeks post cessation of EPO (Ott, Vinberg, Kessing, & Miskowiak, 2016). Systematic review of the literature suggests schizophrenia may also benefit from EPO induced cognitive enhancement (Fountoulakis et al., 2017). In a study of socially defeated mice, a bullying induced model of depression, EPO therapy correlated with symptom improvement as indicated by forced swim testing (Osborn et al., 2013). This was accompanied by maintained hippocampal neurogenesis. In the same study, rapamycin

administration, an mTOR inhibitor, ameliorated EPO's effect on depressive-like symptoms. However, neurogenesis was not assessed in the presence of rapamycin, so it is unclear from this study if mTOR plays a role here. Three-week administration of EPO to young mice led to an increase in hippocampal volume and an approximate 20% higher number of pyramidal neurons in CA1/CA3 hippocampal subregions (Hassouna et al., 2016). This occurred without an increase in BrdU positive cells and therefore irrespective of proliferation and raising the question of the origins of the additional pyramidal neurons. The same study used cultured neurospheres to show that EPO decreased Sox9 and increased miR124, a transcription factor and non-coding RNA associated with neural differentiation. Thus, indicating EPO may induce differentiation of neural progenitor cells *in vivo*.

3.5 N-acetylcysteine

A systematic review of clinical trials involving NAC support its potential role in managing various psychiatric conditions including but not limited to depression, bipolar disorder and schizophrenia (Deepmala et al., 2015). Further analysis of the literature suggests NAC could have a positive impact on human cognition, though further research is required (Skvarc et al., 2017). A preliminary study of patients with psychosis (n=58) found an association between NAC treatment and improved working memory after 24-weeks (Rapado-Castro et al., 2017).

NAC is a glutathione precursor and potent antioxidant; as demonstrated by reversal of a glutathione deficient mouse model of schizophrenia (Duarte et al., 2012). When given to pregnant rats submitted to an endotoxin immune challenge, the antioxidant properties of NAC prevented permanent disorganisation of the CA3 hippocampal subregion in offspring (Rideau Batista Novais et al., 2013). By increasing glutathione and influencing the antioxidant capacity of cells, NAC may augment mitochondrial

respiratory capacity as a means of protecting against disease. NAC improves mitochondrial function in a transgenic mouse model of Huntington's disease (Wright et al., 2015).

NAC may also exert neuroprotection via glutathione independent mechanisms. Cell culture experiments suggest NAC improves heat-shock protein (Hsp) 70 chaperone function in neurones and astrocytes; therefore, protecting cells against proteotoxicity (Gleixner et al., 2017; Jiang et al., 2013). Whether this occurs *in vivo* in neuropsychiatric conditions is unknown. Current NAC formulations have limited bioavailability and short half-life, however, novel nanoparticle prodrug variations of NAC present a potential solution to this issue, nonetheless will require further assessment for indication *in vivo* (Markoutsas & Xu, 2017).

3.6 Leptin

Leptin is an adipokine produced by white adipose tissue with an emerging evidence base in neuropsychiatric disease. A meta-analysis of leptin in bipolar disorder found no association between serum leptin and the mood status (depression, mania or euthymia) of patients (Fernandes et al., 2016). More recently, an eight-year longitudinal study of bipolar patients (n=53) found low leptin and low adiponectin to be predictive of depressive relapse in the next twelve-months (Bond et al., 2017).

Adeno-associated viral vector mediated deletion of the hippocampal leptin receptor induces a depression-like phenotype in adult mice (Guo, Huang, Garza, Chua, & Lu, 2013). Conversely, leptin therapy promotes hippocampal neurogenesis and prevents the depressive phenotype associated with CUMS in rats (Garza, Guo, Zhang, & Lu, 2012).

Leptin levels in schizophrenia patients increase as BMI also increases, possibly representative of leptin-resistance secondary to antipsychotic use (Neelamekam, Nurjono, & Lee, 2014). Data is scarce relating leptin treatment to neuroprotection.

Further studies assessing the direct effect of leptin treatment or the reversal of leptin resistance, on disease related brain structures are required.

3.7 Omega-3 polyunsaturated fatty acids

Dietary omega-3 polyunsaturated fatty acids (PUFAs) may reduce oxidative stress and prevent the onset of mood symptoms. A longitudinal study of middle aged Puerto Ricans (n=787) showed that in people with a high burden of oxidative stress, depressive symptoms were more common when their omega-3 index was low (Bigornia et al., 2016).

Omega-3 index is the combined total of serum eicosapentaenoic acid (EPA) and DHA as looking at omega-3 in isolation is unlikely to reveal the entire story. Recently, more attention has been paid to the ratio of omega-6 to omega-3 PUFAs in relation to psychiatric disease. The baseline ratio of omega-6 fatty acids to omega-3 fatty acids has a positive correlation with the risk for development of mood disorders in high risk young adults (n=69) (Berger et al., 2017). A similar observation was made in young adults at ultra-high risk of psychosis (n=95) (Pawelczyk, Trafalska, Kotlicka-Antczak, & Pawelczyk, 2016).

In light of such evidence, omega-3 PUFA supplementation continues to be tested as augmentation therapy in neuropsychiatric disease with mixed results. In hepatitis C patients receiving interferon alpha (INF α) therapy, depression is a common adverse effect (Machado et al., 2017). EPA but not DHA

supplementation reduced the percentage of patients (n=154) developing depression while on INF α therapy (Su et al., 2014). A randomized trial of older adults at risk of depression (n=51) found omega-3 PUFA supplementation was associated with reduced thalamic oxidative stress and attenuation of symptoms (Duffy et al., 2015). In young psychotic patients, or those at ultra-high risk of psychosis, omega-3 PUFA supplementation increased total antioxidant capacity and reduced symptom severity (Pawelczyk, Grancow-Grabka, Trafalska, Szemraj, & Pawelczyk, 2017; Smesny et al., 2015).

However, multiple other studies have found omega-3 PUFAs to have no, or even negative effects on mental state. A multicentre randomised double-blind clinical placebo control trial showed omega-3 PUFAs plus cognitive behavioural case management in ultra-high risk of psychosis patients (n=301) did not slow conversion to psychosis over a 6-month period (McGorry et al., 2017). In a study of patients at high risk of post-traumatic stress disorder, omega-3 PUFAs had no effect serum BDNF or mental state (Matsuoka et al., 2015). Finally, a randomised double-blind clinical placebo control trial of add on EPA and/or Vitamins C + E in acute schizophrenia (n=99) found these agents to worsen psychosis (Bentsen, Osnes, Refsum, Solberg, & Bohmer, 2013). To explain these discrepancies, groups have postulated omega-3 PUFAs could have opposing neuroprotective or neurotoxic effects depending upon disease stage, for example, prodromal phase versus acute exacerbation of schizophrenia.

3.8 Ebselen

Ebselen is an organoselenium based molecule that mimics the actions of lithium and glutathione peroxidase (endogenous antioxidant), making this currently unmarketed pharmaceutical an attractive

prospect in neuroprotection. Ebselen has been shown in mice to cross the blood brain barrier to inhibit inositol monophosphatase (IMPase), a putative target in bipolar disorder, and exert lithium like effects in animal models of mania (Singh et al., 2013). The same group have shown ebselen to inhibit IMPase in the anterior cingulate cortex of healthy human subjects with significant effects on emotional processing (Singh et al., 2016); findings that have been subsequently replicated (Masaki, Sharpley, Cooper, et al., 2016; Masaki, Sharpley, Godlewska, et al., 2016). Early life treatment with ebselen is comparable to NAC in the prevention of schizophrenia-like symptoms in rats subjected to neonatal hippocampal lesions (Cabungcal et al., 2014). *In vitro* analysis of rat astrocytes indicate ebselen may cause endoplasmic reticulum stress and neurotoxicity, casting some doubt over its neuroprotective potential (Santofimia-Castano et al., 2016). However, ebselen's supposed mechanisms *in vivo* and promising clinical data warrant further investigation as a neuroprotective option.

Conclusion

The evidence presented here is a testament to the significant progress made in neuroprotection. Lithium's maintenance of neurostructural integrity in bipolar disorder is the most pertinent example. However, lithium's story also highlights the limitations in current assessment tools and understanding. MRI results consistent with neuroprotection do not necessarily correlate with functional attenuation of neuroprogression – reduced vulnerability to relapse, increased responsiveness to treatment and delayed cognitive/functional decline. Agents such as aripiprazole or EPO may demonstrate functional improvement in cognition but have little evidence regarding neurostructural benefit. A deeper mechanistic understanding of the agents presented here may explain these discrepancies in future. At this stage, it is also unclear if neuroprotective benefit accrues by directly reversing the operative pathways or via augmentation of state related activation of such pathways, as seem to be the case with antidepressants. Future work may show that agents from multiple classes induce undirected susceptibility to change rather than driving neuroprotective alterations irrespective of environmental factors. Finally, it may be that neuroprotection may require not a single agent but rather multiple adjunctive therapies acting in synchrony to achieve an effect. For example, the use of melatonin or other agents to augment the metabolic side effects of second-generation antipsychotics to improve adherence and maximise benefit. In any case, further validation is ultimately required at this time.

The studies reviewed here were heterogeneous in almost every respect, ranging from clinical trials to cell culture experiments with only some studies having neuroprotection as a primary outcome measure. Furthermore, the scope of this review was limited to neuropsychiatric disorders with reference to neuroprotection in other conditions such as neurodegenerative, cerebrovascular and traumatic brain injuries only where directly relevant to discussion. In order for neuroprotection to be assessed with

confidence, prospective analysis with neuroprotection as the primary outcome and multi-modal analysis including assessment of neurostructural benefit, symptomatic/cognitive effect and biologic/genetic markers of mechanism are required.

Conflicts of Interest

Professor Michael Berk (Co-Author)

Grant/Research Support:

Stanley Medical Research Foundation NIH
MBF
NHMRC

NHMRC Senior Principal Research Fellowship 1059660
Cooperative Research Centre
Simons Autism Foundation

Cancer Council of Victoria, MBF
Rotary Health
Meat and Livestock Board Woolworths

BeyondBlue
Geelong Medical Research Foundation Bristol Myers Squibb
Eli Lilly
Glaxo SmithKline
Organon
Novartis
Mayne Pharma
Servier

Speaker:

Astra Zeneca
Bristol Myers Squibb Eli Lilly
Glaxo SmithKline Lundbeck
Pfizer
Sanofi Synthelabo Servier
Solvay
Wyeth

Consultant:

AstraZeneca
Bristol Myers Squibb Eli Lilly
Bioadvantex
Merck
Glaxo SmithKline Lundbeck
Janssen Cilag Servier

Patents:

Michael Berk is a co-inventor of two provisional patents regarding the use of NAC and related compounds for psychiatric indications, which, while assigned to the Mental Health Research Institute, could lead to personal remuneration upon a commercialisation event.

References

- Abbasi, S. H., Mohammadinejad, P., Shahmansouri, N., Salehiomran, A., Beglar, A. A., Zeinoddini, A., . . . Akhondzadeh, S. (2015). Simvastatin versus atorvastatin for improving mild to moderate depression in post-coronary artery bypass graft patients: A double-blind, placebo-controlled, randomized trial. *J Affect Disord*, 183, 149-155. doi:10.1016/j.jad.2015.04.049
- Ahmed, M., Cannon, D. M., Scanlon, C., Holleran, L., Schmidt, H., McFarland, J., . . . McDonald, C. (2015). Progressive Brain Atrophy and Cortical Thinning in Schizophrenia after Commencing Clozapine Treatment. *Neuropsychopharmacology*, 40(10), 2409-2417. doi:10.1038/npp.2015.90
- Alboni, S., Poggini, S., Garofalo, S., Milior, G., El Hajj, H., Lecours, C., . . . Branchi, I. (2016). Fluoxetine treatment affects the inflammatory response and microglial function according to the quality of the living environment. *Brain Behav Immun*, 58, 261-271. doi:10.1016/j.bbi.2016.07.155
- Alboni, S., van Dijk, R. M., Poggini, S., Milior, G., Perrotta, M., Drenth, T., . . . Branchi, I. (2017). Fluoxetine effects on molecular, cellular and behavioral endophenotypes of depression are driven by the living environment. *Mol Psychiatry*, 22(4), 552-561. doi:10.1038/mp.2015.142
- Amiri, S., Haj-Mirzaian, A., Amini-Khoei, H., Razmi, A., Shirzadian, A., Rahimi-Balaei, M., . . . Ghazi-Khansari, M. (2017). Protective effects of gabapentin against the seizure susceptibility and comorbid behavioral abnormalities in the early socially isolated mice. *European Journal of Pharmacology*, 797, 106-114. doi:10.1016/j.ejphar.2017.01.024
- Arabzadeh, S., Ameli, N., Zeinoddini, A., Rezaei, F., Farokhnia, M., Mohammadinejad, P., . . . Akhondzadeh, S. (2015). Celecoxib adjunctive therapy for acute bipolar mania: a randomized, double-blind, placebo-controlled trial. *Bipolar Disord*, 17(6), 606-614. doi:10.1111/bdi.12324
- Bentsen, H., Osnes, K., Refsum, H., Solberg, D. K., & Bohmer, T. (2013). A randomized placebo-controlled trial of an omega-3 fatty acid and vitamins E plus C in schizophrenia. *Transl Psychiatry*, 3. doi:10.1038/tp.2013.110
- Berger, M. E., Smesny, S., Kim, S. W., Davey, C. G., Rice, S., Sarnyai, Z., . . . Amminger, G. P. (2017). Omega-6 to omega-3 polyunsaturated fatty acid ratio and subsequent mood disorders in young people with at-risk mental states: a 7-year longitudinal study. *Transl Psychiatry*, 7(8), e1220. doi:10.1038/tp.2017.190

- Berk, M., Dandash, O., Daglas, R., Cotton, S. M., Allott, K., Fornito, A., . . . Yucel, M. (2017). Neuroprotection after a first episode of mania: a randomized controlled maintenance trial comparing the effects of lithium and quetiapine on grey and white matter volume. *Transl Psychiatry*, 7(1), e1011. doi:10.1038/tp.2016.281
- Berk, M., Dean, O., Drexhage, H., McNeil, J. J., Moylan, S., O'Neil, A., . . . Maes, M. (2013). Aspirin: a review of its neurobiological properties and therapeutic potential for mental illness. *BMC Med*, 11, 74. doi:10.1186/1741-7015-11-74
- Bigornia, S. J., Harris, W. S., Falcon, L. M., Ordovas, J. M., Lai, C. Q., & Tucker, K. L. (2016). The Omega-3 Index Is Inversely Associated with Depressive Symptoms among Individuals with Elevated Oxidative Stress Biomarkers. *J Nutr*, 146(4), 758-766. doi:10.3945/jn.115.222562
- Bond, D. J., Andreazza, A. C., Hughes, J., Dhanoa, T., Torres, I. J., Kozicky, J. M., . . . Yatham, L. N. (2017). A Longitudinal Study of the Relationships Between Mood Symptoms, Body Mass Index, and Serum Adipokines in Bipolar Disorder. *Journal of Clinical Psychiatry*, 78(4), 441-448. doi:10.4088/JCP.15m10189
- Brambilla, P., Bellani, M., Isola, M., Bergami, A., Marinelli, V., Dusi, N., . . . Furlan, R. (2014). Increased M1/decreased M2 signature and signs of Th1/Th2 shift in chronic patients with bipolar disorder, but not in those with schizophrenia. *Transl Psychiatry*, 4, e406. doi:10.1038/tp.2014.46
- Branchi, I., Santarelli, S., Capoccia, S., Poggini, S., D'Andrea, I., Cirulli, F., & Alleva, E. (2013). Antidepressant treatment outcome depends on the quality of the living environment: a pre-clinical investigation in mice. *PLoS One*, 8(4), e62226. doi:10.1371/journal.pone.0062226
- Brown, B. P., Kang, S. C., Gawelek, K., Zacharias, R. A., Anderson, S. R., Turner, C. P., & Morris, J. K. (2015). In vivo and in vitro ketamine exposure exhibits a dose-dependent induction of activity-dependent neuroprotective protein in rat neurons. *Neuroscience*, 290, 31-40. doi:10.1016/j.neuroscience.2014.12.076
- Brown, N. C., Andreazza, A. C., & Young, L. T. (2014). An updated meta-analysis of oxidative stress markers in bipolar disorder. *Psychiatry Res*, 218(1-2), 61-68. doi:10.1016/j.psychres.2014.04.005
- Brugger, S. P., & Howes, O. D. (2017). Heterogeneity and Homogeneity of Regional Brain Structure in Schizophrenia: A Meta-analysis. *JAMA Psychiatry*, 74(11), 1104-1111. doi:10.1001/jamapsychiatry.2017.2663
- Bumb, J. M., Enning, F., Mueller, J. K., van der List, T., Rohleder, C., Findeisen, P., . . . Leweke, F. M. (2016). Differential melatonin alterations in cerebrospinal fluid and serum of patients with major depressive disorder and bipolar disorder. *Comprehensive Psychiatry*, 68, 34-39. doi:10.1016/j.comppsy.2016.03.005

- Cabungcal, J. H., Counotte, D. S., Lewis, E., Tejeda, H. A., Piantadosi, P., Pollock, C., . . . O'Donnell, P. (2014). Juvenile antioxidant treatment prevents adult deficits in a developmental model of schizophrenia. *Neuron*, 83(5), 1073-1084. doi:10.1016/j.neuron.2014.07.028
- Can, O. D., Ulupinar, E., Ozkay, U. D., Yegin, B., & Ozturk, Y. (2012). The effect of simvastatin treatment on behavioral parameters, cognitive performance, and hippocampal morphology in rats fed a standard or a high-fat diet. *Behavioural Pharmacology*, 23(5-6), 582-592. doi:10.1097/FBP.0b013e328356c3f2
- Chaudhry, I. B., Hallak, J., Husain, N., Laganis, C., Minhas, F., Stirling, J., . . . Deakin, J. F. W. (2012). Minocycline benefits negative symptoms in early schizophrenia; A randomised double-blind placebo-controlled clinical trial in patients on a standard treatment. *Early Intervention in Psychiatry*, 6, 100. doi:10.1111/j.1751-7893.2012.00394.x
- Chaves, C., Marque, C. R., Maia-de-Oliveira, J. P., Wichert-Ana, L., Ferrari, T. B., Santos, A. C., . . . Hallak, J. E. (2015). Effects of minocycline add-on treatment on brain morphometry and cerebral perfusion in recent-onset schizophrenia. *Schizophr Res*, 161(2-3), 439-445. doi:10.1016/j.schres.2014.11.031
- Chen, J., Zuo, S., Wang, J., Huang, J., Zhang, X., Liu, Y., . . . Liu, Z. (2014). Aspirin promotes oligodendrocyte precursor cell proliferation and differentiation after white matter lesion. *Frontiers in Aging Neuroscience*, 6(JAN). doi:10.3389/fnagi.2014.00007
- Cheng, Y., Pardo, M., Armini, R. S., Martinez, A., Mouhsine, H., Zagury, J. F., . . . Beurel, E. (2016). Stress-induced neuroinflammation is mediated by GSK3-dependent TLR4 signaling that promotes susceptibility to depression-like behavior. *Brain Behav Immun*, 53, 207-222. doi:10.1016/j.bbi.2015.12.012
- Chiappini, S., & Schifano, F. (2016). A Decade of Gabapentinoid Misuse: An Analysis of the European Medicines Agency's 'Suspected Adverse Drug Reactions' Database. *CNS Drugs*, 30(7), 647-654. doi:10.1007/s40263-016-0359-y
- Chikama, K., Yamada, H., Tsukamoto, T., Kajitani, K., Nakabeppu, Y., & Uchimura, N. (2017). Chronic atypical antipsychotics, but not haloperidol, increase neurogenesis in the hippocampus of adult mouse. *Brain Res*. doi:10.1016/j.brainres.2017.09.006
- Choudhury, A., Singh, S., Palit, G., Shukla, S., & Ganguly, S. (2016). Administration of N-acetylserotonin and melatonin alleviate chronic ketamine-induced behavioural phenotype accompanying BDNF-independent and dependent converging cytoprotective mechanisms in the hippocampus. *Behavioural Brain Research*, 297, 204-212. doi:10.1016/j.bbr.2015.10.027

- Cousins, D. A., Aribisala, B., Ferrier, I. N., & Blamire, A. M. (2013). Lithium, Gray Matter, and Magnetic Resonance Imaging Signal. *Biological Psychiatry*, 73(7), 652-657. doi:10.1016/j.biopsych.2012.09.029
- Daglas, R., Cotton, S. M., Allott, K., Yucel, M., Macneil, C. A., Hasty, M. K., . . . Berk, M. (2016). A single-blind, randomised controlled trial on the effects of lithium and quetiapine monotherapy on the trajectory of cognitive functioning in first episode mania: A 12-month follow-up study. *Eur Psychiatry*, 31, 20-28. doi:10.1016/j.eurpsy.2015.09.460
- Davis, J., Moylan, S., Harvey, B. H., Maes, M., & Berk, M. (2014). Neuroprogression in schizophrenia: Pathways underpinning clinical staging and therapeutic corollaries. *Aust N Z J Psychiatry*, 48(6), 512-529. doi:10.1177/0004867414533012
- De Sousa, R. T., Zanetti, M. V., Talib, L. L., Gattaz, W. F., & Machado-Vieira, R. (2013). Oxidative stress parameters in recent-onset bipolar disorder and possible lithium antioxidant effect. *Biological Psychiatry*, 73(9), 45S.
- de Sousa, R. T., Zarate, C. A., Jr., Zanetti, M. V., Costa, A. C., Talib, L. L., Gattaz, W. F., & Machado-Vieira, R. (2014). Oxidative stress in early stage Bipolar Disorder and the association with response to lithium. *J Psychiatr Res*, 50, 36-41. doi:10.1016/j.jpsychires.2013.11.011
- Dean, O. M., Kanchanatawan, B., Ashton, M., Mohebbi, M., Ng, C. H., Maes, M., . . . Berk, M. (2017). Adjunctive minocycline treatment for major depressive disorder: A proof of concept trial. *Aust N Z J Psychiatry*, 51(8), 829-840. doi:10.1177/0004867417709357
- Deepmala, Slattey, J., Kumar, N., Delhey, L., Berk, M., Dean, O., . . . Frye, R. (2015). Clinical trials of N-acetylcysteine in psychiatry and neurology: A systematic review. *Neurosci Biobehav Rev*, 55, 294-321. doi:10.1016/j.neubiorev.2015.04.015
- Deiana, S., Watanabe, A., Yamasaki, Y., Amada, N., Kikuchi, T., Stott, C., & Riedel, G. (2015). MK-801-induced deficits in social recognition in rats: reversal by aripiprazole, but not olanzapine, risperidone, or cannabidiol. *Behavioural Pharmacology*, 26(8), 748-765. doi:10.1097/fbp.0000000000000178
- Demirci, K., Özçankaya, R., Yilmaz, H. R., Yiğit, A., Uğuz, A. C., Karakuş, K., . . . Akpınar, A. (2015). Paliperidone regulates intracellular redox system in rat brain: Role of purine mechanism. *Redox Report*, 20(4), 170-176. doi:10.1179/1351000214Y.00000000122
- Demirdağ, A., Nazıroğlu, M., & Övey İ, S. (2017). Duloxetine Reduces Oxidative Stress, Apoptosis, and Ca²⁺ Entry Through Modulation of TRPM2 and TRPV1 Channels

- in the Hippocampus and Dorsal Root Ganglion of Rats. *Molecular Neurobiology*, 54(6), 4683-4695. doi:10.1007/s12035-016-9992-1
- Demirdas, A., Naziroglu, M., & Ovey, I. S. (2017). Short-Term Ketamine Treatment Decreases Oxidative Stress Without Influencing TRPM2 and TRPV1 Channel Gating in the Hippocampus and Dorsal Root Ganglion of Rats. *Cellular and Molecular Neurobiology*, 37(1), 133-144. doi:10.1007/s10571-016-0353-4
- Di Biase, M. A., Zalesky, A., O'Keefe, G., Laskaris, L., Baune, B. T., Weickert, C. S., . . . Cropley, V. (2017). PET imaging of putative microglial activation in individuals at ultra-high risk for psychosis, recently diagnosed and chronically ill with schizophrenia. *Transl Psychiatry*, 7(8), e1225. doi:10.1038/tp.2017.193
- Dominguez-Alonso, A., Valdes-Tovar, M., Solis-Chagoyan, H., & Benitez-King, G. (2015). Melatonin stimulates dendrite formation and complexity in the hilar zone of the rat hippocampus: participation of the Ca⁺⁺/Calmodulin complex. *Int J Mol Sci*, 16(1), 1907-1927. doi:10.3390/ijms16011907
- Dong, D., Wang, Y., Chang, X., Jiang, Y., Klugah-Brown, B., Luo, C., & Yao, D. (2017). Shared abnormality of white matter integrity in schizophrenia and bipolar disorder: A comparative voxel-based meta-analysis. *Schizophr Res*, 185, 41-50. doi:10.1016/j.schres.2017.01.005
- Duarte, J. M. d. N., Kulak, A., Gholam-Razaei, M. M., Cuenod, M., Gruetter, R., & Do, K. Q. (2012). N-Acetylcysteine Normalizes Neurochemical Changes in the Glutathione-Deficient Schizophrenia Mouse Model During Development. *Biological Psychiatry*, 71(11), 1006-1014. doi:10.1016/j.biopsych.2011.07.035
- Duffy, S. L., Lagopoulos, J., Cockayne, N., Lewis, S. J. G., Hickie, I. B., Hermens, D. F., & Naismith, S. L. (2015). The effect of 12-wk ω -3 fatty acid supplementation on in vivo thalamus glutathione concentration in patients "at risk" for major depression. *Nutrition*, 31(10), 1247-1254. doi:10.1016/j.nut.2015.04.019
- Dwivedi, T., & Zhang, H. (2014). Lithium-induced neuroprotection is associated with epigenetic modification of specific BDNF gene promoter and altered expression of apoptotic-regulatory proteins. *Front Neurosci*, 8, 457. doi:10.3389/fnins.2014.00457
- Elnahas, E. M., Abou Zeid, M. S., Kawy, H. S. A., Hendawy, N., & Baher, W. (2016). Celecoxib attenuates depressive-like behavior associated with immunological liver injury in C57BL/6 mice through TNF-alpha and NF-kappa b dependent mechanisms. *Life Sciences*, 163, 23-37. doi:10.1016/j.lfs.2016.08.027
- Fernandes, B. S., Dash, S., Jacka, F., Dodd, S., Carvalho, A. F., Kohler, C. A., . . . Berk, M. (2016). Leptin in bipolar disorder: A systematic review and meta-analysis. *Eur Psychiatry*, 35, 1-7. doi:10.1016/j.eurpsy.2016.02.003

- Fernandes, B. S., Steiner, J., Berk, M., Molendijk, M. L., Gonzalez-Pinto, A., Turck, C. W., . . . Goncalves, C. A. (2015). Peripheral brain-derived neurotrophic factor in schizophrenia and the role of antipsychotics: meta-analysis and implications. *Mol Psychiatry*, 20(9), 1108-1119. doi:10.1038/mp.2014.117
- Flatow, J., Buckley, P., & Miller, B. J. (2013). Meta-analysis of oxidative stress in schizophrenia. *Biol Psychiatry*, 74(6), 400-409. doi:10.1016/j.biopsych.2013.03.018
- Forde, N., Ahmed, M., Holleran, L., Scanlon, C., McDonald, C., & Cannon, D. M. (2012). The effect of clozapine on white matter in schizophrenia: A diffusion imaging and tractography study. *International Journal of Neuropsychopharmacology*, 15, 49. doi:10.1017/S1461145712000508
- Fountoulakis, K. N., Yatham, L., Grunze, H., Vieta, E., Young, A., Blier, P., . . . Moeller, H. J. (2017). The International College of Neuro-Psychopharmacology (CINP) Treatment Guidelines for Bipolar Disorder in Adults (CINP-BD-2017), Part 2: Review, Grading of the Evidence, and a Precise Algorithm. *International Journal of Neuropsychopharmacology*, 20(2), 121-179. doi:10.1093/ijnp/pyw100
- Fusar-Poli, P., Smieskova, R., Kempton, M. J., Ho, B. C., Andreasen, N. C., & Borgwardt, S. (2013). Progressive brain changes in schizophrenia related to antipsychotic treatment? A meta-analysis of longitudinal MRI studies. *Neurosci Biobehav Rev*, 37(8), 1680-1691. doi:10.1016/j.neubiorev.2013.06.001
- Galano, A., Tan, D. X., & Reiter, R. J. (2013). On the free radical scavenging activities of melatonin's metabolites, AFMK and AMK. *J Pineal Res*, 54(3), 245-257. doi:10.1111/jpi.12010
- Galecki, P., Galecka, E., Maes, M., Chamielec, M., Orzechowska, A., Bobinska, K., . . . Szemraj, J. (2012). The expression of genes encoding for COX-2, MPO, iNOS, and sPLA2-IIA in patients with recurrent depressive disorder. *J Affect Disord*, 138(3), 360-366. doi:10.1016/j.jad.2012.01.016
- Garza, J. C., Guo, M., Zhang, W., & Lu, X. Y. (2012). Leptin restores adult hippocampal neurogenesis in a chronic unpredictable stress model of depression and reverses glucocorticoid-induced inhibition of GSK-3B/B-catenin signaling. *Mol Psychiatry*, 17(8), 790-808. doi:10.1038/mp.2011.161
- Georgakis, M. K., Thomopoulos, T. P., Diamantaras, A. A., Kalogirou, E. I., Skalkidou, A., Daskalopoulou, S. S., & Petridou, E. T. (2016). Association of Age at Menopause and Duration of Reproductive Period With Depression After Menopause: A Systematic Review and Meta-analysis. *JAMA Psychiatry*, 73(2), 139-149. doi:10.1001/jamapsychiatry.2015.2653

- Ghanizadeh, A., Rezaee, Z., Dehbozorgi, S., Berk, M., & Akhondzadeh, S. (2014). Lovastatin for the adjunctive treatment of schizophrenia: A preliminary randomized double-blind placebo-controlled trial. *Psychiatry Research*, 219(3), 431-435. doi:10.1016/j.psychres.2014.06.039
- Giakoumatos, C. I., Nanda, P., Mathew, I. T., Tandon, N., Shah, J., Bishop, J. R., . . . Keshavan, M. S. (2015). Effects of lithium on cortical thickness and hippocampal subfield volumes in psychotic bipolar disorder. *J Psychiatr Res*, 61, 180-187. doi:10.1016/j.jpsychires.2014.12.008
- Gildengers, A. G., Butters, M. A., Aizenstein, H. J., Marron, M. M., Emanuel, J., Anderson, S. J., . . . Reynolds, C. F., 3rd. (2015). Longer lithium exposure is associated with better white matter integrity in older adults with bipolar disorder. *Bipolar Disord*, 17(3), 248-256. doi:10.1111/bdi.12260
- Gleixner, A. M., Hutchison, D. F., Sannino, S., Bhatia, T. N., Leak, L. C., Flaherty, P. T., . . . Leak, R. K. (2017). N-Acetyl-L-Cysteine Protects Astrocytes against Proteotoxicity without Recourse to Glutathione. *Mol Pharmacol*, 92(5), 564-575. doi:10.1124/mol.117.109926
- Glombik, K., Stachowicz, A., Trojan, E., Olszanecki, R., Slusarczyk, J., Suski, M., . . . Basta-Kaim, A. (2017). Evaluation of the effectiveness of chronic antidepressant drug treatments in the hippocampal mitochondria - A proteomic study in an animal model of depression. *Prog Neuropsychopharmacol Biol Psychiatry*, 78, 51-60. doi:10.1016/j.pnpbp.2017.05.014
- Goldsmith, D. R., Rapaport, M. H., & Miller, B. J. (2016). A meta-analysis of blood cytokine network alterations in psychiatric patients: comparisons between schizophrenia, bipolar disorder and depression. *Mol Psychiatry*, 21(12), 1696-1709. doi:10.1038/mp.2016.3
- Guan, X. T., Shao, F., Xie, X., Chen, L., & Wang, W. W. (2014). Effects of aspirin on immobile behavior and endocrine and immune changes in the forced swimming test: Comparison to fluoxetine and imipramine. *Pharmacology Biochemistry and Behavior*, 124, 361-366. doi:10.1016/j.pbb.2014.07.002
- Gumuslu, E., Mutlu, O., Ulak, G., Uyar, E., Tatar, O. C., Celikyurt, I. K., . . . Erden, F. (2015). Effects of atypical antipsychotics on emotional memory and locomotion: Altered gene expression levels of BDNF and CREB in the hippocampus of mice. *European Neuropsychopharmacology*, 25, S161-S162.
- Guo, M., Huang, T. Y., Garza, J. C., Chua, S. C., & Lu, X. Y. (2013). Selective deletion of leptin receptors in adult hippocampus induces depression-related behaviours. *International Journal of Neuropsychopharmacology*, 16(4), 857-867. doi:10.1017/s1461145712000703

- Haijma, S. V., Van Haren, N., Cahn, W., Koolschijn, P. C. M. P., Hulshoff Pol, H. E., & Kahn, R. S. (2013). Brain Volumes in Schizophrenia: A Meta-Analysis in Over 18 000 Subjects. *Schizophrenia Bulletin*, 39(5), 1129-1138. doi:10.1093/schbul/sbs118
- Hajek, T., Bauer, M., Simhandl, C., Rybakowski, J., O'Donovan, C., Pfennig, A., . . . Alda, M. (2014). Neuroprotective effect of lithium on hippocampal volumes in bipolar disorder independent of long-term treatment response. *Psychological Medicine*, 44(3), 507-517. doi:10.1017/s0033291713001165
- Hajek, T., Cullis, J., Novak, T., Kopecek, M., Höschl, C., Blagdon, R., . . . Alda, M. (2012). Hippocampal volumes in bipolar disorders: Opposing effects of illness burden and lithium treatment. *Disorders*, 14(3), 261-270. doi:10.1111/j.1399-5618.2012.01013.x
- Halaris, A., Myint, A. M., Meresh, E., Savant, V., Guillemin, G., & Sinacore, J. (2012). The tryptophan-kynurenine pathway in major depression: Neuroprotective effects of treatment. *Neuropsychopharmacology*, 38, S425-S426. doi:10.1038/npp.2012.221
- Hannon, E., Dempster, E., Viana, J., Burrage, J., Smith, A. R., Macdonald, R., . . . Mill, J. (2016). An integrated genetic-epigenetic analysis of schizophrenia: evidence for co-localization of genetic associations and differential DNA methylation. *Genome Biol*, 17(1), 176. doi:10.1186/s13059-016-1041-x
- Hartberg, C. B., Jorgensen, K. N., Haukvik, U. K., Westlye, L. T., Melle, I., Andreassen, O. A., & Agartz, I. (2015). Lithium treatment and hippocampal subfields and amygdala volumes in bipolar disorder. *Bipolar Disord*, 17(5), 496-506. doi:10.1111/bdi.12295
- Hassouna, I., Ott, C., Wüstefeld, L., Offen, N., Neher, R. A., Mitkovski, M., . . . Ehrenreich, H. (2016). Revisiting adult neurogenesis and the role of erythropoietin for neuronal and oligodendroglial differentiation in the hippocampus. *Molecular Psychiatry*, 21(12), 1752-1767. doi:10.1038/mp.2015.212
- Herbet, M., Izdebska, M., Piątkowska-Chmiel, I., Poleszak, E., & Jagiełło-Wójtowicz, E. (2016). Estimation of oxidative stress parameters in rats after simultaneous administration of rosuvastatin with antidepressants. *Pharmacological Reports*, 68(1), 172-176. doi:10.1016/j.pharep.2015.08.004
- Hjelm, B. E., Rollins, B., Mamdani, F., Lauterborn, J. C., Kirov, G., Lynch, G., . . . Vawter, M. P. (2015). Evidence of Mitochondrial Dysfunction within the Complex Genetic Etiology of Schizophrenia. *Mol Neuropsychiatry*, 1(4), 201-219. doi:10.1159/000441252

- Holmes, S. E., Hinz, R., Conen, S., Gregory, C. J., Matthews, J. C., Anton-Rodriguez, J. M., . . . Talbot, P. S. (2017). Elevated Translocator Protein in Anterior Cingulate in Major Depression and a Role for Inflammation in Suicidal Thinking: A Positron Emission Tomography Study. *Biol Psychiatry*. doi:10.1016/j.biopsych.2017.08.005
- Holmes, S. E., Hinz, R., Drake, R. J., Gregory, C. J., Conen, S., Matthews, J. C., . . . Talbot, P. S. (2016). In vivo imaging of brain microglial activity in antipsychotic-free and medicated schizophrenia: a [(11)C](R)-PK11195 positron emission tomography study. *Mol Psychiatry*, 21(12), 1672-1679. doi:10.1038/mp.2016.180
- Hori, H., Yoshimura, R., Katsuki, A., Hayashi, K., Ikenouchi-Sugita, A., Umene-Nakano, W., & Nakamura, J. (2012). The cognitive profile of aripiprazole differs from that of other atypical antipsychotics in schizophrenia patients. *J Psychiatr Res*, 46(6), 757-761. doi:<https://doi.org/10.1016/j.jpsychires.2012.02.013>
- Howes, O., McCutcheon, R., & Stone, J. (2015). Glutamate and dopamine in schizophrenia: An update for the 21st century. *Journal of Psychopharmacology*, 29(2), 97-115. doi:10.1177/0269881114563634
- Huang, N. X., Chen, D., Wu, X. Y., Chen, X. J., Zhang, X. S., Niu, J. Q., . . . Xiao, L. (2016). Aspirin Promotes Oligodendroglial Differentiation Through Inhibition of Wnt Signaling Pathway. *Molecular Neurobiology*, 53(5), 3258-3266. doi:10.1007/s12035-015-9241-z
- Husain, M. I., Chaudhry, I. B., Husain, N., Khoso, A. B., Rahman, R. R., Hamirani, M. M., . . . Young, A. H. (2017). Minocycline as an adjunct for treatment-resistant depressive symptoms: A pilot randomised placebo-controlled trial. *J Psychopharmacol*, 31(9), 1166-1175. doi:10.1177/0269881117724352
- Ibrahim, W. W., Safar, M. M., Khattab, M. M., & Agha, A. M. (2016). 17beta-Estradiol augments antidepressant efficacy of escitalopram in ovariectomized rats: Neuroprotective and serotonin reuptake transporter modulatory effects. *Psychoneuroendocrinology*, 74, 240-250. doi:10.1016/j.psyneuen.2016.09.013
- Ignácio, Z. M., Réus, G. Z., Abelaira, H. M., de Moura, A. B., de Souza, T. G., Matos, D., . . . Quevedo, J. (2017). Acute and chronic treatment with quetiapine induces antidepressant-like behavior and exerts antioxidant effects in the rat brain. *Metabolic Brain Disease*, 32(4), 1195-1208. doi:10.1007/s11011-017-0028-y
- Ignacio, Z. M., Reus, G. Z., Abelaira, H. M., Titus, S. E., Carlessi, A. S., da Luz, J. R., . . . Quevedo, J. (2015). Acute and Chronic Treatments with Quetiapine Increase Mitochondrial Respiratory Chain Complex Activity in the Rat Brain. *Curr Neurovasc Res*, 12(3), 283-292.

- Jiang, Y., Rumble, J. L., Gleixner, A. M., Unnithan, A. S., Pulugulla, S. H., Posimo, J. M., . . . Leak, R. K. (2013). N-Acetyl cysteine blunts proteotoxicity in a heat shock protein-dependent manner. *Neuroscience*, 255, 19-32. doi:10.1016/j.neuroscience.2013.09.049
- Jimenez-Fernandez, S., Gurpegui, M., Diaz-Atienza, F., Perez-Costillas, L., Gerstenberg, M., & Correll, C. U. (2015). Oxidative stress and antioxidant parameters in patients with major depressive disorder compared to healthy controls before and after antidepressant treatment: results from a meta-analysis. *J Clin Psychiatry*, 76(12), 1658-1667. doi:10.4088/JCP.14r09179
- Kanchanatawan, B., Hemrungrojn, S., Thika, S., Sirivichayakul, S., Ruxrungtham, K., Carvalho, A. F., . . . Maes, M. (2017). Changes in Tryptophan Catabolite (TRYCAT) Pathway Patterning Are Associated with Mild Impairments in Declarative Memory in Schizophrenia and Deficits in Semantic and Episodic Memory Coupled with Increased False-Memory Creation in Deficit Schizophrenia. *Mol Neurobiol*. doi:10.1007/s12035-017-0751-8
- Kasahara, T., & Kato, T. (2017). What Can Mitochondrial DNA Analysis Tell Us About Mood Disorders? *Biol Psychiatry*. doi:10.1016/j.biopsych.2017.09.010
- Khan, M. S., Ali, T., Abid, M. N., Jo, M. H., Khan, A., Kim, M. W., . . . Kim, M. O. (2017). Lithium ameliorates lipopolysaccharide-induced neurotoxicity in the cortex and hippocampus of the adult rat brain. *Neurochem Int*, 108, 343-354. doi:10.1016/j.neuint.2017.05.008
- Kim, H.-W., Cheon, Y., Modi, H. R., Rapoport, S. I., & Rao, J. S. (2012). Effects of chronic clozapine administration on markers of arachidonic acid cascade and synaptic integrity in rat brain. *Psychopharmacology*, 222(4), 663-674. doi:10.1007/s00213-012-2671-7
- Kim, S. W., Bae, K. Y., Kim, J. M., Shin, I. S., Hong, Y. J., Ahn, Y., . . . Yoon, J. S. (2015). The use of statins for the treatment of depression in patients with acute coronary syndrome. *Transl Psychiatry*, 5, e620. doi:10.1038/tp.2015.116
- Kohler, O., Benros, M. E., Nordentoft, M., Farkouh, M. E., Iyengar, R. L., Mors, O., & Krogh, J. (2014). Effect of anti-inflammatory treatment on depression, depressive symptoms, and adverse effects: a systematic review and meta-analysis of randomized clinical trials. *JAMA Psychiatry*, 71(12), 1381-1391. doi:10.1001/jamapsychiatry.2014.1611
- Kraus, C., Rabl, U., Vanicek, T., Carlberg, L., Popovic, A., Spies, M., . . . Kasper, S. (2017). Administration of ketamine for unipolar and bipolar depression. *International Journal of Psychiatry in Clinical Practice*, 21(1), 2-12. doi:10.1080/13651501.2016.1254802

- Kraus, M. F., Susmaras, T., Caughlin, B. P., Walker, C. J., Sweeney, J. A., & Little, D. M. (2007). White matter integrity and cognition in chronic traumatic brain injury: a diffusion tensor imaging study. *Brain*, 130(Pt 10), 2508-2519. doi:10.1093/brain/awm216
- Kreisel, T., Frank, M. G., Licht, T., Reshef, R., Ben-Menachem-Zidon, O., Baratta, M. V., . . . Yirmiya, R. (2014). Dynamic microglial alterations underlie stress-induced depressive-like behavior and suppressed neurogenesis. *Mol Psychiatry*, 19(6), 699-709. doi:10.1038/mp.2013.155
- Krivoy, A., Hochman, E., Sendt, K. V., Hollander, S., Vilner, Y., Selakovic, M., . . . Taler, M. (2017). Association between serum levels of glutamate and neurotrophic factors and response to clozapine treatment. *Schizophr Res*. doi:10.1016/j.schres.2017.05.040
- Kudlek Mikulic, S., Mihaljevic-Peles, A., Sagud, M., Bajs Janovic, M., Ganoci, L., Grubisin, J., . . . Bozina, N. (2017). Brain-derived neurotrophic factor serum and plasma levels in the treatment of acute schizophrenia with olanzapine or risperidone: 6-week prospective study. *Nord J Psychiatry*, 71(7), 513-520. doi:10.1080/08039488.2017.1340518
- Levkovitz, Y., Fenchel, D., Kaplan, Z., Zohar, J., & Cohen, H. (2015). Early post-stressor intervention with minocycline, a second-generation tetracycline, attenuates post-traumatic stress response in an animal model of PTSD. *European Neuropsychopharmacology*, 25(1), 124-132. doi:10.1016/j.euroneuro.2014.11.012
- Li, Y. F., Yan, J. C., Zhu, X. L., Zhu, Y., Yao, S., Xu, Y. H., & Ju, S. H. (2017). Dilated Virchow-Robin spaces in the hippocampus impact behaviors and effects of antidepressant treatment in model of depressed rats. *J Affect Disord*, 219, 17-24. doi:10.1016/j.jad.2017.04.035
- Lin, P. Y., Chang, A. Y. W., & Lin, T. K. (2014). Simvastatin treatment exerts antidepressant-like effect in rats exposed to chronic mild stress. *Pharmacology Biochemistry and Behavior*, 124, 174-179. doi:10.1016/j.pbb.2014.06.006
- Liu, D. X., Wang, Z., Gao, Z., Xie, K., Zhang, Q. R., Jiang, H., & Pang, Q. (2014). Effects of curcumin on learning and memory deficits, BDNF, and ERK protein expression in rats exposed to chronic unpredictable stress. *Behavioural Brain Research*, 271, 116-121. doi:10.1016/j.bbr.2014.05.068
- Liu, S. B., Zhang, N., Guo, Y. Y., Zhao, R., Shi, T. Y., Feng, S. F., . . . Zhao, M. G. (2012). G-protein-coupled receptor 30 mediates rapid neuroprotective effects of estrogen via depression of NR2B-containing NMDA receptors. *J Neurosci*, 32(14), 4887-4900. doi:10.1523/jneurosci.5828-11.2012

- Liu, W. X., Wang, J., Xie, Z. M., Xu, N., Zhang, G. F., Jia, M., . . . Yang, J. J. (2016). Regulation of glutamate transporter 1 via BDNF-TrkB signaling plays a role in the anti-apoptotic and antidepressant effects of ketamine in chronic unpredictable stress model of depression. *Psychopharmacology*, 233(3), 405-415. doi:10.1007/s00213-015-4128-2
- Lopresti, A. L., Maes, M., Maker, G. L., Hood, S. D., & Drummond, P. D. (2014). Curcumin for the treatment of major depression: a randomised, double-blind, placebo controlled study. *J Affect Disord*, 167, 368-375. doi:10.1016/j.jad.2014.06.001
- Ludka, F. K., Constantino, L. C., Dal-Cim, T., Binder, L. B., Zomkowski, A., Rodrigues, A. L., & Tasca, C. I. (2016). Involvement of PI3K/Akt/GSK-3 β and mTOR in the antidepressant-like effect of atorvastatin in mice. *J Psychiatr Res*, 82, 50-57. doi:10.1016/j.jpsychires.2016.07.004
- Ludka, F. K., Cunha, M. P., Dal-Cim, T., Binder, L. B., Constantino, L. C., Massari, C. M., . . . Tasca, C. I. (2017). Atorvastatin Protects from A β 1-40-Induced Cell Damage and Depressive-Like Behavior via ProBDNF Cleavage. *Molecular Neurobiology*, 54(8), 6163-6173. doi:10.1007/s12035-016-0134-6
- Luo, Y., Kuang, S. N., Li, H., Ran, D. Z., & Yang, J. Q. (2017). cAMP/PKA-CREB-BDNF signaling pathway in hippocampus mediates cyclooxygenase 2-induced learning/memory deficits of rats subjected to chronic unpredictable mild stress. *Oncotarget*, 8(22), 35558-35572. doi:10.18632/oncotarget.16009
- MacDowell, K. S., Munarriz-Cuezva, E., Caso, J. R., Madrigal, J. L., Zabala, A., Meana, J. J., . . . Leza, J. C. (2017). Paliperidone reverts Toll-like receptor 3 signaling pathway activation and cognitive deficits in a maternal immune activation mouse model of schizophrenia. *Neuropharmacology*, 116, 196-207. doi:10.1016/j.neuropharm.2016.12.025
- Machado, M. O., Oriolo, G., Bortolato, B., Kohler, C. A., Maes, M., Solmi, M., . . . Carvalho, A. F. (2017). Biological mechanisms of depression following treatment with interferon for chronic hepatitis C: A critical systematic review. *J Affect Disord*, 209, 235-245. doi:10.1016/j.jad.2016.11.039
- Markoutsas, E., & Xu, P. (2017). Redox Potential-Sensitive N-Acetyl Cysteine-Prodrug Nanoparticles Inhibit the Activation of Microglia and Improve Neuronal Survival. *Molecular Pharmaceutics*, 14(5), 1591-1600. doi:10.1021/acs.molpharmaceut.6b01028
- Masaki, C., Sharpley, A. L., Cooper, C. M., Godlewska, B. R., Singh, N., Vasudevan, S. R., . . . Cowen, P. J. (2016). Effects of the potential lithium-mimetic, ebselen, on

- impulsivity and emotional processing. *Psychopharmacology (Berl)*, 233(14), 2655-2661. doi:10.1007/s00213-016-4319-5
- Masaki, C., Sharpley, A. L., Godlewska, B. R., Berrington, A., Hashimoto, T., Singh, N., . . . Cowen, P. J. (2016). Effects of the potential lithium-mimetic, ebselen, on brain neurochemistry: a magnetic resonance spectroscopy study at 7 tesla. *Psychopharmacology (Berl)*, 233(6), 1097-1104. doi:10.1007/s00213-015-4189-2
- Matsuoka, Y., Nishi, D., Hamazaki, K., Yonemoto, N., Matsumura, K., Noguchi, H., . . . Hamazaki, T. (2015). Docosahexaenoic Acid for Selective Prevention of Posttraumatic Stress Disorder Among Severely Injured Patients: A Randomized, Placebo-Controlled Trial. *Journal of Clinical Psychiatry*, 76(8), E1015-E1022. doi:10.4088/JCP.14m09260
- Mattei, D., Djodari-Irani, A., Hadar, R., Pelz, A., de Cossio, L. F., Goetz, T., . . . Wolf, S. A. (2014). Minocycline rescues decrease in neurogenesis, increase in microglia cytokines and deficits in sensorimotor gating in an animal model of schizophrenia. *Brain Behav Immun*, 38, 175-184. doi:10.1016/j.bbi.2014.01.019
- Mattei, D., Ivanov, A., Ferrai, C., Jordan, P., Guneykaya, D., Buonfiglioli, A., . . . Wolf, S. A. (2017). Maternal immune activation results in complex microglial transcriptome signature in the adult offspring that is reversed by minocycline treatment. *Transl Psychiatry*, 7. doi:10.1038/tp.2017.80
- McGorry, P. D., Nelson, B., Markulev, C., Yuen, H. P., Schafer, M. R., Mossaheb, N., . . . Amminger, G. P. (2017). Effect of omega-3 Polyunsaturated Fatty Acids in Young People at Ultrahigh Risk for Psychotic Disorders: The NEURAPRO Randomized Clinical Trial. *JAMA Psychiatry*, 74(1), 19-27. doi:10.1001/jamapsychiatry.2016.2902
- Miyaoka, T., Wake, R., Furuya, M., Liaury, K., Ieda, M., Kawakami, K., . . . Horiguchi, J. (2012). Minocycline as adjunctive therapy for patients with unipolar psychotic depression: an open-label study. *Prog Neuropsychopharmacol Biol Psychiatry*, 37(2), 222-226. doi:10.1016/j.pnpbp.2012.02.002
- Modabbernia, A., Heidari, P., Soleimani, R., Sobhani, A., Roshan, Z. A., Taslimi, S., . . . Modabbernia, M. J. (2014). Melatonin for prevention of metabolic side-effects of olanzapine in patients with first-episode schizophrenia: randomized double-blind placebo-controlled study. *J Psychiatr Res*, 53, 133-140. doi:10.1016/j.jpsychires.2014.02.013
- Molendijk, M. L., Spinhoven, P., Polak, M., Bus, B. A., Penninx, B. W., & Elzinga, B. M. (2014). Serum BDNF concentrations as peripheral manifestations of depression:

- evidence from a systematic review and meta-analyses on 179 associations (N=9484). *Mol Psychiatry*, 19(7), 791-800. doi:10.1038/mp.2013.105
- Moller, M., Du Preez, J. L., Emsley, R., & Harvey, B. H. (2012). Social isolation rearing in rats alters plasma tryptophan metabolism and is reversed by sub-chronic clozapine treatment. *Neuropharmacology*, 62(8), 2499-2506. doi:10.1016/j.neuropharm.2012.02.021
- Moller, M., Du Preez, J. L., Viljoen, F. P., Berk, M., Emsley, R., & Harvey, B. H. (2013). Social isolation rearing induces mitochondrial, immunological, neurochemical and behavioural deficits in rats, and is reversed by clozapine or N-acetyl cysteine. *Brain Behav Immun*, 30, 156-167. doi:10.1016/j.bbi.2012.12.011
- Moller, M., Du Preez, J. L., Viljoen, F. P., Berk, M., & Harvey, B. H. (2014). Clozapine alone and in combination with N-acetylcysteine reverses social isolation rearing induced serotonergic metabolism. *European Neuropsychopharmacology*, 24, S563-S564.
- Monroy-Jaramillo, N., Rodriguez-Agudelo, Y., Avina-Cervantes, L. C., Roberts, D. L., Velligan, D. I., & Walss-Bass, C. (2017). Leukocyte telomere length in Hispanic schizophrenia patients under treatment with olanzapine. *J Psychiatr Res*, 90, 26-30. doi:10.1016/j.jpsychires.2017.02.007
- Monte, A. S., De Souza, G. C., McIntyre, R. S., Soczynska, J. K., Dos Santos, J. V., Cordeiro, R. C., . . . Macêdo, D. S. (2013). Prevention and reversal of ketamine-induced schizophrenia related behavior by minocycline in mice: Possible involvement of antioxidant and nitrenergic pathways. *Journal of Psychopharmacology*, 27(11), 1032-1043. doi:10.1177/0269881113503506
- Morais, M., Patricio, P., Mateus-Pinheiro, A., Alves, N. D., Machado-Santos, A. R., Correia, J. S., . . . Bessa, J. M. (2017). The modulation of adult neuroplasticity is involved in the mood-improving actions of atypical antipsychotics in an animal model of depression. *Transl Psychiatry*, 7(6), e1146. doi:10.1038/tp.2017.120
- Morgan, C. J., Dodds, C. M., Furby, H., Pepper, F., Fam, J., Freeman, T. P., . . . Stone, J. M. (2014). Long-Term Heavy Ketamine Use is Associated with Spatial Memory Impairment and Altered Hippocampal Activation. *Front Psychiatry*, 5, 149. doi:10.3389/fpsy.2014.00149
- Morris, G., Carvalho, A. F., Anderson, G., Galecki, P., & Maes, M. (2016). The Many Neuroprogressive Actions of Tryptophan Catabolites (TRYCATs) that may be Associated with the Pathophysiology of Neuro-Immune Disorders. *Curr Pharm Des*, 22(8), 963-977.
- Morris, G., & Maes, M. (2014). Mitochondrial dysfunctions in myalgic encephalomyelitis/chronic fatigue syndrome explained by activated immuno-

- inflammatory, oxidative and nitrosative stress pathways. *Metab Brain Dis*, 29(1), 19-36. doi:10.1007/s11011-013-9435-x
- Motaghinejad, M., Seyedjavadein, Z., Motevalian, M., & Asadi, M. (2016). The neuroprotective effect of lithium against high dose methylphenidate: Possible role of BDNF. *Neurotoxicology*, 56, 40-54. doi:10.1016/j.neuro.2016.06.010
- Mouchlianitis, E., McCutcheon, R., & Howes, O. D. (2016). Brain-imaging studies of treatment-resistant schizophrenia: a systematic review. *Lancet Psychiatry*, 3(5), 451-463. doi:10.1016/s2215-0366(15)00540-4
- Mousavi, S. Y., Khezri, R., Karkhaneh-Yousefi, M. A., Mohammadinejad, P., Gholamian, F., Mohammadi, M. R., . . . Akhondzadeh, S. (2017). A Randomized, Double-Blind Placebo-Controlled Trial on Effectiveness and Safety of Celecoxib Adjunctive Therapy in Adolescents with Acute Bipolar Mania. *J Child Adolesc Psychopharmacol*, 27(6), 494-500. doi:10.1089/cap.2016.0207
- Moylan, S., Berk, M., Dean, O. M., Samuni, Y., Williams, L. J., O'Neil, A., . . . Maes, M. (2014). Oxidative & nitrosative stress in depression: why so much stress? *Neurosci Biobehav Rev*, 45, 46-62. doi:10.1016/j.neubiorev.2014.05.007
- Munkholm, K., Vinberg, M., & Kessing, L. V. (2016). Peripheral blood brain-derived neurotrophic factor in bipolar disorder: a comprehensive systematic review and meta-analysis. *Mol Psychiatry*, 21(2), 216-228. doi:10.1038/mp.2015.54
- Naksuriya, O., Okonogi, S., Schiffelers, R. M., & Hennink, W. E. (2014). Curcumin nanoformulations: a review of pharmaceutical properties and preclinical studies and clinical data related to cancer treatment. *Biomaterials*, 35(10), 3365-3383. doi:10.1016/j.biomaterials.2013.12.090
- Neelamekam, S., Nurjono, M., & Lee, J. (2014). Regulation of Interleukin-6 and Leptin in Schizophrenia Patients: A Preliminary Analysis. *Clinical Psychopharmacology and Neuroscience*, 12(3), 209-214. doi:10.9758/cpn.2014.12.3.209
- Noto, C., Ota, V. K., Gadelha, A., Noto, M. N., Barbosa, D. S., Bonifácio, K. L., . . . Brietzke, E. (2015). Oxidative stress in drug naïve first episode psychosis and antioxidant effects of risperidone. *J Psychiatr Res*, 68, 210-216. doi:10.1016/j.jpsychires.2015.07.003
- Nowakowska, E., Kus, K., Ratajczak, P., Cichocki, M., & Woźniak, A. (2014). The influence of aripiprazole, olanzapine and enriched environment on depressant-like behavior, spatial memory dysfunction and hippocampal level of BDNF in prenatally stressed rats. *Pharmacological Reports*, 66(3), 404-411. doi:10.1016/j.pharep.2013.12.008
- Nucifora, L. G., Tanaka, T., Hayes, L. N., Kim, M., Lee, B. J., Matsuda, T., . . . Sawa, A. (2017). Reduction of plasma glutathione in psychosis associated with

- schizophrenia and bipolar disorder in translational psychiatry. *Transl Psychiatry*, 7(8), e1215. doi:10.1038/tp.2017.178
- Ogłodek, E. A., Just, M. J., Szromek, A. R., & Araszkiewicz, A. (2016). Melatonin and neurotrophins NT-3, BDNF, NGF in patients with varying levels of depression severity. *Pharmacological Reports*, 68(5), 945-951. doi:10.1016/j.pharep.2016.04.003
- Osborn, M., Rustom, N., Clarke, M., Litteljohn, D., Rudyk, C., Anisman, H., & Hayley, S. (2013). Antidepressant-like effects of erythropoietin: a focus on behavioural and hippocampal processes. *PLoS One*, 8(9), e72813. doi:10.1371/journal.pone.0072813
- Ott, C. V., Vinberg, M., Kessing, L. V., & Miskowiak, K. W. (2016). The effect of erythropoietin on cognition in affective disorders - associations with baseline deficits and change in subjective cognitive complaints. *European Neuropsychopharmacology*, 26, S381-S382.
- Paknejad, B., Kebriaeezadeh, A., Ghahremani, M. H., Gharghabi, M., & Ostad, S. N. (2015). Gabapentin Enhances Neurogenesis in E14 Rat Embryonic Neocortex Stem Cells. *Drug Research*, 65(8), 398-402. doi:10.1055/s-0034-1375608
- Pawelczyk, T., Grancow-Grabka, M., Trafalska, E., Szemraj, J., & Pawelczyk, A. (2017). Oxidative stress reduction related to the efficacy of n-3 polyunsaturated fatty acids in first episode schizophrenia: Secondary outcome analysis of the OFFER randomized trial. *Prostaglandins Leukotrienes and Essential Fatty Acids*, 121, 7-13. doi:10.1016/j.plefa.2017.05.004
- Pawelczyk, T., Trafalska, E., Kotlicka-Antczak, M., & Pawelczyk, A. (2016). The association between polyunsaturated fatty acid consumption and the transition to psychosis in ultra-high risk individuals. *Prostaglandins Leukotrienes and Essential Fatty Acids*, 108, 30-37. doi:10.1016/j.plefa.2016.03.010
- Peckham, A. M., Fairman, K. A., & Sclar, D. A. (2017). Prevalence of Gabapentin Abuse: Comparison with Agents with Known Abuse Potential in a Commercially Insured US Population. *Clin Drug Investig*, 37(8), 763-773. doi:10.1007/s40261-017-0530-3
- Pettit, L. K., Varsanyi, C., Tadros, J., & Vassiliou, E. (2013). Modulating the inflammatory properties of activated microglia with Docosahexaenoic acid and Aspirin. *Lipids Health Dis*, 12, 16. doi:10.1186/1476-511x-12-16
- Pfennig, A., Alda, M., Young, T., MacQueen, G., Rybakowski, J., Suwalska, A., . . . Bauer, M. (2014). Prophylactic lithium treatment and cognitive performance in patients with a long history of bipolar illness: no simple answers in complex

- disease-treatment interplay. *Int J Bipolar Disord*, 2, 1. doi:10.1186/s40345-014-0016-7
- Piontkewitz, Y., Bernstein, H., Dobrowolny, H., Bogerts, B., Weiner, I., & Keilhoff, G. (2013). Effects of risperidone treatment in adolescence on hippocampal neurogenesis, parvalbumin expression, and vascularization following prenatal immune activation in rats. *Bulletin*, 39, S123. doi:10.1093/schbul/sbt011
- Poletti, S., Locatelli, C., Radaelli, D., Colombo, C., & Benedetti, F. (2014). Lithium and GSK-3 β promoter gene variants influence gray matter volume in bipolar disorder. *European Neuropsychopharmacology*, 24, S427.
- Rapado-Castro, M., Dodd, S., Bush, A. I., Malhi, G. S., Skvarc, D. R., On, Z. X., . . . Dean, O. M. (2017). Cognitive effects of adjunctive N-acetyl cysteine in psychosis. *Psychol Med*, 47(5), 866-876. doi:10.1017/s0033291716002932
- Reus, G., Abelaira, H., Agostinho, F., Ribeiro, K., Vitto, M., Luciano, T., . . . Quevedo, J. (2012). The administration of olanzapine and fluoxetine has synergistic effects on intracellular survival pathways in the rat brain. *J Psychiatr Res*, 46(8), 1029-1035. doi:10.1016/j.jpsychires.2012.04.016
- Reus, G. Z., Abelaira, H. M., Maciel, A. L., Dos Santos, M. A., Carlessi, A. S., Steckert, A. V., . . . Quevedo, J. (2015). Minocycline protects against oxidative damage and alters energy metabolism parameters in the brain of rats subjected to chronic mild stress. *Metab Brain Dis*, 30(2), 545-553. doi:10.1007/s11011-014-9602-8
- Reus, G. Z., Abelaira, H. M., Tuon, T., Titus, S. E., Ignacio, Z. M., Maciel, A. L., . . . Quevedo, J. (2016). Neuronatomical profile of ketamine after mtor inhibition in the prefrontal cortex. *Biological Psychiatry*, 79(9), 264S. doi:10.1016/j.biopsych.2016.03.1054
- Rideau Batista Novais, A., Guiramand, J., Cohen-Solal, C., Crouzin, N., Ferreira, M. C. D., Vignes, M., . . . Cambonie, G. (2013). N-acetyl-cysteine prevents pyramidal cell disarray and reelin-immunoreactive neuron deficiency in CA3 after prenatal immune challenge in rats. *Pediatr Res*, 73(6), 750-755. doi:10.1038/pr.2013.40
- Romo-Nava, F., Alvarez-Icaza Gonzalez, D., Fresan-Orellana, A., Saracco Alvarez, R., Becerra-Palars, C., Moreno, J., . . . Buijs, R. M. (2014). Melatonin attenuates antipsychotic metabolic effects: an eight-week randomized, double-blind, parallel-group, placebo-controlled clinical trial. *Bipolar Disord*, 16(4), 410-421. doi:10.1111/bdi.12196
- Sachs, B. D., & Caron, M. G. (2014). Chronic fluoxetine increases extra-hippocampal neurogenesis in adult mice. *Int J Neuropsychopharmacol*, 18(4). doi:10.1093/ijnp/pyu029

- Salagre, E., Fernandes, B. S., Dodd, S., Brownstein, D. J., & Berk, M. (2016). Statins for the treatment of depression: A meta-analysis of randomized, double-blind, placebo-controlled trials. *J Affect Disord*, 200, 235-242. doi:10.1016/j.jad.2016.04.047
- Santofimia-Castano, P., Izquierdo-Alvarez, A., de la Casa-Resino, I., Martinez-Ruiz, A., Perez-Lopez, M., Portilla, J. C., . . . Gonzalez, A. (2016). Ebselen alters cellular oxidative status and induces endoplasmic reticulum stress in rat hippocampal astrocytes. *Toxicology*, 357-358, 74-84. doi:10.1016/j.tox.2016.06.002
- Selvaraj, S., Bloomfield, P. S., Cao, B., Veronese, M., Turkheimer, F., & Howes, O. D. (2017). Brain TSPO imaging and gray matter volume in schizophrenia patients and in people at ultra high risk of psychosis: An [11C]PBR28 study. *Schizophr Res*. doi:10.1016/j.schres.2017.08.063
- Sheldrick, A., Camara, S., Ilieva, M., Riederer, P., & Michel, T. M. (2017). Brain-derived neurotrophic factor (BDNF) and neurotrophin 3 (NT3) levels in post-mortem brain tissue from patients with depression compared to healthy individuals - a proof of concept study. *Eur Psychiatry*, 46, 65-71. doi:10.1016/j.eurpsy.2017.06.009
- Simonetti, A., Sani, G., Dacquino, C., Piras, F., De Rossi, P., Caltagirone, C., . . . Spalletta, G. (2016). Hippocampal subfield volumes in short- and long-term lithium-treated patients with bipolar I disorder. *Bipolar Disord*, 18(4), 352-362. doi:10.1111/bdi.12394
- Singh, N., Halliday, A. C., Thomas, J. M., Kuznetsova, O. V., Baldwin, R., Woon, E. C., . . . Churchill, G. C. (2013). A safe lithium mimetic for bipolar disorder. *Nat Commun*, 4, 1332. doi:10.1038/ncomms2320
- Singh, N., Sharpley, A. L., Emir, U. E., Masaki, C., Herzallah, M. M., Gluck, M. A., . . . Churchill, G. C. (2016). Effect of the Putative Lithium Mimetic Ebselen on Brain Myo-Inositol, Sleep, and Emotional Processing in Humans. *Neuropsychopharmacology*, 41(7), 1768-1778. doi:10.1038/npp.2015.343
- Sitges, M., Gómez, C. D., & Aldana, B. I. (2014). Sertraline reduces IL-1 β and TNF- α mRNA expression and overcomes their rise induced by seizures in the rat hippocampus. *PLoS One*, 9(11). doi:10.1371/journal.pone.0111665
- Skvarc, D. R., Dean, O. M., Byrne, L. K., Gray, L., Lane, S., Lewis, M., . . . Marriott, A. (2017). The effect of N-acetylcysteine (NAC) on human cognition - A systematic review. *Neurosci Biobehav Rev*, 78, 44-56. doi:10.1016/j.neubiorev.2017.04.013
- Smesny, S., Milleit, B., Schaefer, M. R., Hipler, U. C., Milleit, C., Wiegand, C., . . . Paul Amminger, G. (2015). Effects of omega-3 PUFA on the vitamin E and glutathione antioxidant defense system in individuals at ultra-high risk of

- psychosis. *Prostaglandins Leukotrienes and Essential Fatty Acids*, 101, 15-21. doi:10.1016/j.plefa.2015.07.001
- Souza, L. C., Wilhelm, E. A., Bortolatto, C. F., Nogueira, C. W., Boeira, S. P., & Jesse, C. R. (2014). The protective effect of melatonin against brain oxidative stress and hyperlocomotion in a rat model of mania induced by ouabain. *Behavioural Brain Research*, 271, 316-324. doi:10.1016/j.bbr.2014.06.030
- Stefanovic, B., Spasojevic, N., Jovanovic, P., Jasnic, N., Djordjevic, J., & Dronjak, S. (2016). Melatonin mediated antidepressant-like effect in the hippocampus of chronic stress-induced depression rats: Regulating vesicular monoamine transporter 2 and monoamine oxidase A levels. *European Neuropsychopharmacology*, 26(10), 1629-1637. doi:10.1016/j.euroneuro.2016.07.005
- Su, K. P., Lai, H. C., Yang, H. T., Su, W. P., Peng, C. Y., Chang, J. P. C., . . . Pariante, C. M. (2014). Omega-3 Fatty Acids in the Prevention of Interferon-Alpha-Induced Depression: Results from a Randomized, Controlled Trial. *Biological Psychiatry*, 76(7), 559-566. doi:10.1016/j.biopsych.2014.01.008
- Tanra, A. J. (2013). The effects of minocycline on microglial cells activation in the dentate gyrus of Gunn rat: A possible animal model of schizophrenia. *Eur Arch Psychiatry Clin Neurosci*, 263(1), S32. doi:10.1007/s00406-013-0433-0
- Toll, A., & Mane, A. (2015). Brain-derived neurotrophic factor levels in first episode of psychosis: A systematic review. *World J Psychiatry*, 5(1), 154-159. doi:10.5498/wjp.v5.i1.154
- Valente, M. M., Bortolotto, V., Cuccurazzu, B., Ubezio, F., Meneghini, V., Francese, M. T., . . . Grilli, M. (2012). alpha2delta ligands act as positive modulators of adult hippocampal neurogenesis and prevent depression-like behavior induced by chronic restraint stress. *Mol Pharmacol*, 82(2), 271-280. doi:10.1124/mol.112.077636
- van der Leeuw, C., Habets, P., Gronenschild, E., Domen, P., Michielse, S., van Kroonenburgh, M., . . . Marcelis, M. (2013). Testing the estrogen hypothesis of schizophrenia: associations between cumulative estrogen exposure and cerebral structural measures. *Schizophr Res*, 150(1), 114-120. doi:10.1016/j.schres.2013.07.033
- van Erp, T. G. M., Thompson, P. M., Kieseppa, T., Bearden, C. E., Marino, A. C., Hoftman, G. D., . . . Cannon, T. D. (2012). Hippocampal morphology in lithium and non-lithium-treated bipolar I disorder patients, non-bipolar co-twins, and control twins. *Human Brain Mapping*, 33(3), 501-510. doi:10.1002/hbm.21239

- Villa, R. F., Ferrari, F., Bagini, L., Gorini, A., Brunello, N., & Tasedda, F. (2017). Mitochondrial energy metabolism of rat hippocampus after treatment with the antidepressants desipramine and fluoxetine. *Neuropharmacology*, 121, 30-38. doi:10.1016/j.neuropharm.2017.04.025
- Vincenzi, B., Stock, S., Borba, C. P., Cleary, S. M., Oppenheim, C. E., Petrucci, L. J., . . . Henderson, D. C. (2014). A randomized placebo-controlled pilot study of pravastatin as an adjunctive therapy in schizophrenia patients: effect on inflammation, psychopathology, cognition and lipid metabolism. *Schizophr Res*, 159(2-3), 395-403. doi:10.1016/j.schres.2014.08.021
- Vita, A., De Peri, L., Deste, G., Barlati, S., & Sacchetti, E. (2015). The Effect of Antipsychotic Treatment on Cortical Gray Matter Changes in Schizophrenia: Does the Class Matter? A Meta-analysis and Meta-regression of Longitudinal Magnetic Resonance Imaging Studies. *Biol Psychiatry*, 78(6), 403-412. doi:10.1016/j.biopsych.2015.02.008
- Wang, H., Liu, S., Tian, Y., Wu, X., He, Y., Li, C., . . . Xiao, L. (2015). Quetiapine Inhibits Microglial Activation by Neutralizing Abnormal STIM1-Mediated Intercellular Calcium Homeostasis and Promotes Myelin Repair in a Cuprizone-Induced Mouse Model of Demyelination. *Frontiers in Cellular Neuroscience*, 9. doi:10.3389/fncel.2015.00492
- Wang, H. N., Liu, G. H., Zhang, R. G., Xue, F., Wu, D., Chen, Y. C., . . . Tan, Q. R. (2016). Quetiapine ameliorates schizophrenia-like behaviors and protects myelin integrity in cuprizone intoxicated mice: The involvement of Notch signaling pathway. *International Journal of Neuropsychopharmacology*, 19(2). doi:10.1093/ijnp/pyv088
- Wang, N., Yu, H. Y., Shen, X. F., Gao, Z. Q., Yang, C., Yang, J. J., & Zhang, G. F. (2015). The rapid antidepressant effect of ketamine in rats is associated with down-regulation of pro-inflammatory cytokines in the hippocampus. *Uppsala journal of medical sciences*, 120(4), 241-248. doi:10.3109/03009734.2015.1060281
- Wehring, H. J., Elsobky, T., McEvoy, J. P., Vyas, G., Richardson, C. M., McMahon, R. P., . . . Kelly, D. L. (2017). Adjunctive Minocycline in Clozapine-Treated Patients with Schizophrenia: Analyzing the Effects of Minocycline on Clozapine Plasma Levels. *Psychiatr Q*. doi:10.1007/s11126-017-9515-x
- Wiedłocha, M., Marcinowicz, P., Krupa, R., Janoska-Jaździk, M., Janus, M., Debowska, W., . . . Szulc, A. (2017). Effect of antidepressant treatment on peripheral inflammation markers - A meta-analysis. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. doi:10.1016/j.pnpbp.2017.04.026

- Wilson, C. B., McLaughlin, L. D., Ebenezer, P. J., Nair, A. R., Dange, R., Harre, J. G., . . . Francis, J. (2014). Differential effects of sertraline in a predator exposure animal model of post-traumatic stress disorder. *Front Behav Neurosci*, 8, 256. doi:10.3389/fnbeh.2014.00256
- Wise, T., Radua, J., Via, E., Cardoner, N., Abe, O., Adams, T. M., . . . Arnone, D. (2017). Common and distinct patterns of grey-matter volume alteration in major depression and bipolar disorder: evidence from voxel-based meta-analysis. *Mol Psychiatry*, 22(10), 1455-1463. doi:10.1038/mp.2016.72
- Wright, D. J., Renoir, T., Smith, Z. M., Frazier, A. E., Francis, P. S., Thorburn, D. R., . . . Gray, L. J. (2015). N-Acetylcysteine improves mitochondrial function and ameliorates behavioral deficits in the R6/1 mouse model of Huntington's disease. *Transl Psychiatry*, 5, e492. doi:10.1038/tp.2014.131
- Wu, A. G., Noble, E. E., Tyagi, E., Ying, Z., Zhuang, Y. M., & Gomez-Pinilla, F. (2015). Curcumin boosts DHA in the brain: Implications for the prevention of anxiety disorders. *Biochimica Et Biophysica Acta-Molecular Basis of Disease*, 1852(5), 951-961. doi:10.1016/j.bbadis.2014.12.005
- Xie, Z. M., Wang, X. M., Xu, N., Wang, J., Pan, W., Tang, X. H., . . . Yang, J. J. (2017). Alterations in the inflammatory cytokines and brain-derived neurotrophic factor contribute to depression-like phenotype after spared nerve injury: improvement by ketamine. *Scientific Reports*, 7. doi:10.1038/s41598-017-03590-3
- Xu, S. X., Zhou, Z. Q., Li, X. M., Ji, M. H., Zhang, G. F., & Yang, J. J. (2013). The activation of adenosine monophosphate-activated protein kinase in rat hippocampus contributes to the rapid antidepressant effect of ketamine. *Behavioural Brain Research*, 253, 305-309. doi:10.1016/j.bbr.2013.07.032
- Yang, S. N., Yang, Y. Y., Wan, F. J., Huang, C. L., & Liu, Y. P. (2017). Effects of isolation rearing and early antipsychotic intervention on oxidative stress-induced apoptosis and brain-derived neurotrophic factor in hippocampus in a rat model of schizophrenia. *Journal of Medical Sciences (Taiwan)*, 37(4), 155-162. doi:10.4103/jmedsci.jmedsci_113_16
- Yoshimura, R., Hori, H., Ikenouchi-Sugita, A., Umene-Nakano, W., Katsuki, A., Hayashi, K., . . . Nakamura, J. (2012). Aripiprazole altered plasma levels of brain-derived neurotrophic factor and catecholamine metabolites in first-episode untreated Japanese schizophrenia patients. *Hum Psychopharmacol*, 27(1), 33-38. doi:10.1002/hup.1257
- Zhang, L., Luo, J. X., Zhang, M. H., Yao, W., Ma, X. L., & Yu, S. Y. (2014). Effects of curcumin on chronic, unpredictable, mild, stress-induced depressive-like

- behaviour and structural plasticity in the lateral amygdala of rats. *International Journal of Neuropsychopharmacology*, 17(5), 793-806.
doi:10.1017/s1461145713001661
- Zhang, Q., He, M., Deng, C., Wang, H., & Huang, X. F. (2014). Effects of olanzapine on the elevation of macrophage infiltration and pro-inflammatory cytokine expression in female rats. *J Psychopharmacol*, 28(12), 1161-1169.
doi:10.1177/0269881114555250
- Zhang, X., Wang, Q., Wang, Y., Hu, J., Jiang, H., Cheng, W., . . . Li, X. (2016). Duloxetine prevents the effects of prenatal stress on depressive-like and anxiety-like behavior and hippocampal expression of pro-inflammatory cytokines in adult male offspring rats. *Int J Dev Neurosci*, 55, 41-48.
doi:10.1016/j.ijdevneu.2016.09.005
- Zhang, Y., Zhang, H., Wang, L., Jiang, W., Xu, H., Xiao, L., . . . Li, X.-M. (2012). Quetiapine enhances oligodendrocyte regeneration and myelin repair after cuprizone-induced demyelination. *Schizophr Res*, 138(1), 8-17.
doi:10.1016/j.schres.2012.04.006
- Zheng, W., Cai, D. B., Yang, X. H., Ungvari, G. S., Ng, C. H., Muller, N., . . . Xiang, Y. T. (2017). Adjunctive celecoxib for schizophrenia: A meta-analysis of randomized, double-blind, placebo-controlled trials. *J Psychiatr Res*, 92, 139-146.
doi:10.1016/j.jpsychires.2017.04.004
- Zheng, X., Zhou, J., & Xia, Y. (2015). The role of TNF- α in regulating ketamine-induced hippocampal neurotoxicity. *Archives of Medical Science*, 11(6), 1296-1302.
doi:10.5114/aoms.2015.56355
- Zhu, F., Zheng, Y., Ding, Y. Q., Liu, Y., Zhang, X., Wu, R., . . . Zhao, J. (2014). Minocycline and risperidone prevent microglia activation and rescue behavioral deficits induced by neonatal intrahippocampal injection of lipopolysaccharide in rats. *PLoS One*, 9(4). doi:10.1371/journal.pone.0093966
- Zung, S., Souza-Duran, F. L., Soeiro-de-Souza, M. G., Uchida, R., Bottino, C. M., Busatto, G. F., & Vallada, H. (2016). The influence of lithium on hippocampal volume in elderly bipolar patients: a study using voxel-based morphometry. *Transl Psychiatry*, 6(6), e846. doi:10.1038/tp.2016.97

	Neurostructural Integrity		Neurogenic Balance		Inflammatory State		Oxidative and Nitrosative Stress	
	<i>Protects cortical volume</i>	<i>Improves white matter</i>	<i>Central</i>	<i>Peripheral</i>	<i>Central</i>	<i>Peripheral</i>	<i>Central</i>	<i>Peripheral</i>
Lithium	Yes	Yes	Rat brain: ↑BDNF, ↓Apoptosis		Rat brain: ↓TLR4, ↓NFκB		↓Oxidative stress	
Clozapine			Mouse brain: ↑BDNF, ↑Proliferation			Rat: ↓toxic TRYCATs, ↓COX signalling	Rat brain: ↑DHA signalling	
Aripiprazole				↑BDNF				
Olanzapine			Rat brain: ↑BDNF, ↓Apoptosis			Rat: ↑TNFα, ↑ IL1β, ↑ IL6		
Risperidone			Rat brain: ↑BDNF, ↓Apoptosis					↓Oxidative stress
Paliperidone					Mouse brain: ↑NRF2, ↓Microglial activity		Rat brain: ↓Oxidative stress	
Quetiapine		Mouse brain: Yes			Mouse brain: ↓Microglial activity		Mouse: ↑mitochondrial activity	
Antidepressants			Mouse brain: ↑BDNF, ↑Hippocampal proliferation	↑BDNF	Rat brain: ↓ IL-β, ↓TNFα	↓IL-4, ↓IL-6, ↓IL-10, ↓ IL-β, ↓toxic TRYCATs	Rat brain: ↑mitochondrial activity	

Table 1: Psychotropic Agents. Evidence of neuroprotective effects and suggested mechanisms of action for psychotropic agents presented in review. Data is divided by mode of detection, either centrally from brain tissue or peripherally from serum. All data represents human clinical studies unless explicitly stated in table. **Key:** ↑ = Increased ↓ = Decreased. For other abbreviations, refer to **Abbreviations**.

	Neurostructural Integrity		Neurogenic Balance		Inflammatory State		Oxidative and Nitrosative Stress	
	<i>Protects cortical volume</i>	<i>Improves white matter</i>	<i>Central</i>	<i>Peripheral</i>	<i>Central</i>	<i>Peripheral</i>	<i>Central</i>	<i>Peripheral</i>
Minocycline	Yes				Mouse brain:↓ IL-β, ↓IL-1, ↓IL-6, ↓TNFα, ↓Microglial activity			
Statins			Rat brain: ↑mTOR				Rat: ↑glutathione	
Aspirin	Rat brain: Yes				↓IL-6, ↓TNFα			
Cyclooxygenase 2 Inhibitors			Rat brain: ↑BDNF, ↓apoptosis		Rat brain: ↓TNFα, ↓NF-κB			
Ketamine			Rat brain: ↑BDNF, ↑ADNP, ↓apoptosis		Rat brain: ↓ IL1-β, ↓IL-6, ↓TNFα			
Pregabalin			Mouse brain: ↑hippocampal neurogenesis					

Table 2: Other Pharmaceutical Agents. Evidence of neuroprotective effects and suggested mechanisms of action for pharmaceutical agents presented in review. Data is divided by mode of detection, either centrally from brain tissue or peripherally from blood. All data represents human clinical studies unless explicitly stated in table. **Key:** ↑ = Increased ↓ = Decreased. For other abbreviations, refer to **Abbreviations**.

	Neurostructural Integrity		Neurogenic Balance		Inflammatory State		Oxidative and Nitrosative Stress	
	<i>Protects cortical volume</i>	<i>Improves white matter</i>	<i>Central</i>	<i>Peripheral</i>	<i>Central</i>	<i>Peripheral</i>	<i>Central</i>	<i>Peripheral</i>
Melatonin			Mouse brain: ↑BDNF, ↑ERK1/2					Rat: ↓Oxidative stress
Oestrogen	Yes		Rat brain: ↑Hippocampal architecture Cortical cells: ↑ERK1/2					
Curcumin			Mouse brain: ↑BDNF, ↑ERK1/2				Rat brain: ↑DHA signaling	
Erythropoietin	Mouse brain: Yes		Mouse brain: ↑proliferation					
N-Acetylcysteine							Mouse brain: ↓Oxidative stress	Mouse:↓Oxidative stress
Leptin			Rat brain: ↑hippocampal neurogenesis					
Omega-3 Fatty Acids							↓Oxidative stress	↑Antioxidant capacity
Ebselen			Mouse brain: ↓IMPase					

Table 2: Nutritional, Hormonal and Other Agents. Evidence of neuroprotective effects and suggested mechanisms of action for nutritional, homonal and other agents presented in review. Data is divided by mode of detection, either centrally from brain tissue or peripherally from blood. All data represents human clinical studies unless explicitly stated in table. **Key:** ↑ = Increased ↓ = Decreased. For other abbreviations, refer to **Abbreviations**.

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Robertson, OD;Coronado, NG;Sethi, R;Berk, M;Dodd, S

Title:

Putative neuroprotective pharmacotherapies to target the staged progression of mental illness

Date:

2019-10-01

Citation:

Robertson, O. D., Coronado, N. G., Sethi, R., Berk, M. & Dodd, S. (2019). Putative neuroprotective pharmacotherapies to target the staged progression of mental illness. *EARLY INTERVENTION IN PSYCHIATRY*, 13 (5), pp.1032-1049. <https://doi.org/10.1111/eip.12775>.

Persistent Link:

<http://hdl.handle.net/11343/285372>