Title: **The Use of the GelPoint Path^R Access Channel for Rectal Trauma**

Running Head: GelPoint^R Path for Rectal Trauma

Names of Authors:

Dr Corina Behrenbruch MBBS(hons), FRACS

Mr Ian Hayes MBBS, MS, FRCS (Gen Surg), FRACS

Mr David Read MBBS, FRACS

The corresponding author is not a recipient of a research scholarship

Figures : 2

Tables : 0

Word count for text : 528

Corresponding author : Dr Corina Behrenbruch

Cori.behrenbruch@gmail.com

Department of General Surgical Specialities

The Royal Melbourne Hospital

Level 6, 300 Grattan St Parkville, 3050, Victoria, Australia

+61393427000

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1111/ans.14997](https://doi.org/10.1111/ans.14997)

The Use of the GelPoint Path^R Access Channel for Rectal Trauma

Behrenbruch C¹, Read D^{1, 2}, Hayes I^{1, 2, 3}

¹Department of General Surgical Specialties, Royal Melbourne Hospital, Victoria, Australia

²Colorectal Surgery Unit, Royal Melbourne Hospital, Victoria, Australia

³Department of Surgery, The University of Melbourne, Victoria, Australia

Extraperitoneal rectal injuries can be difficult to access for repair, and consequently many are diverted. Here the authors report a case of exsanguinating grade 1 rectal injury where repair was greatly facilitated by the use of the access channel component of the GelPoint Path^R; a device designed for transanal excision of polyps. The patient has been consented for publication of the case.

A middle-aged male presented to the emergency department with rectal bleeding following self-digitation. At presentation, he was tachycardic and passing large clots of blood per rectum. A CT angiogram was performed which showed contrast extravastation within the lower rectum in keeping with active haemorrhage (**Figure 1**). There was no evidence of extra-luminal gas to suggest a full-thickness rectal injury. The patient was taken to theatre where flexible sigmoidoscopy demonstrated active arterial bleeding from a mucosal breach in the anterior wall of the lower rectum approximately 5 cm from the anal verge. To gain better access to the area of injury, only the access channel of the GelPoint Path^R platform was inserted (**Figure 2**). This manoeuvre greatly facilitated direct suturing with interrupted 2.0 Vicryl^R, which controlled the haemorrhage.

Extraperitoneal (EP) rectal injuries can be managed by proximal diversion, direct repair, or a combination of both. The current EAST (Eastern Association for the Surgery of Trauma) guidelines cautiously recommend proximal diversion for all EP rectal injuries, but most would interpret that this recommendation applies to full thickness injuries (American Association for the Surge of Trauma (AAST) grades II-V)¹. Despite this, a review of 350 rectal trauma patients from an AAST multi-institutional study showed that 19% of 350 patients diverted for rectal trauma had a grade I injury². The reasons for this are not stated. Access to lower third rectal injuries can be technically difficult but is feasible and without significant septic complications according to a small series by Gonzales *et al.*³. Although conservative management of low-grade rectal injuries is acceptable, the current case presenting with unrelenting haemorrhage excluded this as management option.

Transanal repair of EP rectal injuries is well described using standard proctologic equipment as well as vaginal speculum⁴, Transanal Microsurgery (or Stereoscopic) (TEMS)⁵, and endoscopically with Haemoclips⁶. There have been no published cases detailing the use of Trans Anal Minimally Invasive Surgery (TAMIS) for difficult-to-access low-rectal trauma. TAMIS is an established technique for the resection of rectal polyps, utilising a three-port

system of camera and two working ports, allowing direct suture of the post polypectomy defect with a technique akin to laparoscopic suturing. In our case, the access port for this system was placed in the anus and afforded sufficient view for repair without requiring the gelport and insufflation. The access channel is sutured into position and does not rely upon the steady hand of an assistant, nor does it exert undue stretch upon the anal sphincters. If the defect was not accessible, then the full Gelpoint Path^R platform could potentially be utilised to repair such a defect, although the effects of insufflation on rectal injury remain undescribed. Based on the experience of this case, the authors would recommend considering the use of the TAMIS access channel to facilitate direct transanal repair of rectal injuries.

1. Bosarge PL, Como JJ, Fox N, Falck-Ytter Y, Haut ER, Dorion HA, Patel NJ, Rushing A, Raff LA, McDonald AA, Robinson BR, McGwin G, Jr., Gonzalez RP. Management of penetrating extraperitoneal rectal injuries: An Eastern Association for the Surgery of Trauma practice management guideline. *The journal of trauma and acute care surgery* 2016;**80**(3): 546-551.
2. Brown CVR, Teixeira PG, Furay E, Sharpe JP, Musonza T, Holcomb J, Bui E, Bruns B, Hopper HA, Truitt MS, Burlew CC, Schellenberg M, Sava J, VanHorn J, Eastridge PB, Cross AM, Vasak R, Vercruysse G, Curtis EE, Haan J, Coimbra R, Bohan P, Gale S, Bendix PG. Contemporary management of rectal injuries at Level I trauma centers: The results of an American Association for the Surgery of Trauma multi-institutional study. *The journal of trauma and acute care surgery* 2018;**84**(2): 225-233.
3. Gonzalez RP, Phelan H, 3rd, Hassan M, Ellis CN, Rodning CB. Is fecal diversion necessary for nondestructive penetrating extraperitoneal rectal injuries? *The Journal of trauma* 2006;**61**(4): 815-819.
4. Beatty JS, Atteberry LR, Fasusi OO, Masoumy M, Bates WB, 3rd, Hawkins ML. Transanal repair of traumatic rectal injuries without laparotomy and fecal diversion is feasible and safe in select patients. *The American surgeon* 2014;**80**(2): E56-57.
5. Serra-Aracil X, Gomez-Diaz CJ, Navarro-Soto S, Hidalgo-Rosas JM, Mora-Lopez L. Repair of rectal trauma perforation using transanal endoscopic operation. *Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland* 2012;**14**(7): e427-428.
6. Hahn K-Y, Ko S-Y, Lee W-S, Kim Y-H. Extraperitoneal Rectal Laceration Secondary to Blunt Trauma: Successful Transanal Endoscopic Repair with Hemoclips. *Chinese Medical Journal* 2017;**130**(19): 2384-2385.

Figure 1_ANZJS_Behrenbruch_Rectal Trauma.png

Figure 2_ANZJS_Behrenbruch_Rectal Trauma.png

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Behrenbruch, C;Read, D;Hayes, I

Title:

Use of the GelPoint Path access channel for rectal trauma.

Date:

2019-12

Citation:

Behrenbruch, C., Read, D. & Hayes, I. (2019). Use of the GelPoint Path access channel for rectal trauma.. ANZ J Surg, 89 (12), pp.E589-E590. <https://doi.org/10.1111/ans.14997>.

Persistent Link:

<http://hdl.handle.net/11343/285374>