Phacomatosis pigmentokeratotica: post-zygotic HRAS mutation with malignant degeneration of the sebaceous naevus

Short running title:

PPK with HRAS mutation and multiple BCCs

William C Cranwell, Maie Walsh, Ingrid Winship

Full name of author: Dr William C Cranwell (corresponding author)

Institutional affiliations:

1. Department of Dermatology, The Royal Melbourne Hospital, Parkville
Victoria 3050

Postal address: The Royal Melbourne Hospital, 300 Grattan Street, Parkville, VIC
3050

Email address: william.cranwell@mh.org.au

Telephone number: +613 9342 7000

Full name of author: Dr Maie Walsh

Institutional affiliations:

1. Adult Genetic Medicine, Royal Melbourne Hospital, Melbourne, VIC 3052,
Australia

Postal address: The Royal Melbourne Hospital, 300 Grattan Street, Parkville, VIC
3050

Email address: Maie.Walsh@mh.org.au

Telephone number: +613 9342 7000

Full name of author: Prof Ingrid Winship

Institutional affiliations:

1. Adult Genetic Medicine, Royal Melbourne Hospital, Melbourne, VIC 3052,
Australia
2. Department of Medicine, The University of Melbourne, Melbourne,
Victoria 3010, Australia

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/AJD.13007](https://doi.org/10.1111/AJD.13007)

This article is protected by copyright. All rights reserved

Postal address: The Royal Melbourne Hospital, 300 Grattan Street, Parkville, VIC
3050

Email address: Ingrid.winship@mh.org.au

Telephone number: +613 9342 7000

Acknowledgements: nil

Keywords:

Phacomatosis pigmentkeratolica, Genodermatosis, HRAS mutation, Basal cell carcinoma

Word count: 490

Article type : Case Letter

Phacomatosis pigmentokeratotica: post-zygotic HRAS mutation with malignant degeneration of the sebaceous naevus

Phacomatosis pigmentokeratotica is a rare epidermal naevus syndrome characterised by the coexistence of a sebaceous naevus arranged along Blaschko lines and a speckled lentiginous naevus arranged in a checkerboard pattern. Phacomatosis pigmentokeratotica is a mosaic form of a RASopathy, with recurrent heterozygous *HRAS* variants c.37G>C (p.Gly13Arg) and c.182A>G (p.Gln61Arg) most commonly found in the epidermal and melanocytic naevus¹. A novel *KRAS* c.35G>C (p.G12A) variant has also been described, highlighting the potential for concomitant visceral malignancies associated with phacomatosis pigmentokeratotica². A recent case described phacomatosis pigmentokeratotica with basal cell carcinoma (BCC), syringocystadenoma papilliferum, and trichilemmoma³. Extracutaneous features have been described, including neurologic, ophthalmic, vascular and skeletal defects.

We report a 22-year-old man with a history of phacomatosis pigmentokeratotica, with a pathogenic *HRAS* variant in the sebaceous lesion, mosaic *HRAS* variant in the lentiginous lesions, and wild-type *HRAS* sequence in blood. He was born at 38-weeks' gestation following an unremarkable pregnancy; a congenital sebaceous naevus was evident. This was treated with dermabrasion and multiple excisions. The sebaceous lesion extended to the gingiva on the left side. The lentiginous lesions arose aged 12 years. A left scalp tissue expander was inserted aged 17, but was removed due to expander breakdown and infection. The patient completed High School, but reported significant learning difficulties. He denied any history of seizures, visual symptoms, or neurological symptoms. Multiple BCCs have been excised from the sebaceous naevus over the past two years.

Examination revealed a large sebaceous naevus covering the left scalp, face and neck. Within areas of hyperpigmentation on the back, left arm and left leg, there were multiple lentiginous and

melanocytic naevi. The lesions were clearly demarcated along the midline (Figures 1 & 2). Visual and auditory examinations were normal. A computer tomography of the head and neck showed no central nervous system abnormality.

DNA extracted from punch biopsies from the sebaceous naevus in the left postauricular region showed a pathogenic heterozygous variant, NM 005343.3(HRAS):c.37G>C (p.Gly13Arg) and mosaicism of this variant in the lentiginous naevus on the left forearm. Analysis of peripheral blood showed a wild-type sequence of *HRAS*. These results were consistent with mosaicism. Genetic counselling was provided to the patient and his family.

This case supports the hypothesis that both naevi arose from a single postzygotic *HRAS* mutation in a pluripotent progenitor cell¹. Based on current evidence, the *HRAS* c.37G>C variant is classified as pathogenic. Functional studies show that the c.37G>C variant increases *HRAS* activity.⁴ Post-zygotic somatic mosaicism of an activating *RAS* mutation may compromise an ectodermal, mesodermal, and endodermal differentiation, leading to cutaneous and extracutaneous features of phacomatosis pigmentokeratolica. Generally, the epidermal and melanocytic proliferations remain as benign hamartomas, however basal cell carcinoma and melanoma have been reported in each lesion respectively. The *HRAS* c.37G>C (p.Gly13Arg) mutation has been described in malignant transformation in phacomatosis pigmentokeratolica previously.⁵ It is therefore precautionary to consider that identification of this mutation likely increases the risk of tumorigenesis in either lesion, with consequent appropriate long-term surveillance.

References

1. Groesser L, Herschberger E, Sagrera A, Shwayder T, Flux K, Ehmann L, Wollenberg A, Torrelo A, Bagazgoitia L, Diaz-Ley B. Phacomatosis pigmentokeratolica is caused by a postzygotic *HRAS* mutation in a multipotent progenitor cell. *Journal of Investigative Dermatology*. 2013; 133: 1998-2003.
2. Om A, Cathey SS, Gathings RM, Hudspeth M, Lee JA, Marzolf S, Wine Lee L. Phacomatosis Pigmentokeratolica: A Mosaic *RAS* opathy with Malignant Potential. *Pediatric dermatology*. 2017; 34: 352-5.
3. Loh S-H, Lew B-L, Sim W-Y. A Case of Phacomatosis Pigmentokeratolica Associated With Multiple Basal Cell Carcinomas. *The American Journal of Dermatopathology*. 2018; 40: 131-5.
4. Groesser L, Herschberger E, Ruetten A, Ruivenkamp C, Lopriore E, Zutt M, Langmann T, Singer S, Klingseisen L, Schneider-Brachert W, Toll A. Postzygotic *HRAS* and *KRAS* mutations cause nevus sebaceous and Schimmelpenning syndrome. *Nature genetics*. 2012 Jul;44(7):783.
5. Li JY, Berger MF, Marghoob A, Bhanot UK, Toyohara JP, Pulitzer MP. Combined melanocytic and sweat gland neoplasm: cell subsets harbor an identical *HRAS* mutation in phacomatosis pigmentokeratolica. *Journal of cutaneous pathology*. 2014; 41: 663-71.

Figure Legends:

Figure 1:

- a) Naevus sebaceous extending over the left face, neck, and periauricular regions. Speckled-lentiginous naevus showing a checkerboard appearance on the left neck and upper back. Extensive surgical resection and reconstruction noted.
- b) Linear arrangement of the lentiginous naevus on the right neck.

Figure 2:

- a) Naevus sebaceous affecting the left face and ear.
- b) Multiple lentiginous and melanocytic naevi on the arms and trunk, clearly demarcated along the midline. Epidermoid/sebaceous naevus is seen on the left lower chest.

ajd_13007_f1.jpg

ajd_13007_f2.jpg

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Cranwell, WC;Walsh, M;Winship, I

Title:

Phacomatosis pigmentokeratotica: Postzygotic HRAS mutation with malignant degeneration of the sebaceous naevus

Date:

2019-08-01

Citation:

Cranwell, W. C., Walsh, M. & Winship, I. (2019). Phacomatosis pigmentokeratotica: Postzygotic HRAS mutation with malignant degeneration of the sebaceous naevus. AUSTRALASIAN JOURNAL OF DERMATOLOGY, 60 (3), pp.E245-E246. <https://doi.org/10.1111/ajd.13007>.

Persistent Link:

<http://hdl.handle.net/11343/285457>