Obstructive sleep apnea in multiple pregnancies

Farnaz Dave¹. Honorary research fellow. Department of Obstetric Medicine, The Royal Women's Hospital, 20 Flemington road, Parkville, Victoria 3052, Australia.

Stephen Cole². Obstetrician and Specialist in Maternal Fetal Medicine, Head of Multiple Pregnancy Clinic, The Royal Women's Hospital, 20 Flemington road, Parkville, Victoria 3052, Australia.

Megan Rees³. Consultant Respiratory and Sleep Physician. Department of Respiratory and Sleep Disorders Medicine, The Royal Melbourne Hospital, 300 Grattan street, Parkville, Victoria, 3050, Australia.

Correspondence:

Dr Megan Rees,

Postal Address: PO Box 1060, North Melbourne, Victoria 3051, Australia.

Email: megan.rees@mh.org.au.

Telephone: (+61) 431 110 995.

Conflict of interest: The authors report no conflicts of interest.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/AJO.12958](https://doi.org/10.1111/AJO.12958)

This article is protected by copyright. All rights reserved

DR. MEGAN ANNE REES (Orcid ID : 0000-0002-7736-4000)

Article type : Review

Obstructive sleep apnea in multiple pregnancy

Short title: Obstructive sleep apnea in multiple pregnancy

Word count: Abstract 90, main text 1478

Table: One

Keywords: multiple pregnancy, multifetal pregnancy, preeclampsia, obesity, obstructive sleep apnea

Abstract

Obstructive sleep apnea (OSA) during pregnancy has been associated with gestational diabetes mellitus, hypertension and low birth weight. Multiple pregnancy is associated with similar adverse perinatal outcomes. Multiple pregnancy rates have risen with increased access to assisted conception; simultaneously advancing maternal age and weight are also driving a rise in the incidence of OSA in pregnancy. The intersection of OSA and multiple pregnancy would be presumed to have significant maternal and fetal morbidity, however specific data is sparse. We review the current state of knowledge on OSA in multiple

pregnancy.

Introduction

Obstructive sleep apnea (OSA) involves occlusion of the upper airway causing cessation of breathing numerous times during sleep, this results in oxygen desaturation, transient hypertension and tachycardia, arousal from sleep and sleep fragmentation¹. OSA causes daytime sleepiness, fatigue, and reduction in quality of life¹. OSA is associated with cardiovascular events, insulin resistance and mortality in non-pregnant patients²⁻⁴.

In pregnancy, OSA is associated with gestational diabetes mellitus (GDM), hypertensive disorders, preterm birth and reduced birth weight⁵. In pregnancy pre-existing OSA can worsen, but OSA can also present de novo⁵. Multiple pregnancy is associated with many of the same adverse perinatal outcomes as OSA, including preterm birth, intrauterine growth restriction, GDM and hypertensive disorders^{6,7}. When OSA occurs in a multiple pregnancy these risks may be compounded.

Rates of both OSA and multiple pregnancy have recently increased in parallel with advancing maternal age and greater prevalence of obesity^{8,9}; the intersection of these conditions warrants urgent examination. In this review we describe proposed physiological mechanisms, and present the available data on prevalence, investigation and management of OSA in multiple pregnancy.

Physiology of obstructive sleep apnea in multiple pregnancy

Awake respiratory function is remarkably stable during healthy pregnancy. Any compression of the thoracic cavity by the enlarging uterus appears to be compensated by a reduction in residual lung volume and increased minute ventilation. McAuliffe et al¹⁰, did not show any significant differences in lung function throughout the trimesters between twin or singleton pregnancies in healthy women, suggesting that respiratory compromise from the larger abdominal burden of a multiple pregnancy is unlikely to directly contribute to OSA.

Progesterone has a significant role in sleep physiology, and progesterone levels increase throughout pregnancy. It is likely progesterone levels are higher in multiple pregnancy compared with singleton pregnancy¹¹. Progesterone increases blood flow to pharyngeal tissues predisposing to sleep apnoea. However, progesterone also acts as a respiratory stimulant, increasing minute ventilation and has been associated with increased awakenings from sleep in pregnancy¹² thereby reducing the frequency and severity of pauses in ventilation during sleep. The balance of these effects has not been determined and any subsequent augmentation in multiple pregnancy is undefined.

Prevalence of obstructive sleep apnea in multiple pregnancy

The prevalence of OSA in multiple pregnancy is unknown. In singleton pregnancies in nulliparous women the prevalence is 3.6% in early pregnancy, rising to 8.3% in mid pregnancy¹³. The prevalence is higher in obese pregnant women at 15.4%¹⁴. Furthermore, rates of OSA in pregnancy have been increasing over time. This trend was demonstrated in a retrospective cross sectional analysis⁸, which showed an overall increase in OSA diagnosis from 0.7 to 7.3 per 10,000 from 1998 to 2009, with an average annual increase of 24%

coinciding with a similar rise in rates of obesity (20%).

The increased proportion of multiple pregnancies has been attributed to increased access to assisted conception, as well as increasing maternal age and BMI. In 2016 there were over 4500 multiple births in Australia; over 200 more than in 2015⁹. In addition as spontaneous dizygotic twinning is more likely with an increased BMI, multiple pregnancy rates are likely to continue increasing as obesity rates rise^{15,16}.

Only one study has specifically examined sleep related apneas in multiple pregnancies¹⁷. Nikkola et al (1996) investigated 10 multiple pregnancies; 8 twin, 1 triplet and 1 quadruplet between 30 and 36 weeks' gestation. All subjects had a normal pre-pregnancy BMI, except the mother with quadruplets (BMI 28.4kg/m²). OSA was not identified in any of the patients, although significantly increased respiratory resistance was noted in the patient with quadruplets. This study used a charge-sensitive bed to record nocturnal breathing patterns, oxygenation and leg movements. This methodology underestimates the severity of sleep disordered breathing¹⁸ and has not been validated in pregnancy, furthermore fetal movements may substantially compromise the scoring of events. At the time of writing this remains the largest study investigating sleep apnea in multiple pregnancy.

In 2014, multiple pregnancy was included within a study of OSA in high risk pregnant women²¹. Patients with a BMI >30kg/m², chronic hypertension, pre-gestational diabetes, prior preeclampsia and/or twin pregnancy were included. Of 188 participants 11 were twin gestations, all subjects had an in-home sleep study between 6-20 weeks. A repeated sleep study was performed in 128 in the final trimester; the actual number of multiple pregnancies

in this subsequent group was not reported. In the entire cohort OSA was common in early pregnancy with new-onset OSA in 20% of these high-risk women. In the analysis describing incident OSA in the third trimester, only twin gestation was significantly associated, however the small sample size prevented accurate calculation of risk. This study used a portable wrist worn device (Watch-PAT) which measured multiple variables including oxygen saturation, and estimated sleep time using actigraphy (movement detection). O'Brien et al¹⁹ have demonstrated that results from the Watch-PAT device correlated well with laboratory based polysomnography (PSG) in the third trimester.

Other data on OSA in multiple pregnancies has been published in case reports²⁰⁻²² (see table I). Further studies are required to evaluate the prevalence of OSA in multiple pregnancies.

Given the dynamic physiology of pregnancy screening at a single time point may underestimate the prevalence of this disorder, particularly as OSA is known to be more prevalent in the third trimester of singleton pregnancy²³.

Investigation of obstructive sleep apnea in multiple pregnancy

At present there is no standardised screening tool for assessing risk of OSA in pregnancy although several have been proposed and validated²⁴. Two studies^{25,26} have evaluated OSA assessment tools in high risk cohorts which share many comorbidities with multiple pregnancy. Neither the Epworth Sleep Scale nor the Berlin Questionnaire in these studies could accurately predict OSA risk in women with chronic hypertension, pre-gestational diabetes, obesity and or prior history of preeclampsia²⁵. Tantrakul et al²⁶ also concluded both tools had limited use in the first trimester but improved as the pregnancy progressed.

Any tool developed to screen for OSA in pregnancy needs to assess proven risk factors for

OSA in singleton pregnancies including snoring, elevated BMI and increased neck circumference; features that were also observed in three case reports^{20–22} of OSA in twin pregnancies. Advanced maternal age has also been shown to increase the risk of OSA in singleton pregnancies recently²⁷, and should therefore also be considered in future screening for OSA.

The optimal diagnostic test for women with suspected OSA has not been defined and pregnancy-specific recommendations are absent from guidelines such as the American Academy of Sleep Medicine²⁸. Laboratory PSG is the gold standard measurement of sleep physiology as it obtains continuous electroencephalogram (EEG) data thereby precisely documenting sleep stage, as well as respiratory events and consequences such as oxygen desaturation. The successful use of PSG to diagnose OSA in multiple pregnancies has been documented^{20–22}.

More limited data can be obtained in the home setting; involving either reduced channel EEG or actigraphy (movement detection) as a proxy measure of sleep state. Home based sleep studies may be more convenient, particularly as multiple pregnancies already require supplementary antenatal visits, therefore additional sleep study visits may be burdensome. Conversely, pregnant mothers with young children report a more restful night in a sleep laboratory compared with sleeping at home²⁹.

Management of obstructive sleep apnea in multiple pregnancy

In the non-pregnant population positional therapy, oral devices and continuous positive airway pressure (CPAP) therapy can be used to improve outcomes in OSA³⁰. In singleton

pregnancies CPAP has been shown to improve symptoms³¹ and reduce hypertension³². In two case reports involving CPAP in multiple pregnancies; CPAP controlled or reduced the effects of sleep apnea^{21,22}, whereas in a third case report supplemental oxygen and more complex non-invasive ventilation were required²⁰. In each case clinicians felt nocturnal ventilatory support improved pregnancy outcome, however this limited observational data highlights the need for further research.

Conclusion

Multiple pregnancy and OSA are both separately associated with an increased risk of GDM and hypertensive disorders of pregnancy, including preeclampsia^{13,33,34}. In addition, fetal morbidities such as low birth weight, preterm delivery and neonatal intensive care admission are all more likely with maternal OSA in studies of singleton pregnancies⁵. However, there is little information describing these adverse outcomes in multiple pregnancies complicated by OSA. Investigation or management of OSA is not mentioned in any international obstetric guidelines^{35,36}, and evidence to support intervention is absent. Consequently OSA is rarely considered in routine antenatal practice^{37,38}.

We have described the limited data available regarding OSA in multiple pregnancies. This is a substantial knowledge gap given the rising prevalence of both OSA and multiple pregnancy. The significant maternal and fetal morbidity that can result from OSA in pregnancy should drive further research to provide an evidence-base and improve patient centred care.

References

1. Rowley, J.A., Lareau, S., Fahy, B.F., et al. What Is Obstructive Sleep Apnea in Adults? *Am J Respir Crit Care Med* 2017; **196**: 1–2.. Reichmuth, K.J., Austin,

- D., Skatrud, J.B., Young, T. Association of sleep apnea and type II diabetes: a population-based study. *Am J Respir Crit Care Med* 2005; **172**: 1590–5.
3. Yaggi, H.K., Concato, J., Kernan, W.N., et al. Obstructive sleep apnea as a risk factor for stroke and death. *N Engl J Med* 2005; **353**: 2034–41.
 4. Young, T., Finn, L., Peppard, P.E., et al. Sleep disordered breathing and mortality: eighteen-year follow-up of the Wisconsin sleep cohort. *Sleep* 2008; **31**: 1071–8.
 5. Brown, N.T, Turner, J.M, Kumar, S. The intrapartum and perinatal risks of sleep-disordered breathing in pregnancy: a systematic review and metaanalysis. *Am J Obstet Gynecol* 2018; **219**:147–161.
 6. Fox, N.S., Rebarber, A., Klauser, C.K., et al. Intrauterine growth restriction in twin pregnancies: incidence and associated risk factors. *Am J Perinatol* 2011; **28**: 267–72.
 7. Walker, M.C., Murphy, K.E., Pan, S., et al. Adverse maternal outcomes in multifetal pregnancies. *BJOG An Int J Obstet Gynaecol* 2004; **111**: 1294–6.
 8. Louis, J.M., Mogos, M.F., Salemi, J.L., et al. Obstructive Sleep Apnea and Severe Maternal-Infant Morbidity/Mortality in the United States, 1998-2009. *Sleep* 2014; **37**: 843–9.
 9. Australian Bureau of Statistics. 3301.0 - Births, Australia, 2016. [Cited 2018 June 21]. Available from: <http://www.abs.gov.au/AUSSTATS/abs@.nsf/Latestproducts/3301.0Main%20Features32016>
 10. McAuliffe, F., Kametas, N., Costello, J., et al. Respiratory function in singleton and twin pregnancy. *BJOG An Int J Obstet Gynaecol* 2002; **109**: 765–9.
 11. Lim, H., Powell, S., Mcnamara, H.C., et al. Placental hormone profiles as predictors of preterm birth in twin pregnancy: A prospective cohort study. *PLoS One* 2017; **12**: e0173732.
 12. Wilson, D.L., Barnes, M., Ellett, L., et al. Decreased sleep efficiency, increased wake after sleep onset and increased cortical arousals in late pregnancy. *Aust New Zeal J Obstet Gynaecol* 2011; **51**: 38–46.
 13. Facco, F.L., Parker, C.B., Reddy, U.M., et al. Association between Sleep-Disordered

Breathing and Hypertensive Disorders of Pregnancy and Gestational Diabetes Mellitus. *Obstet Gynecol* 2017; **129**: 31–41.

14. Louis, J., Auckley, D., Miladinovic, B., et al. Perinatal outcomes associated with obstructive sleep apnea in obese pregnant women. *Obs Gynecol* 2012; **120**: 1085–92.
15. Reddy, U.M., Branum, A.M., Klebanoff, M.A. Relationship of Maternal Body Mass Index and Height to Twinning. *Obstet Gynecol* 2005; **105**: 593–7.
16. Hoekstra, C., Willemsen, G., van Beijsterveldt, T., et al. Body composition, smoking, and spontaneous dizygotic twinning. *Fertil Steril*. 2010; **93**: 885–93.
17. Nikkola, E., Ekblad, U., Ekholm, E., et al. Sleep in multiple pregnancy: Breathing patterns, oxygenation, and periodic leg movements. *Am J Obstet Gynecol* 1996; **174**: 1622–5.
18. Svanborg, E., Larsson, H., Carlsson-Nordlander, B., Pirskanen, R. A limited diagnostic investigation for obstructive sleep apnea syndrome. Oximetry and static charge sensitive bed. *Chest* 1990; **98**: 1341–5.
19. O'Brien, L.M., Bullough, A.S., Shelgikar, A.V., et al. Validation of Watch-PAT-200 against polysomnography during pregnancy. *J Clin sleep Med* 2012; **8**: 287–94.
20. Langner, S., Halank, M., Kolditz, M., et al. [Twin pregnancy and severe obstructive sleep apnea]. *Z Geburtshilfe Neonatol* 2007; **211**: 93–7.
21. Gruca-Stryjak, K., Cofta, S., Wysocka, E., et al. [Is there a relationship between pregnancy induced hypertension and obstructive sleep apnea? Case report]. *Pneumonol Alergol Pol* 2014; **82**: 156–62.
22. Canelio, S., Morton, A., McIntyre, H.D. Sleep disordered breathing in pregnancy: the maternal and fetal implications. *J Obstet Gynaecol* 2016; **3615**: 1–9.
23. Facco, F.L., Ouyang, D.W., Zee, P.C., Grobman, W.A. Sleep disordered breathing in a high-risk cohort prevalence and severity across pregnancy. *Am J Perinatol* 2014; **31**: 899–904.
24. Tantrakul, V., Numthavaj, P., Guilleminault, C., et al. Performance of screening questionnaires for obstructive sleep apnea during pregnancy: A systematic review and meta-analysis. *Sleep Med Rev* 2017; **36**: 96–106.

25. Facco, F.L., Ouyang, D.W., Zee, P.C., Grobman, W.A. Development of a pregnancy-specific screening tool for sleep apnea. *J Clin sleep Med* 2012; **8**: 389–94.
26. Tantrakul, V., Sirijanchune, P., Panburana, P., et al. Screening of obstructive sleep apnea during pregnancy: Differences in predictive values of questionnaires across trimesters. *J Clin Sleep Med* 2015; **11**: 157–63.
27. Louis, J.M., Koch, M.A., Reddy, U.M., et al. Predictors of sleep-disordered breathing in pregnancy. *Am J Obstet Gynecol* 2018; **218**: 521.e1-521.e12.
28. Collop, N.A., Anderson, W.M., Boehlecke, B., et al. Clinical guidelines for the use of unattended portable monitors in the diagnosis of obstructive sleep apnea in adult patients. *J Clin sleep Med* 2007; **3**: 737–47.
29. Fung, A.M., Wilson, D.L., Barnes, M., Walker, S.P. Obstructive sleep apnea and pregnancy: The effect on perinatal outcomes. *J Perinatol* 2012; **32**: 399–406.
30. Bratton, D.J., Gaisl, T., Schlatzer, C., Kohler, M. Comparison of the effects of continuous positive airway pressure and mandibular advancement devices on sleepiness in patients with obstructive sleep apnoea: a network meta-analysis. *Lancet Respir Med* 2015; **3**: 869–78.
31. Guilleminault, C., Kreutzer, M., Chang, J.L. Pregnancy, sleep disordered breathing and treatment with nasal continuous positive airway pressure. *Sleep Med* 2004; **5**: 43–51.
32. Edwards, N., Blyton, D.M., Kirjavainen, T., et al. Nasal continuous positive airway pressure reduces sleep-induced blood pressure increments in preeclampsia. *Am J Respir Crit Care Med* 2000; **162**: 252–7.
33. Pamidi, S., Pinto, L.M., Marc, I., et al. Maternal sleep-disordered breathing and adverse pregnancy outcomes: A systematic review and metaanalysis. *Am J Obstet Gynecol* 2014; **210**: 52.e1-52.e14.
34. Edwards, N., Middleton, P.G., Blyton, D.M., Sullivan, C.E. Sleep disordered breathing and pregnancy. *Thorax* 2002; **57**: 555–8.
35. National Institute for Health and Care Excellence, 2011. Multiple pregnancy: antenatal

care for twin and triplet pregnancies. [Cited 2017 June 21] Available from:
<https://www.nice.org.uk/guidance/cg129/resources/multiple-pregnancy-antenatal-care-for-twin-and-triplet-pregnancies-35109458300869>

36. Practice Bulletin No. 169: Multifetal Gestations: Twin, Triplet, and Higher-Order Multifetal Pregnancies. *Obstet Gynecol* 2016; **128**: e131-46.
37. Bourjeily, G., Raker, C., Paglia, M.J., et al. Patient and provider perceptions of sleep disordered breathing assessment during prenatal care: A survey-based observational study. *Ther Adv Respir Dis* 2012; **6**: 211–9.
38. Dominguez, J.E., Lockhart, E.M., Miskovic, A., Bullough, A.S. Recognition of obstructive sleep apnea in pregnancy survey. *International journal of obstetric anesthesia* 2016; **26**: 85–7.

Table I. Published studies of obstructive sleep apnea in multiple pregnancy								
Study	Study design	Participants	Clinical features pre-study	BMI (kg/m2)	Type of Sleep Study	Treatment	Maternal outcomes	Fetal outcomes
Nikkola et al, 1996 ¹⁷	Prospective observational	8 twins 1 triplet 1 quadruplet	-	Mean 22.3 in 8 twin and 1 triplet (range 19.4-24.4)	Laboratory study using static charge sensitive bed with finger oximeter	-	-	8 small for gestational age
				Quadruplets 28.4	Sleep staging estimated by actigraphy			6 growth restricted
Langer et al, 2007 ²⁰	Case report	1 twin	Snoring	-	Laboratory PSG Sleep staging by EEG	BiPAP-ST with oxygen	Progressive preeclampsia	Premature delivery 31/40
Faco et al, 2014 ²³	Prospective cohort	Total 188 high risk cases Only 11 twins	-	32.8	Home sleep evaluation with Watch-PAT100 Sleep staging estimated by actigraphy	-	Preeclampsia in 17.6% (for entire high risk cohort)	Premature birth 9% (for entire high risk cohort)
Gruca-Stryjak et al, 2014 ²¹	Case report	1 twin	Severe daytime sleepiness Snoring	28	Laboratory PSG Sleep staging by EEG	CPAP	Urgent caesarean section at 35/40 due to abnormal CTG of one fetus	Healthy baby

Pauses in breathing during sleep								
Carnelio et al, 2017 ²²	Case reports	Total 3 Only 1 twin	Neck circumference 58cm	48.5	Laboratory PSG	CPAP	Caesarean section after failed induction of labour at 37/40 due to intrahepatic cholestasis	Healthy baby
			Bilateral papillodema	Sleep staging by EEG				
			Crowded oropharynx

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Dave, F;Cole, S;Rees, M

Title:

Obstructive sleep apnoea in multiple pregnancy

Date:

2019-06-01

Citation:

Dave, F., Cole, S. & Rees, M. (2019). Obstructive sleep apnoea in multiple pregnancy. AUSTRALIAN & NEW ZEALAND JOURNAL OF OBSTETRICS & GYNAECOLOGY, 59 (3), pp.341-345. <https://doi.org/10.1111/ajo.12958>.

Persistent Link:

<http://hdl.handle.net/11343/285504>